



HAL
open science

Nicotine enhances alcohol intake and dopaminergic responses through $\beta 2^*$ and $\beta 4^*$ nicotinic acetylcholine receptors.

Stefania Tolu, Fabio Marti, Carole Morel, Carole Perrier, Nicolas Torquet, Stephanie Pons, Renaud de Beaurepaire, Philippe Faure

► To cite this version:

Stefania Tolu, Fabio Marti, Carole Morel, Carole Perrier, Nicolas Torquet, et al.. Nicotine enhances alcohol intake and dopaminergic responses through $\beta 2^*$ and $\beta 4^*$ nicotinic acetylcholine receptors.. Scientific Reports, 2017, 7, pp.45116. <10.1038/srep45116>. <hal-01502363>

HAL Id: hal-01502363

<https://hal.sorbonne-universite.fr/hal-01502363v1>

Submitted on 5 Apr 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

SCIENTIFIC REPORTS



OPEN

Nicotine enhances alcohol intake and dopaminergic responses through $\beta 2^*$ and $\beta 4^*$ nicotinic acetylcholine receptors

Received: 22 November 2016

Accepted: 17 February 2017

Published: 23 March 2017

Stefania Tolu¹, Fabio Marti¹, Carole Morel^{1,†}, Carole Perrier², Nicolas Torquet¹, Stephanie Pons^{3,4}, Renaud de Beaurepaire² & Philippe Faure¹

Alcohol and nicotine are the most widely co-abused drugs. Both modify the activity of dopaminergic (DA) neurons of the Ventral Tegmental Area (VTA) and lead to an increase in DA release in the Nucleus Accumbens, thereby affecting the reward system. Evidences support the hypothesis that distinct nicotinic acetylcholine receptors (nAChRs), the molecular target of acetylcholine (ACh) and exogenous nicotine, are also in addition implicated in the response to alcohol. The precise molecular and neuronal substrates of this interaction are however not well understood. Here we used *in vivo* electrophysiology in the VTA to characterise acute and chronic interactions between nicotine and alcohol. Simultaneous injections of the two drugs enhanced their responses on VTA DA neuron firing and chronic exposure to nicotine increased alcohol-induced DA responses and alcohol intake. Then, we assessed the role of $\beta 4^*$ nAChRs, but not $\beta 2^*$ nAChRs, in mediating acute responses to alcohol using nAChR subtypes knockout mice ($\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice). Finally, we showed that nicotine-induced modifications of alcohol responses were absent in $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice, suggesting that nicotine triggers $\beta 2^*$ and $\beta 4^*$ nAChR-dependent neuroadaptations that subsequently modify the responses to alcohol and thus indicating these receptors as key mediators in the complex interactions between these two drugs.

Alcohol and nicotine are the most commonly abused drugs in the world and their use triggers a broad range of serious negative health consequences with a high cost for the society^{1,2}. Despite dissimilarities in their mechanisms of action and in their behavioural effects, alcohol and tobacco use commonly occur together. A very large majority (80–90%) of dependent drinkers smoke cigarettes³, and alcoholism has been estimated to be 10–14 times more common among smokers than non-smokers⁴.

Multiple factors may contribute to this high comorbidity, including environmental, psychosocial and genetic background^{5,6}. Animal studies also point out the possibility of shared neurobiological mechanisms influencing the development of this co-addiction⁷. Indeed, among others, both alcohol and nicotine act on the mesocorticolimbic dopaminergic (DA) system. This system, originating in the Ventral Tegmental Area (VTA) of the mid-brain and projecting to the Nucleus Accumbens (NAcc) and Prefrontal Cortex (PFC), is involved in reward, motivation, memory and cognition. Both systemic nicotine and alcohol increase synaptic DA release in the NAcc⁸, a key event that is considered to initiate reinforcement.

Increase in DA release is, for both drugs, mainly the consequence of an increase in VTA DA neuron firing rate and bursting activity^{9,10}. However, if nicotine exerts its reinforcing effects by acting on nicotinic acetylcholine receptors (nAChRs), alcohol has been shown to act through a variety of neuronal receptors and ion channels, including different nAChR subtypes¹¹. Several animal studies suggest a role for specific nAChRs in alcohol-elicited DA responses and alcohol reward^{12–15}, thus making nAChRs a potential candidate for the molecular target of alcohol and nicotine interaction.

¹Sorbonne Universités, UPMC Univ Paris 06, INSERM, CNRS, Neurosciences Paris Seine - Institut de Biologie Paris Seine (NPS - IBPS), 75005, Paris, France. ²Groupe Hospitalier Paul Guiraud, BP 20065, F-94806, Villejuif, France. ³Institut Pasteur, Unité Neurobiologie Intégrative des Systèmes Cholinergiques, Département de Neurosciences, F-75724, Paris, France. ⁴CNRS, UMR 3571, F-75724, Paris, France. [†]Present address: Department of Pharmacology and Systems Therapeutics, Icahn School of Medicine at Mount Sinai, New York, NY, USA. Correspondence and requests for materials should be addressed to P.F. (email: phfaure@gmail.com)

nAChRs are a family of pentameric ligand-gated ion channels made up of different α ($\alpha 2$ – $\alpha 10$) and β ($\beta 2$ – $\beta 4$) subunits which can assemble in multiple combinations¹⁶. In this paper, our interest focused on $\beta 2$ -containing ($\beta 2^*$) and $\beta 4^*$ nAChRs as they show specific DA phenotypes with regards to nicotine reinforcement. $\beta 2^{-/-}$ mice lack the ability to self-administer nicotine and do not display nicotine-induced DA release in the NAcc^{17–19}, while $\beta 4^{-/-}$ mice have been recently implicated in the control of nicotine consumption^{20,21}. The interest for $\beta 4^*$ -nAChRs is also supported by human genetic studies assessing the implication of this receptor in the vulnerability to nicotine dependence and in the age of initiation for both tobacco and alcohol consumption^{6,22,23}.

To investigate the multifaceted relationship between alcohol and nicotine, we used *in vivo* electrophysiology and an alcohol-drinking paradigm to analyse acute and chronic interactions between these two drugs. In parallel, we used transgenic mice to investigate the role of specific nAChRs in alcohol-induced responses and alcohol intake in nicotine-naïve and nicotine-exposed mice.

Results

Acute and chronic nicotine enhance responses to alcohol. We first aimed to evaluate *in vivo* the effect of nicotine administration on the evoked response of VTA DA cells to i.v. injections of alcohol. Putative VTA DA neurons were identified according to their location and well-established electrophysiological and pharmacological criteria^{9,17} (see Methods). First, we characterized the evoked response of VTA DA cells to i.v. injections of alcohol. *In vivo*, alcohol injections resulted in a dose-dependent increase of the firing rate and of the bursting activity (%SWB, percentage of spike within a burst) of VTA DA neurons in C57BL/6J mice (Fig. 1a,b and Fig. S1a) (Δ Frequency: One-way ANOVA, $F_{(4,72)} = 10.9$, $p < 0.001$; Δ %SWB: Kruskal–Wallis, $X^2 = 10.5$, $df = 4$, $p < 0.05$). Similarly to what we observed with nicotine²⁴, a fraction of VTA DA cells were inhibited by alcohol, but the current study focuses on cells that were excited by alcohol.

We then performed, on the same neurons, simultaneous injections of the two drugs and compared the responses with those obtained with a single injection of nicotine (30 μ g/kg) or of alcohol (500 mg/kg) alone. Concomitant i.v. injections of nicotine and alcohol resulted in a change in the firing frequency of greater amplitude compared to the two drugs injected separately (Fig. 1c) (Paired Wilcoxon test: double vs single nicotine: $V = 5$, $p < 0.001$; double vs single alcohol: $V = 13$, $p < 0.001$). These data suggest that the two processes, i.e. alcohol- and nicotine-evoked responses, are not in competition and do not saturate the DA cell responses, at least at the tested doses. They are also in line with previous studies showing an additive and/or synergistic effects in the VTA to NAcc pathway^{25–27}. Concurrent injections did not increase the %SWB further compared to single injections of each drug alone (Fig. S1b) (Paired Wilcoxon test: double vs single nicotine: $V = 14$, $p = 1$; double vs single alcohol: $V = 18$, $p = 0.06$). This could be explained by the fact that bursting modifications induced by alcohol are of smaller amplitudes compared to nicotine.

Given that tobacco addiction increases the incidence of alcohol abuse in humans, we then investigated the effect of chronic nicotine on alcohol-induced responses. Mice were exposed to chronic nicotine (10 mg/kg/d) using osmotic Alzet[®] mini-pumps (MPs, see Materials and Methods) for 22–26 days. VTA DA cells spontaneous activity, alcohol-induced responses of DA neurons and alcohol intake were then estimated in the presence of chronic nicotine.

At a cellular level, chronic nicotine pre-exposure induced an increase in the spontaneous bursting activity (%SWB) of VTA DA neurons of mice (nic+) compared to control animals (nic–; a pooled group of naïve mice and mice with mini-pumps delivering saline, see SI and Fig. S2) (Δ %SWB: $W = 2650.5$, $p < 0.05$; but no modification in the firing frequency (Δ Frequency: $t = -1.6$, $df = 113.9$, $p = 0.1$) (Fig. 1d). It also increased the sensitivity of VTA DA neurons to acute injections of alcohol, as indicated by an upward shift of the dose-response curve for both the firing frequency and the %SWB (Fig. 1e and Fig. S1c) (Δ Frequency: Two-way ANOVA: dose: $F_{(3,92)} = 10.0$, $p < 0.001$; nicotine effect: $F_{(1,92)} = 8.2$, $p < 0.01$; Δ %SWB: Kruskal–Wallis: nicotine effect: $X^2 = 6.7$; $df = 3$, $p < 0.01$).

At a behavioural level, this DA cell sensitization matched with an increase in alcohol consumption in a 24-hour voluntary alcohol-drinking paradigm (see Materials and Methods). In this paradigm, mice had to choose between water and increasing concentrations of alcohol during 18 consecutive days. Nic+ mice ingested higher quantities of alcohol compared to their control mice (nic–; Fig. 2a) (repeated measures ANOVA: dose: $F_{(3,93)} = 14.2$, $p < 0.001$; nicotine effect: $F_{(1,31)} = 8.1$; $p < 0.01$). Similarly, the preference ratio for ethanol was higher for nic+ than for nic– mice (Fig. 2b, dose: $F_{(3,93)} = 16.3$, $p < 0.001$; nicotine effect: $F_{(1,31)} = 10.1$, $p < 0.01$). Moreover, the total fluid intake (alcohol+water) was relatively constant across concentration ($F_{(3,93)} = 1.3$; $p = 0.3$) and similar for the two groups ($F_{(1,31)} = 1.7$; $p = 0.2$), thus suggesting that the increased consumption was not due to an increase in thirst, but rather to the increased rewarding properties of the drug (Fig. 2c). Finally, the total alcohol intake during the 18 day-procedure was greater for nic+ than nic– mice (Fig. 2d, Wilcoxon test: $W = 59$, $p < 0.01$). Such results highlighted an adaptation set up by nicotine pre-treatment and are consistent with previous behavioural studies that reported an increase in alcohol intake or preference after nicotine treatment^{28,29}.

Alcohol-induced firing frequency change is modified in $\beta 4^{-/-}$, but not in $\beta 2^{-/-}$ mice. The next step was to investigate the contribution of specific nAChRs in alcohol electrophysiological effects and alcohol intake. The spontaneous activity of DA neurons from $\beta 2^{-/-}$ mice is characterized by a decrease in firing rate ($t = 4.8$, $df = 169.3$, $p < 0.001$) and in bursting activity ($W = 2524$, $p < 0.001$) compared to WT mice, whereas neurons from $\beta 4^{-/-}$ mice display the same spontaneous activity (Frequency: $t = -0.3$, $df = 166.2$, $p = 0.7$; %SWB: $W = 3567$, $p = 0.8$) as WT mice (Fig. S3a,b and refs 9 and 21). Furthermore, VTA DA neurons from $\beta 2^{-/-}$ mice did not respond to acute injections of nicotine (30 μ g/kg) ($W = 148$, $p = 0.9$), whereas $\beta 4^{-/-}$ mice still responded to the drug ($W = 102$, $p < 0.001$) (Fig. S3c and refs 9,17,19 and 21). In response to acute injections of alcohol (500 mg/kg), VTA DA neurons from both $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice displayed a dose-dependent increase of their firing rate (Fig. 3a). ANOVA analysis for the firing frequency modification including WT and both transgenic




Figure 1. Acute and chronic nicotine enhance alcohol responses of VTA DA neurons. (a) (top) Typical electrophysiological recording showing the increase in firing rate of VTA DA cell induced by 500 mg/kg i.v. alcohol injection in WT mice. (bottom) Example responses of a VTA DA neuron to consecutive doses of alcohol. Arrowheads indicate the time of the injection. (b) (left) Mean \pm SEM DA cell firing frequency modification after injection of saline and the indicated ethanol dose (saline, $n = 28$; 125 mg/kg, $n = 13$; 250 mg/kg, $n = 14$; 500 mg/kg, $n = 14$; 750 mg/kg, $n = 7$). Arrowheads indicate the time of the injection. (right) Dose-response curve of ethanol-elicited responses for the same groups of neurons. *** $p < 0.001$, ** $p < 0.01$, Wilcoxon paired test from baseline. Horizontal lines indicate significant dose effect (one-way ANOVA). (c) (top) Example responses of a VTA DA neuron to consecutive injections of alcohol, nicotine and combined alcohol + nicotine. Arrowheads indicate the time of the injections. (bottom) Barplot of the maximum of firing frequency variation from baseline (mean \pm SEM) after alcohol (black), nicotine (white), or combined alcohol + nicotine injections (gray). *** $p < 0.001$, Wilcoxon paired test from baseline is indicated within each vertical bar. Difference between groups is indicated above the horizontal lines. *** $p < 0.001$, Wilcoxon paired test between groups. (d) (left) Examples of electrophysiological recordings of the basal activity of a DA neuron of a nic $-$ (black) and a nic $+$ (gray) mouse. Insets: Enlarged views of two or more action potentials. (right) Barplot of the mean frequency and %SWB for nic $-$ (black, $n = 92$) and nic $+$ mice (gray, $n = 65$). * $p < 0.05$, Wilcoxon test. (e) Dose-response curve of ethanol-elicited DA cell responses for nic $-$ (black) and nic $+$ (gray) mice. Mean \pm SEM of variation from baseline in firing frequency. Horizontal lines indicate significant dose effect and vertical lines indicate treatment effect (*** $p < 0.001$, * $p < 0.05$, two-way ANOVA. Nic $-$: saline: $n = 36$; 125 mg/kg: $n = 19$; 250 mg/kg: $n = 22$; 500 mg/kg: $n = 17$; 750 mg/kg: $n = 9$; nic $+$: saline: $n = 19$; 125 mg/kg: $n = 3$; 250 mg/kg: $n = 9$; 500 mg/kg: $n = 9$; 750 mg/kg: $n = 12$).

mice demonstrated a main effect of alcohol dose (Fig. 3b) ($F_{(3,115)} = 6.5$; $p < 0.001$) and revealed a genotype effect ($F_{(2,115)} = 3.6$, $p < 0.05$). Subsequent ANOVA between WT and $\beta 2^{-/-}$ mice showed only a dose effect ($F_{(3,75)} = 5.1$, $p < 0.01$), but no genotype ($F_{(1,75)} = 0.7$, $p = 0.2$), nor dose-genotype effect ($F_{(3,75)} = 0.1$, $p = 0.9$) was observed, thus indicating that DA neurons from $\beta 2^{-/-}$ mice were not different from those of WT mice in their responses to alcohol. On the contrary, subsequent ANOVA between WT and $\beta 4^{-/-}$ mice revealed a significant dose effect ($F_{(3,81)} = 4.8$, $p < 0.01$) together with a genotype effect ($F_{(1,81)} = 4.8$, $p < 0.05$), but no dose-genotype interaction ($F_{(3,81)} = 0.9$, $p = 0.4$), showing that responses to alcohol were significantly reduced in $\beta 4^{-/-}$ mice. Alcohol-induced %SWB variations in $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice were, contrarily to the variations in firing frequencies, not statistically different from those of WT mice (Fig. S1d) (Kruskal-Wallis: genotype effect: $X^2 = 1.6$, $df = 2$, $p = 0.4$). This could again be explained by the small effect of alcohol on the bursting activity of DA neurons. Having characterized alcohol-elicited VTA DA responses, we tested both transgenic mice for alcohol consumption in comparison to WT mice (Fig. 3c,d). ANOVA between WT and both transgenic mice revealed a main effect of alcohol concentration ($F_{(3,90)} = 27.2$; $p < 0.001$), no genotype effect ($F_{(2,30)} = 2.2$; $p = 0.13$), but a dose-genotype interaction ($F_{(6,90)} = 2.1$; $p = 0.05$). *Post hoc* analysis showed that $\beta 4^{-/-}$ mice consumed significantly more alcohol than WT mice when highest alcohol concentrations were presented (10%: $t = -3.3$, $df = 60.7$, $p < 0.05$; 15%: $t = -3.9$, $df = 47.7$, $p < 0.01$), while $\beta 2^{-/-}$ mice consumed similar amounts as WT mice (Fig. 3c). To further analyse this shift toward higher doses, we measured the intake modification switching from the dose of 10% to that of 15% in WT and mutant mice (Fig. S4a). We found that WT mice did not modify their intake when switching




Figure 2. Chronic nicotine modifies alcohol intake. (a) Mean \pm SEM of ethanol consumption (gr/kg) in nic $^{-}$ (black, $n = 18$) and nic $^{+}$ (gray, $n = 15$) mice during the two bottle choice procedure. Horizontal lines indicate significant dose effect and vertical lines indicate treatment effect. *** $p < 0.001$, ** $p < 0.01$, repeated measures two-way ANOVA. (b) Mean \pm SEM of the preference ratio for alcohol over total fluid intake for the same groups of mice. *** $p < 0.001$, ** $p < 0.01$, repeated measures two-way ANOVA; (c) Mean \pm SEM of the total fluids intake for the same groups of mice. (d) Barplot of the total alcohol intake (within the 18 days procedure) for the same groups. ** $p < 0.01$, Wilcoxon test.

between these two doses (Wilcoxon paired test, 10% vs 15%: $V = 2285$, $p = 0.3$), thus reaching a *plateau* in their consumption. In contrast, $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice kept increasing their consumption (Wilcoxon paired test, 10% vs 15%: $\beta 2^{-/-}$: $V = 340.5$, $p < 0.05$; $\beta 4^{-/-}$: $V = 112$, $p < 0.05$). The comparison of the intake modifications between the three groups revealed a statistical difference between WT and $\beta 4^{-/-}$ mice (Wilcoxon: $W = 921$, $p < 0.05$) but only a tendency between WT and $\beta 2^{-/-}$ mice (Wilcoxon: $W = 1558$, $p = 0.07$), thus confirming a role of $\beta 4$ nAChRs in mediating the acute effects of alcohol. Moreover, the preference ratio for alcohol was also modified in transgenic mice (Fig. 3d). ANOVA comparing the three genotypes showed a main significant dose effect ($F_{(3,90)} = 8.2$, $p < 0.001$), a significant genotype effect ($F_{(2,30)} = 2.6$, $p < 0.05$) but no dose-genotype interaction ($F_{(6,90)} = 0.8$, $p = 0.5$). Subsequent ANOVA between WT and $\beta 2^{-/-}$ mice revealed only a significant dose effect ($F_{(3,75)} = 8.1$, $p < 0.001$), but no genotype effect ($F_{(1,25)} = 1.7$, $p = 0.1$) nor dose-genotype interaction ($F_{(3,75)} = 0.3$, $p = 0.09$). ANOVA between WT and $\beta 4^{-/-}$ mice revealed a significant dose effect ($F_{(3,66)} = 7.4$, $p < 0.001$), a significant genotype effect ($F_{(1,22)} = 7.4$, $p < 0.05$), and no dose-genotype interaction ($F_{(3,66)} = 1.5$, $p = 0.2$).

Thus, these results demonstrated that, compared to WT mice, $\beta 4^{-/-}$, but not $\beta 2^{-/-}$ mice displayed both a modified VTA DA cells evoked response to alcohol and a modified alcohol drinking profile.




Figure 3. Alcohol-induced responses and alcohol reward are modified in $\beta 4^{-/-}$, but not in $\beta 2^{-/-}$ mice. (a) Typical electrophysiological recordings showing the increase in firing rate of VTA DA cells induced by 500 mg/kg i.v. alcohol injection in $\beta 2^{-/-}$ (top) and $\beta 4^{-/-}$ mice (bottom). (b) Dose-response curves of ethanol-elicited DA cell responses for WT (black), $\beta 2^{-/-}$ (red) and $\beta 4^{-/-}$ mice (blue). Horizontal lines indicate significant dose effect and vertical lines indicate strain effect. *** $p < 0.001$, * $p < 0.05$, two-way ANOVA (see Results section for statistical details). WT: saline: $n = 28$; 125 mg/kg: $n = 13$; 250 mg/kg: $n = 14$; 500 mg/kg: $n = 14$; 750 mg/kg: $n = 7$; $\beta 2^{-/-}$: saline: $n = 15$; 125 mg/kg: $n = 5$; 250 mg/kg: $n = 10$; 500 mg/kg: $n = 11$; 750 mg/kg: $n = 8$; $\beta 4^{-/-}$: saline: $n = 23$; 125 mg/kg: $n = 9$; 250 mg/kg: $n = 10$; 500 mg/kg: $n = 14$; 750 mg/kg: $n = 7$. (c) Mean \pm SEM of ethanol intake (gr/kg) in WT (black, $n = 18$), $\beta 2^{-/-}$ (red, $n = 9$) and $\beta 4^{-/-}$ mice (blue, $n = 6$) during the two bottle choice procedure. *** $p < 0.001$, repeated measures two-way ANOVA; ## $p < 0.01$, # $p < 0.05$, multiple comparisons with Bonferroni correction. (d) Mean \pm SEM of the preference ratio for alcohol over total fluid intake for the same groups of mice. *** $p < 0.001$; * $p < 0.05$, repeated measures two-way ANOVA (see Results section for statistical details).

Chronic nicotine has no effect on alcohol-elicited responses and alcohol intake in $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice. We then addressed the question of the impact of chronic nicotine exposure on the alcohol response in these mutant mice. Indeed, chronic exposure to nicotine induces a series of adaptations that particularly implicate heteromeric nAChRs³⁰. We thus asked whether the enhanced responses to alcohol induced by chronic nicotine exposure might rely on modifications of the expression or function of nAChRs in the DA system and thus analysed alcohol-elicited responses in $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice after chronic exposure to nicotine ($\beta 2^{nic+}$ and $\beta 4^{nic+}$ mice).

Surprisingly, in basal conditions, we found that, compared to their respective nic⁻ controls, $\beta 2^{nic+}$ mice displayed an increased basal firing rate and bursting activity (Δ Frequency: $t = -2.9$, $df = 36.4$, $p < 0.01$; Δ %SWB: $W = 558.5$, $p < 0.05$) (Fig. 4a), whereas $\beta 4^{nic+}$ mice showed a decreased firing rate and an unchanged bursting activity (Δ Frequency: $t = 2.7$, $df = 66.2$, $p < 0.01$; Δ %SWB: $W = 862$, $p = 0.9$) (Fig. 4b). However, when we compared the responses evoked by alcohol in nic⁻ and in nic⁺ mice of the two transgenic mouse lines, we




Figure 4. Alcohol-elicited responses and alcohol intake of $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice are not modified by chronic nicotine. (a) Barplot of the mean frequency and %SWB for $\beta 2^{nic-}$ (red, $n = 47$) and $\beta 2^{nic+}$ mice (pink, $n = 33$). $**p < 0.01$, t-test; $*p < 0.05$, Wilcoxon test. (b) Barplot of the mean frequency and %SWB for $\beta 4^{nic-}$ (blue, $n = 41$) and $\beta 4^{nic+}$ mice (light blue, $n = 42$). $*p < 0.05$, t-test. (c) Dose-response curves of ethanol-elicited DA cell responses for $\beta 2^{nic-}$ (red) and $\beta 2^{nic+}$ (pink). $\beta 2^{nic-}$: saline: $n = 15$; 125 mg/kg: $n = 5$; 250 mg/kg: $n = 10$; 500 mg/kg: $n = 11$; 750 mg/kg: $n = 8$; $\beta 2^{nic+}$: saline: $n = 12$; 125 mg/kg: $n = 6$; 250 mg/kg: $n = 7$; 500 mg/kg: $n = 10$; 750 mg/kg: $n = 5$. (d) Dose-response curves of ethanol-elicited DA cell responses for $\beta 4^{nic-}$ (blue) and $\beta 4^{nic+}$ (light blue). $\beta 4^{nic-}$: saline: $n = 23$; 125 mg/kg: $n = 9$; 250 mg/kg: $n = 10$; 500 mg/kg: $n = 14$; 750 mg/kg: $n = 7$; $\beta 4^{nic+}$: saline: $n = 15$; 125 mg/kg: $n = 7$; 250 mg/kg: $n = 10$; 500 mg/kg: $n = 8$; 750 mg/kg: $n = 6$. (e) Mean \pm SEM of ethanol consumption (gr/kg) in $\beta 2^{nic-}$ (red, $n = 9$) and $\beta 2^{nic+}$ (pink, $n = 8$) mice during the two bottle choice procedure. (f) Mean \pm SEM of ethanol consumption (gr/kg) in $\beta 4^{nic-}$ (blue, $n = 6$) and $\beta 4^{nic+}$ (light blue, $n = 6$) mice during the two bottle choice procedure.

observed no modification neither in the variation of the firing frequency (Fig. 4c,d), nor in the variation of %SWB (Fig. S1e,f). VTA DA neurons of $\beta 2\text{nic}+$ and $\beta 4\text{nic}+$ mice (unlike those of $\text{WTnic}+$ mice) showed responses to alcohol of the same amplitude as $\beta 2\text{nic}-$ and $\beta 4\text{nic}-$ mice, respectively ($\Delta\text{Frequency}$: ANOVA: $\beta 2\text{nic}-$ vs $\beta 2\text{nic}+$: dose effect: $F_{(4,78)} = 4.9$, $p < 0.01$; nicotine effect: $F_{(1,78)} = 0.5$, $p = 0.5$; dose \times nicotine effect: $F_{(4,78)} = 0.5$, $p = 0.7$; $\beta 4\text{nic}-$ vs $\beta 4\text{nic}+$: dose effect: $F_{(4,99)} = 8.0$, $p < 0.001$; nicotine effect: $F_{(1,99)} = 2.6$; $p = 0.1$; dose \times nicotine effect: $F_{(4,99)} = 0.2$, $p = 0.9$; $\Delta\% \text{SWB}$: Kruskal-Wallis: $\beta 2\text{nic}-$ vs $\beta 2\text{nic}+$, nicotine effect: $X^2 = 0.01$, $df = 1$, $p = 0.9$; $\beta 4\text{nic}-$ vs $\beta 4\text{nic}+$, nicotine effect: $X^2 = 0.7$, $df = 1$, $p = 0.4$). Similarly, $\beta 2\text{nic}$ and $\beta 4\text{nic}$ mice failed to increase alcohol consumption, showing the same intake profiles as their $\text{nic}-$ controls (Fig. 4e,f) ($\beta 2\text{nic}-$ vs $\beta 2\text{nic}+$: dose effect: $F_{(3,45)} = 33.9$, $p < 0.001$; nicotine effect: $F_{(1,15)} = 0.6$, $p = 0.4$; dose \times nicotine effect: $F_{(3,45)} = 2.3$; $p = 0.1$; $\beta 4\text{nic}-$ vs $\beta 4\text{nic}+$: dose effect: $F_{(3,30)} = 25$; $p < 0.001$; nicotine effect: $F_{(1,10)} = 0.1$; $p = 0.7$; dose \times nicotine effect: $F_{(3,30)} = 0.5$; $p = 0.7$). Interestingly, $\beta 2\text{nic}+$ and $\beta 4\text{nic}+$ kept showing the same increase in alcohol intake when switching from the dose of 10% to that of 15%, contrary to $\text{WT nic}+$ mice, for which the amount of alcohol consumed at the dose of 15% was not statistically different from the intake at the dose of 10% (Fig. S4b) (Wilcoxon paired test, 10% vs 15%: $\text{WT nic}+$: $V = 1647$, $p = 0.2$; $\beta 2\text{nic}+$: $V = 203$, $p < 0.01$; $\beta 4\text{nic}+$: $V = 87$, $p < 0.01$). However, it is important to note that these effects are not dependent on nicotine treatment, since they were already expressed in basal conditions, thus in absence of nicotine.

To summarize, in these mutant mice, despite an effect on the spontaneous VTA DA cells activity, chronic nicotine exposure does not increase VTA DA cells response nor alcohol intake. These results thus suggest that the lack of enhanced responses to alcohol or alcohol intake resulted from the absence of neuroadaptations at the level of $\beta 2$ and/or $\beta 4\text{nAChRs}$ expression and/or function that occurred during nicotine exposure in WT mice.

Discussion

The nicotine-alcohol interactions underlying the high incidence of co-addiction are a very complex phenomenon for which several mechanisms have been proposed. We focused our attention on the mesolimbic DA system that is a potential substrate for mechanistic interaction between these two drugs^{7,8,31}. We first studied the electrophysiological responses of VTA DA neurons to different doses of alcohol in order to establish the dose response curve in C57BL/6J WT mice. To evaluate the short-term interactions between alcohol and nicotine we performed concurrent injections of both drugs and compared them to the responses of each drug alone. A simultaneous injection of nicotine and alcohol led to a summation of the individual effects. Our data are in line with previous *in vitro* and *in vivo* studies showing additive and/or synergistic effects in the VTA to NAcc pathway^{25–27,32,33} and further supports the hypothesis that combined effects of nicotine and alcohol on the DA system may contribute to the high incidence of co-abuse. Moreover, this cumulative effect indicates that, at least at the tested doses, the effect of one does not occlude the other and, more importantly, that the two drugs may cooperate to enhance DA transmission increasing the sensation of pleasure and reward.

One of the possible factors contributing to the development of drug co-abuse is the cross-sensitization, in which the chronic use of one drug induces sensitization to the other. To address this question, we investigated the effects of chronic nicotine pre-exposure on the evoked responses of DA cells to alcohol and on alcohol intake. To achieve a constant level of nicotine, we chose a continuous mode of administration that avoids the need of repetitive injections. Continuous infusion of nicotine has been previously shown to induce behavioural and molecular adaptations, including nAChRs upregulation^{34–38}, which is considered an important feature of nicotine dependence³⁹. Our results showed that chronic nicotine increases the basal bursting activity of DA neurons in WT mice. Previous studies investigating the effects of a passive chronic infusion of nicotine in rodents reported divergent results, showing either no effect of nicotine treatment on the bursting activity or a decrease in the firing rate of DA cells^{36,40–42}. Dissimilar data can be explicated by diversities in nicotine regimen. For example, Besson *et al.*³⁶ and Tan *et al.*⁴⁰ used a different nicotine dose or treatment duration, whereas in Caillé *et al.*⁴¹ and in Grieder *et al.*⁴², MPs were removed 16 or 24 hours before the electrophysiological recordings, thus at the time when animals were already experiencing motivational withdrawal from nicotine. In the same paper, Caillé *et al.* also reported that voluntary nicotine self-administration, but not passive exposure, induced an increase in both the firing rate and bursting activity of VTA DA cells. These adaptations are in part mediated by the BNST (Bed Nucleus Stria Terminalis) glutamatergic drive onto the VTA, which has been suggested to be implicated in learning processes and memory. Nevertheless, the potentiation of glutamatergic signalling on VTA DA cells is not the sole mechanism accounting for bursting activity of DA neurons. We have shown in a previous paper¹⁷ that the cholinergic modulation (through $\beta 2\text{nAChRs}$) of GABA neurons is necessary for the bursting activity of DA neurons. In addition, it has been reported, that passive nicotine infusion through MPs induce an upregulation of nAChRs, in particularly $\alpha 4\beta 2\text{nAChRs}$ expressed on GABA neurons³⁷. In this context, we could speculate that, in mice and at this dose, passive nicotine induces nAChR upregulation, sufficiently to enhance the bursting activity of DA neurons, while the association between active responding and reward delivery is necessary to strengthen glutamatergic inputs from the BNST on DA cells. In addition to DA cell spontaneous firing, chronic nicotine also potentiated VTA DA cell responses to alcohol in WT animals. Such results highlight a neural adaptation, set up by nicotine pre-treatment, which affect alcohol responses and are in line with microdialysis studies showing an increase in alcohol-evoked DA release in the NAcc after repeated nicotine exposure^{43,44}. More interestingly, the sensitization of VTA DA neurons concurred with a sensitization of the animal to the rewarding properties of ethanol, demonstrated by an increased intake. Our results are thus in accordance with previous behavioural studies using different self-administration paradigms which report an increase in alcohol intake or preference after nicotine treatment^{28,29,45}.

Given the increasing evidences suggesting nAChRs as a common molecular substrate for alcohol and nicotine interaction^{12–15,25,26}, we tested $\beta 2-/-$ and $\beta 4-/-$ mice for alcohol responses in order to assess their involvement in alcohol action on the reward system. We found that $\beta 2^*\text{nAChRs}$ are not required for the acute effects of alcohol, given that VTA DA cells of $\beta 2-/-$ mice showed comparable responses to those of WT mice, and that

$\beta 2^{-/-}$ mice ingested unchanged amounts of alcohol. These findings are in agreement with the alcohol-drinking phenotype of $\beta 2^{-/-}$ mice⁴⁶ and confirm studies showing that selective blockade of $\beta 2^*$ nAChR affects neither alcohol consumption^{12,15} nor alcohol-elicited DA release in the NAcc^{47,48}. Interestingly, our data unveil a key role for $\beta 4^*$ nAChRs in mediating alcohol responses and in modulating its reinforcing properties, thereby defining the sensitivity of the reward system to alcohol. This result contrasts with a recent study showing that $\beta 4^{-/-}$ mice consumed similar amount of alcohol as WT mice⁴⁹. But this study not only differed for the concentrations of alcohol presented, but it used a completely different paradigm for alcohol access, the drinking-in-the-dark (DID) procedure, which is a model of binge drinking leading to high alcohol consumptions⁵⁰. The inverted relationship between VTA DA system sensitivity and drug self-administration that we observed in $\beta 4^{-/-}$ mice has been already described for $\alpha 5^{-/-}$ mice for nicotine⁵¹, where the decreased sensitivity of DA cells to nicotine was paralleled by a consumption shift to high doses. Deletion of $\beta 4^*$ nAChRs, which was shown to result in an increased sensitivity for nicotine²¹, results here in an opposite, decreased sensitivity for alcohol. $\beta 4^*$ nAChR subtype thus seems to be implicated, in both alcohol and nicotine responses, leading to a modification of the drug consumption. The downward shift of the dose-response curve of DA neurons found in $\beta 4^{-/-}$ could reflect i) a decreased sensitivity to the reinforcing properties of alcohol so that higher doses are needed to experience the pleasurable effect of alcohol or ii) a decreased sensitivity to the aversive effects. In this latter hypothesis, $\beta 4$ nAChRs could play a regulatory role acting as a “brake” in mediating the negative effects of alcohol, so that mice lacking $\beta 4$ nAChRs may experience fewer signs of aversive effects, which may facilitate alcohol intake at high doses.

The neuronal mechanisms for alcohol and nicotine interaction are not fully understood, yet we clearly show here that they involve the DA system and $\beta 2$ and $\beta 4^*$ nAChRs. Indeed both drugs induce similar responses on DA neurons and their effects are amplified when injected together. Our work also reveal $\beta 4^*$ nAChRs as possible actors implicated in the mediation of the acute effects of alcohol in the VTA, since the lack of this subunit modifies alcohol-evoked responses. Furthermore, it is well known that chronic nicotine exposure triggers a series of changes in nAChRs (distribution, stoichiometry or conformational state) but also various forms of synaptic plasticity, that outlast the presence of the drug and lead to the remodelling of neuronal circuits⁵². Among others, these adaptations underpin (i) modifications of VTA DA cells spontaneous activity and (ii) sensitization in VTA DA neuron responses to alcohol and in alcohol drinking behaviours. Our results suggest that nAChRs contributed to these two phenomena and that they seem to be independent. Indeed, despite chronic nicotine failed to induce any sensitization in DA responses to alcohol and in alcohol drinking behaviours in both $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice, it impacted VTA DA cells spontaneous activity. Yet, the level and the role of nicotine-induced neuroadaptations need further investigation, notably to elucidate whether cross-sensitization is due to molecular changes on nAChRs or rather to synaptic plasticity and to the consecutive reorganization of DA circuits. Understanding this latter point will be of utmost importance to identify specific molecular targets for the development of more effective pharmacological treatments against alcohol and tobacco addiction.

Materials and Methods

Animals. Adult (aged 8–16 weeks) male C57BL/6J wild-type (WT), $\beta 2^{-/-}$ ¹⁸ and $\beta 4^{-/-}$ mice⁵³ were used in this study. Both constitutive KO mice were backcrossed onto C57BL/6J background for at least 20 generation and bred in the same life conditions at Charles River (L'Arbresle, France). Experiments were performed after at least one week of habituation in our animal facility. Animal care and experiments were conducted in accordance with European Ethical Committee guidelines and approved by the Charles Darwin Animal Experimentation Ethical Committee.

In vivo electrophysiology. Single unit extracellular recordings were performed in anesthetized WT, $\beta 2^{-/-}$ and $\beta 4^{-/-}$ mice as detailed in the Supplementary Information. Briefly, glass electrodes containing 1.5% neurobiotin in 0.5% sodium acetate were lowered in the VTA according to stereotaxic coordinates derived from mouse brain atlas, and corrected empirically (antero-posterior: -3 to -4 mm; medio-lateral: 0.3 to 0.7 mm; dorso-ventral: -4 to -4 , 8 mm from bregma). Electrical signals were amplified by a high-impedance amplifier (Axon Instruments) and monitored audibly through an audio monitor (A.M. Systems Inc.). The signal was digitized, sampled at 25 kHz and recorded on a computer using Spike2 software (Cambridge Electronic Design) for later analysis. To distinguish DA from non-DA neurons the following parameters were used: 1) regular firing rate; 2) firing frequency between 1 and 10 Hz; 3) action potential duration between the beginning and the negative trough superior to 1.1 ms. Intravenous injections of nicotine (30 μ g/kg) and alcohol (125 mg/kg, 250 mg/kg, 500 mg/kg and 750 mg/kg) in the saphenous vein were performed in a final volume ranging from 20 to 120 μ l, as function of the administered dose (See SI for drugs specifications). Nicotine dose was chosen according to previous studies showing that nicotine can be intravenously self-administered at this dose in mice and on the base of our previous works^{17,19,24,54}. The range of alcohol doses injected were chosen on the base of previous *in vivo* electrophysiological studies in rats¹⁰. When possible, neurons were electroporated to allow neurobiotin internalization and labelling for neuron identification. D2 receptors pharmacology was performed on the last neuron of the experimental day (see SI and Fig. S5a,b).

Two bottle choice procedure. 24 hours voluntary drinking behaviour was carried out as specified in SI. Briefly, mice were offered water versus increasing concentrations of ethanol (3, 6, 10 and 15% (v/v)) within 18 days.

Osmotic mini-pumps. Surgical implantation of mini-pumps containing nicotine (10 mg/kg/d) or saline solution (0.9% NaCl) is described in SI. This dose was chosen on the base of previous works showing that in mice such dose correspond to nicotine plasma levels comparable to those sampled in smokers, generally ranged from 10 to 50 ng/ml^{35,55}.

Statistical analysis. The analyses were led using the R software (<http://www.r-project.org>). Firing frequency was quantified over 60 s periods, with a 45 s overlapping period. Percentage of spikes within bursts (%SWB) corresponds to the percentage of spikes discharged within bursts in a given time interval. A two-sample t-test was used to compare mean firing rate in two populations while a non-parametric Wilcoxon test was used for %SWB. Shapiro tests were used to test the normality of the data. Firing frequency response was quantified as a percentage of variation from baseline on a 3-min period before and after injection and means were calculated within each dose and each group. One-way or two-way ANOVAs were used to analyse dose-response curves and differences between groups. %SWB variation was calculated as a percentage of variation from baseline on a 3-min period before and after injection. A Kruskal-Wallis test was used to analyse dose-response curves and differences between groups. For alcohol intake and alcohol preference two-way ANOVA with repeated measures were used, followed by Bonferroni tests for *post hoc* analysis, when applicable.

References

- Dani, J. A. & Harris, R. A. Nicotine addiction and comorbidity with alcohol abuse and mental illness. *Nat Neurosci* **8**, 1465–1470 (2005).
- Hendrickson, L. M., Guildford, M. J. & Tapper, A. R. Neuronal nicotinic acetylcholine receptors: common molecular substrates of nicotine and alcohol dependence. *Front Psychiatry* **4**, 29 (2013).
- Falk, D. E., Yi, H.-Y. & Hiller-Sturmhöfel, S. An epidemiologic analysis of co-occurring alcohol and tobacco use and disorders: findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *Alcohol Res Health* **29**, 162–171 (2006).
- DiFranza, J. R. & Guarrera, M. P. Alcoholism and smoking. *J. Stud. Alcohol* **51**, 130–135 (1990).
- Bobo, J. K. & Husten, C. Sociocultural influences on smoking and drinking. *Alcohol Res Health* **24**, 225–232 (2000).
- Schlaepfer, I. R., Hofst, N. R. & Ehringer, M. A. The genetic components of alcohol and nicotine co-addiction: from genes to behavior. *Curr Drug Abuse Rev* **1**, 124–134 (2008).
- Doyon, W. M., Thomas, A. M., Ostroumov, A., Dong, Y. & Dani, J. A. Potential substrates for nicotine and alcohol interactions: a focus on the mesocorticolimbic dopamine system. *Biochemical Pharmacology* **86**, 1181–1193 (2013).
- Di Chiara, G. & Imperato, A. Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc. Natl. Acad. Sci. USA* **85**, 5274–5278 (1988).
- Mameli-Engvall, M. *et al.* Hierarchical Control of Dopamine Neuron-Firing Patterns by Nicotinic Receptors. *Neuron* **50**, 911–921 (2006).
- Gessa, G. L., Muntoni, F., Collu, M., Vargiu, L. & Mereu, G. Low doses of ethanol activate dopaminergic neurons in the ventral tegmental area. *Brain Res.* **348**, 201–203 (1985).
- Narahashi, T., Aistrup, G. L., Marszalec, W. & Nagata, K. Neuronal nicotinic acetylcholine receptors: a new target site of ethanol. *Neurochem. Int.* **35**, 131–141 (1999).
- Lê, A. D., Corrigan, W. A., Harding, J. W., Juzysch, W. & Li, T. K. Involvement of nicotinic receptors in alcohol self-administration. *Alcoholism: Clinical and Experimental Research* **24**, 155–163 (2000).
- Liu, L. *et al.* Nicotinic acetylcholine receptors containing the $\alpha 4$ subunit modulate alcohol reward. *Biol. Psychiatry* **73**, 738–746 (2013).
- Liu, L., Zhao-Shea, R., McIntosh, J. M. & Tapper, A. R. Nicotinic acetylcholine receptors containing the $\alpha 6$ subunit contribute to ethanol activation of ventral tegmental area dopaminergic neurons. *Biochemical Pharmacology* **86**, 1194–1200 (2013).
- Hendrickson, L. M., Zhao-Shea, R. & Tapper, A. R. Modulation of ethanol drinking-in-the-dark by mecamylamine and nicotinic acetylcholine receptor agonists in C57BL/6J mice. *Psychopharmacology* **204**, 563–572 (2009).
- Changeux, J. P. & Edelstein, S. J. Allosteric receptors after 30 years. *Neuron* **21**, 959–980 (1998).
- Tolu, S. *et al.* Co-activation of VTA DA and GABA neurons mediates nicotine reinforcement. *Mol. Psychiatry* **18**, 382–393 (2013).
- Picciozzo, M. R. *et al.* Acetylcholine receptors containing the beta2 subunit are involved in the reinforcing properties of nicotine. *Nature* **391**, 173–177 (1998).
- Maskos, U. *et al.* Nicotine reinforcement and cognition restored by targeted expression of nicotinic receptors. *Nature* **436**, 103–107 (2005).
- Frahm, S. *et al.* Aversion to nicotine is regulated by the balanced activity of $\beta 4$ and $\alpha 5$ nicotinic receptor subunits in the medial habenula. *Neuron* **70**, 522–535 (2011).
- Harrington, L. *et al.* Role of $\beta 4^*$ Nicotinic Acetylcholine Receptors in the Habenulo-Interpeduncular Pathway in Nicotine Reinforcement in Mice. 1–13, doi: 10.1038/npp.2015.346 (2015).
- Bierut, L. J. Convergence of genetic findings for nicotine dependence and smoking related diseases with chromosome 15q24-25. *Trends in Pharmacological Sciences* **31**, 46–51 (2010).
- Sherva, R. *et al.* Variation in nicotinic acetylcholine receptor genes is associated with multiple substance dependence phenotypes. *Neuropsychopharmacology* **35**, 1921–1931 (2010).
- Eddine, R. *et al.* A concurrent excitation and inhibition of dopaminergic subpopulations in response to nicotine. *Sci Rep* **5**, 8184 (2015).
- Tizabi, Y., Copeland, R. L., Louis, V. A. & Taylor, R. E. Effects of combined systemic alcohol and central nicotine administration into ventral tegmental area on dopamine release in the nucleus accumbens. *Alcoholism: Clinical and Experimental Research* **26**, 394–399 (2002).
- Tizabi, Y., Bai, L., Copeland, R. L. & Taylor, R. E. Combined effects of systemic alcohol and nicotine on dopamine release in the nucleus accumbens shell. *Alcohol and Alcoholism* **42**, 413–416 (2007).
- Truitt, W. A. *et al.* Ethanol and nicotine interaction within the posterior ventral tegmental area in male and female alcohol-preferring rats: evidence of synergy and differential gene activation in the nucleus accumbens shell. *Psychopharmacology* **232**, 639–649 (2015).
- Doyon, W. M. *et al.* Nicotine Decreases Ethanol-Induced Dopamine Signaling and Increases Self-Administration via Stress Hormones. *Neuron*, doi: 10.1016/j.neuron.2013.06.006 (2013).
- Leao, R. M. *et al.* Chronic Nicotine Activates Stress/Reward-Related Brain Regions and Facilitates the Transition to Compulsive Alcohol Drinking. *J. Neurosci.* **35**, 6241–6253 (2015).
- Buisson, B. & Bertrand, D. Nicotine addiction: the possible role of functional upregulation. *Trends in Pharmacological Sciences* **23**, 130–136 (2002).
- Funk, D., Marinelli, P. W. & Lê, A. D. Biological processes underlying co-use of alcohol and nicotine: neuronal mechanisms, cross-tolerance, and genetic factors. *Alcohol Res Health* **29**, 186–192 (2006).
- Clark, A. & Little, H. J. Interactions between low concentrations of ethanol and nicotine on firing rate of ventral tegmental dopamine neurones. *Drug and Alcohol Dependence* **75**, 199–206 (2004).
- Engle, S. E., McIntosh, J. M. & Drenan, R. M. Nicotine and ethanol cooperate to enhance ventral tegmental area AMPA receptor function via $\alpha 6$ -containing nicotinic receptors. *Neuropharmacology* **91**, 13–22 (2015).
- Rowell, P. P. & Li, M. Dose-response relationship for nicotine-induced up-regulation of rat brain nicotinic receptors. *Journal of Neurochemistry* **68**, 1982–1989 (1997).

35. Marks, M. J. *et al.* Subsets of acetylcholine-stimulated 86Rb⁺ efflux and [125I]-epibatidine binding sites in C57BL/6 mouse brain are differentially affected by chronic nicotine treatment. *Neuropharmacology* **46**, 1141–1157 (2004).
36. Besson, M. *et al.* Long-term effects of chronic nicotine exposure on brain nicotinic receptors. *Proc. Natl. Acad. Sci. USA* **104**, 8155–8160 (2007).
37. Nashmi, R. *et al.* Chronic nicotine cell specifically upregulates functional alpha 4* nicotinic receptors: basis for both tolerance in midbrain and enhanced long-term potentiation in perforant path. *J. Neurosci.* **27**, 8202–8218 (2007).
38. Metaxas, A. *et al.* Genetic deletion of the adenosine A(2A) receptor prevents nicotine-induced upregulation of $\alpha 7$, but not $\alpha 4\beta 2^*$ nicotinic acetylcholine receptor binding in the brain. *Neuropharmacology* **71**, 228–236 (2013).
39. Govind, A. P., Vezina, P. & Green, W. N. Nicotine-induced upregulation of nicotinic receptors: underlying mechanisms and relevance to nicotine addiction. *Biochemical Pharmacology* **78**, 756–765 (2009).
40. Tan, H., Bishop, S. F., Lauzon, N. M., Sun, N. & Laviolette, S. R. Chronic nicotine exposure switches the functional role of mesolimbic dopamine transmission in the processing of nicotine's rewarding and aversive effects. *Neuropharmacology* **56**, 741–751 (2009).
41. Caille, S., Guillem, K., Cador, M., Manzoni, O. & Georges, F. Voluntary Nicotine Consumption Triggers *In Vivo* Potentiation of Cortical Excitatory Drives to Midbrain Dopaminergic Neurons. *J. Neurosci.* **29**, 10410–10415 (2009).
42. Grieder, T. E. *et al.* Dopaminergic signaling mediates the motivational response underlying the opponent process to chronic but not acute nicotine. *Neuropsychopharmacology* **35**, 943–954 (2010).
43. Blomqvist, O., Ericson, M., Johnson, D. H., Engel, J. A. & Söderpalm, B. Voluntary ethanol intake in the rat: effects of nicotinic acetylcholine receptor blockade or subchronic nicotine treatment. *European Journal of Pharmacology* **314**, 257–267 (1996).
44. Ding, Z.-M. *et al.* Repeated exposure of the posterior ventral tegmental area to nicotine increases the sensitivity of local dopamine neurons to the stimulating effects of ethanol. *Alcohol* **46**, 217–223 (2012).
45. Hauser, S. R. *et al.* Nicotine modulates alcohol-seeking and relapse by alcohol-preferring (P) rats in a time-dependent manner. *Alcoholism: Clinical and Experimental Research* **36**, 43–54 (2012).
46. Kamens, H. M., Andersen, J. & Picciotto, M. R. Modulation of ethanol consumption by genetic and pharmacological manipulation of nicotinic acetylcholine receptors in mice. *Psychopharmacology* **208**, 613–626 (2010).
47. Ericson, M., Molander, A., Löf, E., Engel, J. A. & Söderpalm, B. Ethanol elevates accumbal dopamine levels via indirect activation of ventral tegmental nicotinic acetylcholine receptors. *European Journal of Pharmacology* **467**, 85–93 (2003).
48. Larsson, A., Svensson, L., Söderpalm, B. & Engel, J. A. Role of different nicotinic acetylcholine receptors in mediating behavioral and neurochemical effects of ethanol in mice. *Alcohol* **28**, 157–167 (2002).
49. Patkar, O. L., Belmer, A., Tarren, J. R., Holgate, J. Y. & Bartlett, S. E. The effect of varenicline on binge-like ethanol consumption in mice is $\beta 4$ nicotinic acetylcholine receptor-independent. *Neuroscience Letters* **633**, 235–239 (2016).
50. Leeman, R. F. *et al.* Ethanol consumption: how should we measure it? Achieving concisence between human and animal phenotypes. *Addict Biol* **15**, 109–124 (2010).
51. Morel, C. *et al.* Nicotine consumption is regulated by a human polymorphism in dopamine neurons. *Mol. Psychiatry* **19**, 930–936 (2013).
52. Lüscher, C. & Malenka, R. C. Drug-evoked synaptic plasticity in addiction: from molecular changes to circuit remodeling. *Neuron* **69**, 650–663 (2011).
53. Xu, W. *et al.* Multiorgan autonomic dysfunction in mice lacking the beta2 and the beta4 subunits of neuronal nicotinic acetylcholine receptors. *J. Neurosci.* **19**, 9298–9305 (1999).
54. Picciotto, M. R. *et al.* Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptor in the brain. *Nature* **374**, 65–67 (1995).
55. Matta, S. G. *et al.* Guidelines on nicotine dose selection for *in vivo* research. *Psychopharmacology* **190**, 269–319 (2007).

Acknowledgements

We would like to thank Dr. Alexandre Mourot, Dr. Jérémie Naudé and Dr. Sébastien Valverde for comments on the manuscript. This work was supported by the Centre National de la Recherche Scientifique CNRS UMR 8246, the Foundation for Medical Research (FRM, Equipe FRM 2013 to P.F.), the Bettencourt Schueller Foundation (Coup d'Élan 2012 to P.F.), the Labex Bio-Psy, the Institut de Recherches Scientifiques sur les Boissons (IREB 2013-05) and the Fondation pour la Recherche en alcoologie (FRA 2016-06). The laboratory of P.F. is part of the École des Neurosciences de Paris Ile-de-France RTRA network. P.F. is members of the Laboratory of Excellence, LabEx Bio-Psy, DHU Pepsy.

Author Contributions

S.T. and P.F. designed the study. S.T. performed the electrophysiological recordings and the behavioural experiments. F.M., C.M., C.P. and R.d.B. participated in collecting electrophysiological data. N.T. contributed to behavioural experiments. S.P. provided KO mice. S.T. and P.F. analysed the data and wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing Interests: The authors declare no competing financial interests.

How to cite this article: Tolu, S. *et al.* Nicotine enhances alcohol intake and dopaminergic responses through $\beta 2^*$ and $\beta 4^*$ nicotinic acetylcholine receptors. *Sci. Rep.* **7**, 45116; doi: 10.1038/srep45116 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017