



**HAL**  
open science

## Early born neurons are abnormally positioned in the doublecortin knockout hippocampus

Reham Khalaf-Nazzal, Melissa A. Stouffer, Robert Olaso, Leila Muresan, Audrey Roumegous, Virginie Lavilla, Wassila Carpentier, Imane Moutkine, Sylvie Dumont, Benoit Albaud, et al.

► **To cite this version:**

Reham Khalaf-Nazzal, Melissa A. Stouffer, Robert Olaso, Leila Muresan, Audrey Roumegous, et al.. Early born neurons are abnormally positioned in the doublecortin knockout hippocampus. *Human Molecular Genetics*, 2017, 26 (1), pp.90-108. 10.1093/hmg/ddw370 . hal-01512718

**HAL Id: hal-01512718**

**<https://hal.sorbonne-universite.fr/hal-01512718>**

Submitted on 24 Apr 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Early born neurons are abnormally positioned in the doublecortin knockout hippocampus

#Reham Khalaf-Nazzal<sup>1-3§</sup>, #Melissa A. Stouffer<sup>1-3</sup>, Robert Olasso<sup>4</sup>, Leila Muresan<sup>5-7</sup>, Audrey Roumegous<sup>1-3</sup>, Virginie Lavilla<sup>4</sup>, Wassila Carpentier<sup>8</sup>, Imane Moutkine<sup>1-3</sup>, Sylvie Dumont<sup>9</sup>, Benoit Albaud<sup>10</sup>, Nicolas Cagnard<sup>11</sup>, Hugues Roest Crolius<sup>5-7</sup>, \*Fiona Francis<sup>1-3</sup>.

<sup>1</sup> INSERM UMR-S 839, Paris 75005;

<sup>2</sup> Sorbonne Universités, Université Pierre et Marie Curie, Paris 75005;

<sup>3</sup> Institut du Fer à Moulin, Paris 75005, France

<sup>4</sup> Plateforme de Transcriptomique, Laboratoire de Recherche Translationnelle, CEA/DSV/IG-Centre National de Genotypage, 2 rue Gaston Crémieux, 91057 Evry, France

<sup>5</sup> Ecole Normale Supérieure, Institut de Biologie de l'ENS, IBENS, Paris, F-75005 France.

<sup>6</sup> Inserm, U1024, Paris, F-75005 France.

<sup>7</sup> CNRS, UMR 8197, Paris, F-75005 France.

<sup>8</sup> Plateforme post-génomique de la Pitié-Salpêtrière, Faculty of medicine, 75013, Paris.

<sup>9</sup> Sorbonne Universités, UPMC Paris 06, UMS30 LUMIC, plateforme d'histomorphologie, St Antoine F-75012 PARIS

<sup>10</sup> Plateforme Affymetrix, Institut Curie, Hospital St Louis, Paris

<sup>11</sup> Plateforme Bio-informatique Paris Descartes, Faculté de Necker, 156 rue de Vaugirard, 75730 Paris Cedex 15.

§ Current address, Central research laboratory-A, Developmental Neurosciences and Neurogenetics Group, Faculty of Medicine and Health Sciences, An-Najah National University, Nablus, Palestine

# These authors contributed equally to this work.

\*Correspondance to: Fiona Francis, Institut du Fer à Moulin, 17 rue du Fer à Moulin, 75005 Paris France. Email: [fiona.francis@inserm.fr](mailto:fiona.francis@inserm.fr)

**ABSTRACT**

Human *doublecortin* (*DCX*) mutations are associated with severe brain malformations leading to aberrant neuron positioning (heterotopia), intellectual disability and epilepsy. The *Dcx* protein plays a key role in neuronal migration, and hippocampal pyramidal neurons in *Dcx* knockout (KO) mice are disorganized. The single CA3 pyramidal cell layer observed in wild type (WT) is present as two abnormal layers in the KO, and CA3 KO pyramidal neurons are more excitable than WT. *Dcx* KO mice also exhibit spontaneous epileptic activity originating in the hippocampus. It is unknown however, how hyperexcitability arises and why two CA3 layers are observed.

Transcriptome analyses were performed to search for perturbed postnatal gene expression, comparing *Dcx* KO CA3 pyramidal cell layers with WT. Gene expression changes common to both KO layers indicated mitochondria and Golgi apparatus anomalies, as well as increased cell stress. Intriguingly, gene expression analyses also suggested that the KO layers differ significantly from each other, particularly in terms of maturity. Layer-specific molecular markers and BrdU birthdating to mark the final positions of neurons born at distinct timepoints revealed inverted layering of the CA3 region in *Dcx* KO animals. Notably, many early-born 'outer boundary' neurons are located in an inner position in the *Dcx* KO CA3, superficial to other pyramidal neurons. This abnormal positioning likely affects cell morphology and connectivity, influencing network function. Dissecting this *Dcx* KO phenotype sheds light on coordinated developmental mechanisms of neuronal subpopulations, as well as gene expression patterns contributing to a bi-layered malformation associated with epilepsy.

## INTRODUCTION

The development of neuronal circuits necessitates an intricate sequence of events including cell proliferation, migration, differentiation and synaptogenesis. The hippocampus is a laminated structure that develops following these steps, similar to the neocortex (1). There is evidence for birthdate-dependent layering of principal neurons with deep to superficial molecular and connectivity gradients (2-4). In the rodent, this pattern is induced from mid-embryogenesis, and newly born neurons carry intrinsic properties to help establish their adult phenotypes (5). A population of the earliest born neurons have specifically been shown to function as highly connected hub neurons in juvenile CA3 (6), helping to organize circuit development. Clonal analyses have also shown that related hippocampal pyramidal neurons derived from the same progenitor share interneuron connectivity (7). Thus, the timing and manner that hippocampal neurons are generated are critical for adult function.

Pyramidal cells of the major CA1 and CA3 fields are molecularly distinguishable (3, 8-10). Gene expression boundaries along the radial and septo-temporal axes distinguish hippocampal cell subpopulations related to anatomical position, birthdate and connectivity patterns (3, 5, 11). Subpopulations of interconnected neurons in different hippocampal fields have been found to share similar patterns of gene expression, birthdate and time window of synaptogenesis, these microcircuits being associated with hippocampal function (5, 12).

Mutations in genes associated with neurodevelopmental syndromes often reveal abnormal hippocampi in mouse models. This is characteristic of *doublecortin (Dcx)*, *Lis1*, and *alpha 1 tubulin (Tubal1)* mutations, which in human are associated with cortical malformations featuring neuronal disorganization, including lissencephaly and subcortical band heterotopia (SBH) that present clinically with varying degrees of intellectual disability (ID) and epilepsy (13-16). Mouse models show common but variable defects in hippocampal fields which are often disorganized (17-20). For unknown reasons, hippocampal defects appear to be a signature

1  
2  
3 of type 1 lissencephaly mouse models, more so than neocortical abnormalities (21). Gene  
4  
5 expression patterns that distinguish the features of abnormally positioned neurons have not yet  
6  
7 been reported, which could shed light on their origins and functions.  
8  
9

10  
11 *Dcx* is a developmentally regulated microtubule associated protein (MAP) important for the  
12  
13 stabilization of microtubules in migrating and differentiating neurons (22-25). In *Dcx* KO mice,  
14  
15 hippocampal pyramidal cells are abnormally positioned, mostly in the CA3 region, where the  
16  
17 single pyramidal cell layer observed in wild type (WT) is divided into two layers (18, 19). Video  
18  
19 EEG recordings showed that *Dcx* KO mice have spontaneous epilepsy with the abnormal  
20  
21 electrical activity originating in the hippocampus and secondarily propagating to the cortex  
22  
23 (26). Extracellular and whole cell recordings in the *Dcx* KO CA3 region found that disorganized  
24  
25 cells are more excitable than WT pyramidal cells (26, 27). Field recordings from *Dcx* KO brain  
26  
27 slices suggested that pyramidal cells make a primary contribution to the spontaneous  
28  
29 epileptiform activity, whereas inhibition remained strong during interictal events (27). It is  
30  
31 hence important to further characterize *Dcx* KO CA3 pyramidal cells to investigate the  
32  
33 mechanistic link between mutation of this protein, heterotopic cell positioning and  
34  
35 hyperexcitability.  
36  
37  
38  
39  
40

41  
42  
43 In this study, we assessed global gene expression in the abnormal KO CA3 layers compared  
44  
45 to the WT monolayer at postnatal day 0 (P0). At this stage of development, most CA3 neurons  
46  
47 are expected to have arrived at their final destinations and to be growing axons and dendrites  
48  
49 (2, 28). Gene expression changes at this stage may therefore contribute to abnormal morphology  
50  
51 and circuitry seen in the mature brain (26, 27). We found that both KO layers have perturbed  
52  
53 expression of genes indicating organelle abnormalities, increased cell stress and perturbed cell  
54  
55 metabolism. Comparing KO layers also identified an unexpected inversion of a subpopulation  
56  
57 of early-born KO neurons. Gene ontologies, layer-specific molecular markers and BrdU  
58  
59 birthdating experiments confirmed transcriptome data suggesting that many of the most mature  
60

1  
2  
3 neurons are in an abnormal position and tissue environment at P0. Given the distinct molecular  
4 identities and connectivity patterns conferred by birthdate, this perturbation likely influences  
5  
6 neuron morphology and circuit formation, potentially contributing to the development of  
7  
8 hyperexcitability in this mouse model.  
9  
10  
11

## 12 13 14 RESULTS

### 15 16 17 Laser capture microdissection (LCM) of CA3 cells in *Dcx* KO and WT

18  
19 In the *Dcx* KO hippocampus, the two abnormal bands of pyramidal cells in the CA3 region,  
20 termed here *stratum pyramidale* internal (SPI, closest to the *stratum radiatum*) and *stratum*  
21 *pyramidale* external (SPE, closest to the *stratum oriens*), are visible by late embryonic stages  
22  
23 (Fig. 1). LCM at P0 allowed us to specifically separate SPI from SPE, and a similar region was  
24  
25 excised from WT CA3 (Fig. 1). Reverse transcriptase (RT) PCR from LCM material confirmed  
26  
27 that the microdissected samples expressed the CA3 marker *KAI* (*Grik4*) (9), but did not express  
28  
29 the CA1 marker *Scip* nor the dentate gyrus marker *Prox1* (Supp. Fig. 1).  
30  
31  
32  
33  
34

35 Picochip analyses for RNA quality of CA3 LCM samples (n = 5 per condition) were  
36 compared to analyses of RNAs from whole brain sections that had not been subjected to LCM.  
37  
38 The latter had high RNA integrity number (RIN) values of 9 on average, indicating that the  
39 sectioned and stained brain material was of good quality (Supp. Table 1). The LCM samples  
40 displayed a lower RNA quality, which is expected from this procedure, with RIN values  
41  
42 between 5.2 and 7.2 (Supp. Fig. 2, Supp. Table 1). Overall, RNA integrity showed sufficient  
43  
44 quality and reproducibility to proceed with microarray analyses.  
45  
46  
47  
48  
49  
50  
51  
52

### 53 54 55 Differential gene expression between the WT CA3 region and *Dcx* KO SPI and SPE

56  
57 Reverse transcription, linear amplification and hybridization to microarrays were performed  
58 according to Illumina protocols. Of the 25,600 well-annotated transcript probes on the *Illumina*  
59  
60 *MouseRef-8 v2.0 Expression BeadChip* array, 11,800 (45%) on average demonstrated a

1  
2  
3 detectable level of expression amongst the 15 samples (GEO accession GSE87019). After data  
4 normalization (**Supp. Fig. 3**), 2361 and 1940 transcripts were found to be differentially  
5 expressed in SPI and SPE cells respectively compared to WT when the  $P$  value cut-off was set  
6 to  $< 0.01$ , and 1758 and 1202 transcripts when  $P < 0.005$  (**Fig. 2A, Supp. Table 2**). Considering  
7 all SPI versus WT changes with a  $P$  value  $< 0.005$ , an approximately equal number of transcripts  
8 were up- (52 %) and down-regulated (48 %) (**Fig. 2A, Supp. Table 3A**). For SPE versus WT,  
9 a greater proportion of transcripts were up-regulated (67 %), and only 33 % were down-  
10 regulated (**Fig. 2A, Supp. Table 3A**). Considering genes showing a greater than 1.5 X change  
11 compared to WT ( $P < 0.005$ ), there were 49 up-regulated in SPI and 4 in SPE, and 11 down-  
12 regulated in SPI and 3 in SPE (**Supp. Table 3B**). As expected, *Dcx* appeared amongst the most  
13 significantly down-regulated in both SPI and SPE gene lists.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29  
30 Microarray analyses confirmed that WT and both KO cell populations expressed the  
31 CA3 field-specific markers *KAI* (*Grik4*) (9) and *Bok* (29). Focusing on genes that were  
32 commonly deregulated in both SPI and SPE KO cells compared to WT, 372 and 224 genes with  
33  $P$  values of  $< 0.01$  and  $< 0.005$ , respectively, were identified (**Supp. Table 4**). Some of these  
34 common gene expression changes may be directly and specifically related to the inactivation of  
35 *Dcx* (see below for genes coding protein partners), or due to deregulated cell processes. We  
36 performed DAVID (Database for Annotation, Visualization and Integrated Discovery)  
37 functional annotation clustering analyses for the 372 commonly differentially expressed genes  
38 ( $P < 0.01$ ; **Fig. 2B, Supp. Table 5A**). Significant categories include nucleotide binding and  
39 translation, as well as Golgi apparatus and mitochondrion, the latter likely to indicate organelle  
40 abnormalities, which indeed coincides with our previous observations by electron microscopy  
41 (30). DNA repair and chaperone gene clusters also appeared as significantly different in both  
42 KO layers compared to WT, suggesting abnormalities in cell stress responses.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Indications for increased cell stress in *Dcx* KO layers at P0

Perturbed expression of genes related to organelles can be fundamentally linked with increased cell stress (31, 32). Expression of DNA repair proteins and upregulation of chaperones are known responses to oxidative stress that occur, for example, when the production of reactive oxygen species (ROS) is increased (33, 34). A significant number of genes involved in repairing DNA were found upregulated in SPI and SPE, including genes implicated in excision repair (*Ercc5*) and double-strand break repair (*Hmgb2*) (**Table 1**). Genes involved in protection against oxidative stress, including subunits of glutathione S-transferase (*Gstm1*, *Gstm5*) and sulfiredoxin (*Srxn1*), were also upregulated in SPI and/or SPE (**Table 1**).

Deregulated expression of heat shock protein (HSP) genes was also observed in both KO layers (**Table 1**). HSPs are highly conserved proteins that are often upregulated in response to several types of cell stress, functioning as molecular chaperones to help nascent protein folding or to refold damaged proteins. Two important HSPs upregulated in both SPE and SPI are *Hsp90ab1* and *Hspd1*. *Hsp90ab1* is located in the cytosol and interacts with protein kinases and transcription factors that have crucial roles in developmental processes involved in cell survival upon exposure to various stressful stimuli (34, 35). *Hspd1* (Hsp60) is a mitochondrial matrix chaperone, and deficits in this protein are associated with hypomyelination and neurodegenerative disorders (36). Also related to mitochondrial function, *Dnaja2*, upregulated in both SPE and SPI, functions as a co-chaperone of Hsp70s in protein folding and mitochondrial protein import (37). Other HSPs were conversely downregulated. For example, *Hspa8*, downregulated in SPI, is a constitutively expressed chaperone important for protein import into the endoplasmic reticulum and mitochondria, ubiquitin-proteasome degradation, autophagy, and disassembly of clathrin-coated vesicles (38). Also downregulated in SPI and SPE is *Herpud1*, a component of the endoplasmic reticulum quality control system involved in



1  
2  
3 ubiquitin-dependent degradation of misfolded proteins (39). These proteins, related to organelle  
4  
5 function, would be expected to contribute to cell stress when downregulated.  
6  
7

8         Some changes in cell stress genes were specific to SPI. For example, pleiotrophin (*Ptn*)  
9  
10 was up-regulated in SPI cells. It is a heparin-binding cytokine (40) that acts as a neuroprotective  
11  
12 factor to prevent stress-induced cell death (41). The metallothioneins Mt1 (astrocytes >  
13  
14 neurons) and Mt3 (brain specific isoform, neurons > astrocytes) were also upregulated only in  
15  
16 SPI (**Table 1**). Mts are intracellular proteins that bind metals including zinc, and function in  
17  
18 redox homeostasis (42). Mt overexpression may protect against apoptosis; zinc is a major  
19  
20 inducer of its expression, as is oxidative stress and inflammation (42).  
21  
22  
23

24         Other indications of oxidative stress are related to proline metabolism. The  
25  
26 mitochondrial matrix gene *Aldh4a1* (P5cd) (**Table 1**), critical for proline catabolism, is  
27  
28 downregulated, and mutations in this gene in humans lead to hyperprolinemia, a disease  
29  
30 associated with ID and epilepsy. Additionally, the proline transporter *Slc6a7* is upregulated in  
31  
32 SPI (**Table 1**), further suggesting that intracellular proline could be increased. This Na<sup>+</sup>  
33  
34 dependent transporter is localized to synaptic vesicles and excitatory nerve terminals,  
35  
36 corresponding to the proposed role of proline in ion channel and neurotransmitter receptor  
37  
38 regulation (43). Proline metabolism is connected to the pentose phosphate pathway and redox  
39  
40 homeostasis, and can also serve as a source of glutamate, ATP and ROS (44). On the other  
41  
42 hand, proline might function as an antioxidant by decreasing ROS production (45). Thus several  
43  
44 genes, some clearly linked to organelle function, are indicative of cell stress in P0 *Dcx* KO cells.  
45  
46  
47  
48  
49  
50  
51  
52

### 53 **Metabolism genes are perturbed in KO layers**

54  
55 Potentially also linked to cell stress and/or organelle abnormalities, we observed several  
56  
57 metabolic genes that are commonly deregulated in SPI and SPE compared to WT (**Table 2A**).  
58  
59 For example, phosphofructokinase, platelet type (*Pfkip*), which catalyses the rate-limiting step  
60

1  
2  
3 in glycolysis, phosphoglycerate mutase 1 (*Pgam1*), another glycolytic enzyme, and hexokinase  
4  
5 1 (*Hki*), an enzyme localized to the outer mitochondrial membrane that catalyzes the first step  
6  
7 of glycolysis, are increased in both SPI and SPE. Genes involved in the Krebs cycle are  
8  
9 upregulated in both layers including an ATP synthase subunit (*Atp5a1*), pyruvate  
10  
11 dehydrogenase (*Pdhb*) and isocitrate dehydrogenase 3 (*Idh3g*). The latter enzyme has a role in  
12  
13 redox homeostasis by generation of NADH/NADPH. On the other hand, there is a decrease in  
14  
15 transaldolase 1 (*Taldo1*) in both SPI and SPE, an enzyme of the non-oxidative branch of the  
16  
17 pentose phosphate pathway.  
18  
19  
20  
21

22 Other notable changes in metabolism are related to lipids (**Table 2A**). Downregulation  
23  
24 of genes related to lipid synthesis in SPI and SPE include ATP citrate lyase (*Acly*), acetoacetyl-  
25  
26 CoA synthetase (*Aacs*) and phosphatidylglycerophosphate synthase 1 (*Pgs1*). Two enzymes  
27  
28 involved in biosynthesis of cholesterol, mevalonate decarboxylase (*Mvd*) and hydroxysteroid  
29  
30 17-beta dehydrogenase 7 (*Hsd17b7*), are also downregulated in both KO layers. Furthermore  
31  
32 there is upregulation of three sphingomyelin phosphodiesterase genes coding for enzymes that  
33  
34 break down sphingolipids (*Smpdl3a*, *Smpd4*, *Smpd1*), and upregulation of acyl-CoA  
35  
36 thioesterase 1 (*Acot1*) that hydrolyses lipids (acyl-CoAs). Diazepam binding inhibitor (*Dbi*),  
37  
38 also known as acyl-CoA binding protein, has been shown to regulate long-chain fatty acid  
39  
40 metabolism, and is upregulated to a greater extent in SPI than SPE (46). Plasticity related  
41  
42 protein 3 (*Plppr1*), downregulated in SPI, generates lipids important for cell signaling events  
43  
44 including promotion of neurite growth (47). Overall, perturbed lipid metabolism may have  
45  
46 numerous effects on cell signaling, function of membrane-bound organelles and particularly the  
47  
48 plasma membrane of which cholesterol and sphingolipids are important components.  
49  
50  
51  
52  
53  
54  
55  
56  
57

58 **SPI layer cells are distinguishable from SPE layer cells**  
59  
60

1  
2  
3 We next queried layer-specific gene expression. Many differentially expressed genes were  
4 specific to either SPE versus WT, or SPI versus WT (**Supp. Fig. 4**). DAVID analyses to  
5 compare these up- and down-regulated genes ( $P < 0.005$ ) revealed that functional categories  
6 were indeed different between SPE and SPI (**Supp. Table 5B-E**). The major categories with  
7 the highest enrichment scores are summarized in **Fig. 3**. For SPI up-regulated genes, enriched  
8 categories included several related to synaptic function, cell adhesion and metabolism. Some  
9 of the latter genes were identified in previous analyses (**Table 2B**). Concerning SPE up-  
10 regulated genes, enriched categories included organelle lumen, RNA processing and  
11 mitochondrion. SPE down-regulated genes included membrane, GTP binding and actin  
12 binding, whereas SPI down-regulated gene categories corresponded to those upregulated in  
13 SPE (e.g., RNA processing, organelle lumen). These analyses hence suggest that the SPI layer  
14 of cells differs substantially from SPE at P0.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31  
32 These findings are also supported by Ingenuity Pathway Analyses (IPA), focusing on  
33 processes involved in central nervous system (CNS) development. For genes enriched in SPE  
34 compared to WT, the broad categories predicted to be the most significantly activated (z-score  
35  $> 2$ ) included proliferation of cells, movement of brain cells and development of CNS. For  
36 genes enriched in SPI compared to WT, more than 15 categories were identified as being  
37 significantly activated. The most significant cellular processes included formation of vesicles,  
38 formation of neurites, tubulation of cells, cell movement, long-term potentiation of synapse and  
39 formation of axons (**Supp. Table 6**).  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

50 We next compared SPI and SPE to each other, and identified 2646 transcripts that were  
51 significantly differentially expressed between these two *Dcx* KO cell populations (**Fig. 2A**;  
52 **Supp. Table 4A**). Of these, 97 transcripts showed more than 1.5 X fold change difference ( $P <$   
53  $0.005$ ). The DAVID functional annotation clustering to compare enriched genes in SPI versus  
54 SPE confirmed previous findings of comparing either to WT. Notably, genes preferentially  
55  
56  
57  
58  
59  
60

1  
2  
3 enriched in SPI compared to SPE clustered in processes related to synapses, vesicular transport,  
4 cell adhesion, glycoproteins and cytoskeletal organization (extended list shown in **Supp. Table**  
5 **5F, Supp. Fig. 5A**). SPE gene expression compared to SPI on the other hand, showed clustering  
6 enrichment related to organelle lumen, RNA processing, transcription and chromatin  
7 modification (**Supp. Table 5G, Supp. Fig. 5B**). These clustering analyses are mostly exclusive,  
8 and point to a difference in the maturity status between the two *Dcx* KO CA3 layers. Indeed, a  
9 previous transcriptome study of several mouse lines showed that genes related to transcription  
10 and translation are highly expressed during prenatal development, being turned down  
11 progressively at P0 and P14, whereas expression of genes related to synaptogenesis showed the  
12 opposite regulatory pattern (48). Therefore, based on functional annotation clustering of  
13 differentially regulated genes, SPI cells appear to be more mature than SPE at P0.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

### 32 **Neuronal development genes are differentially expressed between SPI and SPE cells**

33 *Dcx* and many other migration genes continue to be expressed postnatally; therefore, we  
34 compared genes coding for *Dcx* partners and other genes known to be involved in migration  
35 and/or differentiation. Genes potentially related to *Dcx*'s function that showed expression  
36 differences were regulatory subunits of *protein phosphatases* 1 and 2 (either up or down, a  
37 larger number deregulated in SPE) and the close homolog of *Dcx*, *Dclk1* (down in SPE) (**Table**  
38 **3**) (23, 49). *Dclk2*, on the other hand showed no changes. Also down-regulated in SPI compared  
39 to WT, were *Cdk5* and *Mark2*, both kinases that phosphorylate *Dcx* (50, 51), and AP1 (*Ap1m1*)  
40 and AP2 (*Ap2m1*) mu subunit 1 genes of adaptor complexes, which interact with *Dcx* for vesicle  
41 trafficking (52). *Kif1a*, a kinesin shown to be involved with *Dcx* in trafficking cargoes (53) and  
42 *Nfasc*, coding for a cell adhesion molecule of the L1-CAM family interacting with *Dcx*, were  
43 upregulated in SPI (54). If SPI cells are more mature, these gene differences may potentially  
44 distinguish migration from differentiation roles of these partners.  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Several genes known to be involved in development of the cortex and hippocampus,  
4 including *Nde1*, *Dab1* and *Kif2a* (55-57) were up-regulated in either SPI (*Nde1*, *Kif2a*), or both  
5 SPI and SPE when compared to WT (**Table 3**). Three markers for interneurons (*Gad1*, *Sst*,  
6 *Dlx1*) were shown also to be upregulated in both SPI and SPE, which may reflect some  
7 interneuron disorganization in the KO (30).  
8  
9

10  
11  
12  
13  
14  
15 Through IPA functional clustering (e.g., **Supp. Table 6** ‘Cellular movement’  
16 categories), as well as literature searches, we assessed other genes known to be particularly  
17 important for migration (**Table 3**). Notably, SPI showed a significant down-regulation of *Srf*, a  
18 transcription factor regulated by changes in the ratio of polymerized to unpolymerized actin  
19 that is important for neuronal migration (58), and alpha-N-acetyl-neuraminide alpha-2,8-  
20 sialyltransferase 2 (*St8sia2*), a gene which is highly expressed in migrating neurons, but down-  
21 regulated during neuronal differentiation (59). Furthermore, the Rho-GTPase *Rnd2*, a  
22 modulator of radial migration in the developing cortex with a very specific spatiotemporal  
23 expression pattern, being strongly expressed in radial glial progenitors and radially migrating  
24 neurons but sharply down-regulated once neurons finish migration and start differentiation (60),  
25 was similarly down-regulated in SPI.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

41 Thus, although the two *Dcx* KO populations share a common deficit in *Dcx*, and both  
42 differ from WT, they show variable differences in expression of neuronal migration and  
43 differentiation genes. These data may be consistent with the fact that SPI cells are potentially  
44 already in the process of down-regulating a certain number of migration genes, and represent a  
45 subpopulation more advanced in maturity than SPE cells.  
46  
47  
48  
49  
50  
51

52  
53 Further supporting this, changes in genes related to synaptogenesis, ion channels and  
54 neurotransmission are upregulated in SPI (**Table 4**). Some mutant forms of these genes are  
55 known genetic causes of epilepsy (e.g., *Sv2a*, *Lgi1* (61)). In addition, three genes encoding  
56 subunits of the Na<sup>+</sup>/K<sup>+</sup>-ATPase cation transporter (*Atp1a2*, *b1*, *b2*) are upregulated in SPI.  
57  
58  
59  
60

1  
2  
3 These plasma membrane channels regulate excitability by maintaining the resting potential, and  
4 are also a major consumer of the cell's total energy (62). The up-regulation of all of these genes  
5 may point to the advanced maturity of SPI cells. Finally, three categories related to blood vessel  
6 development appeared in the functional annotation clustering for upregulated genes in SPI  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

These plasma membrane channels regulate excitability by maintaining the resting potential, and are also a major consumer of the cell's total energy (62). The up-regulation of all of these genes may point to the advanced maturity of SPI cells. Finally, three categories related to blood vessel development appeared in the functional annotation clustering for upregulated genes in SPI (**Supp. Table 6**), perhaps also suggesting advanced maturity because neurons and blood vessels develop together (63).

### **A partial inversion of the *Dcx* KO hippocampal CA3 field along its radial axis**

The hippocampus is made up of morphologically and functionally distinct pyramidal cells along its radial (deep to superficial), and septal to temporal axes. Molecular differences along these axes may contribute to the acquisition of specific phenotypes (3). We next analysed previously identified layer-specific markers along the radial axis in the *Dcx* KO, since neurons in radial layers are likely to have different maturities at P0, depending on their birthdates. Thompson et al (2008) described 45 molecular markers that specifically label CA3 outer boundary (deep, closest to the *stratum oriens*) neurons in the adult brain. We searched for these molecular markers in our P0 dataset and notably, found 13 genes to be significantly up-regulated in the SPI gene list (**Table 5A**). This suggests that some cells, which would have normally been closest to the *stratum oriens*, are positioned in much more superficial regions closest to the *stratum radiatum* of the *Dcx* KO hippocampus. Significantly, 7 of the 14 genes (*Sema5a*, *Fxyd5*, *Epha7*, *Kcnf1*, *Slc30a3*, *Fxyd7* and *Cdh13*) code for membrane proteins, which may be related to a role forming a boundary at the edge of the pyramidal cell layer. Two further genes down-regulated in outer-boundary cells (*Efnb3*, *Cadps2*) were also lower in expression in SPI than SPE.

Early-born subpopulation markers were likewise identified by Deguchi et al (2011), analysing adult hippocampal cell populations from *Lsi1* and *Lsi2* reporter mice. Despite the age

1  
2  
3 and preparation differences of the starting samples, once again, we identified some genes in  
4  
5 common in our SPI gene lists (**Table 5B**, 6 genes similarly upregulated in SPI vs. WT, of 53;  
6  
7 6 genes similarly down-regulated of 77). Linking the datasets, *Suppression of tumorigenicity*  
8  
9 *18 (St18)* and *stAR-Related Lipid Transfer (START) Domain Containing 13 (Stard13)* are  
10  
11 commonly enriched in SPI, outer boundary (3) and early-born (5) datasets. These comparisons  
12  
13 together indicate that many SPI cells may be misplaced, early-born, outer-boundary cells.  
14  
15

16  
17 The SPI-enriched expression of three of these genes, *St18*, *Cadherin 13 (Cdh13)* and *N-*  
18  
19 *terminal EF-hand calcium binding protein 2 (Necab2)*, was confirmed by qPCR and/or *in situ*  
20  
21 hybridization experiments at P0 (**Fig. 4A,B, Supp. Table 7**). Furthermore, in the adult, *St18*,  
22  
23 *Necab2* and *collagen type VI alpha 1 (Col6a1)*, the latter being an adult CA2/CA3 outer  
24  
25 boundary marker in WT (3, 64), were each found enriched in SPI cells, especially in rostro-  
26  
27 dorsal hippocampal regions (**Fig. 4C**). Outer boundary SPE cells were also labelled in more  
28  
29 temporal regions.  
30  
31

32  
33 To better identify cells, we further checked by qPCR other genes that were differentially  
34  
35 regulated in SPI and/or SPE versus WT transcriptome data (**Supp. Table 3**). Notably, whereas  
36  
37 some genes are specifically upregulated in association with SPI (*Dbi*, *Slc6a7*, *Camk2a*, *Necab2*,  
38  
39 *Npy*, *Atp1b1*), *Adreno-receptor alpha 2A (Adra2a)* and *Gastrin-releasing peptide (Grp)*, were  
40  
41 enriched specifically in SPE (**Fig. 4A**). *In situ* hybridisations of *Grp*, a CA3 gene involved in  
42  
43 cell migration, showed homogenous labelling in the WT CA3 region at P0, as well as the  
44  
45 intermediate migration zone (**Fig. 4B**) (65, 66). In the *Dcx* KO, SPE cells are strongly labelled,  
46  
47 with little expression seen in the SPI layer. We also confirmed that *Cholecystinin B receptor*  
48  
49 (*Cckbr*) and *Cadherin 13 (Cdh13)* were similarly upregulated in SPI and downregulated in SPE  
50  
51 (**Fig. 4A**), which may point to *Cckbr* having an outer-boundary profile.  
52  
53

54  
55 The SPI findings in particular strongly suggest that some cells destined for the outer  
56  
57 boundary layer of the CA3 pyramidal cell layer in the *stratum oriens* region, potentially born  
58  
59  
60



1  
2  
3 at early stages of CA3 neurogenesis (2), are abnormally, superficially positioned in the *Dcx* KO  
4 SPI with respect to SPE. SPE on the other hand, may resemble more generally the WT  
5 pyramidal cell layer organisation, containing CA3 cells born at later time-points.  
6  
7  
8  
9

10 The similar neurodevelopmental outcomes of other hippocampal heterotopia mouse  
11 models (*Lis1*, *Tuba1a*) suggest that the molecular mechanisms mediating these phenotypes may  
12 overlap, at least in the CA3 field. To further investigate this possibility, we performed *in situ*  
13 hybridizations in the *Tuba1a* (+/S140G) mutant mouse at P0. *St18* and *Necab2*, which were  
14 found predominantly in the *Dcx* KO SPI layer (**Fig 4**), also primarily label inner-layer cells in  
15 the CA3 region in the *Tuba1a* mutant compared to a control littermate (**Supp Fig. 6**).  
16 Furthermore, *Grp* expression is similar between *Tuba1a* and *Dcx* mutants, with relatively  
17 homogenous labeling across the major CA3 layer and the migrating cell layer, however with  
18 decreased labeling in the inner-most cells in the mutant. This is intriguing because similar  
19 positioning of early-born CA3 cells across several mouse mutant lines might indicate that these  
20 neurons migrate normally and are less susceptible to cytoskeletal-related gene mutations,  
21 perhaps indicating that migration modes/mechanisms are different between early- and late-born  
22 neurons in this region.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

#### 43 **BrdU birthdating experiments to label early- and late-born cells**

44 Perturbation of layer-specific marker data discussed above suggested that the inside-out  
45 layering of the CA3 region of the hippocampus is abnormal in *Dcx* KO mice. To study cell  
46 birthdates in more detail, bromodeoxyuridine (BrdU) was injected in *Dcx* (+/-) pregnant dams  
47 at either E12.5 or E16.5 to label early-born and late-born neurons, respectively, and brains from  
48 the exposed embryos were analysed at P21-24 to determine the final destinations of CA3 cells.  
49 In WTs, injection of BrdU at E12.5 labelled many neurons (NeuN+) along the border of the  
50 pyramidal cell layer and *stratum oriens*, as expected (2, 5), with 67 +/- 1.3 % of total BrdU-  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 labelled neurons found in the outer half of the CA3 *stratum pyramidale* and the other 33 +/- 1.3  
4  
5 % in the inner half ( $P < 0.001$ , paired  $t$ -tests,  $n = 4$  mice; **Fig. 5**). When looking at each of the  
6  
7 CA3 subfields separately, CA3a and b showed a greater adherence to inside-out labelling  
8  
9 compared to CA3c, in which there was no significant difference between the percentages of  
10  
11 BrdU+ neurons located in the inner or outer halves. By contrast, 56 +/- 1.7% of BrdU-labelled  
12  
13 neurons were found in the SPI of *Dcx* KO mice, with the remaining 46% in SPE ( $P < 0.05$ ,  
14  
15 paired  $t$ -test,  $n = 4$  mice; **Fig. 5**). Comparing the CA3 subfields in KO and WT, the majority of  
16  
17 the lamination defect appears to be in CA3a and b. Overall, the proportion of BrdU+ neurons  
18  
19 labelled at E12.5 in the inner portion of CA3 is greater in KO vs. WT mice ( $P < 0.001$ , Chi-  
20  
21 squared test).  
22  
23  
24  
25

26  
27 Injection of BrdU at E16.5 is expected to label late-born neurons that settle superficially  
28  
29 in the CA3 close to the *stratum radiatum* (2, 5). In the WT CA3, 72 +/- 1.6% of BrdU+ cells  
30  
31 were detected in the inner portion of the pyramidal cell layer ( $P < 0.001$ , paired  $t$ -test,  $n = 4$   
32  
33 mice; **Fig. 5**). Once again, the CA3a and CA3b subfields showed greater adherence to inside-  
34  
35 out labelling, with CA3c actually having more BrdU+ neurons in the inner portion of the  
36  
37 pyramidal layer ( $P < 0.05$ , paired  $t$ -test,  $n = 4$  mice; **Fig. 5**). In the *Dcx* KO CA3 region, BrdU+  
38  
39 neurons were mainly located in the outer (SPE) layer (82 +/- 1.4%,  $P < 0.001$ , paired  $t$ -test,  $n =$   
40  
41 5 mice; **Fig. 5**). There were significantly more BrdU+ neurons in SPE in each CA3 subregion,  
42  
43 even in CA3c. Relative to SPE itself, many of these late-born BrdU+ neurons appeared to be  
44  
45 correctly placed in the internal portion of CA3a and b. These data hence confirm a partial  
46  
47 inversion of birthdate-dependent, inside-out layering, predicted from the transcriptome data,  
48  
49 affecting early-born, outer boundary cells, found in an abnormal superficial position in the *Dcx*  
50  
51 KO hippocampus.  
52  
53  
54  
55  
56  
57  
58  
59  
60

## DISCUSSION

1  
2  
3 In the present study, we queried the genetic content and origin of abnormally positioned  
4 hippocampal neurons in a mouse model of type 1 lissencephaly associated with epilepsy. To  
5 our knowledge, no other studies have attempted to assess global gene expression in heterotopic  
6 neurons. We questioned how the dual CA3 pyramidal cell layers in the *Dcx* KO differ from  
7 each other and from WT, using LCM to separate *Dcx* KO layers at P0. Comparing WT to KO  
8 layers revealed evidence of organelle, cell stress and metabolic gene differences. SPI and SPE  
9 layer-specific gene changes were also identified, suggesting a difference in maturity between  
10 the two layers at P0. Further comparisons with previously published datasets, as well as qPCR,  
11 *in situ* hybridisation and BrdU labelling experiments, confirmed that many superficially-  
12 positioned SPI cells have the identity of normally deeply-positioned, early-born neurons. Later-  
13 born SPE cells had a more similar gene content and arrangement as WT cells. Thus, perturbed  
14 birthdate-dependent organization in addition to separated localisations of SPI and SPE likely  
15 lead to morphological and connectivity abnormalities (27).  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 *Common transcriptome changes in SPI and SPE indicate abnormal organelles, increased cell*  
37 *stress and metabolic abnormalities*  
38  
39  
40

41 The transcriptome experiment performed with laser microdissected material revealed  
42 on average 45 % of the transcripts on the microarray showing a detectable level of expression,  
43 falling into an appropriate range of the number of expressed genes expected per tissue, as shown  
44 by previous RNAseq studies (67). Both KO layers showed appreciable numbers of differentially  
45 expressed genes compared to WT, and these were for the majority layer-specific. This study of  
46 isolated pyramidal cell populations in the hippocampus at P0 may reveal both cell-autonomous  
47 and non-cell autonomous gene expression changes. A lack of *Dcx* has been shown previously  
48 to result in abnormal microtubule bundling (68), as well as abnormal intracellular transport  
49 (53), both intrinsic changes. However, exchanges with the environment (potentially also  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 aberrant due to abnormal cell position) may be different from WT, resulting in either abnormal  
4  
5 cellular responses, and / or abnormal effects on surrounding cells.  
6  
7

8 Of the commonly deregulated genes in SPI and SPE compared to WT, many were  
9  
10 related to cellular response to stress, DNA damage repair pathways and organelles. DNA  
11  
12 damage repair pathways, expressed highly during embryogenesis, were previously shown to be  
13  
14 downregulated postnatally in several mouse lines (48). Two of the most significantly enriched  
15  
16 functional categories in both KO layers were mitochondrion and Golgi apparatus (**Fig. 2, Supp.**  
17  
18 **Table 5A**), corroborating our previous electron microscopy findings that showed ultrastructural  
19  
20 abnormalities in these organelles in both SPI and SPE (30). Perhaps as a result of organelle  
21  
22 dysfunction and/or increased cell stress, apoptotic cell death was also found to be increased in  
23  
24 the hippocampus of *Dcx* KO mice at P2 (30).  
25  
26  
27  
28

29 A direct link between *Dcx*, certain deregulated genes and organelle abnormalities seems  
30  
31 plausible given the role of *Dcx* in microtubular functions (68) endocytosis (69), vesicular  
32  
33 transport (49, 52, 53) and its interaction with molecular motors (49, 53). The misplacement or  
34  
35 impaired transport of particular organelles, and/or intracellular transport in general, due to  
36  
37 abnormal microtubules (e.g. (68)) may affect metabolism, leading to cell stress and abnormal  
38  
39 organelle morphologies. Furthermore, the function and subcellular localisation of mitochondria  
40  
41 (70) and other organelles (71) play important roles in the formation of dendrites, which are  
42  
43 abnormal in the *Dcx* KO (27). Converging data in the literature and the deregulated genes we  
44  
45 identify hence point towards this being a potential mechanism leading to abnormal cell function.  
46  
47  
48  
49

50 Linking metabolism and neuronal migration, oxidative phosphorylation was recently  
51  
52 shown to be necessary for tangential migration of interneurons (72). Although CA3 neurons  
53  
54 migrate radially, they travel further and over a longer period of time relative to CA1 (73), which  
55  
56 could also make them more vulnerable to mitochondrial or metabolic abnormalities.  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Several metabolic genes are similarly deregulated in both KO layers compared to WT, which could also increase cell stress (**Table 2**). Increasing glycolysis in neurons (via genetic mechanisms) has been shown to increase cell stress (74). However, we cannot be certain that metabolic changes, particularly those related to glycolysis, are restricted to pyramidal neurons in our samples. We also found upregulation of genetic markers of interneurons (*Gad1*, *Sst*, *Dlx1*), oligodendrocyte precursor cells (OPCs) (e.g., *Olig1*) and astrocytes (e.g., *Glast*) in both SPI and SPE compared to WT. This may reflect disorganization of other cell types, and potentially non-cell-autonomous effects of *Dcx* mutation. For example, OPCs that do not express *Dcx* may migrate between the two pyramidal cell layers, being attracted to white matter that does not typically form there (75). Interneuron disorganization also seems apparent in the *Dcx* KO hippocampus, but this may be due to mislocalization of the pyramidal cells, invading spaces normally occupied by interneurons (*stratum radiatum* and *stratum oriens*). Alternatively, interneuron disorganization could also be a cell-autonomous effect as they normally express *Dcx*, and KO or knockdown causes slowed tangential migration (22, 23).

There are also several indicators of perturbed lipid metabolism in both SPI and SPE, which be linked to affect plasma membrane and organelle integrity, regardless of the cell types affected. This is interesting in light of previous links between lipid metabolism and epilepsy, and the efficacy of ketogenic diets in treating epilepsy in both mouse models and humans (76). There is however, no major evidence that *Dcx* KO mice experience epilepsy at P0 or *in utero*. Identified defects most likely contribute to the development of the heterotopia and epilepsy in adulthood. Thus, we can link various genetic and cellular defects potentially with epilepsy, and these associations to our knowledge have never previously been made for a heterotopia condition.

*SPI contains more early-born, mature cells than SPE*

1  
2  
3 Differentially regulated genes related to migration, neuronal function and radial position  
4 within the CA3 pyramidal cell layer suggest that SPI and SPE cells have different birthdates  
5 and corresponding maturity statuses. Overall, the formation of two layers from E17.5 is most  
6 probably related to a cell-autonomous lack of *Dcx*. Certain 'migration gene' changes, although  
7 not all, are exclusive to either SPI or SPE. It is possible that the majority of early-born neurons,  
8 which colonise CA3 and eventually become SPI, migrate normally and potentially in a *Dcx*-  
9 independent fashion, whereas later-born neurons that primarily constitute SPE, do not migrate  
10 far enough. It is however, difficult to assess whether SPE is correctly positioned compared to  
11 WT, or if it indicates arrested migration in the intermediate zone. *In utero* genetic interventions  
12 may address this point in the future.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

27 It seems unlikely that early-born neurons over-migrate given that *Dcx* mutations have  
28 only been shown to decrease migration speed (77). SPI cell position close to the *stratum*  
29 *radiatum* likely exposes them to an abnormal environment as they continue to mature, which  
30 may further perturb patterns of gene expression. This could be one reason that the overlap in  
31 previously published, early-born, outer-boundary gene lists was not more complete, although  
32 the differences in age also likely contribute to this. Whether the abnormally positioned early-  
33 born cells represent a specific subset of all early-born cells (e.g. (3, 5, 6) is currently unknown.  
34  
35  
36  
37  
38  
39  
40  
41  
42

43 Later-born neurons may be vulnerable to the loss of *Dcx*; they have farther to travel as  
44 development proceeds, particularly in CA3 due to the curved shape. Furthermore, 'sojourning'  
45 during migration occurs in this hippocampal field (2), and later-born mutant neurons may not  
46 be able to reach earlier-born neurons after this pause. Radial glial cells may also have some  
47 particularity in the CA3 region (7, 78). A decrease in migration speed may hence be revealed  
48 more prominently in this region. As seen in migrating neuroblasts from the cortex (79) or in the  
49 postnatal rostral migratory stream (RMS), *Dcx* knockdown slowed migration speed, and the  
50 RMS neurons started to mature before reaching their normal location (80). The same situation,  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 presumably for cell-autonomous reasons, could produce SPE in the lengthy trajectory of CA3,  
4  
5 and these particular features of CA3 may also help explain why the entire hippocampus is not  
6  
7 affected in the *Dcx* KO as it is in other mutants (*Lis1*, *Tuba1a*). Following this logic however,  
8  
9 a complete inversion of all layers might be expected, which is not the case as revealed by the  
10  
11 BrdU and *in situ* hybridisation data. Further studies are required to understand which  
12  
13 subpopulations of early- and later-born neurons correctly or under-migrate and why. In the  
14  
15 future it will be interesting to compare embryonic expression patterns of e.g. up- or down-  
16  
17 regulated SPE genes, which may reveal promising candidates for *in utero* genetic intervention.  
18  
19  
20  
21

22 Similar disruption of hippocampal birthdate-dependent organization was also observed  
23  
24 in the *reeler* mutant, even though the extent of disorganization was more severe (81). Also, our  
25  
26 *in situ* hybridization experiments in the *Tuba1a* mutant were comparable to the *Dcx* KO,  
27  
28 demonstrating perturbed inside-out lamination in CA3. This suggests overlapping mechanisms  
29  
30 or increased vulnerability of particular neuronal populations depending on birthdate. It will be  
31  
32 interesting in the future to also compare to *Lis1* mutants that have similar hippocampal  
33  
34 phenotypes and to further focus on the differences between these models.  
35  
36  
37  
38

39 To speculate further on this issue, we refer to the study by Pramparo et al. (48), who  
40  
41 analyzed gene transcription changes in whole brains from four lissencephaly mouse models  
42  
43 (*Dcx*, *Lis1*, *Ndel1* and *14-3-3epsilon*) at three developmental time points. They observed that  
44  
45 each mouse model had unique enrichment of particular gene clusters in addition to those that  
46  
47 were commonly perturbed. We might expect similar findings if analogous hippocampal LCM  
48  
49 transcriptome studies were performed in *Lis1* and *Tuba1a* mutant mice. However, these models  
50  
51 have more severe lamination defects that are present throughout the hippocampus, including  
52  
53 two or more layers and fragmentation. *Lis1* mutants also exhibit variable neocortical defects  
54  
55 depending on the gene dosage (17, 82). Unlike *Lis1*, where gene dosage appears critical for all  
56  
57 migrating neurons, the decreased severity of the *Dcx* KO phenotype demonstrates that *Dcx* is  
58  
59  
60

1  
2  
3 not essential for most of the mouse hippocampus and the neocortex to develop normally, and  
4  
5 that the CA3 later-born (SPE) cells are perhaps the most vulnerable to *Dcx* mutation. Our gene  
6  
7 expression data in different cell populations for various cellular processes (e.g. upregulation of  
8  
9 cell adhesion in SPI versus WT, and deregulated morphology and cytoskeleton gene clusters in  
10  
11 SPI or SPE versus WT) show concordance with the Pramparo et al. study, however, with the  
12  
13 caveat that differences are also expected when comparing whole brain to microdissected  
14  
15 hippocampus. It remains to be seen which other cellular processes may be uniquely deregulated  
16  
17  
18  
19  
20 in *Dcx* KO mice compared to the other models.  
21  
22  
23

#### 24 *Dcx* KO network formation

25  
26  
27 The disorganization of pyramidal cells and/or interneurons in the *Dcx* KO may impair network  
28  
29 formation and lead to epileptogenesis. A subset of early-born GABAergic neurons function as  
30  
31 hubs, generating giant depolarizing potentials (GDPs) that synchronize neural firing across the  
32  
33 postnatal hippocampus *in vitro* (83, 84). Potential interneuron disorganization associated with  
34  
35 both SPI and SPE (as suggested by the transcriptome data) might change their function, or have  
36  
37 an effect on developing networks.  
38  
39  
40

41  
42 Pyramidal cells with the same birthdate show selective interconnectivity in the  
43  
44 trisynaptic circuit (5), with some early-born CA3a/b pyramidal neurons functioning as highly  
45  
46 interconnected hub neurons in juvenile mice *in vitro* (6). With a large number of early-born  
47  
48 neurons being misplaced into a distinct layer (SPI), it is likely that these normal developmental  
49  
50 patterns are perturbed. Another important aspect of circuit formation depends on the clonal  
51  
52 relationship of sister pyramidal cells, which share interneuron connectivity (7). We do not know  
53  
54 if neurons derived from the same progenitor cell distribute across SPI and SPE, or whether they  
55  
56 tend to group together into one of these layers, but either formation could be pathological in  
57  
58 terms of network formation. The time of onset of abnormal electrical activity (or seizures) is  
59  
60



1  
2  
3 currently unknown in this model, but given the finding that early-born neurons are abnormally  
4 positioned and considering the important role of birthdate-dependent network synchronization  
5 outlined in the studies above, it seems likely that postnatal neuronal activity will be  
6  
7  
8  
9  
10 subsequently perturbed.  
11

### 12 13 14 15 *Conclusion*

16  
17 We show here an advanced molecular analysis of the heterotopic CA3 layers in *Dcx* KO  
18 mice which reveals organelle, cell stress and metabolism abnormalities as well as abnormal  
19 positioning of SPI and SPE. It remains to be seen what repercussions the relative displacement  
20 of early-born, outer boundary cells might have on neuronal differentiation, cell-to-cell  
21 communication, connectivity and function. It remains possible that SPI cells, exhibiting a  
22 unique molecular identity, still become connected appropriately, even though they are separated  
23 from the environment of SPE cells. The mechanisms leading to misplaced early-born versus  
24 late-born neurons still remain unknown, but may be related to different migration mechanisms,  
25 especially in the complex CA3 region (85). The abnormal neuronal position for rostral-dorsal  
26 SPE cells, lacking an outer-boundary with the *stratum oriens*, could also have consequences on  
27 connectivity. Further studies are required, including investigating synaptogenesis during  
28 postnatal stages, and performing gene expression analyses of more mature *Dcx* KO cells, to  
29 help answer these questions concerning the consequences of abnormal neuronal position on  
30 neuronal and network function.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## MATERIALS AND METHODS

### *Animals*

*Dcx* KO and WT mice were maintained on the *Sv129Pas* background with more than ten generations of backcrosses (19, 22). Genotyping was performed by PCR to verify the inactivation of the *Dcx* gene in KO animals (22). All experiments were performed in accordance with institutional, national and international guidelines (EC directive 86/609) and were approved by the local ethical committees (French MESR N°: 00984\_02). The day of confirmation of vaginal plug was defined as embryonic day zero (E0.5), and the day of birth was defined as postnatal day zero (P0). We considered this latter timepoint to be suitably late to reduce the study of mixed populations of migrating and differentiating neurons, in favour of studying just differentiating neurons, and suitably early to be before epileptic seizures begin. P0 animals were anesthetized by placing on ice for 5 minutes before decapitation. For laser capture microdissection (LCM), the whole head was immediately frozen in isopentane at -35°C for 1 minute before storing at -80°C for future cryostat cutting. For *in situ* hybridization and immunohistochemistry experiments, animals were perfused, and brains were fixed using 4% paraformaldehyde (PFA) in phosphate buffered saline (PBS). Brains were post-fixed in the same solution overnight. Embryonic and early postnatal brains were cryoprotected in 30% sucrose, embedded and frozen in OCT tissue freezing medium, and cut using a cryostat (Leica) at 20 µm thickness, and spread on SuperFrost II slides.

### *Laser capture microdissection (LCM)*

Coronal brain sections (12 µm) containing the rostro-caudal hippocampus were prepared using a cryostat (Leica) maintained at -20°C and mounted on PENmembrane slides (1440–1000, PALM, Bernried, Germany) which were pre-treated by RNase ZAP (Ambion) and UV irradiated in a cell culture hood for 30 min at 254 nm. After sectioning, the slides were stored

1  
2  
3 in a -80°C freezer for use within 1 week. On the day of LCM, the slides were removed from the  
4  
5 freezer, and fixed in 70% ethanol, prepared in RNase free water for 2 min and then in 50%  
6  
7 ethanol for 5 seconds, and stained with 1% cresyl violet (Sigma) for 1 min. Subsequently, slides  
8  
9 were rinsed and dehydrated using serial dilutions of ethanol (50% for 5 sec, 75% for 5 sec,  
10  
11 100% for 30 sec). Sections were air-dried and subjected to LCM within the next 30 minutes.  
12  
13 Samples were cut using a Zeiss LCM system (PALM Microbeam). In WT animals, the  
14  
15 hippocampal CA3 region was microdissected in one piece. In *Dcx* KO animals, internal *stratum*  
16  
17 *pyramidale* (SPI) and external *stratum pyramidale* (SPE) CA3 cells in the two layers were  
18  
19 separated. Per brain, the entire CA3 structure (bilateral) from the dorsal hippocampus was  
20  
21 excised; for each condition 60 sections per animal (n=5) were pooled.  
22  
23  
24  
25  
26  
27

#### 28 *RNA isolation, quantity and quality assessment*

29  
30 Total RNA was isolated from pooled microdissected material using an RNA isolation kit  
31  
32 (Arcturus picopure) according to manufacturer instructions. RNA for each brain was eluted in  
33  
34 13 µl of the elution buffer provided in the kit. RNA quantity was measured using a nanodrop  
35  
36 spectrophotometer. RNA quality was checked using an Agilent 2100 Bioanalyzer with the RNA  
37  
38 6000 Pico LabChip Kit (5065-4473, Agilent Technologies, Palo Alto) according to the  
39  
40 manufacturer's instructions (**Supp Fig 2**). No statistical differences in RNA qualities ('RIN'  
41  
42 values) were seen between the samples.  
43  
44  
45  
46  
47

#### 48 *Linear RNA amplification and GeneChip hybridization*

49  
50 15 samples, corresponding to n=5 samples for each condition were processed using the Illumina  
51  
52 TotalPrep RNA Amplification Kit (Ambion, Life Technologies) and the Whole-genome Gene  
53  
54 Expression Direct Hybridization Assay (Illumina) according to manufacturers' instructions.  
55  
56 Briefly, 150-200 ng total RNA was used to prepare double-stranded cDNA using a T7 oligo  
57  
58 (dT) primer (Illumina protocol). Reverse transcription was followed by *in vitro* transcription in  
59  
60

1  
2  
3 the presence of biotinylated nucleotides. cRNA samples were hybridized to the Illumina  
4  
5 Mouse-Ref-8 expression beadchip arrays in the appropriate buffer overnight at 58 °C. After  
6  
7 hybridization and washes, fluorescent tagging was achieved by incubation with streptavidin-  
8  
9 Cy3. Each array contains approximately 25,600 well-annotated RefSeq transcripts, represented  
10  
11 as oligonucleotides attached to beads (average of 30 beads/per transcript), corresponding to  
12  
13 over 19,100 unique genes.  
14  
15

### 16 17 18 *Data analysis*

#### 19 20 21 *Intra-sample normalisation*

22  
23 In order to compare gene expression intensities between samples, expression values within  
24  
25 samples were first normalised to account e.g. for differences in mRNA library preparation. To  
26  
27 this end, each log-transformed expression value was corrected by subtracting the median of the  
28  
29 distribution of log-transformed expression values and dividing by the standard deviation of this  
30  
31 distribution of log-transformed expression values and dividing by the standard deviation of this  
32  
33 distribution (**Supp Figure 3A**).  
34

#### 35 36 37 *Inter-sample normalisation*

38  
39 Samples were distributed on two different slides, requiring a correction for potential  
40  
41 experimental or technical differences between slides. A principal component analysis of the  
42  
43 expression values of the 15 samples (**Supp Figure 3B, C**) showed that the principal eigenvector  
44  
45 ( $Z[1]$ ) can separate the samples according to their slide of origin. To normalize, we averaged  
46  
47 the expression values of each gene across the four SPI and the four SPE samples of slide 1  
48  
49 respectively. We subtracted this average value from the value of the single SPI and SPE samples  
50  
51 on slide 2 (respectively  $\Delta_I$  and  $\Delta_E$  in **Supp Figure 3D**). When data points are close to zero on  
52  
53 both axes, this indicates that slides 1 and 2 show the same expression values. While this appears  
54  
55 to be true for many genes, we also observed a distribution of points along a slope. For genes  
56  
57 lying on this diagonal, this indicates a bias of the same magnitude for both SPI and SPE samples.  
58  
59  
60

We first performed a gene-wise correction of the SPI samples on slide 1 to be compatible with corresponding samples on slide 2. The corrected value for each gene  $j$  is obtained as the sum of the original value and an estimate of the bias error. The estimate of the bias is based on the bias of SPE (since we assume this to be independent of the sample, i.e. affecting intensities in the same way on the two slides) weighted with a constant  $w_I$ .

The correction for SPI samples is:

$$I_j^{corr,slide_1} = I_j^{slide_1} + w_I \cdot \Delta_{E,E,j}$$

$$\text{with: } \Delta_{E,j} = \text{average}(E_j^{slide_1}) - E_j^{slide_2}$$

where  $I$  and  $E$  represent the intensity of gene  $j$  on the slide indicated by the superscript (the subscript of the sample on the slide is omitted for better readability). The constant  $w_I$  was determined by minimizing the error between the intensities of the genes on two different slides according to a least square criteria (as in the equations below).

$$\text{For SPI: } \min_{w_I} \left[ \sum_j \left( I_j^{slide_2} - \text{average}(I_j^{corr,slide_1}) \right)^2 \right]$$

$$\text{For SPE: } \min_{w_E} \left[ \sum_j \left( E_j^{slide_2} - \text{average}(E_j^{corr,slide_1}) \right)^2 \right]$$

Similarly, the SPE samples were corrected on slide 1 with the bias estimated from SPI and thus all samples were aligned to the samples as found on slide 2.

After corrections, principal component analysis showed that SPI and SPE samples from slide 2 respectively cluster with the SPI and SPE samples from slide 1.

Finally the linear model fit of the LIMMA package (86) was applied on these bias-corrected expression values in order to detect the differentially expressed genes pertaining to the three conditions, using a  $t$ -test. The Benjamini-Hochberg false discovery rate control (with  $P = 0.05$ ) was applied to correct for multiple testing. Results are presented as ratios between the conditions.

### *Pathway analysis and functional category clustering*

A global overview of the biological processes and enriched functional clusters in SPI and SPE lists was obtained using a Database for Visualization and Integrative Discovery (DAVID) (Dennis et al., 2003). The following sources were used: COG-Ontology, SP-PIR-Keywords, UP-SEQ-Feature, GOterm-BP-FAT, GOterm-CC-FAT GOterm-MF-FAT, KEGG pathways, and INTERPRO protein information resources. Analysis was performed using the default settings (medium stringency); enrichment scores  $> 1.5$  were considered significant.

For further specific questions, Ingenuity Systems Pathway Analysis (IPA) (Ingenuity Systems, Mountain View, CA; [www.ingenuity.com](http://www.ingenuity.com)) was used to define enriched functional categories and pathways specific to the developing central nervous system.

### *Quantitative PCR (qPCR) validation*

Linear amplified LCM RNAs (Ambion Life Technologies or Nugen Ovation kit) were used to prepare cDNA. Real time qPCR assays using the SYBRgreen method followed MIQE guidelines (87). Gene-specific primers were designed using Primer Express Software (PE Applied Biosystems, **Supp Table 8**), except for *Atp1b1*, which were purchased from Sigma (KSPQ12012, primer set 1). Standard curves were generated from assays made with serial dilutions of cDNA to calculate PCR efficiencies ( $90\% < \text{efficiency} < 105\%$ , with  $r^2 \geq 0.998$ ). Threshold cycles ( $C_t$ ) were transformed into quantity values using the formula  $(1 + \text{Efficiency})^{-C_t}$ . Only means of triplicates with a coefficient of variation of less than 10 % were analysed. Inter-plate variation was below 10 %. Values were normalized to the geometric mean of 3 Normalization Factors (NF) found to be the most stable through all samples using the geNorm approach. These were ATP synthase, H<sup>+</sup> transporting mitochondrial F1 complex, beta subunit (*Atp5b*), eukaryotic translation initiation factor 4A2 (*Eif4a2*) and prosaposin (*Psap*). Average

1  
2  
3 values  $\pm$  standard deviations were compared for 3-5 animals of each genotype. Ratios were  
4  
5  
6 calculated and *P* values using the Student *t*-test.  
7

#### 8 9 *mRNA in situ hybridization*

10  
11 Specific antisense RNA probes (see **Supp Table 8**) for *Col6a1*, *Necab2*, *St18* and *Grp* genes  
12  
13 were used for *in situ* hybridization analyses. Equivalent sense probes were also generated for  
14  
15 comparison. Digoxigenin (DIG) probes were synthesized with a labeling kit according to the  
16  
17 manufacturer's instructions (Roche Diagnostics). Following a protocol adapted from Bally-Cuif  
18  
19 and Wassef (1994) (88), frozen cryostat sections were rinsed 3  $\times$  5 min in PBS, postfixed in 4%  
20  
21 PFA, rinsed in PBS 2  $\times$  5 min, and treated with proteinase K (10  $\mu$ g/ml) for 10 min. Sections  
22  
23 were incubated in PBS + glycine (2 mg/ml), rinsed in PBS, and postfixed in a mixture of 4%  
24  
25 PFA, followed by a rinse in PBS. Tissue sections were hybridized at 70°C overnight with the  
26  
27 DIG-labeled probes diluted 1/100 in hybridization buffer (50% deionised formamide, 10%  
28  
29 dextran sulphate, 1 mg/ml Yeast RNA, 1x Denhardt's solution). The next day, sections were  
30  
31 sequentially washed in 2X saline sodium citrate (SSC) Tween 0.1% at 70°C then in maleate  
32  
33 buffer (Maleic acid 100 mM, NaCl 150 mM, 0.1% Tween20, pH 7.5) at room temperature. For  
34  
35 immunological detection of DIG-labeled hybrids, sections were first blocked (2% blocking  
36  
37 reagent-Roche Applied Science, cat. 1096176, 20% sheep serum in maleate buffer) and then  
38  
39 incubated overnight at 4°C in the same solution containing sheep anti-DIG-alkaline  
40  
41 phosphatase-conjugated Fab fragments (Roche Diagnostics) diluted 1/2000. The following day,  
42  
43 sections were washed 4  $\times$  15 min in maleate buffer and 30 min in NTMT buffer (100 mm NaCl,  
44  
45 100 mm Tris-HCl, pH 9.5, 50 mm MgCl<sub>2</sub>, 0.1% Tween 20). The alkaline phosphatase  
46  
47 chromogen reaction was performed in NTMT buffer containing 100 mg/ml nitroblue  
48  
49 tetrazolium (Roche Diagnostics) and 50 mg/ml 5-bromo-4-chloro-3-indolyl phosphate (Roche)  
50  
51 at room temperature for 1-7 days and stopped with PBS. Sections were mounted on glass slides,  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 dried, counter-colored using nuclear fast red and dehydrated in graded ethanol solutions, and  
4  
5 coverslipped with Vectamount (Vector Laboratories).  
6  
7

#### 8 9 *Birthdate studies*

10  
11 Pregnant mice were injected intraperitoneally (i.p) four times with 2 hour intervals at E12.5  
12  
13 with 100 mg/kg bromo deoxy uridine (BrdU), or once with 200 mg/kg BrdU at E16.5. Animals  
14  
15 were sacrificed at P21-24. To determine the number of BrdU-positive cells, every tenth section  
16  
17 of 20  $\mu$ m (200  $\mu$ m intervals) of the dorsal hippocampus from both cerebral hemispheres was  
18  
19 processed for immunohistochemistry (n = 3-5 animals per genotype per BrdU injection date).  
20  
21  
22  
23  
24  
25  
26

#### 27 *Immunohistochemistry*

28  
29 Cryostat sections at 20  $\mu$ m thickness were washed in PBS 1X, and incubated in 2 N HCl at  
30  
31 37°C for 40 min to denature DNA. Sections were thoroughly washed in PBS 1X to neutralize  
32  
33 the acid, and incubated in blocking solution (0.1 M PBS, 0.3% Triton X-100, 2% normal goat  
34  
35 serum) for two hours at room temperature. This was followed by incubation with the primary  
36  
37 antibodies; anti-BrdU (Rat, clone ICR1, ABD serotec 1:100), anti-NeuN (Mouse, clone A60  
38  
39 Millipore, 1:200); diluted in blocking solution, overnight at 4°C. Brain sections were then  
40  
41 washed three times in PBS 1X and incubated with fluorochrome-conjugated secondary  
42  
43 antibodies (1:400, Invitrogen) diluted in blocking solution containing Hoechst fluorescent  
44  
45 nuclear counterstain (1:5000). Sections were mounted using fluoromount G (Electron  
46  
47 Microscopy Devices) and cover-slipped.  
48  
49  
50  
51  
52

#### 53 *Image acquisition and quantification*

54  
55  
56  
57  
58  
59  
60



1  
2  
3 Bright-field *in situ* hybridization images were acquired with a Coolsnap CCD camera fitted to  
4  
5 a Provis Olympus Microscope using 4X, 10X and 20X objectives (magnification/numerical  
6  
7 aperture).  
8  
9

10  
11 Fluorescent BrdU and NeuN immunostained sections were acquired using SP5II Leica  
12  
13 confocal microscope using a 40X objective. BrdU-labelled cell quantification was performed  
14  
15 using Icy software (Version 1.5.4.2) and the Spot Detector module (89). BrdU-labelled cells  
16  
17 were counted using the following criteria: 1. Threshold parameters were optimised for inclusion  
18  
19 of only strongly labelled BrdU cells, and identical settings were applied for counting of all  
20  
21 images. 2. Co-immunostaining for both BrdU and NeuN was required to exclude astrocytes.  
22  
23  
24 Division of the hippocampal CA3 subfields in coronal brain sections into CA3a, CA3b and  
25  
26 CA3c was performed as described previously (90). The WT pyramidal cell layer was divided  
27  
28 in half on the radial axis, described as “inner” and “outer”, for comparison to the KO bilayers.  
29  
30  
31  
32  
33

### 34 **Acknowledgements**

35  
36 We thank Nicolas Narboux-Neme, Sylvie Thomasseau, Katia Boutourlinsky, Richard  
37  
38 Belvindrah and Gael Grannec for technical help and advice, Evelyne Souil for initial advice  
39  
40 concerning LCM, David Gentien for advice and use of the Institut Curie platform for Picochip  
41  
42 analyses and Dominique Wendum for the use of the Saint Antoine LCM platform. We thank  
43  
44 the Inserm Avenir program, the French Agence National de la Recherche (ANR- 08-MNP-013),  
45  
46 the European Union's Seventh Framework Programme for research, technological development  
47  
48 and demonstration under grant agreement no: Health-F2-602531-2013, the CNRS, the  
49  
50 Fondation Bettencourt Schueller and the Fondation Jérôme Lejeune for grant support to FF. We  
51  
52 thank University Paris 6 and An-Najah University, Palestine for support to RKN. We thank the  
53  
54 IFM imaging and animal house facilities. We thank the Ile de France region for support of  
55  
56 imaging and animal house facilities and the CNRS TAAM for mouse housing. MS was funded  
57  
58  
59  
60



1  
2  
3 by the European project DESIRE. FF and MS are associated with the BioPsy Labex project and  
4  
5 the Ecole des Neurosciences de Paris Ile-de-France network.  
6  
7  
8  
9

10 **Conflict of Interest Statement.** None declared.  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

## References

- 1 Forster, E., Zhao, S. and Frotscher, M. (2006) Laminating the hippocampus. *Nat. Rev. Neurosci.*, **7**, 259-267.
- 2 Altman, J. and Bayer, S.A. (1990) Prolonged sojourn of developing pyramidal cells in the intermediate zone of the hippocampus and their settling in the stratum pyramidale. *J. Comp. Neurol.*, **301**, 343-364.
- 3 Thompson, C.L., Pathak, S.D., Jeromin, A., Ng, L.L., MacPherson, C.R., Mortrud, M.T., Cusick, A., Riley, Z.L., Sunkin, S.M., Bernard, A. *et al.* (2008) Genomic anatomy of the hippocampus. *Neuron*, **60**, 1010-1021.
- 4 Slomianka, L., Amrein, I., Knuesel, I., Sorensen, J.C. and Wolfer, D.P. (2011) Hippocampal pyramidal cells: the reemergence of cortical lamination. *Brain Struct. Funct.*, **216**, 301-317.
- 5 Deguchi, Y., Donato, F., Galimberti, I., Cabuy, E. and Caroni, P. (2011) Temporally matched subpopulations of selectively interconnected principal neurons in the hippocampus. *Nat. Neurosci.*, **14**, 495-504.
- 6 Marissal, T., Bonifazi, P., Picardo, M.A., Nardou, R., Petit, L.F., Baude, A., Fishell, G.J., Ben-Ari, Y. and Cossart, R. (2012) Pioneer glutamatergic cells develop into a morpho-functionally distinct population in the juvenile CA3 hippocampus. *Nat. Commun.*, **3**, 1316.
- 7 Xu, H.T., Han, Z., Gao, P., He, S., Li, Z., Shi, W., Kodish, O., Shao, W., Brown, K.N., Huang, K. *et al.* (2014) Distinct lineage-dependent structural and functional organization of the hippocampus. *Cell*, **157**, 1552-1564.
- 8 Datson, N.A., Morsink, M.C., Steenbergen, P.J., Aubert, Y., Schlumbohm, C., Fuchs, E. and de Kloet, E.R. (2009) A molecular blueprint of gene expression in hippocampal subregions CA1, CA3, and DG is conserved in the brain of the common marmoset. *Hippocampus*, **19**, 739-752.
- 9 Tole, S., Christian, C. and Grove, E.A. (1997) Early specification and autonomous development of cortical fields in the mouse hippocampus. *Development*, **124**, 4959-4970.
- 10 Tole, S. and Grove, E.A. (2001) Detailed field pattern is intrinsic to the embryonic mouse hippocampus early in neurogenesis. *J. Neurosci.*, **21**, 1580-1589.
- 11 Haussler, U., Bielefeld, L., Frioriep, U.P., Wolfart, J. and Haas, C.A. (2012) Septotemporal position in the hippocampal formation determines epileptic and neurogenic activity in temporal lobe epilepsy. *Cereb. Cortex*, **22**, 26-36.
- 12 Galimberti, I., Bednarek, E., Donato, F. and Caroni, P. (2010) EphA4 signaling in juveniles establishes topographic specificity of structural plasticity in the hippocampus. *Neuron*, **65**, 627-642.
- 13 des Portes, V., Francis, F., Pinard, J.M., Desguerre, I., Moutard, M.L., Snoeck, I., Meiners, L.C., Capron, F., Cusmai, R., Ricci, S. *et al.* (1998) doublecortin is the major gene causing X-linked subcortical laminar heterotopia (SCLH). *Hum. Mol. Genet.*, **7**, 1063-1070.
- 14 Gleeson, J.G., Allen, K.M., Fox, J.W., Lamperti, E.D., Berkovic, S., Scheffer, I., Cooper, E.C., Dobyns, W.B., Minnerath, S.R., Ross, M.E. *et al.* (1998) Doublecortin, a brain-specific gene mutated in human X-linked lissencephaly and double cortex syndrome, encodes a putative signaling protein. *Cell*, **92**, 63-72.
- 15 Barkovich, A.J., Guerrini, R., Kuzniecky, R.I., Jackson, G.D. and Dobyns, W.B. (2012) A developmental and genetic classification for malformations of cortical development: update 2012. *Brain*, **135**, 1348-1369.
- 16 Bahi-Buisson, N., Souville, I., Fourniol, F.J., Toussaint, A., Moores, C.A., Houdusse, A., Lemaitre, J.Y., Poirier, K., Khalaf-Nazzal, R., Hully, M. *et al.* (2013) New insights into

- 1  
2  
3 genotype-phenotype correlations for the doublecortin-related lissencephaly spectrum. *Brain*,  
4 **136**, 223-244.
- 5  
6 17 Hirotsune, S., Fleck, M.W., Gambello, M.J., Bix, G.J., Chen, A., Clark, G.D.,  
7 Ledbetter, D.H., McBain, C.J. and Wynshaw-Boris, A. (1998) Graded reduction of Pafah1b1  
8 (Lis1) activity results in neuronal migration defects and early embryonic lethality. *Nat.*  
9 *Genet.*, **19**, 333-339.
- 10  
11 18 Corbo, J.C., Deuel, T.A., Long, J.M., LaPorte, P., Tsai, E., Wynshaw-Boris, A. and  
12 Walsh, C.A. (2002) Doublecortin is required in mice for lamination of the hippocampus but  
13 not the neocortex. *J. Neurosci.*, **22**, 7548-7557.
- 14  
15 19 Kappeler, C., Dhenain, M., Phan Dinh Tuy, F., Saillour, Y., Marty, S., Fallet-Bianco,  
16 C., Souville, I., Souil, E., Pinard, J.M., Meyer, G. *et al.* (2007) Magnetic resonance imaging  
17 and histological studies of corpus callosal and hippocampal abnormalities linked to  
18 doublecortin deficiency. *J. Comp. Neurol.*, **500**, 239-254.
- 19  
20 20 Keays, D.A., Tian, G., Poirier, K., Huang, G.J., Siebold, C., Cleak, J., Oliver, P.L.,  
21 Fray, M., Harvey, R.J., Molnar, Z. *et al.* (2007) Mutations in alpha-tubulin cause abnormal  
22 neuronal migration in mice and lissencephaly in humans. *Cell*, **128**, 45-57.
- 23  
24 21 Belvindrah, R., Nosten-Bertrand, M. and Francis, F. (2014) Neuronal migration and its  
25 disorders affecting the CA3 region. *Front. Cell. Neurosci.*, **8**, 63.
- 26  
27 22 Kappeler, C., Saillour, Y., Baudoin, J.P., Tuy, F.P., Alvarez, C., Houbron, C., Gaspar,  
28 P., Hamard, G., Chelly, J., Metin, C. *et al.* (2006) Branching and nucleokinesis defects in  
29 migrating interneurons derived from doublecortin knockout mice. *Hum. Mol. Genet.*, **15**,  
30 1387-1400.
- 31  
32 23 Koizumi, H., Tanaka, T. and Gleeson, J.G. (2006) Doublecortin-like kinase functions  
33 with doublecortin to mediate fiber tract decussation and neuronal migration. *Neuron*, **49**, 55-  
34 66.
- 35  
36 24 Francis, F., Koulakoff, A., Boucher, D., Chafey, P., Schaar, B., Vinet, M.C., Friocourt,  
37 G., McDonnell, N., Reiner, O., Kahn, A. *et al.* (1999) Doublecortin is a developmentally  
38 regulated, microtubule-associated protein expressed in migrating and differentiating neurons.  
39 *Neuron*, **23**, 247-256.
- 40  
41 25 Gleeson, J.G., Lin, P.T., Flanagan, L.A. and Walsh, C.A. (1999) Doublecortin is a  
42 microtubule-associated protein and is expressed widely by migrating neurons. *Neuron*, **23**,  
43 257-271.
- 44  
45 26 Nosten-Bertrand, M., Kappeler, C., Dinocourt, C., Denis, C., Germain, J., Phan Dinh  
46 Tuy, F., Verstraeten, S., Alvarez, C., Metin, C., Chelly, J. *et al.* (2008) Epilepsy in Dcx  
47 knockout mice associated with discrete lamination defects and enhanced excitability in the  
48 hippocampus. *PLoS One*, **3**, e2473.
- 49  
50 27 Bazelot, M., Simonnet, J., Dinocourt, C., Bruel-Jungerman, E., Miles, R., Fricker, D.  
51 and Francis, F. (2012) Cellular anatomy, physiology and epileptiform activity in the CA3  
52 region of Dcx knockout mice: a neuronal lamination defect and its consequences. *Eur. J.*  
53 *Neurosci.*, **35**, 244-256.
- 54  
55 28 Angevine, J.B., Jr. and Sidman, R.L. (1961) Autoradiographic study of cell migration  
56 during histogenesis of cerebral cortex in the mouse. *Nature*, **192**, 766-768.
- 57  
58 29 Newrzella, D., Pahlavan, P.S., Kruger, C., Boehm, C., Sorgenfrei, O., Schrock, H.,  
59 Eisenhardt, G., Bischoff, N., Vogt, G., Wafzig, O. *et al.* (2007) The functional genome of  
60 CA1 and CA3 neurons under native conditions and in response to ischemia. *BMC Genomics*,  
**8**, 370.
- 61  
62 30 Khalaf-Nazzal, R., Bruel-Jungerman, E., Rio, J.P., Bureau, J., Irinopoulou, T., Sumia,  
63 I., Roumegous, A., Martin, E., Olasso, R., Parras, C. *et al.* (2013) Organelle and cellular  
64 abnormalities associated with hippocampal heterotopia in neonatal doublecortin knockout  
65 mice. *PLoS One*, **8**, e72622.

- 1  
2  
3 31 Youle, R.J. and van der Blik, A.M. (2012) Mitochondrial fission, fusion, and stress.  
4 *Science*, **337**, 1062-1065.
- 5 32 Jiang, Z., Hu, Z., Zeng, L., Lu, W., Zhang, H., Li, T. and Xiao, H. (2011) The role of  
6 the Golgi apparatus in oxidative stress: is this organelle less significant than mitochondria?  
7 *Free Radic. Biol. Med.*, **50**, 907-917.
- 8 33 Raza, M.U., Tufan, T., Wang, Y., Hill, C. and Zhu, M.Y. (2016) DNA Damage in  
9 Major Psychiatric Diseases. *Neurotox. Res.*, in press.
- 10 34 Arya, R., Mallik, M. and Lakhota, S.C. (2007) Heat shock genes - integrating cell  
11 survival and death. *J. Biosci.*, **32**, 595-610.
- 12 35 Nathan, D.F., Vos, M.H. and Lindquist, S. (1997) In vivo functions of the  
13 *Saccharomyces cerevisiae* Hsp90 chaperone. *Proc. Natl. Acad. Sci. USA*, **94**, 12949-12956.
- 14 36 Magnoni, R., Palmfeldt, J., Hansen, J., Christensen, J.H., Corydon, T.J. and Bross, P.  
15 (2014) The Hsp60 folding machinery is crucial for manganese superoxide dismutase folding  
16 and function. *Free Radic. Res.*, **48**, 168-179.
- 17 37 Terada, K. and Mori, M. (2000) Human DnaJ homologs dj2 and dj3, and bag-1 are  
18 positive cochaperones of hsc70. *J. Biol. Chem.*, **275**, 24728-24734.
- 19 38 Stricher, F., Macri, C., Ruff, M. and Muller, S. (2013) HSPA8/HSC70 chaperone  
20 protein: structure, function, and chemical targeting. *Autophagy*, **9**, 1937-1954.
- 21 39 Paredes, F., Parra, V., Torrealba, N., Navarro-Marquez, M., Gatica, D., Bravo-Sagua,  
22 R., Troncoso, R., Pennanen, C., Quiroga, C., Chiong, M. *et al.* (2016) HERPUD1 protects  
23 against oxidative stress-induced apoptosis through downregulation of the inositol 1,4,5-  
24 trisphosphate receptor. *Free Radic. Biol. Med.*, **90**, 206-218.
- 25 40 Li, Y.S., Milner, P.G., Chauhan, A.K., Watson, M.A., Hoffman, R.M., Kodner, C.M.,  
26 Milbrandt, J. and Deuel, T.F. (1990) Cloning and expression of a developmentally regulated  
27 protein that induces mitogenic and neurite outgrowth activity. *Science*, **250**, 1690-1694.
- 28 41 Asai, H., Morita, S. and Miyata, S. (2011) Effect of pleiotrophin on glutamate-induced  
29 neurotoxicity in cultured hippocampal neurons. *Cell Biochem. Funct.*, **29**, 660-665.
- 30 42 Coyle, P., Philcox, J.C., Carey, L.C. and Rofe, A.M. (2002) Metallothionein: the  
31 multipurpose protein. *Cell. Mol. Life Sci.*, **59**, 627-647.
- 32 43 Renick, S.E., Kleven, D.T., Chan, J., Stenius, K., Milner, T.A., Pickel, V.M. and  
33 Fremeau, R.T., Jr. (1999) The mammalian brain high-affinity L-proline transporter is enriched  
34 preferentially in synaptic vesicles in a subpopulation of excitatory nerve terminals in rat  
35 forebrain. *J. Neurosci.*, **19**, 21-33.
- 36 44 Phang, J.M., Liu, W. and Zabirnyk, O. (2010) Proline metabolism and  
37 microenvironmental stress. *Annu. Rev. Nutr.*, **30**, 441-463.
- 38 45 Chen, C. and Dickman, M.B. (2005) Proline suppresses apoptosis in the fungal  
39 pathogen *Colletotrichum trifolii*. *Proc. Natl. Acad. Sci. USA*, **102**, 3459-3464.
- 40 46 Bouyakdan, K., Taib, B., Budry, L., Zhao, S., Rodaros, D., Neess, D., Mandrup, S.,  
41 Faergeman, N.J. and Alquier, T. (2015) A novel role for central ACBP/DBI as a regulator of  
42 long-chain fatty acid metabolism in astrocytes. *J. Neurochem.*, **133**, 253-265.
- 43 47 Savaskan, N.E., Brauer, A.U. and Nitsch, R. (2004) Molecular cloning and expression  
44 regulation of PRG-3, a new member of the plasticity-related gene family. *Eur. J. Neurosci.*,  
45 **19**, 212-220.
- 46 48 Pramparo, T., Libiger, O., Jain, S., Li, H., Youn, Y.H., Hirotsune, S., Schork, N.J. and  
47 Wynshaw-Boris, A. (2011) Global developmental gene expression and pathway analysis of  
48 normal brain development and mouse models of human neuronal migration defects. *PLoS*  
49 *Genet.*, **7**, e1001331.
- 50 49 Liu, J.S., Schubert, C.R., Fu, X., Fourniol, F.J., Jaiswal, J.K., Houdusse, A., Stultz,  
51 C.M., Moores, C.A. and Walsh, C.A. (2012) Molecular basis for specific regulation of  
52 neuronal kinesin-3 motors by doublecortin family proteins. *Mol. Cell*, **47**, 707-721.
- 53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 50 Tanaka, T., Serneo, F.F., Higgins, C., Gambello, M.J., Wynshaw-Boris, A. and  
4 Gleeson, J.G. (2004) Lis1 and doublecortin function with dynein to mediate coupling of the  
5 nucleus to the centrosome in neuronal migration. *J. Cell Biol.*, **165**, 709-721.
- 6  
7 51 Schaar, B.T., Kinoshita, K. and McConnell, S.K. (2004) Doublecortin microtubule  
8 affinity is regulated by a balance of kinase and phosphatase activity at the leading edge of  
9 migrating neurons. *Neuron*, **41**, 203-213.
- 10 52 Friocourt, G., Chafey, P., Billuart, P., Koulakoff, A., Vinet, M.C., Schaar, B.T.,  
11 McConnell, S.K., Francis, F. and Chelly, J. (2001) Doublecortin interacts with mu subunits of  
12 clathrin adaptor complexes in the developing nervous system. *Mol. Cell. Neurosci.*, **18**, 307-  
13 319.
- 14  
15 53 Deuel, T.A., Liu, J.S., Corbo, J.C., Yoo, S.Y., Rorke-Adams, L.B. and Walsh, C.A.  
16 (2006) Genetic interactions between doublecortin and doublecortin-like kinase in neuronal  
17 migration and axon outgrowth. *Neuron*, **49**, 41-53.
- 18 54 Kizhatil, K., Wu, Y.X., Sen, A. and Bennett, V. (2002) A new activity of doublecortin  
19 in recognition of the phospho-FIGQY tyrosine in the cytoplasmic domain of neurofascin. *J.*  
20 *Neurosci.*, **22**, 7948-7958.
- 21  
22 55 Homma, N., Takei, Y., Tanaka, Y., Nakata, T., Terada, S., Kikkawa, M., Noda, Y. and  
23 Hirokawa, N. (2003) Kinesin superfamily protein 2A (KIF2A) functions in suppression of  
24 collateral branch extension. *Cell*, **114**, 229-239.
- 25 56 Rice, D.S., Sheldon, M., D'Arcangelo, G., Nakajima, K., Goldowitz, D. and Curran, T.  
26 (1998) Disabled-1 acts downstream of Reelin in a signaling pathway that controls laminar  
27 organization in the mammalian brain. *Development*, **125**, 3719-3729.
- 28 57 Pawlisz, A.S., Mutch, C., Wynshaw-Boris, A., Chenn, A., Walsh, C.A. and Feng, Y.  
29 (2008) Lis1-Nde1-dependent neuronal fate control determines cerebral cortical size and  
30 lamination. *Hum. Mol. Genet.*, **17**, 2441-2455.
- 31  
32 58 Pinheiro, E.M., Xie, Z., Norovich, A.L., Vidaki, M., Tsai, L.H. and Gertler, F.B.  
33 (2011) Lpd depletion reveals that SRF specifies radial versus tangential migration of  
34 pyramidal neurons. *Nat. Cell Biol.*, **13**, 989-995.
- 35 59 Rieger, S., Volkmann, K. and Koster, R.W. (2008) Polysialyltransferase expression is  
36 linked to neuronal migration in the developing and adult zebrafish. *Dev. Dyn.*, **237**, 276-285.
- 37 60 Heng, J.I., Nguyen, L., Castro, D.S., Zimmer, C., Wildner, H., Armant, O.,  
38 Skowronska-Krawczyk, D., Bedogni, F., Matter, J.M., Hevner, R. *et al.* (2008) Neurogenin 2  
39 controls cortical neuron migration through regulation of Rnd2. *Nature*, **455**, 114-118.
- 40 61 Noebels, J. (2015) Pathway-driven discovery of epilepsy genes. *Nat. Neurosci.*, **18**,  
41 344-350.
- 42  
43 62 Howarth, C., Gleeson, P. and Attwell, D. (2012) Updated energy budgets for neural  
44 computation in the neocortex and cerebellum. *J. Cereb. Blood Flow Metab.*, **32**, 1222-1232.
- 45 63 Carmeliet, P. and Tessier-Lavigne, M. (2005) Common mechanisms of nerve and  
46 blood vessel wiring. *Nature*, **436**, 193-200.
- 47 64 Lein, E.S., Hawrylycz, M.J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe, A.F.,  
48 Boguski, M.S., Brockway, K.S., Byrnes, E.J. *et al.* (2007) Genome-wide atlas of gene  
49 expression in the adult mouse brain. *Nature*, **445**, 168-176.
- 50 65 Paul, P., Qiao, J., Kim, K.W., Romain, C., Lee, S., Volny, N., Mobley, B., Correa, H.  
51 and Chung, D.H. (2013) Targeting gastrin-releasing peptide suppresses neuroblastoma  
52 progression via upregulation of PTEN signaling. *PLoS One*, **8**, e72570.
- 53 66 Patel, M., Kawano, T., Suzuki, N., Hamakubo, T., Karginov, A.V. and Kozasa, T.  
54 (2014) Galpha13/PDZ-RhoGEF/RhoA signaling is essential for gastrin-releasing peptide  
55 receptor-mediated colon cancer cell migration. *Mol. Pharmacol.*, **86**, 252-262.
- 56  
57  
58  
59  
60



- 1  
2  
3 67 Ramskold, D., Wang, E.T., Burge, C.B. and Sandberg, R. (2009) An abundance of  
4 ubiquitously expressed genes revealed by tissue transcriptome sequence data. *PLoS Comput.*  
5 *Biol.*, **5**, e1000598.  
6  
7 68 Bielas, S.L., Serneo, F.F., Chechlac, M., Deerinck, T.J., Perkins, G.A., Allen, P.B.,  
8 Ellisman, M.H. and Gleeson, J.G. (2007) Spinophilin facilitates dephosphorylation of  
9 doublecortin by PP1 to mediate microtubule bundling at the axonal wrist. *Cell*, **129**, 579-591.  
10 69 Yap, C.C., Vakulenko, M., Kruczek, K., Motamedi, B., Digilio, L., Liu, J.S. and  
11 Winckler, B. (2012) Doublecortin (DCX) mediates endocytosis of neurofascin independently  
12 of microtubule binding. *J. Neurosci.*, **32**, 7439-7453.  
13 70 Kimura, T. and Murakami, F. (2014) Evidence that dendritic mitochondria negatively  
14 regulate dendritic branching in pyramidal neurons in the neocortex. *J Neurosci*, **34**, 6938-  
15 6951.  
16 71 Jan, Y.N. and Jan, L.Y. (2010) Branching out: mechanisms of dendritic arborization.  
17 *Nat. Rev. Neurosci.*, **11**, 316-328.  
18 72 Lin-Hendel, E.G., McManus, M.J., Wallace, D.C., Anderson, S.A. and Golden, J.A.  
19 (2016) Differential Mitochondrial Requirements for Radially and Non-radially Migrating  
20 Cortical Neurons: Implications for Mitochondrial Disorders. *Cell Rep.*, **15**, 229-237.  
21 73 Khalaf-Nazzal, R. and Francis, F. (2013) Hippocampal development - old and new  
22 findings. *Neuroscience*, **248**, 225-242.  
23 74 Herrero-Mendez, A., Almeida, A., Fernandez, E., Maestre, C., Moncada, S. and  
24 Bolanos, J.P. (2009) The bioenergetic and antioxidant status of neurons is controlled by  
25 continuous degradation of a key glycolytic enzyme by APC/C-Cdh1. *Nat. Cell Biol.*, **11**, 747-  
26 752.  
27 75 Rogister, B., Ben-Hur, T. and Dubois-Dalcq, M. (1999) From neural stem cells to  
28 myelinating oligodendrocytes. *Mol. Cell. Neurosci.*, **14**, 287-300.  
29 76 Ruskin, D.N. and Masino, S.A. (2012) The nervous system and metabolic  
30 dysregulation: emerging evidence converges on ketogenic diet therapy. *Front Neurosci*, **6**, 33.  
31 77 Friocourt, G., Liu, J.S., Antypa, M., Rakic, S., Walsh, C.A. and Parnavelas, J.G.  
32 (2007) Both doublecortin and doublecortin-like kinase play a role in cortical interneuron  
33 migration. *J. Neurosci.*, **27**, 3875-3883.  
34 78 Barry, G., Piper, M., Lindwall, C., Moldrich, R., Mason, S., Little, E., Sarkar, A.,  
35 Tole, S., Gronostajski, R.M. and Richards, L.J. (2008) Specific glial populations regulate  
36 hippocampal morphogenesis. *J. Neurosci.*, **28**, 12328-12340.  
37 79 Pramparo, T., Youn, Y.H., Yingling, J., Hirotsune, S. and Wynshaw-Boris, A. (2010)  
38 Novel embryonic neuronal migration and proliferation defects in Dcx mutant mice are  
39 exacerbated by Lis1 reduction. *J. Neurosci.*, **30**, 3002-3012.  
40 80 Belvindrah, R., Nissant, A. and Lledo, P.M. (2011) Abnormal neuronal migration  
41 changes the fate of developing neurons in the postnatal olfactory bulb. *J. Neurosci.*, **31**, 7551-  
42 7562.  
43 81 Boyle, M.P., Bernard, A., Thompson, C.L., Ng, L., Boe, A., Mortrud, M., Hawrylycz,  
44 M.J., Jones, A.R., Hevner, R.F. and Lein, E.S. (2011) Cell-type-specific consequences of  
45 Reelin deficiency in the mouse neocortex, hippocampus, and amygdala. *J. Comp. Neurol.*,  
46 **519**, 2061-2089.  
47 82 Youn, Y.H., Pramparo, T., Hirotsune, S. and Wynshaw-Boris, A. (2009) Distinct  
48 dose-dependent cortical neuronal migration and neurite extension defects in Lis1 and Ndel1  
49 mutant mice. *J. Neurosci.*, **29**, 15520-15530.  
50 83 Bonifazi, P., Goldin, M., Picardo, M.A., Jorquera, I., Cattani, A., Bianconi, G.,  
51 Represa, A., Ben-Ari, Y. and Cossart, R. (2009) GABAergic hub neurons orchestrate  
52 synchrony in developing hippocampal networks. *Science*, **326**, 1419-1424.  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 84 Picardo, M.A., Guigue, P., Bonifazi, P., Batista-Brito, R., Allene, C., Ribas, A.,  
4 Fishell, G., Baude, A. and Cossart, R. (2011) Pioneer GABA cells comprise a subpopulation  
5 of hub neurons in the developing hippocampus. *Neuron*, **71**, 695-709.  
6  
7 85 Stouffer, M.A., Golden, J.A. and Francis, F. (2016) Neuronal migration disorders:  
8 Focus on the cytoskeleton and epilepsy. *Neurobiol. Dis.*, **92**, 18-45.  
9  
10 86 Smyth, G.K., Michaud, J. and Scott, H.S. (2005) Use of within-array replicate spots  
11 for assessing differential expression in microarray experiments. *Bioinformatics*, **21**, 2067-  
12 2075.  
13  
14 87 Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M., Mueller,  
15 R., Nolan, T., Pfaffl, M.W., Shipley, G.L. *et al.* (2009) The MIQE guidelines: minimum  
16 information for publication of quantitative real-time PCR experiments. *Clin. Chem.*, **55**, 611-  
17 622.  
18  
19 88 Bally-Cuif, L. and Wassef, M. (1994) Ectopic induction and reorganization of Wnt-1  
20 expression in quail/chick chimeras. *Development*, **120**, 3379-3394.  
21  
22 89 Olivo-Marin, J.-C. (2002) Extraction of spots in biological images using multi scale  
23 products. *Pattern Recognition* **35**, 1989-1996.  
24  
25 90 Li, X.G., Somogyi, P., Ylinen, A. and Buzsaki, G. (1994) The hippocampal CA3  
26 network: an in vivo intracellular labeling study. *J. Comp. Neurol.*, **339**, 181-208.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## FIGURE LEGENDS

### Figure 1

WT versus *Dcx* KO hippocampus at late embryonic and early postnatal stages. (A, B) Cresyl violet stained coronal sections of the hippocampus at E18.5 and P0, prior to laser capture microdissection cutting. The CA1 and CA3 fields of the pyramidal cell layer are indicated on the WT section. Arrows on the *Dcx* KO section indicate the double pyramidal cell layer. So, *stratum oriens*, sr, *stratum radiatum*. (C) Laser capture microdissection from cresyl violet stained P0 brain sections. Hippocampal CA3 WT and *Dcx* KO internal (SPI) and external (SPE) layers are shown, with schematized indications of areas selected for excision. Scale bars, 300  $\mu\text{m}$  E18.5; 300  $\mu\text{m}$  (B) and 70  $\mu\text{m}$  (C) P0.

### Figure 2

Global gene expression changes in the two *Dcx* KO layers, compared to WT. (A) Increasing numbers of differentially expressed genes as *P* value cutoffs are lowered. SPE versus WT, SPI versus WT, and SPI versus SPE comparisons are shown. The y-axis indicates numbers of genes. The red color indicates up-regulated, and the green color indicates down-regulated genes. (B) Genes commonly de-regulated in SPI and SPE were subjected to DAVID functional clustering analyses. Each significant functional category (y-axis), is plotted against the value of enrichment score (x-axis) obtained using the DAVID pathway analysis software. DAVID analyses are also resumed in Supp Table 5.

### Figure 3

DAVID functional clustering of SPE and SPI layer-specific gene expression differences (A-D). Functional clustering of up-regulated and down-regulated differentially expressed genes from comparisons of SPI cells to WT (A,B), and SPE cells to WT (C,D), respectively. Each



1  
2  
3 significant functional category (y-axis), is plotted against the value of enrichment score (x-  
4 axis).  
5  
6  
7  
8  
9

#### 10 **Figure 4**

11 Validation of transcriptome data with quantitative RT-PCR (qPCR) and *in situ* hybridizations  
12 (ISH). (A) qPCR analyses for selected genes with comparison to transcriptome data. Y-axis  
13 corresponds to relative fluorescence units. Up-regulation compared to WT is indicated as  
14 positive values and down-regulation as negative ( $P < 0.05$ , Student's *t*-tests). Results are  
15 presented as the ratio of means  $\pm$  SEM ( $n = 3-5$  samples per genotype). (B,C) ISH in coronal  
16 sections showing layer-specific CA3 gene expression patterns. In the *Dcx* KO CA3 at P0 (B),  
17 *Necab2* and *St18* are primarily expressed in SPI, whereas *Grp* is primarily expressed in SPE,  
18 as well as in late-migrating neurons in the intermediate zone (IZ) in both the KO and WT. (C)  
19 ISH in coronal sections (R, rostral, C, caudal) in adults. The WT section is equivalent to the  
20 most rostral KO section. *Col6A1* and *St18* label outer-boundary cells closest to the *stratum*  
21 *oriens* in WT, whereas mainly SPI cells are labeled in the *Dcx* KO in septal regions, as well as  
22 some outer boundary cells in more temporal regions. *Necab2* labels CA2 cells and CA3 outer-  
23 boundary cells in WT, and similar to *Col6A1* and *St18*, shows a predominant expression in SPI  
24 cells in septal regions.  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47

#### 48 **Figure 5**

49 BrdU labeling of early- and late-born neurons. Representative immunohistochemistry images  
50 of dorsal hippocampi from WT (left) and *Dcx* KO males exposed to BrdU at E12.5 (A) and  
51 E16.5 (B) show BrdU (green) and NeuN (red) labeling. White arrow heads indicate co-labeled  
52 cells. Bar graphs depict the average percent ( $\pm$  SEM) of total BrdU<sup>+</sup> neurons (NeuN<sup>+</sup>) found  
53 in SPI (inner) and SPE (outer) heterotopic layers of *Dcx* KO mice, or the inner and outer halves  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 of the single WT pyramidal layer. The proportions of BrdU+ neurons located in the inner and  
4  
5 outer layers, born at either E12.5 or E16.5, were significantly different from WT in the CA3  
6  
7 region and each subregion (CA3a, b, c; Chi-squared tests,  $p < 0.05$ ,  $n = 4-5$  mice per genotype  
8  
9 per injection time). Significant differences between the percentages of total BrdU+ neurons  
10  
11 located in SPI/inner and SPE/outer are indicated (paired  $t$ -tests,  $*p < 0.05$ ,  $**p < 0.01$ ,  $***p <$   
12  
13  $0.001$ ).  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

**Table 1. Stress-response genes significantly ( $P < 0.01$ ) changed in SPI and/or SPE vs. WT. Values are ratios. Values in parentheses,  $P < 0.05$ .**

	SPI vs. WT	SPE vs. WT
<b>DNA stress response</b>		
High mobility group box 2 (Hmgb2)	1,332	1,185
Structural maintenance of chromosomes 6 (Smc6)	1,202	1,209
Transformation related protein 53 binding protein 1 (Trp53bp1)	1,195	1,320
Excision repair cross-complementing rodent repair deficiency, complementation group 5 (Ercc5)	1,157	1,202
FGF receptor activating protein 1 (Frag1)	1,155	1,191
Myeloid differentiation primary response gene 116 (Myd116)	1,133	1,074
Structural maintenance of chromosomes 3 (Smc3)	1,100	
Polynucleotide kinase 3'-phosphatase (Pnkp)	0,913	0,874
Serum/glucocorticoid regulated kinase 1 (Sgk1)	0,875	0,900
Structural maintenance of chromosomes 1A (Smc1a)	0,772	0,907
<b>Antioxidants</b>		
Glutathione S-transferase, mu 1 (Gstm1)	1,287	1,181
Glutathione S-transferase, mu 5 (Gstm5)	1,259	
Sulfiredoxin 1 homolog ( <i>S. cerevisiae</i> ) (Srxn1)	1,145	
<b>Chaperones/Heat shock proteins</b>		
Heat shock protein 90kDa alpha (cytosolic), class B member 1 (Hsp90ab1)	1,281	1,553
Heat shock protein 1 (chaperonin) (Hspd1)	1,271	1,235
DnaJ (Hsp40) homolog, subfamily A, member 2 (Dnaja2)	1,162	1,165
Heat shock protein 2 (Hspa2), transcript variant 1	1,159	
Heat shock protein 1 (Hspb1)		0,874
Homocysteine-inducible, endoplasmic reticulum stress-inducible, ubiquitin-like domain member 1 (Herpud1)	0,894	0,906
Hspb associated protein 1 (Hspbap1)	0,872	
Heat shock protein 8 (Hspa8)	0,773	
<b>Other</b>		
Pleiotrophin (Ptn)	1,941	
Metallothionein 1 (Mt1)	1,814	
Metallothionein 3 (Mt3)	1,646	(0,784)
Aldehyde dehydrogenase 1 family, member L1 (Aldh111)	1,331	1,123
Aldehyde dehydrogenase family 3, subfamily A2 (Aldh3a2)	0,891	
Aldehyde dehydrogenase 4 family, member A1 (Aldh4a1)	0,819	
Aldehyde dehydrogenase 1 family, member L2 (Aldh112)	0,774	
Solute carrier family 6 (neurotransmitter transporter, L-proline), member 7 (Slc6a7)	1,544	0,821
<b>Metabolic stress response</b>		
Aldolase C, fructose-bisphosphate (Aldoc)	2,116	
Enolase 2, gamma neuronal (Eno2)	1,281	

**Table 2. Metabolic genes significantly deregulated in KO layers (P < 0.01).** Values are ratios. Values in parentheses, P < 0.05**A. Genes up- or downregulated in SPI and SPE vs. WT**

<b>Definition</b>	<b>SPI vs. WT</b>	<b>SPE vs. WT</b>
Aldolase C, fructose-bisphosphate (Aldoc)	2,116	1,326
Solute carrier family 1 (glial high affinity glutamate transporter), member 3 (Slc1a3); Glast	1,638	(1,199)
Phosphofruktokinase, platelet (Pfkp)	1,639	1,288
Diazepam binding inhibitor (Dbi), transcript variant 2; Acap	1,509	1,163
Isocitrate dehydrogenase 3 (NAD+), gamma (Idh3g)	1,148	(1,124)
ATP synthase, H <sup>+</sup> transporting, mitochondrial F1 complex, alpha subunit, isoform 1 (Atp5a1)	1,282	1,272
Enolase 2, gamma neuronal (Eno2)	1,281	0,838
Acyl-CoA thioesterase 1 (Acot1)	1,280	1,213
Sphingomyelin phosphodiesterase, acid-like 3A (Smpd13a)	1,269	1,162
Cytochrome c oxidase subunit VIIa polypeptide 2-like (Cox7a2l)	1,254	(1,091)
Cytochrome C oxidase assembly factor 5 (Coa5)	1,210	1,160
Phosphoglycerate mutase 1 (Pgam1)	1,192	1,128
Pyruvate dehydrogenase (lipoamide) beta (Pdhb)	1,172	1,269
Sphingomyelin phosphodiesterase 1, acid lysosomal (Smpd1)	1,156	1,170
Hexokinase 1 (Hk1)	(1,108)	1,214
Sphingomyelin phosphodiesterase 4 (Smpd4)	(1,068)	1,120
Acetoacetyl-CoA synthetase (Aacs)	0,906	0,907
Phosphatidylglycerophosphate synthase 1 (Pgs1)	0,867	0,827
Mitochondrial Translocator Assembly And Maintenance Protein, Homolog (S. Cerevisiae) (Tam41)	0,836	1,125
Nitric oxide associated 1 (Noa1)	0,834	0,894
ATP citrate lyase (Acly)	0,831	0,867
Hydroxysteroid (17-beta) dehydrogenase 7 (Hsd17b7)	0,823	0,862
Transaldolase 1 (Taldo1)	0,806	0,847
Plasticity-related protein 3/phospholipid phosphatase related 1 (Plppr1)	0,796	(1,177)
Mevalonate (diphospho) decarboxylase (Mvd)	0,780	0,759

**B. Genes up- or downregulated in SPI vs. WT**

<b>Definition</b>	<b>SPI vs. WT</b>	<b>SPE vs. WT</b>
Aldehyde dehydrogenase 1 family, member L1 (Aldh1l1)	1,391	
Aldolase A, fructose-bisphosphate (Aldoa)	1,360	
Enolase 1, alpha non-neuron (Eno1)	1,275	
Solute carrier family 2 (facilitated glucose transporter), member 1 (Slc2a1); Glut1	1,190	
Malate dehydrogenase 2, NAD (mitochondrial) (Mdh2)	1,160	
Solute carrier family 2 (facilitated glucose transporter), member 3 (Slc2a3); Glut3	1,113	
6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 2 (Pfkfb2)	0,877	
6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 4 (Pfkfb4)	0,835	

**Table 3. Neurodevelopmental genes significantly deregulated in KO layers ( $P < 0.01$ ).**  
 Genes highlighted in dark gray and light gray were upregulated and downregulated, respectively, in both SPI and SPE vs. WT.

SPI vs WT				SPE vs WT			
Upregulated		Downregulated		Upregulated		Downregulated	
Gene	Ratio	Gene	Ratio	Gene	Ratio	Gene	Ratio
Ptn	1,941	Dcx	0,521	Ppp1cb	1,408	Dcx	0,577
Rasgrp1	1,825	Rnd2	0,625	Dlx1	1,372	Rasgrp1	0,771
Slc1a3	1,638	Tiam2	0,669	Capn2	1,282	Ppp1r1a	0,872
Pdgfra	1,540	Efnb1	0,781	Rasgrf1	1,271	Ppp1ca	0,885
Wnt5a	1,522	St8sia2	0,802	Neurod1	1,245	Gas6	0,888
Gad1	1,490	Nr2f2	0,809	Nckap1	1,229	Ppp2ca	0,888
Ppp1r3c	1,489	Cdk5	0,841	Ppp2r2b	1,227	Ap2a2	0,896
Reln	1,487	Ap1m1	0,851	Gad1	1,225	Robo1	0,899
Dlx1	1,482	Mark2	0,852	Ncam1*	1,224	Dcl1	0,899
Il16	1,480	Katna1	0,856	Sst	1,216	Gdnf	0,913
Mapt*	1,443	Srf	0,859	Ctnnb1*	1,207	Pou4f1	0,944
Ccnd1	1,316	Lmnb2	0,865	Lmnb1	1,196		
Ppp2r2c	1,309	Ptk2	0,874	Tpm3	1,195		
Rasgrf1	1,309	Ap2m1	0,897	Pafah1b1*	1,174		
Ctnnb1*	1,301	Pak4	0,907	Ppp2r1b	1,171		
Enpp2	1,290	Itgb1bp1	0,910	Rhob	1,170		
Robo1	1,266	Prkcz*	0,914	Tubgcp2	1,166		
Sst	1,264	Pou4f1	0,947	Dab1	1,144		
Rhob	1,251			Prkcz*	1,103		
Ncam1*	1,235						
Nckap1	1,217						
Prkaca	1,215						
Cav1	1,214						
Kif2a	1,206						
Ppp1r14c	1,205						
Cx3cl1	1,200						
Flt1	1,195						
Nde1*	1,194						
Kif1a	1,192						
Stat3*	1,189						
Ednrb	1,185						
Unc5c	1,170						
Nfasc	1,155						
Nos3	1,153						
Gas6	1,150						
Ap2a2	1,149						
Map2k4	1,143						
Dab1	1,132						
Cdkn2c	1,127						
Angpt2	1,121						
Chi3l1	1,117						
Pdgfrb	1,108						

\*Probe dependent

**Table 4. Genes related to epilepsy, excitability, and/or neurotransmission upregulated in SPI vs. WT ( $P < 0.01$ ).** Values are fold changes. Values in parentheses,  $P < 0.05$

Definition	SPI vs. WT
ATPase, Na <sup>+</sup> /K <sup>+</sup> transporting, alpha 2 polypeptide (Atp1a2)	1,657
ATPase, Na <sup>+</sup> /K <sup>+</sup> transporting, beta 1 polypeptide (Atp1b1)	1,283
ATPase, Na <sup>+</sup> /K <sup>+</sup> transporting, beta 2 polypeptide (Atp1b2)	1,350
Leucine-rich repeat LGI family, member 1 (Lgi1)	1,313
Synaptic vesicle glycoprotein 2 a (Sv2a)	1,218
Potassium large conductance calcium-activated channel, subfamily M, beta member 4 (Kcnmb4)	1,213
Potassium voltage-gated channel, shaker-related subfamily, member 1 (Kcna1)	1,157
Potassium voltage-gated channel, shaker-related subfamily, member 2 (Kcna2)	1,125

**Table 5. SPI gene expression compared to outer-boundary and early-born CA3 transcriptome studies ( $P < 0.01$ ).** Values are ratios. Genes common in A and B are underlined.

**A. Genes commonly up- or down-regulated in outer-boundary pyramidal neurons of adult CA3 and SPI (Thompson et al., 2008; Table S2)**

<b>Upregulated</b>	<b>SPI vs. WT</b>	<b>Upregulated</b>	<b>SPI vs. SPE</b>
Necab2	1,636	Necab2	1,697
<u>St18</u>	1,634	<u>St18</u>	1,641
Lypd1	1,405	Fxyd5	1,639
Fxyd5	1,367	Fxyd7	1,639
Fxyd7	1,367	Sema5a	1,471
Epha7	1,344	Lypd1	1,417
Sema5a	1,330	<u>Stard13</u>	1,226
Cdh13	1,222	Cdh13	1,211
<u>Stard13</u>	1,220	Slc30a3	1,112
Kcnip3	1,155	Klotho	1,103
Thrsp	1,143	<b>Downregulated</b>	<b>SPI vs. SPE</b>
Kcnf1	1,105	Efnb3	0,922
Klotho	1,088	Cadps2	0,924

**B. Genes upregulated in *Lsi1* and/or *Lsi2* pyramidal neurons of adult CA3 compared to SPI (Deguchi et al., 2011; Supp. Fig. 3)**

<b>Upregulated</b>	<b>SPI vs. WT</b>	<b>Upregulated</b>	<b>SPI vs. SPE</b>
<u>St18</u>	1,634	<u>St18</u>	1,641
Rfx4	1,260	Slc40a1	1,313
Slc40a1	1,242	<u>Stard13</u>	1,226
<u>Stard13</u>	1,220	Rfx4	1,120
Dleu7	1,139		
Wwtr1	1,116		
<b>Downregulated</b>		<b>Downregulated</b>	
Fzd7	0,687	Fzd7	0,633
D16Ert472e	0,756	Odz4	0,684
Odz4	0,820	D16Ert472e	0,787
Ptprd	0,911		

**Genes downregulated in *Lsi1* and/or *Lsi2* pyramidal neurons of adult CA3 compared to SPI (Deguchi et al., 2011; Supp Fig 3)**

<b>Upregulated</b>	<b>SPI vs. WT</b>	<b>Upregulated</b>	<b>SPI vs. SPE</b>
Hba-a1	1,874	Lmna	1,240
Epha5	1,226	Gabbr3	1,158
Haghl	1,193	Lrrc33	1,128
Prss35	1,153	Ube3a	1,069
Dlgap2	1,110		
Lrrc33	1,089		
<b>Downregulated</b>		<b>Downregulated</b>	
Zfp277	0,821	Mid1	0,762
Csnk1e	0,885	Use1	0,825
Ap2m1	0,897	Zfp277	0,853
Entpd5	0,906		
Creb3	0,932		
Rad50	0,939		



**ABBREVIATIONS**

SPI- *stratum pyramidale* internal

WT- wildtype

KO- knockout

SPE- *stratum pyramidale* external

RT-PCR- Reverse transcriptase PCR

qPCR- quantitative RT-PCR

ISH- *in situ* hybridisation

LCM- laser capture microdissection

RIN- RNA integrity number

DAVID- Database for Annotation, Visualization and Integrated Discovery

IPA- Ingenuity Systems Pathway Analysis

P0- postnatal day 0

BrdU- bromodeoxyuridine

MAP- microtubule associated protein

EEG- electroencephalogram

SBH- subcortical band heterotopia

Dcx- doublecortin

ID- intellectual disability

ROS- reactive oxygen species

HSP- heat shock protein

NADH (NADPH)- Nicotinamide adenine dinucleotide (phosphate)

CNS- central nervous system


GDPs- giant depolarizing potentials

GABA- gamma-aminobutyric acid

PFA- paraformaldehyde

PBS- phosphate buffered saline



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Khalaf\_Stouffer\_Fig1

111x122mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Khalaf\_Stouffer\_Fig2

173x135mm (300 x 300 DPI)


Review

Figure 3. DAVID functional clustering of SPE and SPI layer-specific gene expression differences.



Khalaf\_Stouffer\_Fig3

173x228mm (300 x 300 DPI)




Khalaf\_Stouffer\_Fig4

173x108mm (300 x 300 DPI)

Review

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Khalaf\_Stouffer\_Fig5

86x173mm (300 x 300 DPI)