Preparation of Substituted 2 H -Pyrans via a Cascade Reaction from Methyl Coumalate and Activated Methylene Nucleophiles

Liang Chang, Kristína Plevová, Serge Thorimbert, Luc Dechoux

► To cite this version:

Liang Chang, Kristína Plevová, Serge Thorimbert, Luc Dechoux. Preparation of Substituted 2 H -Pyrans via a Cascade Reaction from Methyl Coumalate and Activated Methylene Nucleophiles. Journal of Organic Chemistry, 2017, 82 (10), pp.5499-5505. 10.1021/acs.joc.7b00761 . hal-01520277

HAL Id: hal-01520277

<https://hal.sorbonne-universite.fr/hal-01520277>

Submitted on 10 May 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Preparation of Substituted 2*H*-Pyrans *via* a Cascade Reaction from Methyl Coumalate and Activated Methylene Nucleophiles

Liang Chang, Kristína Plevová, Serge Thorimbert * and Luc Dechoux *

Sorbonne Universités, UPMC Univ Paris 06, Institut Parisien de Chimie Moléculaire, UMR CNRS 8232, 4 place Jussieu, F-75005 Paris, France.

Supporting Information Placeholder

ABSTRACT: The reaction of methyl coumalate with a wide range of methylene active compounds such as keto-esters or keto-sulfones and cyclic or acyclic diketones afforded more than thirty 2,3,5,6-tetrasubstituted 2*H*-pyrans. The reaction proceeds via a cascade reaction involving a Michael addition - 6*π*-electrocyclic ring opening - Proton transfer and 6*π* Electrocyclization in which a variety of functional groups were tolerated.

The 2*H*-pyran core is widely encountered in natural products¹ and is often used as a key intermediate in total synthesis of natural products.² Since 2*H*-pyrans are in equilibrium with 1-oxatrienes by a 6*π* electrocyclic process, 1-oxatrienes generally afford an entry into the 2*H*-pyran skeleton (Figure 1). The 6*π* disrotatory process has been already discussed at different levels of theory.³ The equilibrium between 2*H*-pyrans **A** and 1-oxatrienes **B** depends on many factors such as the presence of electron-withdrawing substituents or increased steric interactions. The most common entry into the 1-oxatriene core is the Knoevenagel reaction (most generally condensation between an enal and a cyclic 1,3-dicarbonyl compound). From a synthetic point of view, the tandem Knoevenagel / 6*π*-electrocyclization (Figure 1) can be considered as a formal [3+3] cycloaddition.^{4,5} Conjugated 1-oxatrienes can also be obtained through a Claisen rearrangement of propargyl vinyl ethers⁶ or cycloisomerization of diynols.⁷ Other strategies towards 2*H*-pyrans that do not involve the “1-oxatriene pathway” but rather rely on formal [4+2]- or [3+3]-cycloaddition have been reported.^{8,9}

We present a study enabling a general synthesis of tetrasubstituted 2*H*-pyrans by addition of methylene active nucleophiles **D** to methyl coumalate (**MC**). We have recently demonstrated that Grignard reagents add regioselectively on the C6 of **MC**.¹⁰ Our hypothesis was that **MC** should react with the same regiochemistry with stabilized nucleophiles **D** like keto-esters or diketones in a tandem 1,6-Michael addition - 6*π*-electrocyclic ring opening - 1,5 H shift reaction offering an entry to 1-oxatrienes **B'**. This highly strained 1-oxatriene bearing two electron-withdrawing groups should undergo a final 6*π*-electrocyclization, deliver the 2*H*-pyran systems **A'** (Figure 1).

Figure 1. Approaches to 2*H*-Pyrans

This hypothesis was also based on a report of the literature¹¹ in which the aceto-phenylsulfone reacted with **MC** at high temperature to give the corresponding decarboxylated 2*H*-pyran in 67% yield. Only one example was described and we decided to reinvestigate the reaction conditions and to study the scope and limitations. We demonstrate here that careful control of the temperature is crucial and, in addition to the previously reported 2*H*-pyran, that 2-(2*H*-pyran-2-yl)acetic acids are also available in good yields. We also proposed a modified mechanism that takes into account some of our previous results.

We started our investigation with methyl acetoacetate (MAA) as a model nucleophile in the cascade reaction (Table 1). The first attempt was carried out with 20 mol % of cesium carbonate in DCM at room temperature for 15 h. The expected 2*H*-pyran **1a** was obtained in low yield due to low conversion (MC recovered in 75 mol %, see SI, Table S1). A stoichiometric amount of cesium carbonate gave compound **1a** with moderate yield (60%) (Table 1, entry 1).

Table 1. Optimization of the Reaction Conditions^a

Entry	Base	Solvent	Temp	Time (h)	Products (yield)
1	Cs ₂ CO ₃	DCM	RT	3	1a (60%)
2	Cs ₂ CO ₃	EtOH	RT	3	1a (80%)
3	Na ₂ CO ₃	EtOH	RT	1	1a (90%)
4	Et ₃ N	EtOH	RT	3	1a (46%) ^b
5	Et ₃ N	DCM	RT	3	1a (55%)
6	Et ₃ N	DCM	reflux	15	2a (87%)
7	Et ₃ N	EtOH	reflux	15	2a (58%)
8	Et ₃ N	THF	reflux	6	2a (54%) ^c
9	Et ₃ N	Toluene	reflux	6	2a (80%)
10	DBU	DCM	reflux	1	2a (32%) ^c
11	Et ₃ N ^d	DCM	reflux	24	2a (77%) ^e
12	Et ₃ N ^d	DCM	reflux	48	2a (86%)
13	Et ₃ N	Neat	50°C	15	2a (60%)

^a Reaction conditions: All reactions were performed with MC (1 mmol) and MAA (1 mmol), in 5 mL of organic solvent. ^b 54% of MC was observed in the crude reaction mixture, determined by ¹H NMR. ^c Degradation was observed. ^d 20 mol % of Et₃N. ^e 18% of MC was observed in the crude reaction mixture, determined by ¹H NMR.

2*H*-Pyran **1a** was obtained with better yield when the reaction was performed in EtOH with either cesium or sodium carbonate (entries 2-3). At rt NEt₃ gave moderate conversions and yields using either EtOH or DCM as solvents (entries 4-5). Interestingly, after prolonged reaction time, some traces of the decarboxylated 2*H*-pyran **2a** were observed (see SI, Table S1). In order to favour the complete decarboxylation, the reaction mixture was heated to reflux for 15h in DCM. Product **2a** was thus isolated in good yield (87%, entry 6). Changing DCM for toluene, EtOH or THF did not improve the yield of **2a** (entries 7-9) and the use of DBU instead of NEt₃ in DCM gave only modest results (entry 10). Interestingly, the reaction was successfully tested with a catalytic amount of Et₃N. Complete conversion of MC into the expected decarboxylated 2*H*-pyran **2a** was reached in 48 h with 86% yield (entry 12, see also SI, Table S1 for shorter reaction times). A solvent-free reaction led to the formation of the expected compound **2a**, however in a moderate 60% yield after 15 hours at 50°C (entry 13).

Having in hand the best conditions to synthesize either 2*H*-pyran **1a** (entry 3) or its corresponding decarboxylated analogue **2a** (entry 6), we explored the scope and limitations of these reactions.

The data in Table 2 illustrate the reaction with Na₂CO₃ in EtOH for 1h at room temperature. 1,3-Ketoesters gave the expected 2*H*-pyrans **1a-c,e,f** in yields ranging from 86% to 92%. Diverse functions such as methylether (**1c**), halogen (**1e**) or a second ester function (**1f**) gave the expected products whereas 1,3-diketones gave moderate to good yields. For example, the symmetrical 1,3-pentanedione gave the 2*H*-pyran **1g** in 87% yield whereas the non-symmetrical acetoacetophenone gave the 2*H*-pyrans **1h** and **1h'** in a 3/1 mixture, which could not be purified by flash chromatography, reflecting the two diastereoisomeric 1-oxatriene intermediates **5h** (EWG=COPh R=Me) and **5h'** (EWG=COMe and R=Ph) (see Scheme 1).

The purifications of some of 2*H*-pyrans **1** were not satisfying. We surmised that direct esterification of the carboxylic acid function following Steglich procedure¹² could solve the problem. Compounds **3d,3i,3j** were efficiently prepared in a one-pot manner with yields from 52% to 72% (Table 2).

Table 2. Substrate Scope of the Tandem Michael / 6π / 1,5H shift / 6π Reaction at Room Temperature^a

^a Reaction conditions : MC/Methylene active nucleophile/Na₂CO₃ (1/1/1) in EtOH for 1 h. ^b 5mL of DCM, 1 equiv. of EDC.HCl and 20 mol% of DMAP were directly added to the reaction mixture after 1h and stirred 15h.

We next examined the outcome of the cascade reactions in the presence of NEt₃ in DCM under reflux. Under these conditions, more than 20 examples of 2*H*-pyrans **2** were synthesized. Results are reported in Table 3. The decarboxylated 2*H*-pyrans **2a-f** were prepared from the 1,3-ketoesters in excellent yields. Again, ether, halogen or ester moieties were tolerated in this cascade reaction. Satisfyingly and in contrast to the first conditions, the crowded 2*H*-pyran **2k** and the aromatic 2*H*-pyrans **2l,m,s** were synthesized with up to 97% yield (*vide supra* for mechanistic explanations). The 4-hydroxycoumarin gave the expected tricyclic product **2n** in 82% yield. In these conditions, β-oxosulfones led to 2*H*-pyrans **2j,t** in almost quantitative yield.¹¹

Table 3. Substrates Scope of the Decarboxylative Tandem Reaction^a

^a Reaction conditions : MC/Methylene active nucleophile/Et₃N (1/1/1) in DCM reflux for 15 h.

Various 1,3-diketones could also lead to the expected 2H-Pyrans in moderate to excellent yields. Even aromatic diketones, which were unproductive at rt produced the expected pyrans **2p** albeit in moderate yields. Again, acetoacetophenone gave a 1.1/1 mixture of isomers **2h** and **2h'** in good yield (82%). This result seems to indicate that there is no stereoselectivity in the proton shift step (see *vide supra* for mechanism). Finally 2H-pyrans **2i,q,r** resulting from the addition of 5- and 6-membered cyclic diketones were isolated in good yields (84-96 %).

Our proposal for the mechanism of this cascade reaction might involve a 1,6-addition of the enolate on **MC** followed by a 6- π -electrocyclic ring opening¹³ leading to the dienolic carboxylates **4** (Scheme 1) followed by a [1,5]-H shift through a deprotonation-reprotonation sequence. We ruled out a [1,5]-H sigmatropic shift due to high steric constraint in the required planar transition state due to the *cis,cis* configuration of the dienolic system **4**.¹⁴ A further rapid 6 π electrocyclicization would lead to the 2H-pyran **1'** from oxatriene **5**. This last electrocyclic reaction seemed to be a rapid process since oxatriene intermediate **5** was never detected. This observation suggests that even during the formation of 2H-pyran **2**, the mechanism involves intermediate **1'**. This proposition is confirmed by the fact that 2H-pyran **1a**, treated with NEt₃ in DCM delivered, after 15h at reflux, the corresponding decarboxylated 2H-pyran **2a** in 95 %. The mechanistic pathway to 2H-pyrans **2** implies

that decarboxylation may operate on the transient 1-oxatriene **5** giving the stabilized enolate **6** which, after reprotonation from ammonium, gives the oxatriene **7** and regenerates NEt₃ (Table 1, entry 12). The mechanistic pathway ends with the rapid 6 π electrocyclicization delivering the stable 2H-pyran **2**.

Scheme 1. Proposed Unified Mechanism for Cascade Reaction Leading to 2H-pyrans 1 and 2

A possible rationale for the modest yields obtained with aromatic di-ketones at rt could be explained by steric factors. The planarity of the substituents of the double bonds (EWG=COAr and R=Ar) in the 6 π transition state is required for the transformation of 1-oxatriene **5** to 2H-pyran **1'**. Since the 6 π electrocyclicization was inhibited and all the steps are reversible, starting materials were recovered accompanied by complex mixture of by-products. In contrast, by heating the reaction mixture, intermediate **5** was irreversibly converted to enolate **6** which after reprotonation and 6- π electrocyclicization gave the stable 2H-pyran **2**.

In summary, we described here a rapid and efficient synthesis of more than thirty stable tetrasubstituted 2H-pyrans **1**, **2** and **3**. The reactions to **1** or **2** proceed through a cascade 1,6-Michael / 6 π electrocyclic ring opening / [1,5]-H transfer / (decarboxylation) / 6 π electrocyclicization reaction. In addition to the broad substrate scope, this methodology has the advantage of using readily available starting materials and mild conditions. The resulting 2H-pyrans bear functional groups, which should allow chemoselective transformations useful in total synthesis.² Secoiridoid monoterpenes such as secologanin or gentiopicroside are also potential targets with biological activities of interest.¹⁵

EXPERIMENTAL SECTION

General. All reactions were carried out under argon atmosphere with magnetic stirring in dry and distilled solvents. ¹H and ¹³C NMR spectra were recorded at 400 MHz for ¹H nuclei, 100 MHz for ¹³C nuclei. Chemical shifts are reported in δ units, parts per million (ppm) using, for ¹H and ¹³C, solvent residual peak as internal standard references: chloroform (7.26 ppm for

¹H NMR and 77.16 ppm for ¹³C NMR). Coupling constants (*J*) are given in Hz, multiplicities are abbreviated as: s (singlet), bs (broad singlet), d (doublet), dd (doublet of doublets), ddd (doublet of doublets of doublets), t (triplet), m (multiplet). High resolution mass spectra (HR-MS) were recorded on a LTQ-Orbitrap Mass Spectrometer [Thermo Scientific]. IR: Shimadzu IRAffinity-1CE spectrometer, wavenumbers in cm⁻¹. Reactions were monitored by thin-layer chromatography carried out on silica plates (silica gel 60 F254, Merck) using UV-light for visualization. Column chromatography was performed on silica gel 60 (0.040–0.063 mm, Merck) using the indicated eluent given in volume ratio. All starting materials were purchased at the highest commercial quality and used without further purification unless otherwise stated.

Procedure A: General procedure for the synthesis of compounds 1a-c, 1e-j. The corresponding methylene active compound (1 mmol, 1 equiv) was added at r.t. to a solution of methyl coumalate (154 mg, 1 mmol, 1 equiv) in EtOH (5 mL). After Na₂CO₃ (105 mg, 1 mmol, 1 equiv) was added to this colorless solution, it turned to orange. Thus prepared solution was stirred for 1 h at r.t. under Argon. Then the reaction mixture was diluted with H₂O (20 mL) and extracted with CH₂Cl₂ (3 × 20 mL). Then the aqueous layer, containing the product as sodium salt, was acidified with 1M HCl (5 mL, until pH = 2) and extracted with DCM (3 × 20 mL). Collected organic layers were dried over anhydrous MgSO₄, filtrated and evaporated under reduced pressure to afford the target products.

Procedure B: General procedure for the synthesis of compounds 2a-u. To a solution of methyl coumalate (154 mg, 1 mmol, 1 equiv) in CH₂Cl₂ (5 mL) at r.t. was added the corresponding methylene active compound (1 mmol, 1 equiv) followed by triethylamine (101 mg, 1 mmol, 1 equiv). The solution was stirred for 15 h under reflux, cool to r.t. then quenched with 1M HCl (10 mL). The mixture was extracted with DCM (3 × 20 mL). The organic layers were washed with brine and dried over anhydrous MgSO₄, filtrated and evaporated and the residue was subjected to column chromatography on silica yielding the respective title compounds.

Procedure C: General procedure for the synthesis of compounds 3d, 3i, 3j. The corresponding methylene active compound (1 mmol, 1 equiv) was added at r.t. to a solution of methyl coumalate (154 mg, 1 mmol, 1 equiv) in EtOH (5 mL). After Na₂CO₃ (105 mg, 1 mmol, 1 equiv) was added to this colorless solution, it turned to orange. Thus prepared solution was stirred for 1h at r.t. under Argon. Then N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (192 mg, 1 mmol, 1 equiv) and DMAP (24 mg, 0.2 mmol, 0.2 equiv) was added in CH₂Cl₂ (5mL) solution. Once reaction was completed, the reaction mixture was quenched with 1M HCl (10 mL) and mixture was extracted into CH₂Cl₂ (3 × 20 mL). Collected organic layers were washed with brine and dried over anhydrous MgSO₄, filtrated and evaporated and the residue was subjected to column chromatography on silica yielding the respective title compounds.

2-(3,5-bis(methoxycarbonyl)-6-methyl-2H-pyran-2-yl)acetic acid (1a) : The title compound was prepared according to procedure A, 243 mg; 90% yield; yellow oil; ¹H NMR (400 MHz, CDCl₃) δ 10.65 (br s, 1H), 7.56 (s, 1H), 5.73 (dd, *J* = 10.1, 3.1 Hz, 1H), 3.77 (s, 6H), 2.84 (dd, *J* = 15.2, 10.1 Hz, 1H), 2.56 (dd, *J* = 15.2, 3.1 Hz, 1H), 2.37 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 174.9, 170.6, 165.4, 164.6, 132.3, 114.8, 104.4, 72.2, 51.9, 51.6, 37.9, 20.2; IR (film, cm⁻¹): ~3600-

2600 (br), 2976, 2868, 1716, 1436, 1362, 1244, 1182, 1112, 933, 790; HRMS (ESI) Found *m/z* 293.0632 calcd for [M+Na⁺] C₁₂H₁₄NaO₇ 293.0632.

2-(6-butyl-3,5-bis(methoxycarbonyl)-2H-pyran-2-yl)acetic acid (1b): The title compound was prepared according to procedure A, 268 mg; 90% yield; yellow oil; ¹H NMR (400 MHz, CDCl₃) δ 10.43 (br s, 1H), 7.57 (s, 1H), 5.75 (dd, *J* = 10.1, 3.0 Hz, 1H), 3.77 (s, 3H), 3.77 (s, 3H), 2.98 – 2.76 (m, 2H), 2.55–2.57 (m, 2H), 1.66 – 1.52 (m, 2H), 0.95 (t, *J* = 7.4 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 175.2, 174.0, 165.3, 164.7, 132.6, 114.6, 104.1, 72.0, 51.9, 51.6, 37.7, 34.6, 20.7, 14.0; IR (film, cm⁻¹): ~3600–2600 (br), 2960, 2873, 1714, 1558, 1436, 1232, 1112, 1060, 977, 756; HRMS (ESI) Found *m/z* 321.0957 calcd for [M+Na⁺] C₁₄H₁₈NaO₇ 321.0945

2-(3,5-bis (methoxycarbonyl) -6- (methoxymethyl)-2H-pyran-2-yl) acetic acid (1c): The title compound was prepared according to procedure A, 273 mg; 91% yield; colorless oil; ¹H NMR (400 MHz, CDCl₃) δ 9.63 (br s, 1H), 7.53 (s, 1H), 5.83 (dd, *J* = 9.9, 3.2 Hz, 1H), 4.65 (d, *J* = 14.0 Hz, 1H), 4.44 (d, *J* = 14.0 Hz, 1H), 3.79 (s, 3H), 3.78 (s, 3H), 3.40 (s, 3H), 2.90 (dd, *J* = 15.6, 9.9 Hz, 1H), 2.57 (dd, *J* = 15.6, 3.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 173.9, 167.4, 164.8, 164.3, 131.5, 116.6, 105.9, 72.3, 69.3, 59.3, 52.1, 51.9, 37.6; IR (film, cm⁻¹): ~3700–2500 (br), 2953, 1710, 1641, 1570, 1242, 1112, 1064, 977, 756; HRMS (ESI) Found *m/z* 323.0748 calcd for [M+Na⁺] C₁₃H₁₆NaO₈ 323.0737.

2-(6-(chloromethyl)-3,5-bis(methoxycarbonyl)-2H-pyran-2-yl) acetic acid (1e): The title compound was prepared according to procedure A, 255 mg; 84% yield; colorless oil; ¹H NMR (400 MHz, CDCl₃) δ 9.72 (br s, 1H), 7.52 (s, 1H), 5.81 (dd, *J* = 9.9, 3.1 Hz, 1H), 4.90 (d, *J* = 11.3 Hz, 1H), 4.29 (d, *J* = 11.3 Hz, 1H), 3.81 (s, 3H), 3.78 (s, 3H), 2.93 (dd, *J* = 16.2, 9.9 Hz, 1H), 2.56 (dd, *J* = 16.2, 3.1 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 174.7, 164.2, 164.2, 164.1, 131.1, 117.9, 106.1, 72.0, 52.2 (2C), 39.8, 37.4; IR (film, cm⁻¹): ~3700–2400 (br), 2978, 1716, 1436, 1384, 1251, 1238, 1193, 1180, 1097, 1070, 1002, 933, 833, 756; HRMS (ESI) Found *m/z* 303.0266, calcd for [M-H⁻] C₁₂H₁₂ClO₇ 303.0277.

2-(6-(2-methoxy-2-oxoethyl)-3,5-bis(methoxycarbonyl)-2H-pyran-2-yl) acetic acid (1f): The title compound was prepared according to procedure A, 302 mg; 92% yield; colorless oil; ¹H NMR (400 MHz, CDCl₃) δ 9.16 (br s, 1H), 7.55 (s, 1H); 5.80 (dd, *J* = 10.1, 2.7 Hz, 1H); 4.14 (d, *J* = 16.2 Hz, 1H); 3.79 (s, 3H); 3.77 (s, 3H); 3.71 (s, 3H); 3.48 (d, *J* = 16.2 Hz, 1H); 3.07 (dd, *J* = 16.4, 10.1 Hz, 1H); 2.53 (dd, *J* = 16.4, 2.7 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 174.8; 168.7; 164.9; 164.6; 164.4; 131.4; 116.5; 106.4; 72.4; 52.6; 52.1; 52.0; 39.0; 37.4; IR (film, cm⁻¹): ~3500–2400 (br), 2960, 1714, 1697, 11435, 1357, 1247, 1093, 997, 877, 775; HRMS (ESI) Found *m/z* 351.0687, calcd for [M+Na⁺] C₁₄H₁₆NaO₉ 351.0691.

2-(5-acetyl-3-(methoxycarbonyl)-6-methyl-2H-pyran-2-yl)acetic acid (1g): The title compound was prepared according to procedure A, 228 mg; 90% yield; colorless oil; ¹H NMR (400 MHz, CDCl₃) δ 10.42 (br s, 1H), 7.46 (s, 1H), 5.74 – 5.54 (m, 1H), 3.73 (s, 3H), 2.77 (m, 1H), 2.50 (m, 1H), 2.28 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ 195.6, 174.4, 171.1, 164.5, 132.3, 114.8, 112.9, 72.3, 52.0, 37.9, 28.9, 21.0; IR (film, cm⁻¹): ~3500–2400 (br), 2976, 2870, 1712, 1624, 1436, 1382, 1238, 1105, 1070, 1018, 935, 840; HRMS (ESI) Found *m/z* 277.0784, calcd for [M+Na⁺] C₁₂H₁₄NaO₆ 277.0683.

Dimethyl-2,6-dimethyl-2H-pyran-3,5-dicarboxylate (2a): The title compound was prepared according to procedure B,

196 mg; 87% yield; colorless oil; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.47 (s, 1H), 5.34 (q, $J = 6.5$ Hz, 1H), 3.74 (s, 3H), 3.74 (s, 3H), 2.34 (s, 3H), 1.30 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 170.8, 165.8, 165.0, 130.8, 117.2, 103.5, 72.5, 51.6, 51.4, 20.5, 19.2; **IR** (film, cm^{-1}): 2951, 2918, 1716, 1689, 1637, 1433, 1379, 1309, 1226, 1128, 1045, 910, 752; **HRMS** (ESI) Found m/z 233.0996, calcd for $[\text{M}+\text{Li}^+]$ $\text{C}_{11}\text{H}_{14}\text{LiO}_5$ 233.0996.

Dimethyl-2-methyl-6-propyl-2H-pyran-3,5-dicarboxylate (2b): The title compound was prepared according to procedure B, 224 mg; 88% yield; white solid, m.p. 123–126°C; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.47 (s, 1H), 5.34 (q, $J = 6.5$ Hz, 1H), 3.73 (s, 3H), 3.73 (s, 3H), 3.04–2.84 (m, 1H), 2.51–2.42 (m, 1H), 1.67–1.50 (m, 2H), 1.27 (d, $J = 6.5$ Hz, 3H), 0.95 (t, $J = 7.4$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 174.1, 165.6, 165.0, 131.0, 117.2, 103.4, 72.2, 51.6, 51.3, 34.7, 20.6, 19.0, 13.9; **IR** (film, cm^{-1}): 2974, 2872, 1701, 1639, 1556, 1435, 1307, 1224, 1122, 1068, 914, 756; **HRMS** (ESI) Found m/z 277.1051, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{13}\text{H}_{18}\text{NaO}_5$ 277.1046.

Dimethyl-6-(methoxymethyl)-2-methyl-2H-pyran-3,5-dicarboxylate (2c): The title compound was prepared according to procedure B, 208 mg, 81% yield, white solid, m.p. 140–141°C, $R_f = 0.3$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.44 (s, 1H), 5.46 (q, $J = 6.5$ Hz, 1H), 4.67 (d, $J = 14.3$ Hz, 1H), 4.40 (d, $J = 14.3$ Hz, 1H), 3.76 (s, 3H), 3.75 (s, 3H), 3.39 (s, 3H), 1.35 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 168.0, 165.1, 164.7, 130.1, 118.8, 104.9, 72.8, 69.6, 59.1, 51.8, 51.6, 19.0; **IR** (film, cm^{-1}): 2951, 2931, 1699, 1639, 1566, 1435, 1309, 1226, 1124, 1076, 991, 912, 752; **HRMS** (ESI) Found m/z 279.0835, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{12}\text{H}_{16}\text{NaO}_6$ 279.0839.

Dimethyl-6-cyclopropyl-2-methyl-2H-pyran-3,5-dicarboxylate (2d): The title compound was prepared according to procedure B, 237 mg; 94% yield; white solid, m.p. 101–103°C; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.52 (s, 1H), 5.27 (q, $J = 6.5$ Hz, 1H), 3.77 (s, 3H), 3.73 (s, 3H), 3.17–3.10 (m, 1H), 1.35–1.24 (m, 1H), 1.21 (d, $J = 6.5$ Hz, 3H), 1.09–0.97 (m, 1H), 0.97–0.88 (m, 1H), 0.88–0.74 (m, 1H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 173.8, 166.2, 165.1, 131.4, 116.3, 103.5, 72.0, 51.5, 51.4, 18.6, 13.1, 10.1, 7.1; **IR** (film, cm^{-1}): 2951, 2929, 1687, 1635, 1539, 1301, 1222, 1062, 1043, 931, 875, 815; **HRMS** (ESI) Found m/z 275.0885, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{13}\text{H}_{16}\text{NaO}_5$ 275.0890.

Dimethyl-6-(chloromethyl)-2-methyl-2H-pyran-3,5-dicarboxylate (2e): The title compound was prepared according to procedure B, 179 mg; 71% yield; colorless oil; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.45 (d, $J = 0.4$ Hz, 1H), 5.43 (q, $J = 6.5$ Hz, 1H), 4.97 (d, $J = 11.2$ Hz, 1H), 4.24 (d, $J = 11.2$ Hz, 1H), 3.81 (s, 3H), 3.78 (s, 3H), 1.36 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 164.6, 164.1, 129.6, 120.4, 105.5, 72.6, 52.0, 51.9, 40.1, 18.9; **IR** (film, cm^{-1}): 2953, 2926, 1703; 1643, 1573, 1435, 1307, 1228, 1124, 1076, 939, 796, 754; **HRMS** (ESI) Found m/z 267.0610, calcd for $[\text{M}+\text{Li}^+]$ $\text{C}_{11}\text{H}_{13}\text{ClLiO}_5$ 267.0606.

Dimethyl-6-(2-methoxy-2-oxoethyl)-2-methyl-2H-pyran-3,5-dicarboxylate (2f): The title compound was prepared according to procedure B, 233 mg, 82% yield, colorless oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.41 (s, 1H), 5.35 (q, $J = 6.5$ Hz, 1H), 4.05 (d, $J = 16.2$ Hz, 1H), 3.71 (s, 3H), 3.71 (s, 3H), 3.67 (s, 3H), 3.44 (d, $J = 16.2$ Hz, 1H), 1.32 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz,

CDCl_3) δ 168.7, 165.2, 164.8, 164.7, 129.8, 118.8, 105.4, 73.0, 52.2, 51.7, 51.6, 39.0, 19.1; **IR** (film, cm^{-1}): 3001, 2954, 1739, 1712, 1689, 1641, 1573, 1463, 1332, 1122, 1047, 948, 871; **HRMS** (ESI) Found m/z 307.0784, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{13}\text{H}_{16}\text{NaO}_7$ 307.0788.

Methyl-5-acetyl-2,6-dimethyl-2H-pyran-3-carboxylate (2g): The title compound was prepared according to procedure B, 199 mg; 95% yield, colorless oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.40 (s, 1H), 5.32 (q, $J = 6.5$ Hz, 1H), 3.73 (s, 3H), 2.29 (s, 3H), 2.28 (s, 3H), 1.27 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 195.2, 171.1, 164.9, 131.0, 117.1, 112.3, 72.5, 51.6, 28.9, 21.3, 19.2; **IR** (film, cm^{-1}): 2976, 1703, 1633, 1558, 1436, 1274, 1228, 1147, 1122, 954, 756; **HRMS** (ESI) Found m/z 233.0783, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{11}\text{H}_{14}\text{NaO}_4$ 233.0784.

Methyl-5-acetyl-2-methyl-6-phenyl-2H-pyran-3-carboxylate (2h): The title compound was prepared according to procedure B, 117 mg, 43% yield, yellow oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.69 (s, 1H), 7.60–7.30 (m, 5H), 5.53 (q, $J = 6.5$ Hz, 1H), 3.78 (s, 3H), 1.88 (s, 3H), 1.48 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 196.8, 165.6, 165.0, 133.7, 131.7, 131.1, 129.9, 128.5, 119.0, 115.8, 72.6, 51.8, 29.8, 18.3; **IR** (film, cm^{-1}): 2976, 1705, 1625, 1435, 1363, 1283, 1119, 875, 771, 793; **HRMS** (ESI) Found m/z 295.0942, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{16}\text{H}_{16}\text{NaO}_4$ 295.0941.

Methyl-5-benzoyl-2,6-dimethyl-2H-pyran-3-carboxylate (2h'): The title compound was prepared according to procedure B, 106 mg, 39% yield, colorless oil, $R_f = 0.3$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.72–7.38 (m, 5H), 7.32 (s, 1H), 5.43 (q, $J = 6.5$ Hz, 1H), 3.72 (s, 3H), 2.09 (s, 3H), 1.44 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 193.7, 169.2, 165.1, 139.1, 132.3, 131.6, 128.9, 128.6, 116.8, 112.2, 72.4, 51.6, 26.9, 20.7, 19.3; **IR (film, cm^{-1}): 2976, 1703, 1624, 1435, 1381, 1238, 1147, 1043, 939, 877, 725; **HRMS** (ESI) Found m/z 311.0675, calcd for $[\text{M}+\text{K}^+]$ $\text{C}_{16}\text{H}_{16}\text{KO}_4$ 311.0680.**

Methyl-2-methyl-5-oxo-5,6,7,8-tetrahydro-2H-chromene-3-carboxylate (2i): The title compound was prepared according to procedure B, 188 mg, 85% yield, colorless oil, $R_f = 0.3$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.50 (s, 1H), 5.39 (q, $J = 6.5$ Hz, 1H), 3.72 (s, 3H), 2.49–2.43 (m, 2H), 2.43–2.36 (m, 2H), 2.05–1.88 (m, 2H), 1.33 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 194.1, 175.0, 165.1, 126.7, 119.4, 110.9, 73.5, 51.6, 36.4, 28.8, 20.3, 20.2; **IR** (film, cm^{-1}): 2954, 2927, 1703, 1633, 1556, 1435, 1381, 1274, 1145, 1122, 1076, 954, 754; **HRMS** (ESI) Found m/z 245.0789, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{12}\text{H}_{14}\text{NaO}_4$ 245.0784.

Methyl-2,6-dimethyl-5-(methylsulfonyl)-2H-pyran-3-carboxylate (2j): The title compound was prepared according to procedure B, 223 mg, 91% yield, colorless oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 7.28 (s, 1H), 5.36 (q, $J = 6.5$ Hz, 1H), 3.73 (s, 3H), 2.95 (s, 3H), 2.33 (s, 3H), 1.34 (d, $J = 6.5$ Hz, 3H); $^{13}\text{C NMR}$ (100 MHz, CDCl_3) δ 168.0, 164.3, 128.0 (2C), 118.4, 113.8, 73.1, 51.9, 44.2, 19.3; **IR** (film, cm^{-1}): 2980, 2954, 1703, 1633, 1556, 1435, 1381, 1274, 1228, 1145, 1122, 1043, 954, 754; **HRMS** (ESI) Found m/z 269.0458, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{10}\text{H}_{14}\text{NaO}_5\text{S}$ 269.0454.

Dimethyl-6-(tert-butyl)-2-methyl-2H-pyran-3,5-dicarboxylate (2k): The title compound was prepared according to Procedure B, 71 mg, 27% yield, colorless oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; $^1\text{H NMR}$ (400 MHz, CDCl_3) δ

7.36 (s, 1H), 5.36 (q, $J = 6.5$ Hz, 1H), 3.76 (s, 3H), 3.75 (s, 3H), 1.27 (d, $J = 6.5$ Hz, 3H), 1.25 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 174.8, 167.4, 164.9, 132.7, 116.6, 105.2, 71.3, 51.7, 51.6, 38.2, 27.6, 17.5; **IR** (film, cm^{-1}): 2951, 1701, 1637, 1529, 1435, 1384, 1301, 1230, 1215, 1107, 933, 914, 756; **HRMS** (ESI) Found m/z , 291.1214, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{14}\text{H}_{20}\text{NaO}_5$ 291.1203

5-ethyl-3-methyl-2-methyl-6-phenyl-2H-pyran-3,5-dicarboxylate (2l): The title compound was prepared according to procedure B, 262 mg, 80% yield, yellow oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.65 (s, 1H), 7.53 – 7.30 (m, 5H), 5.53 (q, $J = 6.5$ Hz, 1H), 4.07 (q, $J = 7.1$, 2H), 3.78 (s, 3H), 1.49 (d, $J = 6.5$ Hz, 3H), 1.05 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.8, 165.7, 164.9, 133.9, 131.6, 130.8(2C), 129.5(2C), 127.7, 118.5, 105.3, 72.7, 60.4, 51.7, 18.5, 13.8; **IR** (film, cm^{-1}): 2978, 1697, 1637, 1548, 1487, 1436, 1375, 1300, 1230, 1124, 1080, 989, 920, 754; **HRMS** (ESI) Found m/z 325.1071, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{17}\text{H}_{18}\text{NaO}_5$ 325.1052.

5-ethyl-3-methyl-2-methyl-6-(4-nitrophenyl)-2H-pyran-3,5-dicarboxylate (2m): The title compound was prepared according to procedure B, 336 mg, 97% yield, yellow oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 8.24 (d, $J = 8.9$ Hz, 2H), 7.62 (s, 1H), 7.61 (d, $J = 8.9$ Hz, 2H), 5.56 (q, $J = 6.5$ Hz, 1H), 4.11 (q, $J = 7.1$ Hz, 2H), 3.82 (s, 3H), 1.52 (d, $J = 6.5$ Hz, 3H), 1.12 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 164.7, 164.6, 163.0, 148.8, 140.1, 130.6, 130.5, 123.0, 120.0, 106.7, 73.1, 60.9, 52.0, 18.9, 13.9; **IR** (film, cm^{-1}): 2989, 1693, 1633, 1546, 1435, 1371, 1298, 1236, 1134, 1018, 1991, 856, 752; **HRMS** (ESI) Found m/z 370.0910, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{17}\text{H}_{17}\text{NNaO}_7$ 370.0903.

Methyl-2-methyl-5-oxo-2,5-dihydropyrano-[3,2-c]-chromene-3-carboxylate (2n): The title compound was prepared according to procedure B, 222 mg, 82% yield, yellow oil; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.86 – 7.81 (m, 1H), 7.66 (s, 1H), 7.64 – 7.55 (m, 1H), 7.36 – 7.28 (m, 2H), 5.71 (q, $J = 6.5$ Hz, 1H), 3.82 (s, 3H), 1.50 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 164.5, 161.4, 160.1, 153.9, 133.7, 127.4, 124.4, 123.6, 122.3, 117.1, 114.8, 100.4, 74.0, 52.0, 20.5; **IR** (film, cm^{-1}): 2926, 2854, 1722, 1699, 1637, 1606, 1556, 1490, 1296, 1242, 1205, 1029, 975, 935, 752; **HRMS** (ESI) Found m/z 295.0584, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{15}\text{H}_{12}\text{NaO}_5$ 295.0577.

Methyl-6-ethyl-2-methyl-5-propionyl-2H-pyran-3-carboxylate (2o): The title compound was prepared according to procedure B, 216 mg, 91% yield, colorless oil; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.42 (s, 1H), 5.32 (q, $J = 6.5$ Hz, 1H), 3.72 (s, 3H), 2.98 – 2.77 (m, 1H), 2.73 – 2.50 (m, 2H), 2.48 – 2.29 (m, 1H), 1.25 (d, $J = 6.5$ Hz, 3H), 1.10 (t, $J = 7.5$ Hz, 3H), 1.06 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 197.8, 174.7, 164.9, 130.7, 117.1, 111.4, 71.9, 51.6, 33.5, 26.9, 18.7, 11.1, 8.1. **IR** (film, cm^{-1}): 2980, 2941, 1707, 1678, 1624, 1531, 1436, 1377, 1290, 1220, 1195, 1068, 898, 761; **HRMS** (ESI) Found m/z 277.0840, calcd for $[\text{M}+\text{K}^+]$ $\text{C}_{13}\text{H}_{18}\text{KO}_4$ 277.0837.

Methyl-5-benzoyl-2-methyl-6-phenyl-2H-pyran-3-carboxylate (2p): The title compound was prepared according to procedure B, 166 mg, 50% yield, yellow oil, $R_f = 0.3$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.64 (s, 1H), 7.53 – 7.46 (m, 2H), 7.26 – 7.14 (m, 3H), 7.15 – 6.97 (m, 5H), 5.57 (q, $J = 6.5$ Hz, 1H), 3.73 (s, 3H), 1.57 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 193.9, 165.0, 164.0,

138.1, 133.0, 132.2, 132.1, 131.1, 130.1(2C), 129.3(2C), 128.0(2C), 127.9(2C), 119.0, 113.4, 72.4, 51.8, 18.6; **IR** (film, cm^{-1}): 2974, 2852, 1695, 1672, 1543, 1446, 1435, 1240, 1176, 1053, 1020 941, 904, 771; **HRMS** (ESI) Found m/z 357.1107, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{21}\text{H}_{18}\text{NaO}_4$ 357.1097.

Methyl-2,7,7-trimethyl-5-oxo-5,6,7,8-tetrahydro-2H-chromene-3-carboxylate (2q): The title compound was prepared according to procedure B, 210 mg, 84% yield, colorless oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.51 (s, 1H), 5.41 (q, $J = 6.5$ Hz, 1H), 3.72 (s, 3H), 2.43 – 2.31 (m, 2H), 2.30 – 2.11 (m, 2H), 1.33 (d, $J = 6.5$ Hz, 3H), 1.08 (s, 3H), 1.00 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 193.8, 173.7, 165.1, 126.6, 119.1, 109.9, 73.6, 51.6, 50.3, 42.5, 32.3, 29.4, 27.2, 20.2; **IR** (film, cm^{-1}): 2965, 2872, 1703, 1660, 1571, 1398, 1307, 1217, 1145, 1068, 939, 761; **HRMS** (ESI) Found m/z 273.1099, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{14}\text{H}_{18}\text{NaO}_4$ 273.1097.

Methyl-2-methyl-5-oxo-2,5,6,7-tetrahydrocyclopenta-[b]-pyran-3-carboxylate (2r): The title compound was prepared according to procedure B, 201 mg, 96% yield, colorless oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.29 (s, 1H), 5.65 (q, $J = 6.5$ Hz, 1H), 3.76 (s, 3H), 2.81 – 2.57 (m, 2H), 2.54 (ddd, $J = 12.1$, 9.2, 6.4 Hz, 2H), 1.46 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 199.1, 187.9, 165.0, 125.2(2C), 120.3, 113.7, 51.9, 34.1, 26.7, 21.0; **IR** (film, cm^{-1}): 2989, 2953, 1693, 1633, 1546, 1344, 991, 856, 752; **HRMS** (ESI) Found m/z 231.0637, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{11}\text{H}_{12}\text{NaO}_4$ 231.0628.

Dimethyl-6-(4-fluorophenyl)-2-methyl-2H-pyran-3,5-dicarboxylate (2s): The title compound was prepared according to procedure B, 251 mg, 82% yield, yellow oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.64 (s, 1H), 7.52 – 7.42 (m, 2H), 7.12 – 7.03 (m, 2H), 5.54 (q, $J = 6.5$ Hz, 1H), 3.80 (s, 3H), 3.64 (s, 3H), 1.47 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.9, 165.6, 164.8, 131.9, 131.8, 131.4(2C), 118.7, 115.1(2C), 114.9, 104.7, 72.8, 51.8, 51.5, 18.5; **IR** (film, cm^{-1}): 2976, 1699, 1637, 1602, 1595, 1367, 1083, 1049, 1024, 952, 914, 840, 754; **HRMS** (ESI) Found m/z 329.0807, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{16}\text{H}_{15}\text{FNaO}_5$ 329.0796.

Methyl-2,6-dimethyl-5-(phenylsulfonyl)-2H-pyran-3-carboxylate (2t): The title compound was prepared according to procedure B, 300 mg, 97% yield, colorless oil, $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.90 – 7.79 (m, 2H), 7.63 – 7.48 (m, 3H), 7.41 (s, 1H), 5.34 (q, $J = 6.5$ Hz, 1H), 3.75 (s, 3H), 2.29 (s, 3H), 1.28 (d, $J = 6.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 168.1, 164.4, 142.5, 133.0 (2C), 129.3, 128.3(2C), 126.5, 118.4, 114.4, 73.1, 51.9, 19.5, 19.5; **IR** (film, cm^{-1}): 2980, 1703, 1633, 1556, 1435, 1274, 999, 927, 756; **HRMS** (ESI) Found m/z 331.0600, calcd for $[\text{M}+\text{Na}^+]$ $\text{C}_{15}\text{H}_{16}\text{NaO}_5\text{S}$ 331.0611.

Dimethyl 6-cyclopropyl-2-(2-ethoxy-2-oxoethyl)-2H-pyran-3,5-dicarboxylate (3d): The title compound was prepared according to procedure C, 220 mg; 68% yield; colorless oil; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; ^1H NMR (400 MHz, CDCl_3) δ 7.61 (s, 1H), 5.67 (dd, $J = 10.2$, 3.0 Hz, 1H), 4.27 – 4.09 (m, 2H), 3.79 (s, 3H), 3.77 (s, 3H), 3.12 (tt, $J = 8.2$, 4.9 Hz, 1H), 2.74 (dd, $J = 15.4$, 10.2 Hz, 1H), 2.43 (dd, $J = 15.4$, 3.0 Hz, 1H), 1.34 – 1.31 (m, 1H), 1.28 (t, $J = 7.1$ Hz, 3H), 1.05 – 1.00 (m, 1H), 0.95 – 0.80 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 173.9, 169.4, 165.8, 164.7, 132.8, 113.7, 103.8, 72.2, 60.9, 51.7, 51.47, 37.6, 14.1, 13.2, 10.8, 7.7. **IR** (film, cm^{-1}): 2968, 2954, 1732, 1707, 1693, 1547, 1449, 1393, 1175, 1114, 985,

908, 877, 752; **HRMS** (ESI) Found m/z 347.1107 calcd for $[M+Na^+]$ $C_{16}H_{20}NaO_7$ 347.1101.

Methyl-2-(2-ethoxy-2-oxoethyl)-5-oxo-5,6,7,8-tetrahydro-2H-chromene-3-carboxylate (3i): The title compound was prepared according to procedure C, 211 mg; 72% yield; colorless oil; $R_f = 0.2$ [1:4 (v/v) EA/cyclohexane]; 1H NMR (400 MHz, $CDCl_3$) δ 7.63 (s, 1H), 5.79 (dd, $J = 9.8, 3.2$ Hz, 1H), 4.20 (q, $J = 7.1$ Hz, 2H), 3.79 (s, 3H), 2.78 (dd, $J = 15.0, 9.8$ Hz, 1H), 2.61 (dd, $J = 15.0, 3.2$ Hz, 1H), 2.55 – 2.37 (m, 4H), 2.10 – 1.95 (m, 2H), 1.29 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 194.0, 174.8, 169.2, 164.7, 128.1, 117.1, 111.4, 73.4, 61.0, 51.9, 39.3, 36.4, 28.4, 20.2, 14.2; **IR** (film, cm^{-1}): 2983, 2871, 1726, 1709, 1633, 1592, 1445, 1383, 1271, 1229, 759, 726; **HRMS** (ESI) Found m/z 317.1005, calcd for $[M+Na^+]$ $C_{15}H_{18}NaO_6$: 317.0996.

Methyl-2-(2-ethoxy-2-oxoethyl)-6-methyl-5-(phenylsulfonyl)-2H-pyran-3-carboxylate (3j): The title compound was prepared according to procedure C, 193 mg; 52 % yield; yellow oil; $R_f = 0.2$ [1:5 (v/v) EA/cyclohexane]; 1H NMR (400 MHz, $CDCl_3$) δ 7.95 – 7.81 (m, 2H), 7.68 – 7.52 (m, 3H), 7.50 (d, $J = 0.5$ Hz, 1H), 5.72 (dd, $J = 9.9, 3.3$ Hz, 1H), 4.15 (q, $J = 7.1$ Hz, 2H), 3.80 (s, 3H), 2.73 (dd, $J = 15.0, 9.9$ Hz, 1H), 2.50 (dd, $J = 15.0, 3.3$ Hz, 1H), 2.33 (s, 3H), 1.28 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ 168.9, 168.0, 164.0, 142.2, 133.2, 129.6 (2C), 129.4 (2C), 126.7, 116.3, 115.2, 73.1, 61.0, 52.1, 38.5, 19.2, 14.2; **IR** (film, cm^{-1}): 3061, 2847, 2360, 2341, 1663, 1645, 1620, 1459, 1386, 1276, 1150, 883, 759; **HRMS** (ESI) Found m/z 403.0823, calcd for $[M+Na^+]$ $C_{18}H_{20}NaO_7S$ 403.0822.

Supporting Information

Supporting Information with full analysis of products **1**, **2** and **3** is available free of charge on the ACS Publications website.

Corresponding Authors

serge.thorimbert@upmc.fr; luc.dechoux@upmc.fr.
http://www.ipcm.fr/presentation-682?lang=en.

ACKNOWLEDGMENT

We thank University P. et M. Curie (UPMC) and CNRS for fundings. The Fédération de Recherche (FR2769) provided technical access for analysis. Acknowledgements to the French Embassy in Slovakia and to China Scholarship Council (CSC) for financial supports to K.P. and L.C. respectively.

REFERENCES

- (1) Xie P.; Yang, J.; Zheng, J.; Huang, Y. *Eur. J. Org. Chem.* **2014**, 1189-1194 and references cited herein.
- (2) (a) For a review see: Hsung, R.P.; Kurdyumov, A.V.; Sydorenko, N. *Eur. J. Org. Chem.* **2005**, 23-44. For selected recent examples see: (b) Luo, G.Y.; Wu, H.; Tang, Y.; Li, H.; Yeom, H.S.; Yang, K.; Hsung, R.P. *Synthesis*, **2015**, 47, 2713-2720. (c) Li, K.; Ou, J.; Gao,

S. *Angew. Chem. Int. Ed.* **2016**, 55, 14778-14783. (d) Suzuki, T.; Watanabe, S.; Kobayashi, S.; Tanino, K. *Org. Lett.* **2017**, 19, 922-925. (e) Takeda, Y.; Narita, K.; Katoh, T. *Eur. J. Org. Chem.* **2017**, 901-907.

(3) (a) Gosink, T.A. *J. Org. Chem.* **1974**, 39, 1942-1944. (b) Rodriguez-Otero, J. *J. Org. Chem.* **1999**, 64, 6842-6848. (c) Krasnaya, Zh.A. *Chemistry of Heterocyclic Compounds* **1999**, 35, 1255-1271. (d) Riveira, M.J.; Quiroga, G.N.; Mata, E.G.; Gandon, V.; Mischne, M.P. *J. Org. Chem.* **2015**, 80, 6515-6519.

(4) For selected examples see: (a) Peng, W.; Hirabaru, T.; Kawafuchi, H.; Inokuchi, T. *Eur. J. Org. Chem.* **2011**, 5469-5474. (b) Fotiadou, A.D.; Zografos, A.L. *Org. Lett.* **2012**, 14, 5464-5667. (c) Riveira, M.J.; Mischne, M.P. *Chem. Eur. J.* **2012**, 18, 2382-2388. (d) Edayadulla, N.; Lee, Y.R. *Bull. Korean Chem. Soc.* **2013**, 34, 2963-2967. (e) Narayana, V.R.; Pudukulathan, Z.; Varala, R. *Org. Commun.* **2013**, 6, 110-119.

(5) For other activated methylene groups involved in formal [3+3]-CA see for hydroxyquinolones: (a) Riveira, M.J.; Mischne, M.P. *Synth. Commun.* **2013**, 43, 208-220. (b) For β -oxosulfones: Pena, J.; Moro, R.F.; Basabe, P.; Marcos, I.S.; Diez, D. *RSC Advances*, **2012**, 2, 8041-8049. (c) From dienyl-diketones: Jacob, S.D.; Brooks, J.L.; Frontier, A. J. *J. Org. Chem.* **2014**, 79, 10296-10302.

(6) (a) Menz, H.; Kirsch, S.F. *Org. Lett.* **2006**, 8, 4795-4797. (b) Zhu, Z.B.; Kirsch, S.F. *Chem. Commun.* **2013**, 49, 2272-2283. (c) Vidhani, D.V.; Krafft, M.E.; Alabugin, I.V. *Org. Lett.* **2013**, 15, 4462-4465. (d) Vidhani, D.V.; Krafft, M.E.; Alabugin, I.V. *J. Org. Chem.* **2014**, 79, 352-364. (e) Chong, Q.; Wang, C.; Wang, D.; Wang, H.; Wu, F.; Xin, X.; Wan, B. *Tetrahedron Lett.* **2015**, 56, 401-403.

(7) Rauser, M.; Schroeder, S.; Niggeman, M. *Chem. Eur. J.* **2015**, 21, 15929-15933.

(8) Sun, J.; Zhu, D.; Gong, H.; Yan, C.G. *Tetrahedron* **2013**, 69, 10565-10572.

(9) (a) Hu, J.; Dong, W.; Wu, X.Y.; Tong, X. *Org. Lett.* **2012**, 14, 5530-5533. (b) Ni, C.; Tong, X. *J. Am. Chem. Soc.* **2016**, 138, 7872-7875.

(10) (a) Agarwal, J.; Bayounes, O.; Thorimbert, S.; Dechoux, L. *RSC Advances*, **2014**, 4, 2772-2775. (b) Plevová, K.; Chang, L.; Martin, E.; Llopis, Q.; Dechoux, L.; Thorimbert, S. *Adv. Synth. Catal.* **2016**, 358, 3293-3297.

(11) Addition of aceto-phenylsulfone to methyl coumalate, using DBU in DCM, was described giving one 2H-pyran structure in 67%. See: Frisbee, A.R.; Nantz, M.H.; Kramer, G.W. Fuchs, P.L. *J. Am. Chem. Soc.* **1984**, 106, 7143-7145.

(12) Neises, B.; Steglich, W. *Angew. Chem. Int. Ed.* **1978**, 17, 522-524.

(13) This hypothesis is supported by two results: a first experiment using stoichiometric amount of NaH as base in DCM with 2,4-pentanedione gave **4g** in the enol form after acidic quench. When the reaction was followed by 1H NMR, this compound was detected as an intermediate.

(14) Alabugin, I.V.; Manoharan, M.; Breiner, B.; Lewis, F.D. *J. Am. Chem. Soc.* **2003**, 125, 9329-9342.

(15) (a) Wu, S.; Zhang, Y.; Agarwal, J.; Mathieu, E.; Thorimbert, S.; L. Dechoux. *Tetrahedron*, **2015**, 71, 7663-7669. (b) Wu, S.; Yang, L.; Sun, W.; Si, W.; Xiao, S.; Wang, Q.; Dechoux, L.; Thorimbert, S.; Sollogoub, M.; Zhou, D.; Zhang, Y. *Eur. J. Med. Chem.* **2017**, 130, 308-319.