



HAL
open science

Coastal ocean acidification and increasing total alkalinity in the northwestern Mediterranean Sea

Lydia Kapsenberg, Samir Alliouane, Frédéric Gazeau, Laure Mousseau,
Jean-Pierre Gattuso

► To cite this version:

Lydia Kapsenberg, Samir Alliouane, Frédéric Gazeau, Laure Mousseau, Jean-Pierre Gattuso. Coastal ocean acidification and increasing total alkalinity in the northwestern Mediterranean Sea. *Ocean Science*, 2017, 13 (3), pp.411-426. 10.5194/os-13-411-2017 . hal-01534516

HAL Id: hal-01534516

<https://hal.sorbonne-universite.fr/hal-01534516>

Submitted on 7 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Coastal ocean acidification and increasing total alkalinity in the northwestern Mediterranean Sea

Lydia Kapsenberg¹, Samir Alliouane¹, Frédéric Gazeau¹, Laure Mousseau¹, and Jean-Pierre Gattuso^{1,2}

¹Sorbonne Universités, Université Pierre et Marie Curie-Paris 6, CNRS-INSU, Laboratoire d’Océanographie de Villefranche, 06230 Villefranche-sur-Mer, France

²Institute for Sustainable Development and International Relations, Sciences Po, 27 rue Saint Guillaume, 75007 Paris, France

Correspondence to: Jean-Pierre Gattuso (gattuso@obs-vlfr.fr)

Received: 30 August 2016 – Discussion started: 13 September 2016

Revised: 10 April 2017 – Accepted: 14 April 2017 – Published: 15 May 2017

Abstract. Coastal time series of ocean carbonate chemistry are critical for understanding how global anthropogenic change manifests in near-shore ecosystems. Yet, they are few and have low temporal resolution. At the time series station Point B in the northwestern Mediterranean Sea, seawater was sampled weekly from 2007 through 2015, at 1 and 50 m, and analyzed for total dissolved inorganic carbon (C_T) and total alkalinity (A_T). Parameters of the carbonate system such as pH (pH_T , total hydrogen ion scale) were calculated and a deconvolution analysis was performed to identify drivers of change. The rate of surface ocean acidification was -0.0028 ± 0.0003 units $pH_T \text{ yr}^{-1}$. This rate is larger than previously identified open-ocean trends due to rapid warming that occurred over the study period (0.072 ± 0.022 °C yr^{-1}). The total pH_T change over the study period was of similar magnitude as the diel pH_T variability at this site. The acidification trend can be attributed to atmospheric carbon dioxide (CO_2) forcing (59 %, 2.08 ± 0.01 ppm $\text{CO}_2 \text{ yr}^{-1}$) and warming (41 %). Similar trends were observed at 50 m but rates were generally slower. At 1 m depth, the increase in atmospheric CO_2 accounted for approximately 40 % of the observed increase in C_T (2.97 ± 0.20 $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$). The remaining increase in C_T may have been driven by the same unidentified process that caused an increase in A_T (2.08 ± 0.19 $\mu\text{mol kg}^{-1} \text{ yr}^{-1}$). Based on the analysis of monthly trends, synchronous increases in C_T and A_T were fastest in the spring–summer transition. The driving process of the interannual increase in A_T has a seasonal and shallow component, which may indicate riverine or groundwater influence. This study exemplifies the importance of understanding changes in coastal carbonate chemistry through the

lens of biogeochemical cycling at the land–sea interface. This is the first coastal acidification time series providing multi-year data at high temporal resolution. The data confirm rapid warming in the Mediterranean Sea and demonstrate coastal acidification with a synchronous increase in total alkalinity.

1 Introduction

Maintaining time series of oceanographic data is essential for understanding anthropogenic changes in the ocean (Tanhua et al., 2013). On land, fossil fuel burning, cement production, and land use changes have contributed ~ 600 Gt carbon to the atmosphere during the period 1750–2015 (Le Quéré et al., 2016). In the recent decade 2006–2015, an estimated 25 % of this anthropogenic carbon has been absorbed by the ocean in the form of carbon dioxide (CO_2 ; Le Quéré et al., 2016), and is causing global changes to the ocean carbonate system. Absorption of CO_2 by seawater produces carbonic acid, which decreases seawater pH, and is of great concern for biological processes and marine ecosystems (Doney et al., 2009; Gattuso and Hansson, 2011; Pörtner et al., 2014). Since the preindustrial era, global mean ocean pH has declined by 0.1 (Rhein et al., 2013). Due to the declining trend of ocean pH with increasing anthropogenic CO_2 , the process is termed “ocean acidification”. This expression represents a suite of chemical changes, including increases in total dissolved inorganic carbon (C_T) and partial pressure of CO_2 ($p\text{CO}_2$) and decrease in calcium carbonate saturation states (Ω , aragonite and calcite; Dickson, 2010). Rates of ocean acidification differ by ocean region and range from

–0.0026 (Irminger Sea, North Atlantic) to –0.0013 (South Pacific) units pH yr^{-1} (Bates et al., 2014). Such time series remain spatially limited, especially in coastal regions, which provide valuable ecosystem services (Barbier et al., 2011; Costanza et al., 1997) and are under high anthropogenic impact (Halpern et al., 2008). Here, we present the first coastal ocean acidification time series at high temporal resolution.

Compared to the global ocean, marginal seas play a critical role in anthropogenic CO_2 storage via enhanced CO_2 uptake and export to the ocean interior (Lee et al., 2011). As a marginal sea, the Mediterranean Sea has a naturally high capacity to absorb but also buffer anthropogenic CO_2 (Álvarez et al., 2014; Palmiéri et al., 2015). This is primarily due to the high total alkalinity (A_T) of Mediterranean waters and overturning circulation (Lee et al., 2011; Palmiéri et al., 2015; Schneider et al., 2010). In the Mediterranean Sea, the salinity– A_T relationship is driven by the addition of river discharge and Black Sea input, which are generally high in A_T (Copin-Montégut, 1993; Schneider et al., 2007). Combined with evaporation, this results in higher A_T and salinity in the Mediterranean Sea compared to the Atlantic Mediterranean source water (Jiang et al., 2014). On average, Mediterranean Sea A_T is 10 % higher than in the global ocean (Palmiéri et al., 2015). The surface ocean acidification rate, estimated at ΔpH_T (total hydrogen ion scale) of –0.08 since 1800, is comparable to that of the global ocean despite a 10 % greater anthropogenic carbon inventory (Palmiéri et al., 2015). Due to its important role in carbon sequestration and ecological sensitivity to anthropogenic change with economic consequences (Lacoue-Labarthe et al., 2016), the Mediterranean Sea could provide insight to global trends (Lejeune et al., 2010).

Over the last few years, numerous studies have estimated ocean acidification rates across the Mediterranean Sea (Table 1). Together, these studies cover various periods with a range of techniques yielding different results. For example, estimates of change in pH of bottom waters since the preindustrial era range between –0.005 and –0.06 (Palmiéri et al., 2015) and up to as much as –0.14 for full profile estimates (Touratier and Goyet, 2011). Techniques for estimating ocean acidification in the Mediterranean Sea thus far include the following: (1) hind-casting, using high-resolution regional circulation models (Palmiéri et al., 2015), the TrOCA approach as applied to cruise-based profile data (Krasakopoulou et al., 2011; Touratier and Goyet, 2011; Touratier et al., 2016), and others (Howes et al., 2015); (2) partially reconstructed time series (Marcellin Yao et al., 2016); (3) comparative study periods (Luchetta et al., 2010; Meier et al., 2014); and (4) sensor-based observations over a short study period (Flecha et al., 2015). Ocean acidification time series of consistent sampling over many years are lacking for the Mediterranean Sea (The MerMex Group, 2011), particularly along the coast where river discharge influences the carbonate system (Ingrosso et al., 2016).

Table 1. Previous estimates or documentation of pH change (ΔpH) in the Mediterranean Sea. “Total” indicates estimates made for the whole Mediterranean Sea. TrOCA is the “Tracer combining Oxygen, inorganic Carbon, and total Alkalinity” method, NR means “not reported”, and PI is “preindustrial era”.

Region	Site	Method	Study period	pH scale	°C	$\Delta\text{pH yr}^{-1} \pm \text{SE}$	Total ΔpH	Reference
NW	Point B, 1 m	time series, anomaly	2007–2015	total	in situ	–0.0028 \pm 0.0003	–0.0252	This study
NW	Point B, 1 m	time series, anomaly	2007–2015	total	25	–0.0017 \pm 0.0002	–0.0153	This study
NW	Point B	model	1967–2003	total	in situ	–0.0014	–0.05	Howes et al. (2015)
NW	DYFAMED	time series, observed	1995–2011	seawater	in situ	–0.003 \pm 0.001	–0.051	Marcellin Yao et al. (2016)
NW	DYFAMED	time series comparison	1998–2000, 2003–2005	seawater	in situ*	–	–0.02	Meier et al. (2014)
NW	Gulf of Lion	TrOCA	PI–2011	NR	in situ*	–	–0.15 to –0.11	Touratier et al. (2016)
East	N Adriatic Sea	cruise comparison	1983, 2008	total	25	–0.0025	–0.063	Luchetta et al. (2010)
East	Otranto Strait	TrOCA	PI–1995	seawater	25	–	< –0.1 to –0.05, \pm 0.014	Krasakopoulou et al. (2011)
Total	Full profile	TrOCA	PI–2001	NR	in situ*	–	–0.14 to –0.05	Touratier and Goyet (2011)
Total	Bottom waters	model	1800–2001	total	in situ*	–	–0.06 to –0.005	Palmiéri et al. (2015)
Total	Surface waters	model	1800–2001	total	in situ*	–	–0.084 \pm 0.001	Palmiéri et al. (2015)
Gibraltar Strait	Espartel sill	pH, μCO_2 sensors	2012–2015	total	25	–0.0044 \pm 0.00006	–	Flecha et al. (2015)

* Indicates studies where the reported pH change was assumed to be at in situ temperatures.

Compared to the open ocean, shallow coastal sites exhibit natural variability in carbonate chemistry over annual time frames (Hofmann et al., 2011; Kapsenberg and Hofmann, 2016; Kapsenberg et al., 2015), complicating the detection and relevance of open ocean acidification in isolation of other processes (Duarte et al., 2013). In the northwestern (NW) Pacific coast, rapid acidification of surface waters ($\Delta\text{pH}_T - 0.058$ units yr^{-1}) at Tatoosh Island was documented in the absence of changes in known drivers of local pH variability (e.g., upwelling, eutrophication, and more; Wootton and Pfister, 2012; Wootton et al., 2008). Further inshore, in the Hood Canal subbasin of the Puget Sound, only 24–49 % of the estimated pH decline from preindustrial values could be attributed to anthropogenic CO_2 (Feely et al., 2010). The excess decrease in pH was attributed to increased remineralization (Feely et al., 2010). Acidification rates documented along the North Sea Dutch coastline and inlets were highly variable in space, with some exceeding the expected anthropogenic CO_2 rate by an order of magnitude while others exhibited an increase in pH (Provoost et al., 2010).

Variability in coastal carbonate chemistry stems from both physical (e.g., upwelling, river discharge; Feely et al., 2008; Vargas et al., 2016) and biological processes (e.g., primary production, respiration, net calcification). Within watersheds, coastal carbonate chemistry is affected by eutrophication (Borges and Gypens, 2010; Cai et al., 2011), groundwater supply (Cai et al., 2003), and land use and rain influence on river alkalinity (Raymond and Cole, 2003; Stets et al., 2014). Over longer periods, pH can also be influenced by atmospheric deposition (Omstedt et al., 2015). Through primary production and respiration, coastal ecosystems produce pH fluctuations over timescales ranging from hours (e.g., sea-grass, kelp) to months (e.g., phytoplankton blooms; Kapsenberg and Hofmann, 2016). Due to existing pH variability in coastal seas, it is necessary to quantify high-frequency trends in order to interpret the pH changes inferred from lower-frequency sampling.

In this study, we present the first complete time series data quantifying the present-day ocean acidification rate for a coastal site in the Mediterranean Sea, based on weekly measurements of A_T and C_T sampled from 2007 through 2015. For a subset of this time series, pH variability was documented using a SeaFETTM Ocean pH Sensor in order to assess hourly pH variability. For comparison and consistency with other ocean acidification time series around the world, we report rates of change based on anomalies (Bates et al., 2014) and identify drivers of change.

2 Materials and methods

2.1 Site description

A carbonate chemistry time series was initiated in 2007 and maintained through 2015 in the NW Mediterranean Sea at the

entrance of the Bay of Villefranche-sur-Mer, France (Fig. 1): Point B station (43.686° N, 7.316° E, 85 m bottom depth). A second site, the Environment Observable Littoral buoy (EOL, 43.682° N, 7.319° E, 80 m bottom depth), was used for pH sensor deployment starting in 2014. These two sites are 435 m apart. Point B has been an oceanographic station since 1957. A full site description and research history has been detailed by De Carlo et al. (2013). Briefly, the Bay is a narrow north–south facing inlet with steep bathymetry and estimated volume of 310 million m^3 . The surrounding region is predominantly composed of limestone with a series of shallow, submarine groundwater karst springs (Gilli, 1995). The north current, a major and structuring counterclockwise current in the Ligurian Sea, can sometimes flow close to Point B. The Bay can also be, on occasion, influenced by local countercurrents. Both of these hydrodynamic movements have signatures of river discharge. Limestone erosion can be observed in the A_T of rivers nearest to Point B (Paillon, due 4 km west; Var due 10 km west; and Roya due 26 km east). River A_T ranges between 1000 and 2000 $\mu\text{mol kg}^{-1}$ (data from Agence de l'Eau Rhône-Méditerranée-Corse, <http://sierm.eaurmc.fr>), and is lower than seawater A_T . The Paillon River, whose plume on occasion reaches into the Bay (L. Mousseau, personal observation, 2013, 2014), was sampled on 18 August 2014 and had a A_T of $1585 \pm 0.1 \mu\text{mol kg}^{-1}$ ($N = 2$, Gattuso, unpublished). Due to low primary productivity, seasonal warming drives the main annual variability in carbonate chemistry at this location (De Carlo et al., 2013).

2.2 Point B data collection, processing, and analysis

To document long-term changes in ocean carbonate chemistry at Point B, seawater was sampled weekly from 9 January 2007 to 22 December 2015. Samples were collected at 1 and 50 m, using a 12 L Niskin bottle at 09:00 LT. Seawater was transferred from the Niskin bottle to 500 mL borosilicate glass bottles and fixed within an hour via addition of saturated mercuric chloride for preservation of carbonate parameters, following recommendations by Dickson et al. (2007). Duplicate samples were collected for each depth. For each sampling event, CTD casts were performed either with a Seabird 25 or Seabird 25+ profiler whose sensors are calibrated at least every 2 years. Accuracy of conductivity (SBE4 sensor) and temperature (SBE3 sensor) measurements from CTD casts were 0.0003 S m^{-1} and $0.001 \text{ }^\circ\text{C}$, respectively.

Within 6 months of collection, bottle samples were analyzed for C_T and A_T via potentiometric titration following methods described by Edmond (1970) and DOE (1994), by Service National d'Analyse des Paramètres Océaniques du CO_2 , at the Université Pierre et Marie Curie in Paris, France. Average accuracy of C_T and A_T measurements was 2.6 and $3 \mu\text{mol kg}^{-1}$, respectively, compared to seawater certified reference material (CRM) provided by A. Dickson (Scripps Institution of Oceanography). Repeatability of replicate samples was better than $3 \mu\text{mol kg}^{-1}$.




Figure 1. Map of study region in the NW Mediterranean Sea (a), along the north current (b) in the Bay of Villefranche-sur-Mer, France (c). Point B, the EOL buoy, and the offshore time series station DYFAMED are marked. Bathymetric line units are measured in meters (m) (c).

Only obvious outliers were omitted from the analyses: three C_T values at 1 m ($>2300 \mu\text{mol kg}^{-1}$), one A_T value at 1 m ($>2900 \mu\text{mol kg}^{-1}$), and one A_T value at 50 m ($<2500 \mu\text{mol kg}^{-1}$). The C_T and A_T measurements on replicate bottle samples were averaged for analyses.

Calculations of the carbonate system parameters were performed using the R package seacarb version 3.1 with C_T , A_T , in situ temperature, and salinity as inputs (Gattuso et al., 2016). Total concentrations of silicate (SiOH_4) and phosphate (PO_4^{3-}) were used, when available from Point B (Mousseau, unpublished, <http://somlit.epoc.u-bordeaux1.fr/fr/>). Detection limits for nutrients were $0.03 \mu\text{M}$ for SiOH_4 and 0.003 to $0.02 \mu\text{M}$ for PO_4^{3-} ; relative precision of these analyses is 5–10 % (Aminot and K erouel, 2007). Total boron concentration was calculated from salinity using the global ratio determined by Lee et al. (2010). The following constants were used: K_1 and K_2 from Lueker et al. (2000), K_f from Perez and Fraga (1987), and K_s from Dickson (1990). Reported measured parameters are temperature, salinity, A_T , and C_T , and derived parameters are pH_T (total hydrogen ion scale), pH_T normalized to 25°C (pH_{T25}), $p\text{CO}_2$, and aragonite (Ω_a) and calcite (Ω_c) saturation states. Salinity-normalized changes in A_T (nA_T) and C_T (nC_T) were calculated by dividing by in situ salinity and multiplying by 38. Except for pH_{T25} , all parameters are reported at in situ temperatures.

The average uncertainties of the derived carbonate parameters were calculated according to the Gaussian method (Dickson and Riley, 1978) implemented in the “errors” function of the R package seacarb 3.1 (Gattuso et al., 2016). The uncertainties are $\pm 2.7 \times 10^{-10} \text{ mol H}^+$ (about 0.015 units pH_T), $\pm 15 \mu\text{atm } p\text{CO}_2$, and ± 0.1 unit of the aragonite and calcite saturation states.

To quantify interannual changes in carbonate parameters, the data were detrended for seasonality by subtracting the respective climatological monthly means computed for the period 2009–2015 from the time series (hereafter “monthly means”). The resulting residuals were analyzed using a linear regression to compute anomaly trends. This approach follows methods from Bates et al. (2014) to allow for comparisons of trends observed at different time series stations. All analyses were performed in R (R Core Team, 2016).

2.3 Deconvolution of pH_T and $p\text{CO}_2$

To identify proportional contributions of various drivers to ocean acidification trends at Point B, deconvolution of time series pH_T and $p\text{CO}_2$ was performed following methods from Garc a-Ib a ez et al. (2016) for observations at 1 and 50 m. The equation is described below for pH_T , where changes in pH_T are driven by changes in temperature (T), salinity (S), A_T , and C_T , over time (t).

$$\frac{d\text{pH}_T}{dt} = \frac{\partial \text{pH}_T}{\partial T} \frac{dT}{dt} + \frac{\partial \text{pH}_T}{\partial S} \frac{dS}{dt} + \frac{\partial \text{pH}_T}{\partial A_T} \frac{dA_T}{dt} + \frac{\partial \text{pH}_T}{\partial C_T} \frac{dC_T}{dt} \quad (1)$$

Here, $\frac{\partial \text{pH}_T}{\partial \text{var}} \frac{d\text{var}}{dt}$ represents the slope contribution of changing “var” to the estimated change in pH_T ($\frac{d\text{pH}_T}{dt}$), where var is either temperature (T), salinity (S), A_T , or C_T . The sensitivity of pH to var ($\frac{\partial \text{pH}_T}{\partial \text{var}}$) was estimated by calculating pH_T using the true observations of var and holding the other three variables constant (mean value of the time series) and regressing it to var. Sensitivity ($\frac{\partial \text{pH}_T}{\partial \text{var}}$) was then multiplied by the anomaly rate of var (Table 2). The calculation was repeated for $p\text{CO}_2$ ($\frac{dp\text{CO}_2}{dt}$) in order to compare the rate of increase with that of atmospheric CO_2 .

Table 2. Time series anomaly regression analyses on seawater carbonate chemistry at Point B for salinity (S), temperature (T), dissolved inorganic carbon (C_T), total alkalinity (A_T), pH_T , pH_T normalized to 25 °C (pH_{T25}), $p\text{CO}_2$, calcite (Ω_c) and aragonite (Ω_a) saturation state, and salinity-normalized A_T (nA_T) and C_T (nC_T), at 1 and 50 m. Slopes represent the change in the variable unit per year. $P \ll 0.001$ indicates p values far smaller than 0.001. SE stands for standard error. Degrees of freedom are denoted by df.

Depth (m)	Variable	Slope \pm SE	Intercept \pm SE	N	F	df	Slope P	R^2	
1	S	-0.0017 ± 0.0044	3.38 ± 8.82	417	0.147	1, 415	0.702	0	
	T (°C)	0.072 ± 0.022	-145 ± 44	413	10.999	1, 411	0.001	0.026	
	C_T ($\mu\text{mol kg}^{-1}$)	2.97 ± 0.20	-5965 ± 400	416	221.87	1, 414	$\ll 0.001$	0.349	
	A_T ($\mu\text{mol kg}^{-1}$)	2.08 ± 0.19	-4189 ± 379	417	122.429	1, 415	$\ll 0.001$	0.228	
	pH_T	-0.0028 ± 0.0003	5.72 ± 0.66	412	74.205	1, 410	$\ll 0.001$	0.153	
	pH_{T25}	-0.0017 ± 0.0002	3.46 ± 0.43	412	64.204	1, 410	$\ll 0.001$	0.1354	
	$p\text{CO}_2$ (μatm)	3.53 ± 0.39	-7105 ± 776	412	83.927	1, 410	$\ll 0.001$	0.17	
	Ω_c	-0.0109 ± 0.0022	22.0 ± 4.5	412	24.08	1, 410	$\ll 0.001$	0.055	
	Ω_a	-0.0064 ± 0.0015	12.9 ± 3.1	412	17.33	1, 410	$\ll 0.001$	0.041	
	nA_T ($\mu\text{mol kg}^{-1}$)	2.20 ± 0.28	-4425 ± 560	412	62.34	1, 410	$\ll 0.001$	0.132	
	nC_T ($\mu\text{mol kg}^{-1}$)	3.12 ± 0.29	-6275 ± 579	412	117.486	1, 410	$\ll 0.001$	0.223	
	50	S	0.0063 ± 0.0020	-12.8 ± 4.1	412	9.858	1, 410	0.002	0.0235
		T (°C)	0.088 ± 0.019	-177 ± 38	408	21.927	1, 406	$\ll 0.001$	0.0512
C_T ($\mu\text{mol kg}^{-1}$)		2.16 ± 0.21	-4344 ± 418	411	108.105	1, 409	$\ll 0.001$	0.2091	
A_T ($\mu\text{mol kg}^{-1}$)		1.59 ± 0.15	-3192 ± 309	412	106.947	1, 410	$\ll 0.001$	0.2069	
pH_T		-0.0026 ± 0.0002	5.28 ± 0.50	407	112.111	1, 405	$\ll 0.001$	0.2168	
pH_{T25}		-0.0013 ± 0.0003	2.55 ± 0.54	407	21.863	1, 405	$\ll 0.001$	0.0512	
$p\text{CO}_2$ (μatm)		2.79 ± 0.25	-5603 ± 501	407	125.1	1, 405	$\ll 0.001$	0.236	
Ω_c		-0.0070 ± 0.0027	14.0 ± 5.4	407	6.648	1, 405	0.01	0.0162	
Ω_a		-0.0038 ± 0.0019	7.6 ± 3.7	407	4.155	1, 405	0.042	0.0102	
nA_T ($\mu\text{mol kg}^{-1}$)		1.15 ± 0.13	-2309 ± 254	407	82.309	1, 405	$\ll 0.001$	0.1689	
nC_T ($\mu\text{mol kg}^{-1}$)		1.82 ± 0.19	-3661 ± 376	407	94.98	1, 405	$\ll 0.001$	0.19	

As a subcomponent of $\frac{\partial p\text{CO}_2}{\partial C_T} \frac{dC_T}{dt}$, the rate of anthropogenic CO_2 increase was estimated from atmospheric CO_2 concentrations nearest to Point B (Plateau Rosa, Italy, courtesy of the World Data Center for Greenhouse Gases, <http://ds.data.jma.go.jp/gmd/wdcgg/>). For these data, missing daily values were linearly interpolated, climatological monthly means were calculated and subtracted from the time series to generate an anomaly time series. A linear regression was performed on anomalies where the slope represents the rate of anthropogenic CO_2 increase in the atmosphere. Finally, to help identify different processes that might have contributed to the observed trends, linear regressions were performed on changes in A_T and C_T per month (mean value of observations within 1 month) from 2009 through 2015 and on the salinity– A_T relationship by year.

2.4 SeaFET data collection, processing, and analysis

To capture pH variability at higher-than-weekly sampling frequencies, a SeaFET™ Ocean pH sensor (Satlantic) was deployed on the EOL buoy (435 m from the Point B sampling site) starting in June 2014, at 2 m depth. Autonomous sampling was hourly and deployment periods averaged 58 ± 25 days with 5 ± 2 calibration samples per deployment.

Field calibration samples for pH were collected using a Niskin bottle next to SeaFET within 15 min of measurement. This sampling scheme was sufficient for this site as there is no large high-frequency pH variability. Unlike Point B sampling, SeaFET calibration samples were processed for pH using the spectrophotometric method (Dickson et al., 2007) with purified m-cresol purple (purchased from Robert H. Byrne's laboratory, University of South Florida). In situ temperature, salinity, and A_T measured at Point B, within 30 min of the SeaFET sampling, were used to calculate in situ pH_T of the calibration samples.

SeaFET voltage was converted to pH_T using the respective calibration samples for each deployment period, following the methods and code described in Bresnahan et al. (2014) but adapted for use in R. Only 5 % of the data was removed during quality control, due to biofouling in one deployment and battery exhaustion in another, yielding 610 days of data. The mean offset between calibration samples and the calibrated SeaFET pH time series was ± 0.007 , indicating a high-quality pH dataset (data shown in Fig. 7c).

The estimated standard uncertainty in SeaFET pH_T is ± 0.01 and was calculated as the square root of the sum of each error squared. The sources of errors are measurement error of spectrophotometric pH (± 0.004 , $N = 68$ mean SD of 5 replicate measurements per calibration sample for samples




Figure 2. Time series observations (a–g) and anomaly trends (h–n) for temperature, salinity, and seawater carbonate chemistry at Point B, 1 m. Regression slopes are drawn \pm SE (standard error, in grey) and noted with a star for significance at $\alpha = 0.05$. Variable abbreviations are the same as in Table 2.

collected between 16 July 2014 and 3 May 2016), spatiotemporal mismatch sampling at EOL (± 0.007 , mean offset of pH_T of the calibration samples from calibrated time series), and variability in purified m-cresol dye batch accuracy compared to tris buffer CRM pH (± 0.006 , mean offset of pH_T of the spectrophotometric measurement of tris buffer from the CRM value).

3 Results

3.1 Time series trends

At Point B from January 2007 to December 2015, more than 400 samples were collected for carbonate chemistry at both 1 and 50 m. Anomaly trends detected at 1 m (Fig. 2) were also significant at 50 m (Fig. 3, Table 2), with the exception that salinity only increased at 50 m (0.0063 ± 0.0020 units yr^{-1}). At 1 m, trends were significant for pH_T (-0.0028 units yr^{-1}),

A_T ($2.08 \mu\text{mol kg}^{-1} \text{yr}^{-1}$), C_T ($2.97 \mu\text{mol kg}^{-1} \text{yr}^{-1}$), pCO_2 ($3.53 \mu\text{atm yr}^{-1}$), and Ω_a (-0.0064 units yr^{-1}). At the same time, temperature anomaly increased (0.072 °C yr^{-1}). Changes in carbonate chemistry parameters were greater at 1 compared to 50 m, with the exception of salinity and temperature. The warming rate at 50 m was slightly greater compared to 1 m, mostly due to increasing summer temperatures since 2007. Time series data are available online (Gattuso et al., 2014).

Strong seasonal cycles of carbonate chemistry parameters were present at Point B at 1 m (Fig. 4). Climatological monthly means (2007–2015) are described briefly and listed in Supplement Table S1. Mean temperature range was 11.2 °C with a maximum of $24.77 \pm 1.35 \text{ °C}$ in August and minimum of $13.58 \pm 0.41 \text{ °C}$ in February. The range in A_T was $19 \mu\text{mol kg}^{-1}$ from June to September. The C_T range was $33 \mu\text{mol kg}^{-1}$ with a peak in late winter and minimum values in August and October. Due to summer warming co-

inciding with the period of peak primary productivity (De Carlo et al., 2013), warming countered the influence of low C_T on pH. As a result, pH_T reached minimum values in summer (8.02 ± 0.03 , July and August) and peaked in late winter (8.14 ± 0.01 , February and March), for an overall annual pH range of 0.12. The corresponding pCO_2 range was $128 \mu\text{atm}$ from February to August. Seasonal cycles were smaller at 50 compared to 1 m (Table S1).

3.2 Deconvolution of pH_T and pCO_2

Deconvolutions of pH and pCO_2 are presented in Tables 3 and 4, respectively. The estimated anomaly trends $\left(\frac{dpH_T}{dt}, \frac{dpCO_2}{dt}\right)$ from the deconvolution fall within the error of the observed anomaly trends (Table 2). The contribution of warming to the pH_T anomaly ($-0.0011 \text{ units yr}^{-1}$, at 1 m) matched the difference between the trends of pH_T and pH_{T25C} (Table 2), which verifies that the deconvolution reproduced influences of temperature sensitivity well. Overall, these results indicate that the deconvolution analyses represent the observed trends well.

At both depths, the predominant driver of $\frac{dpH_T}{dt}$ and $\frac{dpCO_2}{dt}$ was the increase in C_T . Increasing A_T countered 66–69 and 60 % of the influence of increasing C_T on $\frac{dpH_T}{dt}$ and $\frac{dpCO_2}{dt}$, respectively. At 1 m, warming accounted for 41 and 37 % of $\frac{dpH_T}{dt}$ and $\frac{dpCO_2}{dt}$, respectively. Since warming was slightly greater at 50 compared to 1 m, warming accounted for a larger proportional influence on $\frac{dpH_T}{dt}$ and $\frac{dpCO_2}{dt}$ at 50 compared to 1 m. Increasing salinity at 50 m contributed slightly to $\frac{dpH_T}{dt}$ (4 %) and $\frac{dpCO_2}{dt}$ (2 %).

The atmospheric CO_2 anomaly at Plateau Rosa increased by $2.08 \pm 0.01 \text{ ppm yr}^{-1}$ ($F_{1,3285} = 4664$, $P \ll 0.001$, $R^2 = 0.93$) during the study period 2007–2015, and represents the anthropogenic CO_2 forcing on seawater pH. To estimate the influence of anthropogenic CO_2 forcing at Point B, we assume air–sea CO_2 equilibrium (e.g., an increase in atmospheric CO_2 causes an equal increase in seawater pCO_2) for the water mass at 1 m. This assumption is based on evidence that Point B is a weak sink for atmospheric CO_2 with near-balanced air–sea CO_2 flux on an annual time frame (De Carlo et al., 2013). Considering the error associated with deconvolution of pCO_2 at 1 m, the atmospheric CO_2 increase can, at most, represent 38–43 % of the total C_T contribution $\left(\frac{\partial pCO_2}{\partial C_T} \frac{dC_T}{dt}\right)$ to $\frac{dpCO_2}{dt}$. This leaves 57–62 % of the total C_T contribution to pCO_2 trends unaccounted for.

As A_T is not influenced by addition of anthropogenic CO_2 to seawater, but it did increase, the next question was whether or not the changes in A_T and C_T were process-linked. At 1 m, regressions of annual monthly trends of A_T and C_T revealed similar seasonal cycles for both parameters (Fig. 5, Table S2). The fastest increases in A_T and C_T occurred from May through July. The smallest (non-significant) changes occurred in January. The synchronicity between monthly trends

of A_T and C_T was also observed at 50 m, but the rates were slower (analysis not shown).

3.3 Salinity and A_T relationships

Over an annual observation period at 1 m, salinity was a poor predictor of A_T , with the exception of 2007 (Fig. 6). The R^2 value for each annual salinity– A_T relationship at 1 m ranged from 0.00 (in 2013) to 0.87 (in 2007) with y intercepts (A_{T0} , total alkalinity of the freshwater end-member) ranging between $-176 \mu\text{mol kg}^{-1}$ (in 2007) and $2586 \mu\text{mol kg}^{-1}$ (in 2013). The interannual variability of the salinity– A_T relationship was driven by the variability in A_T observed at salinity < 38.0 that was present from November through July.

Changes in salinity (based on monthly means) at Point B were small and ranged from 37.64 ± 0.26 to 38.21 ± 0.11 from May to September, following freshwater input in winter and spring and evaporation throughout summer and fall (Fig. 4). The highest (> 38.0) and most stable salinity observations were made in August through October and coincided with the period of maximum A_T (2562 and $2561 \pm 9 \mu\text{mol kg}^{-1}$ in September and October, respectively). Minimum A_T ($2543 \pm 14 \mu\text{mol kg}^{-1}$) was observed in June, 1 month after minimum salinity. To capture this seasonality without the interannual variation of A_T , the salinity– A_T relationship at 1 m was estimated from climatological monthly means (cA_T and cS , $N = 12$) where cA_T units are measured in micromoles per kilogram ($\mu\text{mol kg}^{-1}$) and error terms are standard errors ($R^2 = 0.74$):

$$cA_T = 1554.9 (\pm 185.9) + 26.3 (\pm 4.9) \times cS. \quad (2)$$

At 50 m, monthly salinity and A_T were less correlated over an annual cycle (analysis not shown). Salinity remained stable at 38.0 from January through September while A_T declined by $13 \mu\text{mol kg}^{-1}$. In general, seasonal changes were dampened at 50 compared to 1 m.

3.4 High-frequency pH data

To verify the weekly sampling scheme at Point B, a continuous record of high-frequency pH observations was obtained via SeaFET deployments from June 2014 to April 2016 (Fig. 7). Sensor data corroborated the seasonal pH and temperature cycle observed at Point B (Fig. 7a–b). Event-scale effects (e.g., pH_T change ≥ 0.1 for days to weeks, sensu Kapsenberg and Hofmann, 2016) were absent at this site, suggesting that weekly sampling was sufficient to describe seasonal and interannual changes in carbonate chemistry at Point B. The magnitude of diel pH_T variability was small: the 2.5th to 97.5th percentiles ranged between 0.01 and 0.05 units pH_T (Fig. 7d–e). Variability increased from winter to spring with the greatest variations in April, May, and June ($\Delta pH_T > 0.035$). The magnitude of diel pH variability was not correlated with temperature or the concentration of chlorophyll *a* (data not shown).

Table 3. Deconvolution of pH_T anomalies ($\frac{d\text{pH}_T}{dt}$, units $\text{pH}_T\text{yr}^{-1}$) at 1 and 50 m. Sensitivity of pH_T with respect to variables ($\frac{\partial\text{pH}_T}{\partial\text{var}}$), where the variable var is either temperature (T), salinity (S), total alkalinity (A_T), or dissolved inorganic carbon (C_T), was multiplied by the anomaly of var ($\frac{d\text{var}}{dt}$, Table 2). SE is standard error and RMSE is root mean square error. Rounding was performed at the end of the calculations, prior to the estimation of the percentage of contributions.

Depth (m)	var	$\frac{\partial\text{pH}_T}{\partial\text{var}} \pm \text{SE}$	$\frac{\partial\text{pH}_T}{\partial\text{var}} \frac{d\text{var}}{dt} \pm \text{RMSE}$	Contribution (%)	$\frac{d\text{pH}_T}{dt} \pm \text{RMSE}$
1	T ($^{\circ}\text{C}$)	$-0.0153 \pm <0.0001$	-0.0011 ± 0.0003	41	-0.0027 ± 0.0005
	S	$-0.0117 \pm <0.0001$	$<0.0001 \pm 0.0001$	0	
	A_T ($\mu\text{mol kg}^{-1}$)	$0.0015 \pm <0.0001$	0.0031 ± 0.0003	-115	
	C_T ($\mu\text{mol kg}^{-1}$)	$-0.0016 \pm <0.0001$	-0.0047 ± 0.0003	174	
50	T ($^{\circ}\text{C}$)	$-0.0154 \pm <0.0001$	-0.0014 ± 0.0003	54	-0.0026 ± 0.0005
	S	$-0.0116 \pm <0.0001$	$-0.0001 \pm <0.0001$	4	
	A_T ($\mu\text{mol kg}^{-1}$)	$0.0015 \pm <0.0001$	0.0024 ± 0.0002	-92	
	C_T ($\mu\text{mol kg}^{-1}$)	$-0.0016 \pm <0.0001$	-0.0035 ± 0.0003	135	

Table 4. Deconvolution of $p\text{CO}_2$ anomalies ($\frac{dp\text{CO}_2}{dt}$, $\mu\text{atm yr}^{-1}$) at 1 and 50 m. Details are the same as in Table 3.

Depth (m)	var	$\frac{\partial p\text{CO}_2}{\partial\text{var}} \pm \text{SE}$	$\frac{\partial p\text{CO}_2}{\partial\text{var}} \frac{d\text{var}}{dt} \pm \text{RMSE}$	Contribution (%)	$\frac{dp\text{CO}_2}{dt} \pm \text{RMSE}$
1	T ($^{\circ}\text{C}$)	16.49 ± 0.05	1.19 ± 0.36	37	3.23 ± 0.57
	S	$10.14 \pm <0.01$	-0.02 ± 0.05	-1	
	A_T ($\mu\text{mol kg}^{-1}$)	-1.478 ± 0.005	-3.08 ± 0.28	-95	
	C_T ($\mu\text{mol kg}^{-1}$)	1.735 ± 0.006	5.14 ± 0.35	159	
50	T ($^{\circ}\text{C}$)	15.55 ± 0.03	1.37 ± 0.29	48	2.84 ± 0.49
	S	$9.355 \pm <0.001$	0.06 ± 0.02	2	
	A_T ($\mu\text{mol kg}^{-1}$)	-1.327 ± 0.002	-2.11 ± 0.20	-74	
	C_T ($\mu\text{mol kg}^{-1}$)	1.629 ± 0.005	3.52 ± 0.34	124	

4 Discussion

High-resolution time series are necessary to document coastal ocean acidification. Based on weekly measurements from 2007 through 2015, near-surface pH_T decreased by 0.0028 ± 0.0003 units yr^{-1} at Point B in the NW coastal Mediterranean Sea. Temperature increased by 0.072 ± 0.022 $^{\circ}\text{C yr}^{-1}$. In addition, A_T increased by 2.08 ± 0.19 $\mu\text{mol kg}^{-1} \text{yr}^{-1}$, a change that is unrelated to direct effects of CO_2 absorption by seawater. In less than a decade, the total change in pH at Point B (ΔpH_T : 0.0252, Table 1) was of the same magnitude as the diel pH variability (ΔpH_T : 0.01–0.05) and one-fifth of the magnitude of the seasonal pH cycle (ΔpH_T : 0.12) found at this site.

We can identify the drivers of ocean acidification at Point B using the deconvolution of pH and $p\text{CO}_2$ trends and by assuming that the increase in A_T was due to increases in its carbon constituents, bicarbonate (HCO_3^-) and carbonate (CO_3^{2-}) ions. This assumption is based on the fact that (1) HCO_3^- and CO_3^{2-} ions together make up 96% of A_T at Point B, (2) increases in both HCO_3^- and CO_3^{2-} would contribute to A_T and C_T and thereby explain the observed synchronicity of monthly trends in A_T and C_T (Fig. 5). Us-

ing the $p\text{CO}_2$ deconvolution, we can then sum the contributions to $p\text{CO}_2$ from A_T (-3.08 $\mu\text{atm } p\text{CO}_2 \text{ yr}^{-1}$) and C_T (5.14 $\mu\text{atm } p\text{CO}_2 \text{ yr}^{-1}$) to determine the proportional contribution of ΔC_T to $\Delta p\text{CO}_2$ that is unrelated to changes in C_T brought on by A_T . This remaining 2.06 $\mu\text{atm } p\text{CO}_2 \text{ yr}^{-1}$ increase due to increasing C_T closely matched the magnitude of atmospheric CO_2 increase during the observation period (2.08 ± 0.01 ppm yr^{-1} at Plateau Rosa, Italy). The closeness of these numbers does not imply causation. However, given that surface waters at Point B exhibit a near-zero air–sea CO_2 flux over annual cycles (De Carlo et al., 2013), the evidence supports the conclusion that the ocean acidification trend at Point B closely follows the rate of atmospheric CO_2 increase. The influence of atmospheric CO_2 can also be observed in the monthly changes in C_T . Monthly C_T trends are positive and statistically significant over more months than A_T trends (8 vs. 3 months), which are more seasonally restricted (Fig. 5).

Applying this simple model to pH_T , assuming changes in C_T are due to both increasing A_T (a neutralizing effect) and atmospheric CO_2 forcing, the surface ocean acidification trend at Point B can be attributed primarily to atmospheric CO_2 forcing (59%) and secondarily to warming (41%).




Figure 3. Time series observations (a–g) and anomaly trends (h–n) for temperature, salinity, and seawater carbonate chemistry at Point B, 50 m. Regression slopes are drawn \pm SE (standard error, in grey) and noted with a star for significance at $\alpha = 0.05$. Variable abbreviations are the same as in Table 2.

Using this same approach for observations at 50 m, the C_T contribution to $p\text{CO}_2$ trends that was unrelated to A_T change was $1.41 \mu\text{atm } p\text{CO}_2 \text{ yr}^{-1}$, which is 68 % of the rate of atmospheric CO_2 increase, in contrast to 99 % at 1 m. Changes in pH_T can then be attributed to atmospheric CO_2 forcing (42 %), warming (54 %), and increasing salinity (3 %). Due to slightly enhanced warming and reduced CO_2 penetration observed at 50 m, warming had a greater effect on the ocean acidification trend than atmospheric CO_2 forcing at depth compared to the surface.

The acidification rate at Point B (-0.0028 units $\text{pH}_T \text{ yr}^{-1}$) is larger than those reported for other ocean time series sites (-0.0026 to -0.0013 units $\text{pH} \text{ yr}^{-1}$, Bates et al., 2014), likely due to differences in warming rates. The observed warming from 2007 through 2015 (0.72 to $0.88 \pm 0.2 \text{ }^\circ\text{C decade}^{-1}$) is extremely rapid relative to global trends in the upper 75 m from 1971 to 2010 ($0.11 \text{ }^\circ\text{C decade}^{-1}$, Rhein et al., 2013). The coastal region of

Point B has warmed steadily since 1980 and with periods of rapid warming (Parravicini et al., 2015). Exacerbated warming may be related to the positive phases of the Atlantic Multidecadal Oscillation (AMO) and North Atlantic Oscillation (NAO), both of which are associated with episodic warming of the Mediterranean Sea (Lejeune et al., 2010). The AMO has been positive since the 1990s¹ and positive NAO phases were prevalent during the second half of our study².

Nearest to Point B, the acidification rate at DYFAMED, an open-sea site about 50 km offshore from Point B (Fig. 1), was estimated at -0.003 ± 0.001 units $\text{pH}_{\text{SW}} \text{ yr}^{-1}$ from 1995 to 2011 (Marcellin Yao et al., 2016). At DYFAMED, warming contributed approximately 30 % to the acidification rate and the remaining 70 % was attributed to anthropogenic CO_2 (Marcellin Yao et al., 2016). While the uncertainty for DY-

¹<http://www.cgd.ucar.edu/cas/catalog/climind/AMO.html>

²http://www.cpc.ncep.noaa.gov/products/precip/CWlink/pna/month_ao_index.shtml




Figure 4. Monthly distribution of seawater carbonate chemistry at Point B, 1 m, using a combination of a violin plot showing the relative frequency of the observations (shaded blue area) and a box plot showing the median, first, and third quartiles, as well as outliers (blue).

FAMED pH data is large, the trends are comparable to those observed at Point B. However, A_T at the DYFAMED did not change significantly from 2007 through 2014 ($F_{1,51} = 3.204$, $P = 0.0794$, $R^2 = 0.08$; data from Coppola et al., 2016). This probably indicates that the processes driving changes in A_T at Point B are unique to the coastal environment.




Figure 5. Monthly trends of total alkalinity (A_T , blue) and dissolved inorganic carbon (C_T , green) for the period 2007–2015. Error bars are \pm SE of the slope estimate and significance is noted (*) at $\alpha = 0.05$.




Figure 6. Salinity and total alkalinity (A_T) relationships at Point B for the period 2007–2015, by year, at 1 m. Data points are colored by month.

Similar changes in coastal carbonate chemistry were observed elsewhere in the near-shore Mediterranean Sea. In the northern Adriatic Sea, Luchetta et al. (2010) determined an acidification rate of -0.0025 units $\text{pH}_T \text{ yr}^{-1}$ and an increase in A_T of $2.98 \mu\text{mol kg}^{-1} \text{ yr}^{-1}$ at depths shallower than 75 m, by comparing cruise data between the winters of 1983 and 2008. Point B and Adriatic Sea observations are independent but reflect changes in seawater chemistry that may be occurring in more coastal regions of the Mediterranean. Additional time series would help resolve the spatial extent of these observations.

While the trends in atmospheric CO_2 forcing and temperature account for the ocean acidification trend at Point B, the increase in A_T and C_T beyond what can be attributed to changes in atmospheric CO_2 was unexpected. The fastest




Figure 7. Time series pH_T (a) and temperature (b) from SeaFET pH sensor deployments at EOL, 2 m. Discrete calibration samples are noted in green, and grey vertical lines bracket deployment periods (a). Calibration sample offsets from processed pH time series are shown in (c). Violin and box plots (see Fig. 4) show diel pH range by month (d), and an example of this pH variability is shown for May 2015 (e).

increases in A_T and C_T occurred from May through July (Fig. 5), when the thermal stratification settled. In the NW Mediterranean, the main processes governing seasonal variability in A_T are evaporation, which increases A_T , in summer (i.e., June through September at Point B) and, to a lesser extent, phytoplankton uptake of nitrate (NO_3^-) and phosphate (PO_4^{3-}), which increase A_T from January through

March (Cossarini et al., 2015). During the transition of these processes, salinity decreases to a minimum in May, reflecting freshwater input that dilutes A_T to minimum values at the start of summer. For C_T , peak values occur in winter when the water column is fully mixed. For reference, at DYFAMED, mixing occurs down to more than 2000 m depth and C_T is up to $100 \mu\text{mol kg}^{-1}$ higher in deep waters (Copin-Montégut and Bégovic, 2002). Notably, monthly trends of C_T at Point B were not statistically significant from November through January for the period 2007–2015. Following winter, C_T declines due to a combination of phytoplankton bloom carbon uptake and freshwater dilution (assuming river $C_T < \text{seawater } C_T$), until the onset of summer stratification. Summer warming leads to $p\text{CO}_2$ outgassing (De Carlo et al., 2013), thereby further decreasing C_T . In addition, the increases in A_T and C_T from 2007 through 2015 were more pronounced at 1 compared to 50 m. Thus, the processes driving A_T and C_T trends are stronger at the surface and affect carbonate chemistry primarily during the spring–summer transition from May through July.

Identifying the processes causing an increase in A_T and C_T beyond what can be attributed to changes in atmospheric CO_2 requires some speculation. Some biogeochemical processes can be ruled out as drivers. For example, changes in benthic processes are very unlikely. Reduced calcium carbonate (CaCO_3) precipitation rates would increase A_T but would increase C_T . Even so, the dominant ecosystem in the Bay of Villefranche-sur-Mer is seagrass meadows, which harbor relatively few calcifying organisms. Dissolution of calcium carbonate sediment would contribute to A_T and C_T increase in the water column. However, as the water was supersaturated with respect to both aragonite and calcite, this could only be mediated by biological processes. Carbonate dissolution following CO_2 production via oxic degradation of organic matter releases A_T and C_T in a 1 : 1 ratio (Moulin et al., 1985). Likewise, anaerobic remineralization produces alkalinity (Cai et al., 2011). In the sediment of the Bay of Villefranche, sulfate reduction coupled with precipitation of sulfide minerals is the dominant anoxic mineralization pathway (Gaillard et al., 1989). An increase in these processes would explain the observed increase in A_T and C_T , but as trends were slower at 50 compared to 1 m, this would suggest the dominance of a process taking place in surface waters.

In the upper water column at Point B, changes in biological processes are unlikely to explain the observed trends in A_T and C_T . For example, the concentration of chlorophyll *a* (a proxy of primary production) has decreased since 1995, nutrients have increased, and phytoplankton blooms have shifted towards earlier dates in the year (Irisson et al., 2012). While a decrease in net primary production could drive C_T trends, the observed increase in NO_3^- and PO_4^{3-} would cause a small decrease in A_T (Wolf-Gladrow et al., 2007). Stimulated community respiration could result from warming waters but enhanced remineralization would cause a decrease in A_T (Wolf-Gladrow et al., 2007).

The lack of salinity change at the surface excludes additional processes as drivers of the A_T and C_T increase at Point B. For example, increased summertime evaporation (concentration effect) and reduced freshwater input (decreased dilution effect) would both be expected to cause an increase in salinity, which was not observed. Increased input of eastern Mediterranean Sea waters could increase A_T , but this is unlikely as this water mass flows much deeper than Point B (Millot and Taupier-Letage, 2005).

Instead, the observed changes in A_T and C_T could be due to increased limestone weathering, which would increase the input of A_T from land to the sea via rivers and groundwater. Rivers contribute both A_T and C_T to the Mediterranean Sea (Copin-Montégut, 1993; Tamše et al., 2015). River A_T originates from erosion and is correlated with bedrock composition (e.g., McGrath et al., 2016). Positive trends in river A_T have been documented in North America and occur via a number of processes including (1) the interplay of rainfall and land use (Raymond and Cole, 2003), (2) anthropogenic limestone addition used to enhance agricultural soil pH (Oh and Raymond, 2006; Stets et al., 2014) and freshwater pH (Clair and Hindar, 2005), and (3) potentially indirect effects of anthropogenic CO_2 on groundwater CO_2 acidification and weathering (Macpherson et al., 2008). These and other processes were hypothesized to have driven A_T changes in the Baltic Sea (Müller et al., 2016). There, an increase in A_T of $3.4 \mu\text{mol kg}^{-1} \text{yr}^{-1}$ was observed from 1995 to 2014 (mean salinity = 7). In contrast to Point B, the increase in Baltic Sea A_T was not noticeable at salinity > 30 (Müller et al., 2016).

Given the above speculations, the simplest plausible mechanisms causing the unexpected A_T and C_T trends would be through increasing A_T of the freshwater end-member of Point B. Local precipitation, however, did not have an influential effect and was not correlated with salinity or A_T (Fig. S1 in the Supplement). While submarine groundwater springs can be a significant source of nutrients, A_T , and C_T for the ocean (Cai et al., 2003; Slomp and Van Cappellen, 2004), carbonate chemistry contributions of local submarine springs are currently unknown (Gilli, 1995). Signatures of limestone erosion can be observed in A_T of nearby rivers (Var, Paillon, and Roya) but detailed time series are not available. Likewise, riverine influence at Point B has not been quantified. If river runoff exerts a dominant control on Point B carbonate chemistry, there is a lag effect, as freshwater influence peaked in May but A_T and C_T increased fastest from May through July. Consequently, this hypothesis needs further investigation. Until the source of A_T increase is properly identified, use of this observation in modeling should be implemented with caution.

5 Conclusion

Predictions of coastal ocean acidification remain challenging due the complexity of biogeochemical processes occurring

at the ocean–land boundary. At the Point B coastal oceanographic station in the NW Mediterranean Sea, surface ocean acidification was due to atmospheric CO_2 forcing and rapid warming over the observation period 2007–2015. However, additional trends in A_T and C_T were observed and remain unexplained, but these trends could relate to riverine and groundwater input. The influence of coastal boundary processes influencing seawater A_T and C_T presents a potentially major difference between coastal and offshore changes in ocean chemistry. This study highlights the importance of considering other anthropogenic influences in the greater land–sea region that may contribute to coastal biogeochemical cycles (sensu Duarte et al., 2013) and alter projections of anthropogenic change in near-shore waters.

Data availability. Time series data from Point B are available at PANGAEA® (doi:10.1594/PANGAEA.727120)

The Supplement related to this article is available online at doi:10.5194/os-13-411-2017-supplement.

Author contributions. Jean-Pierre Gattuso initiated the study, Laure Mousseau supervised data collection, Samir Alliouane performed SeaFET deployments and calibration, Jean-Pierre Gattuso and Lydia Kapsenberg designed and Jean-Pierre Gattuso conducted statistical analyses, and Lydia Kapsenberg prepared the paper with contributions from all authors.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. Thanks are due to the Service d'Observation Rade de Villefranche (SO-Rade) of the Observatoire Océanologique and the Service d'Observation en Milieu Littoral (SOMLIT/CNRS-INSU) for their kind permission to use the Point B data. Discrete samples were analyzed for C_T and A_T by the Service National d'Analyse des Paramètres Océaniques du CO_2 . The authors thank Jean-Yves Carval, Anne-Marie Corre, Maïa Durozier, Ornella Passafiume and Frank Petit for sampling assistance, Steeve Comeau and Alice Webb for help with data analysis, and Bernard Gentili for producing Fig. 1. Atmospheric CO_2 data from Plateau Rosa were collected by Ricerca sul Sistema Energetico (RSE S.p.A.); we are grateful for their contribution. We acknowledge L. Coppola for providing DYFAMED data (Coppola et al., 2016), Météo-France for supplying the meteorological data, and the HyMeX database teams (ESPRI/IPSL and SEDOO/Observatoire Midi-Pyrenees) for their help in accessing them. The Agence de l'Eau Rhône-Méditerranée-Corse kindly provided data on the chemistry of local rivers. Alexandre Dano, Gilles Dandec, and Dominique Chassagne provided the high-resolution bathymetric data for the volume estimate of the bay. We are grateful for helpful comments from Nicolas Metzl on the manuscript and those from two anonymous reviewers. This work is a contribution to the European Project on

Ocean Acidification (EPOCA; contract no. 211384) and the Med-Sea project (contract no. 265103), which received funding from the European Community's Seventh Framework Programme, and to the United States National Science Foundation Ocean Sciences Postdoctoral Research Fellowship (OCE-1521597), awarded to LK.

Edited by: M. Hoppema

Reviewed by: two anonymous referees

References

- Álvarez, M., Sanleón-Bartolomé, H., Tanhua, T., Mintrop, L., Luchetta, A., Cantoni, C., Schroeder, K., and Civitarese, G.: The CO₂ system in the Mediterranean Sea: a basin wide perspective, *Ocean Sci.*, 10, 69–92, doi:10.5194/os-10-69-2014, 2014.
- Aminot, A. and Kérouel, R.: Dosage automatique des nutriments dans les eaux marines: méthodes d'analyse en milieu marin, edited by: Aminot, A. and Kérouel, R., Ifremer, 188 pp., 2007.
- Barbier, E. B., Hacker, S. D., Kennedy, C., Koch, E. W., Stier, A. C., and Silliman, B. R.: The value of estuarine and coastal ecosystem services, *Ecol. Monogr.*, 81, 169–193, doi:10.1890/10-1510.1, 2011.
- Bates, N. R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., González-Dávila, M., Lorenzoni, L., Muller-Karger, F., Olafsson, J., and Santana-Casiano, J. M.: A time-series view of changing ocean chemistry due to ocean uptake of anthropogenic CO₂ and ocean acidification, *Oceanography*, 27, 126–141, 2014.
- Borges, A. V. and Gypens, N.: Carbonate chemistry in the coastal zone responds more strongly to eutrophication than ocean acidification, *Limnol. Oceanogr.*, 55, 346–353, doi:10.4319/lo.2010.55.1.0346, 2010.
- Bresnahan, P. J., Martz, T. R., Takeshita, Y., Johnson, K. S., and LaShomb, M.: Best practices for autonomous measurement of seawater pH with the Honeywell Durafet, *Methods Oceanogr.*, 9, 44–60, 2014.
- Cai, W.-J., Wang, Y., Krest, J., and Moore, W. S.: The geochemistry of dissolved inorganic carbon in a surficial groundwater aquifer in North Inlet, South Carolina, and the carbon fluxes to the coastal ocean, *Geochim. Cosmochim. Ac.*, 67, 631–639, doi:10.1016/S0016-7037(02)01167-5, 2003.
- Cai, W.-J., Hu, X., Huang, W.-J., Murrell, M. C., Lehrter, J. C., Lohrenz, S. E., Chou, W.-C., Zhai, W., Hollibaugh, J. T., Wang, Y., Zhao, P., Guo, X., Gundersen, K., Dai, M., and Gong, G.-C.: Acidification of subsurface coastal waters enhanced by eutrophication, *Nat. Geosci.*, 4, 766–770, doi:10.1038/ngeo1297, 2011.
- Clair, T. A. and Hindar, A.: Liming for the mitigation of acid rain effects in freshwaters: a review of recent results, *Environ. Rev.*, 13, 91–128, doi:10.1139/a05-009, 2005.
- Copin-Montégut, C.: Alkalinity and carbon budgets in the Mediterranean Sea, *Global Biogeochem. Cy.*, 7, 915–925, doi:10.1029/93GB01826, 1993.
- Copin-Montégut, C. and Bégovic, M.: Distributions of carbonate properties and oxygen along the water column (0–2000 m) in the central part of the NW Mediterranean Sea (Dyfamed site): influence of winter vertical mixing on air–sea CO₂ and O₂ exchanges, *Deep-Sea Res. Pt. II*, 49, 2049–2066, doi:10.1016/S0967-0645(02)00027-9, 2002.
- Coppola, L., Diamond Riquier, E., and Carval, T.: Dyfamed observatory data, SEANOE, doi:10.17882/43749, 2016.
- Cossarini, G., Lazzari, P., and Solidoro, C.: Spatiotemporal variability of alkalinity in the Mediterranean Sea, *Biogeosciences*, 12, 1647–1658, doi:10.5194/bg-12-1647-2015, 2015.
- Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R. V., Paruelo, J., Raskin, R. G., Sutton, P., and van den Belt, M.: The value of the world's ecosystem services and natural capital, *Nature*, 387, 253–260, 1997.
- De Carlo, E. H., Mousseau, L., Passafiume, O., Drupp, P. S., and Gattuso, J.-P.: Carbonate chemistry and air–sea CO₂ flux in a NW Mediterranean bay over a four-year period: 2007–2011, *Aquat. Geochem.*, 19, 399–442, doi:10.1007/s10498-013-9217-4, 2013.
- Dickson, A.: The carbon dioxide system in seawater: equilibrium chemistry and measurements, in: Guide to best practices for ocean acidification research and data reporting, edited by: Fabry, V. J., Hansson, L., and Gattuso, J.-P., Publications Office of the European Union, Luxembourg, 17–40, 2010.
- Dickson, A. G.: Standard potential of the reaction: AgCl(s) + 1/2 H₂(g) = Ag(s) + HCl(aq), and the standard acidity constant of the ion HSO₄[−] in synthetic sea water from 273.15 to 318.15 K, *J. Chem. Thermodyn.*, 22, 113–127, doi:10.1016/0021-9614(90)90074-Z, 1990.
- Dickson, A. G. and Riley, J. P.: The effect of analytical error on the evaluation of the components of the aquatic carbon-dioxide system, *Mar. Chem.*, 6, 77–85, doi:10.1016/0304-4203(78)90008-7, 1978.
- Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practices for ocean CO₂ measurements, PICES Special Publication, 3, 191 pp., 2007.
- DOE: Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, 1994.
- Doney, S. C., Fabry, V. J., Feely, R. A., and Kleypas, J. A.: Ocean acidification: the other CO₂ problem, *Annu. Rev. Mar. Sci.*, 1, 169–192, doi:10.1146/annurev.marine.010908.163834, 2009.
- Duarte, C. M., Hendriks, I. E., Moore, T. S., Olsen, Y. S., Steckbauer, A., Ramajo, L., Carstensen, J., Trotter, J. A., and McCulloch, M.: Is ocean acidification an open-ocean syndrome? Understanding anthropogenic impacts on seawater pH, *Estuar. Coast.*, 36, 221–236, doi:10.1007/s12237-013-9594-3, 2013.
- Edmond, J. M.: High precision determination of titration alkalinity and total carbon dioxide content of sea water by potentiometric titration, *Deep-Sea Res.*, 17, 737–750, doi:10.1016/0011-7471(70)90038-0, 1970.
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for upwelling of corrosive “acidified” water onto the continental shelf, *Science*, 320, 1490–1492, doi:10.1126/science.1155676, 2008.
- Feely, R. A., Alin, S. R., Newton, J., Sabine, C. L., Warner, M., Devol, A., Krembs, C., and Maloy, C.: The combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an urbanized estuary, *Estuar. Coast. Shelf S.*, 88, 442–449, doi:10.1016/j.ecss.2010.05.004, 2010.
- Flecha, S., Pérez, F. F., García-Lafuente, J., Sammartino, S., Ríos, A. F., and Huertas, I. E.: Trends of pH decrease in the Mediter-

- anean Sea through high frequency observational data: indication of ocean acidification in the basin, *Sci. Rep.*, 5, 16770, doi:10.1038/srep16770, 2015.
- Gaillard, J.-F., Pauwels, H., and Michard, G.: Chemical diagenesis in coastal marine sediments, *Oceanol. Acta*, 12, 175–187, 1989.
- García-Ibáñez, M. I., Zunino, P., Fröb, F., Carracedo, L. I., Ríos, A. F., Mercier, H., Olsen, A., and Pérez, F. F.: Ocean acidification in the subpolar North Atlantic: rates and mechanisms controlling pH changes, *Biogeosciences*, 13, 3701–3715, doi:10.5194/bg-13-3701-2016, 2016.
- Gattuso, J. P. and Hansson, L.: Ocean acidification, Oxford University Press, Oxford, 2011.
- Gattuso, J.-P., Epitalon, J.-M., and Lavigne, H.: seacarb: Seawater Carbonate Chemistry. R package version 3.1.1 <https://cran.r-project.org/package=seacarb> (last access: 3 April 2017), 2016.
- Gattuso, J.-P., Alliouane, S., and Mousseau, L.: Seawater carbonate chemistry in the Bay of Villefranche, Point B (France), January 2007 – December 2015, doi:10.1594/PANGAEA.727120, 2014.
- Gilli, E.: Etude des sources karstiques sous-marines et littorales des Alpes Maritimes entre Menton et Nice, 41, http://www.side.developpement-durable.gouv.fr/EXPLOITATION/DEFAULT/doc/IFD/I_IFD_REFDOC_0113893 (last access: 12 May 2017), 1995.
- Halpern, B. S., Walbridge, S., Selkoe, K. A., Kappel, C. V., Micheli, F., D'Agrosa, C., Bruno, J. F., Casey, K. S., Ebert, C., and Fox, H. E.: A global map of human impact on marine ecosystems, *Science*, 319, 948–952, 2008.
- Hofmann, G. E., Smith, J. E., Johnson, K. S., Send, U., Levin, L. A., Micheli, F., Paytan, A., Price, N. N., Peterson, B., Takeshita, Y., Matson, P. G., Crook, E. D., Kroeker, K. J., Gambi, M. C., Rivest, E. B., Frieder, C. A., Yu, P. C., and Martz, T. R.: High-frequency dynamics of ocean pH: a multi-ecosystem comparison, *PLoS One*, 6, e28983, doi:10.1371/journal.pone.0028983, 2011.
- Howes, E. L., Stemann, L., Assailly, C., Irisson, J. O., Dima, M., Bijma, J., and Gattuso, J. P.: Pteropod time series from the North Western Mediterranean (1967–2003): impacts of pH and climate variability, *Mar. Ecol.-Prog. Ser.*, 531, 193–206, 2015.
- Ingrassio, G., Giani, M., Comici, C., Kralj, M., Piacentino, S., De Vittor, C., and Del Negro, P.: Drivers of the carbonate system seasonal variations in a Mediterranean gulf, *Estuar. Coast. Shelf S.*, 168, 58–70, doi:10.1016/j.ecss.2015.11.001, 2016.
- Irisson, J.-O., Webb, A., Passafiume, O., and Mousseau, L.: Detecting hydrologic variations in a long term monitoring time series, *Europole Mer Gordon-like conference “Time-series analysis in marine science and application for industry”*, Brest, France, 17–21 September 2012.
- Jiang, Z.-P., Tyrrell, T., Hydes, D. J., Dai, M., and Hartman, S. E.: Variability of alkalinity and the alkalinity-salinity relationship in the tropical and subtropical surface ocean, *Global Biogeochem. Cy.*, 28, 729–742, doi:10.1002/2013GB004678, 2014.
- Kapsenberg, L. and Hofmann, G. E.: Ocean pH time-series and drivers of variability along the northern Channel Islands, California, USA, *Limnol. Oceanogr.*, 61, 953–968, doi:10.1002/lno.10264, 2016.
- Kapsenberg, L., Kelley, A. L., Shaw, E. C., Martz, T. R., and Hofmann, G. E.: Near-shore Antarctic pH variability has implications for biological adaptation to ocean acidification, *Sci. Rep.*, 5, 9638, doi:10.1038/srep09638, 2015.
- Krasakopoulou, E., Souvermezoglou, E., and Goyet, C.: Anthropogenic CO₂ fluxes in the Otranto Strait (E. Mediterranean) in February 1995, *Deep-Sea Res. Pt. I*, 58, 1103–1114, doi:10.1016/j.dsr.2011.08.008, 2011.
- Lacoue-Labarthe, T., Nunes, P. A. L. D., Ziveri, P., Cinar, M., Gazeau, F., Hall-Spencer, J. M., Hilmi, N., Moschella, P., Safa, A., Sauzade, D., and Turley, C.: Impacts of ocean acidification in a warming Mediterranean Sea: An overview, *Regional Studies in Marine Science*, 5, 1–11, doi:10.1016/j.rsma.2015.12.005, 2016.
- Lee, K., Kim, T.-W., Byrne, R. H., Millero, F. J., Feely, R. A., and Liu, Y.-M.: The universal ratio of boron to chlorinity for the North Pacific and North Atlantic oceans, *Geochim. Cosmochim. Ac.*, 74, 1801–1811, doi:10.1016/j.gca.2009.12.027, 2010.
- Lee, K., Sabine, C. L., Tanhua, T., Kim, T.-W., Feely, R. A., and Kim, H.-C.: Roles of marginal seas in absorbing and storing fossil fuel CO₂, *Energ. Environ. Sci.*, 4, 1133–1146, doi:10.1039/C0EE00663G, 2011.
- Lejeune, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C. F., and Pérez, T.: Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea, *Trends Ecol. Evol.*, 25, 250–260, doi:10.1016/j.tree.2009.10.009, 2010.
- Le Quéré, C., Andrew, R. M., Canadell, J. G., Sitch, S., Korsbakken, J. I., Peters, G. P., Manning, A. C., Boden, T. A., Tans, P. P., Houghton, R. A., Keeling, R. F., Alin, S., Andrews, O. D., Anthoni, P., Barbero, L., Bopp, L., Chevallier, F., Chini, L. P., Ciais, P., Currie, K., Delire, C., Doney, S. C., Friedlingstein, P., Gkritzalis, T., Harris, I., Hauck, J., Haverd, V., Hoppema, M., Klein Goldewijk, K., Jain, A. K., Kato, E., Körtzinger, A., Landschützer, P., Lefèvre, N., Lenton, A., Lienert, S., Lombardozzi, D., Melton, J. R., Metzl, N., Millero, F., Monteiro, P. M. S., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-I., O'Brien, K., Olsen, A., Omar, A. M., Ono, T., Pierrot, D., Poulter, B., Rödenbeck, C., Salisbury, J., Schuster, U., Schwinger, J., Séférian, R., Skjelvan, I., Stocker, B. D., Sutton, A. J., Takahashi, T., Tian, H., Tilbrook, B., van der Laan-Luijkx, I. T., van der Werf, G. R., Viovy, N., Walker, A. P., Wiltshire, A. J., and Zaehle, S.: Global Carbon Budget 2016, *Earth Syst. Sci. Data*, 8, 605–649, doi:10.5194/essd-8-605-2016, 2016.
- Luchetta, A., Cantoni, C., and Catalano, G.: New observations of CO₂-induced acidification in the northern Adriatic Sea over the last quarter century, *Chem. Ecol.*, 26, 1–17, doi:10.1080/02757541003627688, 2010.
- Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for *K*₁ and *K*₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium, *Mar. Chem.*, 70, 105–119, doi:10.1016/S0304-4203(00)00022-0, 2000.
- Macpherson, G. L., Roberts, J. A., Blair, J. M., Townsend, M. A., Fowle, D. A., and Beisner, K. R.: Increasing shallow groundwater CO₂ and limestone weathering, *Konza Prairie, USA, Geochim. Cosmochim. Ac.*, 72, 5581–5599, doi:10.1016/j.gca.2008.09.004, 2008.
- Marcellin Yao, K., Marcou, O., Goyet, C., Guglielmi, V., Touratier, F., and Savy, J.-P.: Time variability of the north-western Mediterranean Sea pH over 1995–2011, *Mar. Environ. Res.*, 116, 51–60, doi:10.1016/j.marenvres.2016.02.016, 2016.
- McGrath, T., McGovern, E., Cave, R. R., and Kivimäe, C.: The inorganic carbon chemistry in coastal and shelf waters around Ire-

- land, *Estuar. Coast.*, 39, 27–39, doi:10.1007/s12237-015-9950-6, 2016.
- Meier, K. J. S., Beaufort, L., Heussner, S., and Ziveri, P.: The role of ocean acidification in *Emiliania huxleyi* coccolith thinning in the Mediterranean Sea, *Biogeosciences*, 11, 2857–2869, doi:10.5194/bg-11-2857-2014, 2014.
- Millot, C. and Taupier-Letage, I.: Circulation in the Mediterranean Sea, in: *The Mediterranean Sea*, edited by: Saliot, A., Springer, Berlin, Heidelberg, 29–66, 2005.
- Moulin, E., Jordens, A., and Wollast, R.: Influence of the aerobic bacterial respiration on the early dissolution of carbonates in coastal sediments, in: *Progress in Belgian Oceanographic Research: Proceedings of a Symposium Held at the Palace of Academies Brussels*, edited by: Van Grieken, R. and Wollast, R., Brussels, 196–208, 1985.
- Müller, J. D., Schneider, B., and Rehder, G.: Long-term alkalinity trends in the Baltic Sea and their implications for CO₂-induced acidification, *Limnol. Oceanogr.*, 61, 1984–2002, doi:10.1002/lno.10349, 2016.
- Oh, N.-H. and Raymond, P. A.: Contribution of agricultural liming to riverine bicarbonate export and CO₂ sequestration in the Ohio River basin, *Global Biogeochem. Cy.*, 20, GB3012, doi:10.1029/2005GB002565, 2006.
- Omstedt, A., Edman, M., Claremar, B., and Rutgersson, A.: Modelling the contributions to marine acidification from deposited SO_x, NO_x, and NH_x in the Baltic Sea: Past and present situations, *Cont. Shelf Res.*, 111, 234–249, doi:10.1016/j.csr.2015.08.024, 2015.
- Palmiéri, J., Orr, J. C., Dutay, J.-C., Béranger, K., Schneider, A., Beuvier, J., and Somot, S.: Simulated anthropogenic CO₂ storage and acidification of the Mediterranean Sea, *Biogeosciences*, 12, 781–802, doi:10.5194/bg-12-781-2015, 2015.
- Parravicini, V., Mangialajo, L., Mousseau, L., Peirano, A., Morri, C., Montefalcone, M., Francour, P., Kulbicki, M., and Bianchi, C. N.: Climate change and warm-water species at the north-western boundary of the Mediterranean Sea, *Mar. Ecol.*, 36, 897–909, doi:10.1111/maec.12277, 2015.
- Perez, F. F. and Fraga, F.: The pH measurements in seawater on the NBS scale, *Mar. Chem.*, 21, 315–327, doi:10.1016/0304-4203(87)90054-5, 1987.
- Pörtner, H.-O., Karl, D., Boyd, P. W., Cheung, W., Lluch-Cota, S. E., Nojiri, Y., Schmidt, D. N., and Zavialov, P.: Ocean systems, in: *Climate Change 2014, Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Field, C. B., Barros, V. R., Dokken, D. J., Mach, K. J., Mastrandrea, M. D., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., Genova, R. C., Girma, B., Kissel, E. S., Levy, A. N., MacCracken, S., Mastrandrea, P. R., and White, L. L., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 411–484, 2014.
- Provoost, P., van Heuven, S., Soetaert, K., Laane, R. W. P. M., and Middelburg, J. J.: Seasonal and long-term changes in pH in the Dutch coastal zone, *Biogeosciences*, 7, 3869–3878, doi:10.5194/bg-7-3869-2010, 2010.
- Raymond, P. A. and Cole, J. J.: Increase in the export of alkalinity from North America's largest river, *Science*, 301, 88–91, 2003.
- R Core Team: R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria, <https://www.r-project.org/> (last access: 3 April 2017), 2016.
- Rhein, M., Rintoul, S. R., Aoki, S., Campos, E., Chambers, D., Feely, R. A., Gulev, S., Johnson, G. C., Josey, S. A., Kostianoy, Mauritzen, C., Roemmich, D., Talley, L. D., and Wang, F.: Observations: Ocean, in: *Climate Change 2013, The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Schneider, A., Wallace, D. W. R., and Körtzinger, A.: Alkalinity of the Mediterranean Sea, *Geophys. Res. Lett.*, 34, L15608, doi:10.1029/2006GL028842, 2007.
- Schneider, A., Tanhua, T., Körtzinger, A., and Wallace, D. W. R.: High anthropogenic carbon content in the eastern Mediterranean, *J. Geophys. Res.*, 115, C12050, doi:10.1029/2010JC006171, 2010.
- Slomp, C. P. and Van Cappellen, P.: Nutrient inputs to the coastal ocean through submarine groundwater discharge: controls and potential impact, *J. Hydrol.*, 295, 64–86, doi:10.1016/j.jhydrol.2004.02.018, 2004.
- Stets, E. G., Kelly, V. J., and Crawford, C. G.: Long-term trends in alkalinity in large rivers of the conterminous US in relation to acidification, agriculture, and hydrologic modification, *Sci. Total Environ.*, 488–489, 280–289, doi:10.1016/j.scitotenv.2014.04.054, 2014.
- Tamše, S., Ogrinc, N., Walter, L. M., Turk, D., and Faganeli, J.: River sources of dissolved inorganic carbon in the Gulf of Trieste (N Adriatic): stable carbon isotope evidence, *Estuar. Coast.*, 38, 151–164, doi:10.1007/s12237-014-9812-7, 2015.
- Tanhua, T., Bates, N. R., and Körtzinger, A.: The marine carbon cycle and ocean anthropogenic CO₂ inventories, in: *Ocean Circulation and Climate: A 21st Century Perspective, 2nd Ed.*, edited by: Siedler, G., Griffies, S., Gould, J., and Church, J., 103, Academic Press, 787–816, 2013.
- The MerMEX Group: Durrieu de Madron, X., Guieu, C., Sempéré, R., Conan, P., Cossa, D., D'Ortenzio, F., Estournel, C., Gazeau, F., Rabouille, C., Stemmann, L., Bonnet, S., Diaz, F., Koubbi, P., Radakovitch, O., Babin, M., Baklouti, M., Bancon-Montigny, C., Belviso, S., Bensoussan, N., Bonsang, B., Bouloubassi, I., Brunet, C., Cadiou, J. F., Carlotti, F., Chami, M., Charmasson, S., Charrière, B., Dachs, J., Doxaran, D., Dutay, J. C., Elbaz-Poulichet, F., Eléaume, M., Eyrolles, F., Fernandez, C., Fowler, S., Francour, P., Gaertner, J. C., Galzin, R., Gasparini, S., Ghiglione, J. F., Gonzalez, J. L., Goyet, C., Guidi, L., Guizien, K., Heimbürger, L. E., Jacquet, S. H. M., Jeffrey, W. H., Joux, F., Le Hir, P., Leblanc, K., Lefèvre, D., Lejeusne, C., Lemé, R., Loÿe-Pilot, M. D., Mallet, M., Méjanelle, L., Mélin, F., Mellon, C., Mériçot, B., Merle, P. L., Migon, C., Miller, W. L., Mortier, L., Mostajir, B., Mousseau, L., Moutin, T., Para, J., Pérez, T., Petrenko, A., Poggiale, J. C., Prieur, L., Pujopay, M., Pulido, V., Raimbault, P., Rees, A. P., Ridame, C., Rontani, J. F., Ruiz Pino, D., Sicre, M. A., Taillandier, V., Tamburini, C., Tanaka, T., Taupier-Letage, I., Tedetti, M., Testor, P., Thébault, H., Thouvenin, B., Touratier, F., Tronczynski, J., Ulses, C., Van Wambeke, F., Vantrepotte, V., Vaz, S., and Ver-

- ney, R.: Marine ecosystems' responses to climatic and anthropogenic forcings in the Mediterranean, *Prog. Oceanogr.*, 91, 97–166, doi:10.1016/j.pocean.2011.02.003, 2011.
- Touratier, F. and Goyet, C.: Impact of the Eastern Mediterranean Transient on the distribution of anthropogenic CO₂ and first estimate of acidification for the Mediterranean Sea, *Deep-Sea Res. Pt. I*, 58, 1–15, doi:10.1016/j.dsr.2010.10.002, 2011.
- Touratier, F., Goyet, C., Houpert, L., de Madron, X. D., Lefèvre, D., Stabholz, M., and Guglielmi, V.: Role of deep convection on anthropogenic CO₂ sequestration in the Gulf of Lions (north-western Mediterranean Sea), *Deep-Sea Res. Pt. I*, 113, 33–48, doi:10.1016/j.dsr.2016.04.003, 2016.
- Vargas, C. A., Contreras, P. Y., Pérez, C. A., Sobarzo, M., Saldías, G. S., and Salisbury, J.: Influences of riverine and upwelling waters on the coastal carbonate system off Central Chile and their ocean acidification implications, *J. Geophys. Res.-Biogeo.*, 121, 1468–1483, doi:10.1002/2015JG003213, 2016.
- Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A., and Dickson, A. G.: Total alkalinity: The explicit conservative expression and its application to biogeochemical processes, *Mar. Chem.*, 106, 287–300, doi:10.1016/j.marchem.2007.01.006, 2007.
- Wootton, J. T. and Pfister, C. A.: Carbon system measurements and potential climatic drivers at a site of rapidly declining ocean pH, *PLoS One*, 7, e53396, doi:10.1371/journal.pone.0053396, 2012.
- Wootton, J. T., Pfister, C. A., and Forester, J. D.: Dynamic patterns and ecological impacts of declining ocean pH in a high-resolution multi-year dataset, *P. Natl. Acad. Sci. USA*, 105, 18848–18853, 2008.