



HAL
open science

Geographically distinct patterns of reproductive isolation and hybridisation in two sympatric species of the *Jaera albifrons* complex (marine isopods)

Ambre Ribardière, Claire Daguin--thiébaud, Céline Houbin, Jérôme Coudret, Caroline Broudin, Olivier Timsit, Thomas Broquet

► To cite this version:

Ambre Ribardière, Claire Daguin--thiébaud, Céline Houbin, Jérôme Coudret, Caroline Broudin, et al.. Geographically distinct patterns of reproductive isolation and hybridisation in two sympatric species of the *Jaera albifrons* complex (marine isopods). 2017. hal-01534577

HAL Id: hal-01534577

<https://hal.sorbonne-universite.fr/hal-01534577v1>

Preprint submitted on 7 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Geographically distinct patterns of reproductive isolation and hybridisation in two sympatric species
2 of the *Jaera albifrons* complex (marine isopods)

3

4 Ambre Ribardière^{1,2}, Claire Daguin-Thiébaud^{1,2}, Céline Houbin^{2,3}, Jérôme Coudret^{1,2}, Caroline
5 Broudin^{2,3}, Olivier Timsit⁴, Thomas Broquet^{1,2}

6

7 1- Centre National de la Recherche Scientifique, Team Diversity and Connectivity of Coastal Marine
8 Landscapes, Unité Mixte de Recherche 7144, Station Biologique de Roscoff, 29680 Roscoff, France.

9 2- Sorbonne Universités, UPMC Univ Paris 06, Station Biologique de Roscoff, 29680 Roscoff, France.

10 3- Centre National de la Recherche Scientifique, FR 2424, Station Biologique de Roscoff, 29680
11 Roscoff, France.

12 4- Groupe d'Etude des Milieux Estuariens et Littoraux de Normandie, Centre Régional d'Etudes
13 Côtières, 54 rue du Docteur Charcot, 14530 Luc-sur-Mer, France.

14

15 Running title: Isolation and hybridization in *J. albifrons*

16

17 Corresponding author: Ambre Ribardière, ambre.ribardiere@sb-roscoff.fr, tel +33 298 292 544, fax
18 +33 298 292 324

19

20 **Abstract**

21 Sympatric species that in some populations hybridize and in other populations remain reproductively
22 isolated open interesting research possibilities for the study of hybridization and speciation. Here we
23 test for such a situation in two littoral isopods (*Jaera albifrons* and *J. praehirsuta*) that occur in mixed
24 populations and where past morphological descriptions suggested that the two species are generally
25 reproductively isolated except in rare populations where hybridization may be happening. Using field
26 surveys and microsatellite genetic structure analyses in two regions from France (Normandy and
27 Brittany), we confirmed that introgressive hybridization occurs in a subset of mixed *J. albifrons* / *J.*
28 *praehirsuta* populations (region Normandy) where the two species are found in the same habitat
29 (pebbles on the shore). Moreover, we found that introgression in these populations is differential, 21
30 out of 23 microsatellite markers showing little genetic divergence between species (hierarchical
31 analysis of molecular variance $F_{CT}=0.017$) while the remaining two loci were strongly differentiated
32 ($F_{CT}=0.428$). By contrast, *J. albifrons* and *J. praehirsuta* in mixed populations from region Brittany
33 occupied distinct habitats (pebbles and seaweeds, respectively) with little overlap and showed
34 stronger genetic divergence ($F_{CT}=0.132$). In hybridizing populations, the majority of individuals show
35 morphological traits that are characteristic of one or the other species. This raises the question of the
36 forces that act to maintain this polymorphism, noting that hybridizing populations seem to be
37 geographically isolated from potential source parental populations and show no detectable habitat
38 divergence between species.

39

40 **Keywords**

41 Mosaic hybrid zone, isolating barriers, sexual isolation, genetic incompatibilities, introgression,
42 crustaceans

43 **Introduction**

44 Natural hybridization events inform our understanding of isolating barriers between species,
45 the conditions of species coexistence despite hybridization, and the mechanisms of speciation. The
46 archetypal hybrid zone structure is a region of contact between two otherwise allopatric species. In
47 such hybrid zones, flanked on one side by populations of one species and on the other side by the
48 other species, the dynamics of the system is most often driven by a balance between immigration
49 from pure parental populations and selection against hybrids (the tension zone model, Barton &
50 Hewitt, 1985). In such systems, individuals freely hybridize in the contact zone and hybrids have
51 reduced fitness due to the segregation of genetic incompatibilities. Studies of naturally hybridizing
52 populations have also increasingly highlighted the role of other isolating barriers, including
53 environmentally-induced selection (Endler, 1977, Moore, 1977, Arnold, 1997) and sexual isolation
54 (Seehausen et al., 1997, Poelstra et al., 2014).

55 Hybridizing populations vary not only in the nature of isolating mechanisms that are involved
56 but also in geographic structure. Hybrid zones are typically characterized by a clinal structure
57 (gradients of allelic frequencies between pure parental populations). More complex spatial
58 structures are found when the environment induces differential selection on hybridizing species and
59 the distribution of habitats is discrete (e.g. islands, lakes, host plants) or otherwise heterogeneous,
60 leading to patchy hybrid zones (mosaic hybrid zones, Harrison & Rand, 1989, and other types of
61 replicated hybridizing populations, reviewed e.g. in Harrison & Larson, 2016). Variable degrees of
62 patchiness can also be induced by colonization history or population stochasticity (Gompert et al.,
63 2010). Whatever causes patchiness, patchy systems allow us to compare multiple, potentially
64 independent contact zones (McKinnon & Rundle, 2002, Bierne et al., 2003, Butlin et al., 2014). Such
65 comparisons are also possible in hybrid zones that have a simpler spatial structure but that can be
66 sampled along replicated transects (e.g. Teeter *et al.*, 2010), and, notably, in experimental
67 populations (Pritchard & Edmands, 2013). These comparative analyses may increase our

68 understanding of isolation mechanisms, their associated genomic architecture, and, promisingly,
69 speciation (Harrison & Larson, 2016, Westram et al., 2016).

70 Particularly intriguing are the situations where one can compare populations composed by a
71 mixture of individuals of two species that in some instances hybridize and in other instances remain
72 strongly reproductively isolated. That is, sympatric or mixed populations that may or may not be
73 reproductively isolated; hereafter we will use the term "mixed populations", defined as populations
74 where individuals of two species are close enough so that they can meet and interact frequently. An
75 illustrative case in point is the lake Victoria cichlid 'speciation transect' (Seehausen, 2009) where
76 mixed populations of *Pundamilia pundamilia* and *P. nyererei* show more or less hybridization
77 depending on variations in premating behavioral mechanisms themselves linked with variations in
78 habitat (water clarity). Fish studies have provided a few other related examples where a pair of
79 species shows contrasted levels of reproductive isolation when in sympatry (benthic and limnetic
80 three-spined sticklebacks, Taylor et al., 2006, swordtail fish, Culumber et al., 2011, and lake
81 whitefish, Gagnaire et al., 2013, river and blueback herring, Hasselman et al., 2014, and river and
82 brook lampreys, Rougemont et al., 2015). Comparing sympatric non-hybridizing / hybridizing
83 populations provides power to interpret admixture patterns (e.g. shared ancestral polymorphism vs
84 current gene flow) or assess whether differential introgression patterns are due to heterogeneous
85 recombination, selection, or gene flow (Gagnaire et al., 2013, Powell et al., 2013, Rougemont et al.,
86 2016).

87 Here we focus on the *Jaera albifrons* group, a complex of five marine isopod species that live
88 on the shores of the temperate and cold waters of the North-Atlantic Ocean. It includes *J. albifrons*, *J.*
89 *praehirsuta*, *J. ischiosetosa*, *J. forsmanni*, and *J. posthirsuta* (Bocquet, 1953, Naylor & Haahtela, 1966,
90 Bocquet, 1972). Note that *Jaera albifrons* designates one of the five species of the *Jaera albifrons*
91 group (the distinction will be noted using the words "complex" or "group" throughout). All five
92 species occupy a narrow but geographically extended belt in the intertidal zone and they have largely
93 overlapping distribution ranges. In short, individuals from one species frequently coexist with

94 individuals from at least one other species throughout their distribution range, and mixed
95 populations are the rule rather than the exception. In this context, the five species of the *Jaera*
96 *albifrons* group were shown to be reproductively isolated by at least three types of barriers: i)
97 ecological isolation (variations in local habitat preferences along the seashore), ii) sexual isolation
98 (differences in male secondary sexual traits used in tactile courtship, and strong female-driven mate
99 choice), and iii) genetic incompatibilities (reviewed in Solignac, 1978, 1981, Mifsud, 2011). The
100 reproductive isolation resulting from the combination of these pre- and post-zygotic barriers is
101 thought to be very strong in nature.

102 However, intermediate male sexual traits have been reported in a few populations, suggesting
103 that hybridization may happen in some rare places (Solignac, 1978). One such potentially hybridizing
104 population has been intensively studied by Charles Bocquet and Michel Solignac between 1965 and
105 1970. They described a *Jaera albifrons* / *J. praeheirsuta* mixed population located in Luc-sur-Mer,
106 Normandy (France) where the analysis of male secondary sexual traits and experimental crosses led
107 them to conclude that this population contained an exceptional proportion of hybrids (15 to 32%
108 depending on sampling event and classification thresholds, Bocquet & Solignac, 1969, Solignac,
109 1969a, b, 1978). Based on morphological descriptions for a large number of individuals sampled or
110 raised in the lab from this population (nearly 2000 ind., Solignac, 1978) and comparison with
111 experimental crosses (Bocquet & Solignac, 1969), their conclusion on hybridization between species
112 seems very strong. This past work convincingly suggests that hybridization was occurring in at least
113 one particular site in 1965-70 while all other known *J. albifrons* / *J. praeheirsuta* sympatric populations
114 were reproductively isolated. As discussed above, such a situation seems interesting for the study of
115 isolating barriers and speciation.

116 With this study, our objectives were i) to test for hybridization between *J. albifrons* and *J.*
117 *praeheirsuta* using genetic tools, ii) to investigate the geographic structure of hybridizing populations
118 and the nature of the isolating mechanisms, and iii) to compare genetic patterns within hybridizing vs
119 non-hybridizing mixed populations. For this purpose, we searched for mixed populations and

120 morphologically intermediate individuals as described in Normandy ca. 50 years ago, analyzed the
121 genetic structure of local populations using a panel of 23 microsatellite loci, and compared it with
122 mixed populations from another French region (Brittany) where the two species had been described
123 as reproductively isolated.

124

125 **Materials and methods**

126 Model species

127 Members of the *Jaera albifrons* complex are small marine crustaceans (2-5 mm total adult
128 length, Fig. 1). Out of the five species comprising the *Jaera albifrons* complex, two are restricted to
129 the temperate waters of the North-American east coast (*J. posthirsuta*) or the European coast (*J.*
130 *forsmani*), while the three others are more widely distributed on both sides of the Northern Atlantic
131 (Bocquet, 1972). These five species are found in abundance in the intertidal zone, where they can
132 show local habitat preferences involving variations in micro-habitat (under rocks or on seaweeds),
133 level on the intertidal zone, and salinity (Naylor & Haahtela, 1966, Jones, 1972). However, these
134 habitat preferences vary widely, meaning that ecological isolation is also very variable (Veuille, 1976,
135 Solignac, 1981).

136 The identification of species within the *Jaera albifrons* complex is based on male secondary
137 sexual traits (Fig. 1). Mating is preceded by a courtship behavior whereby males mount females in a
138 head-to-tail position and used different parts of their pereopods to brush or press the female's
139 back. Males of the five species differ in the distribution of setae and spines borne by the pereopods
140 used to court females (Jones & Fordy, 1971, Solignac, 1978), and a female's acceptance or rejection is
141 a major driver of reproductive isolation between species.

142

143 Species survey and sampling

144 We sampled *J. albifrons* and *J. prae-hirsuta* in two regions. First we focused on the area where
145 Michel Solignac had described hybridization between these two species in 1965 and 1970 (Solignac,

146 1978). For that, we surveyed strictly all potential habitats on a 25 Km portion of the coast around this
147 original site, finding the population studied by M. Solignac to be extinct (Luc-sur-Mer, Fig. 2), possibly
148 due to the regular removal of pebbles from the beach for touristic activities. We extended this survey
149 35 Km East and 35 Km West by visiting a large number of (but not strictly all) potential habitats,
150 where we found three sites with a mixture of *J. albifrons* and *J. prae-hirsuta* (sites 7-9, see Results,
151 Table 1, and Fig. 2). This gave us a 95 Km continuous portion of the coastline where we have a
152 precise, although not strictly exhaustive, view of the distribution of species (from Grandcamp-Maisy
153 to Honfleur, second world war landing beaches, highlighted in yellow in Fig. 2).

154 Less intensive surveys were more recently conducted further West and East in order to check
155 for additional mixed or pure *J. albifrons* / *J. prae-hirsuta* populations (such populations were found
156 but not analyzed in this study, see Results).

157 Second, we searched for similarly mixed populations of the same pair of species in a region
158 where no hybridization had been found despite extensive field studies (area around Roscoff biology
159 station, Brittany, France, Fig. 2, Bocquet & Solignac, 1969, Solignac, 1969b). In this region we selected
160 6 sampling sites where the two species were found at the same location (Fig. 2 and Table 1).

161 In both regions (Brittany and Normandy) we looked for individuals of the *Jaera albifrons*
162 complex under rocks and on seaweeds (*Ascophyllum nodosum*, *Fucus vesiculosus* and *Fucus serratus*
163 essentially) in the intertidal zone. Animals found on rocks were collected in the field using a small
164 brush. By contrast, samples of seaweeds were brought back to the lab where we checked for the
165 presence of *Jaera* individuals by shaking algae repeatedly in freshwater (Solignac, 1978). All
166 individuals were kept alive until identification based upon observation of male secondary sexual
167 traits. Because females of all species are morphologically identical, this study is based on male
168 individuals only. As adult females are larger than males, many females could be left alive in the field.
169 All collected individuals were fixed in ethanol after species identification.

170 Finally, one sampling site (Ste-Honorine-des-Pertes, site 8 in figure 2) was selected for a
171 detailed analysis of the micro-distribution of individuals. In this site we performed an exhaustive

172 survey on a large portion of the beach, recording the precise localization of each individual with a
173 Trimble GeoExplorer 6000 GPS (average horizontal accuracy 55 mm). There we also collected
174 females, which were kept in the lab until they produced offspring (sperm storage allows females to
175 produce offspring in absence of males). These offspring were reared in the lab for at least six weeks,
176 until each individual could be sexed and each male could be identified based on secondary sexual
177 traits. This identification of series of male sibs gave a presumptive species identification for their
178 mother (e.g. a female could be classified as *J. albifrons*, *J. praehirsuta* or "hybrid" if it gave birth to a
179 series of males bearing *J. albifrons*, *J. praehirsuta* or intermediate morphological traits, respectively).

180

181 Genetic analyses

182 All genetic analyses are based on individual genotypes obtained at 23 microsatellite loci (all
183 loci described in Ribardière et al., 2015, except Ja01). DNA extraction and amplification followed the
184 protocols described in Ribardière *et al.* (2015). Pairwise linkage disequilibrium between loci was
185 tested in FSTAT version 2.9.3.2 (Goudet, 2001) in order to avoid redundant information. Departure
186 from Hardy-Weinberg equilibrium was also tested in FSTAT in order to detect technical artifacts (null
187 alleles or allelic dropouts) or departure from random mating within sampling sites. The occurrence of
188 null alleles, already detected in the *Jaera albifrons* complex with these microsatellites (Ribardière *et*
189 *al.*, 2015) was specifically investigated with the software MICROCHECKER (Van Oosterhout *et al.*, 2004).
190 About 10% of the genotypes were randomly replicated in order to evaluate the frequency of
191 genotyping errors and for each locus genotyping error rate was calculated as error rate = (number of
192 false genotypes) / (total number of repeated genotypes). The level of polymorphism was estimated
193 by measuring observed and expected heterozygosity in FSTAT.

194 Our test of reproductive isolation or hybridization between *J. albifrons* and *J. praehirsuta*
195 within our two sampling regions (Brittany and Normandy) is based upon estimates of genetic
196 differentiation. We estimated the distribution of genetic variance among sampling sites within a
197 species (F_{SC}) and between species (F_{CT}) in a hierarchical analysis of molecular variance (AMOVA,

198 Excoffier et al., 1992) implemented in ARLEQUIN version 3.5 (Excoffier & Lischer, 2010). We performed
199 this analysis independently within each region. The between-species component F_{CT} will thus inform
200 us on the strength of genetic differentiation between *J. albifrons* and *J. praehirsuta* within each
201 region. Small samples were not included in these analyses (that is, individuals with an intermediate
202 phenotype, and individuals from site 9, where one of the two species was represented by only one
203 individual). These analyses were performed using both allelic frequencies (F_{ST} -like) and the distance
204 between alleles (R_{ST} -like) and significance was assessed using 10100 permutations as implemented in
205 ARLEQUIN. We ran these analyses first using all markers (global AMOVA) and then for each
206 microsatellite locus independently (locus-by-locus AMOVA).

207 Because some loci showed strongly contrasted levels of between-species genetic structure
208 (F_{CT}) in Normandy vs Brittany (see results), the significance of the difference between $F_{CT-Normandy}$
209 against $F_{CT-Brittany}$ was tested by bootstrapping individuals 1000 times in R version 3.2.3 (R Core Team,
210 2016). This allowed us to estimate how often the two F_{CT} values obtained from a given resampled
211 dataset overlapped, and thus whether $F_{CT-Normandy}$ differed significantly from $F_{CT-Brittany}$ at the locus
212 tested.

213 Pairwise estimates of genetic differentiation between samples were also obtained in a non-
214 hierarchical model in ARLEQUIN (that is, F_{ST} between all pairs of populations, where a population is
215 defined by a given species in a given sampling site, Table 1). This is useful i) to evaluate whether the
216 global differentiation between species is consistent across sampling sites (i.e. using F_{ST} between
217 species within each site separately), and ii) to investigate genetic structure within each species
218 separately, in particular by testing for isolation by distance between sampling sites with SPAGEDI 1.4
219 (Hardy & Vekemans, 2002) and GENEPOP 4.2.2 (Rousset, 2008).

220 Finally, the distribution of genetic variance was also investigated using individual analyses
221 without a priori grouping of samples. We first ran a clustering analysis within each region using
222 STRUCTURE 2.3.4 (Pritchard *et al.*, 2000) with an admixture model (10 independent repetitions, burn-in
223 period= 50000, MCMC= 300000). The most likely number of clusters (K) was determined *via*

224 HARVESTER v.0.6.1 (Earl, 2012) using ΔK as described by Evanno *et al.* (2005). The *Ancestdist* option
225 was used in STRUCTURE to calculate 95% probability intervals for an individual's membership q to
226 each cluster. The width of such intervals (difference between upper and lower bounds) gave us an
227 estimate of the precision of membership values. Second, we performed a Principal Component
228 Analysis (PCA) based on all individual genotypes using the R package ade4 (Dray & Dufour, 2007).
229 Because the locus-by-locus AMOVA showed that two loci had a striking behavior (see results), we ran
230 both analyses (PCA and STRUCTURE) with and without these two loci.

231

232 **Results**

233 **Population survey**

234 In Brittany we sampled 142 *J. albifrons* and 80 *J. praeheirsuta* in six sites where the two species
235 were co-occurring (Fig. 2 and Table 1, only males are considered throughout the paper, unless stated
236 otherwise). In these sites, all *J. albifrons* but two were found under rocks, while all *J. praeheirsuta*
237 except three were found on seaweeds (located directly above rocks or within a radius of a few
238 meters). In this region, all individuals could be morphologically identified at the species level without
239 any overlap of traits (i.e. no individuals with intermediate morphology).

240 In Normandy we sampled 44 *J. albifrons* and 57 *J. praeheirsuta* from the three mixed
241 populations that we found (sites 7-9, Fig. 2). Contrary to the Brittany situation, all individuals were
242 collected under rocks, while no individuals could be found on seaweeds. In addition, we found 9
243 individuals clearly showing intermediate morphological traits as described by Solignac (Fig. 1, "hybrid
244 types" 5 to 13 in Solignac 1978, p.172-177). These individuals will be referred to as "intermediate
245 phenotypes" hereafter. The portion of the coast that was intensively surveyed revealed a single pure
246 *J. albifrons* population located >70 Km East of mixed populations (empty circle in figure 2).

247 Later, less intensive surveys revealed another pure *J. albifrons* population 30 Km West (Fig. 2),
248 and interestingly, one additional mixed *J. albifrons* / *J. praeheirsuta* population (with intermediate

249 phenotypes) further East (location Yport, Fig. 2). No *J. praehirsuta* individuals were found outside of
250 mixed populations anywhere in Normandy.

251 The fine-scale distribution of individuals at site 8 (Ste-Honorine-des-Pertes, Normandy)
252 showed that individuals (61 males and 138 females) of the two species and intermediate phenotypes
253 were largely intermingled, with *J. praehirsuta* being distributed all along the shore while the
254 distribution of *J. albifrons* was more irregular (supplementary material Fig. S1).

255

256 **Genetic diversity**

257 All microsatellite loci could be amplified in individuals of the two species (consistent with
258 Ribardière *et al.*, 2015) as well as in individuals with intermediate phenotypes. There was no linkage
259 disequilibrium after Bonferroni correction between all pairs of loci in each population, so that all 23
260 markers were kept for further analyses. The level of polymorphism was globally consistent across
261 species (Table 1). Unless stated otherwise, individuals with intermediate phenotypes were removed
262 from the following analyses for we had too few of them (2 in Grandcamp, site 7, and 7 in Ste-
263 Honorine-des-Pertes, site 8, Fig. 2). Observed heterozygosity H_o (0.45 in *J. albifrons* and 0.51 in *J.*
264 *praehirsuta*) was on average lower than within population gene diversity H_e (0.51 and 0.59,
265 respectively), resulting in a significant departure from Hardy-Weinberg equilibrium (F_{IS} values in
266 Table 1). Departure from HWE was driven in Brittany by loci Ja37, Ja39, Ja55 and Ja94, and in
267 Normandy by locus Ja55 (loci with significant positive F_{IS} in one to six samples, data not shown).
268 These loci, except Ja94, showed signs of a null allele in more than half of populations, as tested using
269 MICROCHECKER. While we did not detect locus-specific HW disequilibrium patterns that were
270 consistent across populations, the main downstream quantitative analysis (analysis of molecular
271 variance, see below) was run with and without the four loci cited above (and we report locus-specific
272 results as well). Genotyping error rates estimated from replicated individuals ranged from 0 to 5.88%
273 per locus (average 1.7%), and were due roughly equally to allelic dropouts and false alleles.

274

275 **Genetic differentiation between *J. albifrons* and *J. praehirsuta***

276 The hierarchical analysis of molecular variance based on 23 microsatellite loci (Table 2) showed
277 that the between-species differentiation was higher in Brittany ($F_{CT-Brittany} = 0.132$, $p < 0.005$) than in
278 Normandy ($F_{CT-Normandy} = 0.074$, $p = 0.34$). Accordingly, the locus-by-locus AMOVA showed that most
279 loci were less differentiated between species in Normandy than in Brittany (Fig. 3). However, two loci
280 (Ja41 and Ja64) revealed a strikingly different pattern: these two loci showed a very strong level of
281 differentiation (Ja41, $F_{CT-Normandy} = 0.462$; Ja64, $F_{CT-Normandy} = 0.384$) in Normandy region while the
282 remaining 21 loci showed no or little differentiation (locus specific $F_{CT-Normandy}$ ranged from -0.05 to
283 0.079], Fig. 3). Without these two peculiar loci, there is thus no genetic differentiation between
284 species in Normandy ($F_{CT-Normandy} = 0.017$, $F_{CT-Brittany} = 0.125$, table 2). Note that this result is unchanged
285 when removing the four loci that showed a departure from HW equilibrium in some populations (not
286 shown).

287 Moreover, the differentiation between species at loci Ja41 and Ja64 in Normandy was not only
288 much stronger than at other loci but it was also stronger than the differentiation observed at the
289 same two loci in Brittany (Ja41, $F_{CT-Brittany} = 0.199$, Ja64, $F_{CT-Brittany} = 0.194$, Fig. 3), and this difference
290 was significantly different from 0 (bootstrap p -value < 0.001 for Ja41 and $p = 0.001$ for Ja64,
291 supplementary material Figs. S2 and S3). The distribution of allelic frequencies at these two loci is
292 presented in supplementary material figures S4 and S5.

293 Although the heterogeneity across loci appeared somewhat lower in Brittany (Fig. 3), there
294 was nonetheless some variation, with two other loci showing a particularly strong level of
295 differentiation (Ja66, $F_{CT-Brittany} = 0.605$; Ja80, $F_{CT-Brittany} = 0.37$). The difference in F_{CT} between regions
296 was significant at these two loci (bootstrap p -value < 0.001 for Ja66 and Ja80, supplementary material
297 figure S3).

298 Because we studied only three mixed populations in Normandy, and one of them contained
299 nearly only *J. albifrons* (site 9, Longues-sur-Mer, Fig. 2, Table 1), the permutation procedure
300 implemented in ARLEQUIN to test for the significance of $F_{CT-Normandy}$ is essentially powerless (the

301 between-species component of genetic variation is tested by permuting populations within species).
302 We therefore checked if the patterns found in the AMOVA (which considers all sites simultaneously)
303 were consistent across sites. Tables 3 and 4 show pairwise F_{ST} values calculated in a simple non-
304 hierarchical framework. Most importantly, it shows that the between-species differentiation was
305 consistent across sites, both in Normandy (F_{ST} between species equal to 0.104 at Grandcamp, site 7,
306 and 0.101 at Ste-Honorine-des-Pertes, site 8) and in Brittany (F_{ST} between species within sites in [0.1;
307 0.19]). These results consider all loci, but the same geographical consistency is observed when
308 considering the locus-specific patterns described above (data not shown). That is, the global AMOVA
309 results are repeatable across sites (e.g. F_{ST} at the two sites from Normandy = 0.546 (site 7) and 0.396
310 (site 8) when considering only loci Ja41 and Ja64, and F_{ST} = 0.05 (site 7) and 0.056 (site 8) with all
311 other loci).

312 Individual analyses bring some complementary information, in particular because the
313 individuals with intermediate morphology could be included in spite of their low abundance (as well
314 as individuals from site 9). Running STRUCTURE with $K=2$, we found that *J. albifrons* and *J. praehirsuta*
315 cluster into two clearly identified groups both in Brittany and Normandy using a panel of 23 loci (Fig.
316 4). However, while an individual's membership q to its assigned cluster was similar for both species in
317 Brittany and Normandy (membership averaged over all individuals and 10 STRUCTURE runs, Brittany:
318 $\bar{q}_{albifrons} = 0.99$, $\bar{q}_{praehirsuta} = 0.97$, and Normandy: $\bar{q}_{albifrons} = 0.97$, $\bar{q}_{praehirsuta} = 0.95$), the
319 uncertainty associated with q was larger in Normandy (average width of 95% probability interval
320 $w_{albifrons}^{95} = 0.24$, $w_{praehirsuta}^{95} = 0.32$) than Brittany ($w_{albifrons}^{95} = 0.12$, $w_{praehirsuta}^{95} = 0.19$). As it
321 turned out, the apparent genetic clustering in Normandy was almost entirely due to the effect of two
322 loci only (Ja41 and Ja64), while a STRUCTURE analysis using the remaining 21 loci showed that the two
323 species were genetically homogeneous (Fig. 4B).

324 The results from molecular analyses of variance and clustering analyses could be well
325 visualized using PCA performed with all 23 loci (Figure 5). The two isopod species were clearly
326 differentiated in one region (Brittany) and less so in the other one (Normandy). We see also that

327 individuals with intermediate phenotypes were genetically indistinguishable from individuals with *J.*
328 *prae-hirsuta* traits.

329 The particular geographical distribution of individuals (replicates of mixed populations
330 comprising intermediate phenotypes and absence of clinal structure, see discussion) precluded the
331 use of genetic tools dedicated to the analysis of hybridization in clinal hybrid zones.

332

333 **Genetic structure within species**

334 Because there is ongoing hybridization and introgression between the two species in
335 Normandy (see discussion), the within-species genetic structure is best investigated using samples
336 from Brittany. Brittany is also the region where we have more sampling locations. The genetic
337 structure among *J. albifrons* samples appeared slightly higher than within *J. prae-hirsuta*. This is visible
338 from pairwise F_{ST} estimates (Table 3), ranging from 0.005 to 0.066 (13 out of 15 pairs significantly
339 different) when considering only *J. albifrons* within Brittany, and -0.075 to 0.017 (1 out of 15 pairs
340 significant) when considering only *J. prae-hirsuta*.

341 In addition, a significant pattern of isolation-by-distance (sup. Fig. S6) was observed in Brittany
342 both in *J. albifrons* ($R^2 = 0.73$, Mantel test p -value < 0.01) and *J. prae-hirsuta* ($R^2 = 0.11$, p -value = 0.02).
343 In line with the pairwise F_{ST} results, genetic differentiation increased more rapidly with distance in *J.*
344 *albifrons* than in *J. prae-hirsuta*, although 95% confidence intervals calculated in GENEPOP overlapped
345 (10000 permutations, *J. albifrons* [0.013, 0.037], *J. prae-hirsuta* [-0.0005, 0.018], Fig. S6).

346

347 **Discussion**

348 The first result of this study is the clear-cut confirmation that introgressive hybridization is
349 happening between *J. albifrons* and *J. prae-hirsuta* in at least two mixed populations from Normandy,
350 France. As developed below, this opens interesting questions regarding the conditions of coexistence
351 of the two parental morphs in hybridizing populations that seem to receive no influx from pure
352 parental populations and shows no detectable ecological heterogeneity.

353

354 1- Hybridization between *Jaera albifrons* and *J. prae-hirsuta*

355 Analyses of molecular variance (Table 2 and Fig. 3) and admixture analyses (Figs. 4 and 5) both
356 showed that mixed populations from Normandy have a homogeneous genetic structure at 21 out of
357 23 multi-allelic loci. Critically, this genetic homogeneity contrasts with the differentiation observed in
358 mixed populations from Brittany, where individuals bearing sexual traits specific to *J. albifrons* or *J.*
359 *prae-hirsuta* cluster into clearly marked genetic groups (Figs. 4A and 5). Hence shared ancestral
360 polymorphism cannot explain the lack of differentiation between species in Normandy, which
361 therefore supports the hypothesis of ongoing hybridization.

362 These findings agree with the conclusions reached by C. Bocquet and M. Solignac nearly 50
363 years ago, who studied the morphological variation of secondary sexual traits in a population from
364 the same region (Luc-sur-Mer, Fig. 2, Bocquet & Solignac, 1969, Solignac, 1969a, b, 1978 chapter 6).
365 Similarly to the results reported by these authors, we found that in Normandy several males have
366 intermediate phenotypes and the two species occupy the same habitat (under stones and pebbles on
367 the shore) while in Brittany we did not detect any intermediate phenotypes and the two species
368 occupy two different habitats, with some overlap; *J. albifrons* lives primarily under stones, while *J.*
369 *prae-hirsuta* is found primarily on seaweeds.

370 We conclude from these observations and the contrast in species divergence in the two
371 regions that the two species are currently hybridizing in populations from Normandy, but not in
372 Brittany.

373

374 2- A semi-permeable barrier to gene flow

375 The genetic homogeneity observed across species in Normandy further shows that
376 hybridization has been introgressive, as correctly concluded by Solignac (1969b) from the continuous
377 range of morphological characteristics observed in natural populations and by comparison with
378 experimental crosses (Bocquet & Solignac, 1969). While it is now clearly established that

379 introgression proceeds differentially across loci in hybrid zones (with no known exceptions, Harrison
380 & Larson, 2014), investigating this variation was not part of our original plan with this study given
381 that we were using a panel of only 23 loci. However, the locus-by-locus AMOVA analyses revealed a
382 surprisingly trenchant pattern, whereby 21 loci showed no differentiation at all between species in
383 populations from Normandy (F_{CT} in [-0.05; 0.08]) and the two remaining loci where strongly
384 differentiated ($F_{CT} = 0.384$ and 0.462). Moreover, these two loci were also significantly more
385 differentiated in the hybridizing populations (Normandy) than in reproductively isolated ones
386 (Brittany) while showing no differentiation within each species (see F_{SC} values in Table 2). This
387 strongly suggests that there is a semi-permeable barrier to gene flow between *J. albifrons* and *J.*
388 *praeheirsuta* in hybridizing populations from Normandy. Hypotheses other than a reduction in gene
389 flow at these loci seem impossible to reconcile with the fact that the same two loci are significantly
390 less differentiated in non-hybridizing sympatric populations. Alternate hypotheses such as the
391 differential sorting of ancestral polymorphism or reduced variability at these loci due to a locally low
392 recombination rate would require a history of differentiation between species whereby ancestral
393 polymorphism or recombination have evolved differentially in the two regions studied (ca 250km
394 apart). While this is theoretically possible, a more parsimonious hypothesis is that the two loci are
395 encompassed in one or two genomic regions where inter-specific gene flow is hampered because
396 these regions are linked with one or several isolating barriers.

397 Additional indicators of a semi-permeable barrier to gene flow are two other markers (Ja66
398 and Ja80) also showing a heterogeneous pattern. As most markers they are more differentiated in
399 Brittany than in Normandy, but, interestingly, they show a stronger differentiation than the other loci
400 in Brittany (Fig. 3). We cannot currently make assumptions based only on these results, which
401 emphasize the necessity to study the heterogeneity of genome with an extended set of genetic
402 markers.

403

404 3- Is introgression symmetrical?

405 Interspecific crosses are generally not equally likely in both directions, especially when
406 behavioral isolation is involved (e.g. Coyne & Orr, 2004, p. 226). Such asymmetries leave specific
407 signatures in the genome that are most easily detected by comparing uni- and bi-parentally inherited
408 genetic variation (typically, markers from the mitochondrial and nuclear DNA, e.g. Toews & Brelsford,
409 2012). It would be interesting to test for asymmetric introgression of mtDNA in our system given that
410 Bocquet and Solignac (1969) have suggested that interspecific crosses may occur more easily in one
411 direction (female *J. praehirsuta* x male *J. albifrons*) than the other. This result was obtained from
412 experimental crosses with individuals from a hybridizing population (Luc-sur-Mer, Normandy), and
413 the asymmetry was further confirmed by Solignac (1981) using individuals from other origins (non-
414 hybridizing populations). Yet we could not use this approach here because mtDNA analyses so far
415 have indicated that most of the genetic variation is shared by all species of the *Jaera albifrons*
416 complex (perhaps excluding the American species *J. posthirsuta*, which has not been included in
417 these analyses). That is, the four European species form a polyphyletic clade (16S rDNA, Mifsud,
418 2011, and COI, Ribardièere and Broquet unpublished), in strong contrast with the patterns obtained
419 with nuclear data (AFLP, Mifsud, 2011, and microsatellites, this study). There are no mitochondrial
420 haplotypes or clades that are specific to *J. albifrons* or *J. praehirsuta* (not shown), and the cyto-
421 nuclear discordance, also certainly interesting in its own right, is not informative of recent
422 introgression directionality. The symmetry of introgression can in some cases be evaluated by taking
423 advantage of differences between nuclear loci (differential introgression), but our microsatellite
424 dataset is too limited for this approach. Nevertheless, results of admixture analyses show that
425 individuals showing an intermediate phenotype share more genetic background with *J. praehirsuta*
426 (Figs. 4 and 5) which suggests that introgression is asymmetric (genetic variation from *J. albifrons*
427 introgressing into the genetic background of *J. praehirsuta*). This can also be seen by looking at allelic
428 frequencies at loci Ja41 and Ja64 for the phenotypically intermediate individuals, which are similar to
429 that of individuals bearing *J. praehirsuta* traits and different from the frequencies observed in *J.*
430 *albifrons* (supplementary material Figs. S4 and S5).

431

432 4- Geographical structure and persistence of hybridizing populations

433 In a 1969 paper, C. Bocquet and M. Solignac reported that many "morphological hybrids" have
434 been observed during the preceding 15 years in their study area of Luc-Sur-Mer (Bocquet & Solignac,
435 1969). There is no more suitable habitat at this site, but Solignac (1978, p. 171) mentioned that
436 "hybrids" had been found in Ste-Honorine-des-Pertes, which is one of the two sites sampled in the
437 present study (site 8). This means that hybridizing populations have persisted in Normandy for at
438 least several decades. Moreover, during a recent additional survey aiming at extending the 95 Km
439 coastline region studied here, we found intermediate phenotypes in a population located roughly
440 100 Km East of the mixed populations studied here (location "Yport", Fig. 2). Even more
441 unexpectedly, we detected a *J. albifrons* – *J. praeheirsuta* mixed population with some intermediate
442 male phenotypes in the Isles of Scilly, UK, an archipelago that is located more than 400 Km away
443 across the English Channel (Fig. 2). This means that hybridization between these two species is
444 probably much more widespread than previously thought (Solignac, 1969a, b, 1978). Perhaps more
445 importantly, this also strongly suggests that hybridizing populations have been persisting for a long
446 time.

447 In this study we identified 9 individuals (out of 110 males found in Normandy) showing
448 intermediate morphological traits. While this figure depends on what one recognizes as
449 morphologically pure or intermediate individuals, the majority of males clearly show strict *J. albifrons*
450 or *J. praeheirsuta* sexual traits despite the extensive genetic introgression demonstrated here
451 (bimodal hybrid zone, Jiggins & Mallet, 2000). We concur with Solignac's observation (1978, p. 188)
452 that the coexistence of the two morphs in spite of introgressive hybridization is of great interest, and
453 we discuss below the mechanisms that may allow this coexistence in the long term and in repeated
454 areas. This part of the discussion will focus on hybridizing populations only (that is, results from
455 Normandy).

456 The closest population containing only *J. albifrons* that we found was located at more than 30
457 Km from the hybridizing populations. Moreover, the mixed populations that we found in this region
458 are geographically isolated from one another (there is most likely a discontinuity at least between
459 sites 7 and 8, Fig. 2). All species of the *Jaera albifrons* group have a direct development without a
460 dispersive larval phase, and in Normandy they do not live on seaweeds, which could potentially drift
461 across populations. Gravid females caught in the water column could occasionally be moved over a
462 great distance, but we did not find any *J. praeirsuta* outside of the mixed *J. albifrons* / *J. praeirsuta*
463 populations in Normandy. It is difficult to conduct an exhaustive survey over large areas for such
464 small species and we could have missed pure *J. albifrons* (and perhaps even *J. praeirsuta*)
465 populations at dispersal distance from our hybridizing populations, but we feel that it is unlikely.
466 Given that these crustaceans do not have a dispersive larval phase, and given the patchy distribution
467 of habitats, we infer from our surveys that hybridizing populations are replicated and patchily
468 distributed.

469 The hybridizing populations analysed in this study do not seem to be flanked by -or otherwise
470 functionally connected to- pure parental populations. They seem to be independent replicates of
471 hybridizing populations potentially distributed on a much larger geographical area than the one
472 studied here (e.g. on the UK coast). The influx of individuals from pure parental populations of *J.*
473 *praeirsuta* (and probably *J. albifrons*) is thus most likely not one of the forces acting to stabilize the
474 system. This interpretation needs further testing (e.g. from additional surveys and analyses of spatial
475 genetic structure), since if this hypothesis is confirmed, it would exclude dispersal-dependent models
476 of hybrid zones, chief among them the tension zone model (Barton & Hewitt, 1985), which relies on a
477 balance between immigration of parental genotypes and selection against hybrids. An alternative
478 hybrid zone model without immigration from parental populations involves ecological variation and
479 an advantage of hybrids in intermediate habitats (Moore, 1977). This hypothesis seem also be
480 excluded in our case because we were unable to detect any variation in habitat within hybridizing
481 populations (there was no identifiable variation in the distribution of *J. albifrons*-like, *J. praeirsuta*-

482 like, and morphologically intermediate individuals within a site). Other classical models are also
483 inappropriate, for they combine the tension zone balance with ecological variation, either through
484 environmentally-induced selection against hybrids (Endler, 1977) or a patchy distribution of habitats
485 favouring one or the other species (mosaic hybrid zones, Harrison & Rand, 1989). The literature is
486 also rich in empirical studies of hybridizing populations that do not fit one of these classical models,
487 but it seems that situations where species coexist in spite of extensive introgression (i.e. bimodal
488 hybrid zones) most often involve either an income of individuals from pure parental populations or
489 ecological variation and habitat specialization within hybrid zones (or both). When incompletely
490 isolated species occupy different ecological niches, comparative analyses of replicate hybrid zones
491 have made quite clear that the maintenance of bimodality is correlated with the opportunity for
492 ecological specialization (e.g. Culumber et al., 2011, Gagnaire et al., 2013). In the *Jaera albifrons* / *J.*
493 *praehirsuta* system, the two species are more differentiated in our populations from Brittany where
494 they specialize in two different habitats (rocks vs seaweeds) but, interestingly, parental forms coexist
495 despite introgression in *hybridizing populations* in Normandy where there is probably no habitat
496 specialization.

497 Assuming that there is no immigration from pure parental populations and no ecological
498 variability, what evolutionary forces would allow pure *J. albifrons* and *J. praehirsuta* phenotypes to
499 coexist in hybridizing populations? Past work suggests two strong candidates. First, the strongest
500 isolating barrier between species of the *Jaera albifrons* complex is sexual isolation (Solignac, 1981).
501 The courtship behavior of males (plus perhaps unknown male characteristics such as pheromone
502 production or other unnoticed phenotypic variation) and female preference may still partially isolate
503 *J. albifrons* from *J. praehirsuta* in hybridizing populations. One compelling hypothesis in this regard is
504 that females of one species accept heterospecific mating more readily than females of the alternate
505 species. This is nearly the rule in case of behavioral isolation (e.g. Coyne & Orr, 2004) and empirical
506 tests suggest that this happens in the hybridizing populations studied here (although with a limited
507 sample size, Bocquet & Solignac, 1969). The hypothesis that sexual isolation is a strong component is

508 also in line with other examples where hybrid zones remain bimodal (reviewed in Jiggins & Mallet,
509 2000).

510 The second candidate is selection against hybrids. This hypothesis is supported by Solignac
511 (1978, p. 186), who reported a strong hybrid breakdown in experimental F2 and backcrosses using
512 individuals from the hybridizing population of Luc-sur-Mer, Normandy. This suggests that the *J.*
513 *albifrons* / *J. praeheirsuta* hybridizing populations documented in this study persist through a balance
514 between hybridization *versus* partial (perhaps asymmetrical) sexual isolation and selection against
515 certain recombined genotypes. This situation seems to be infrequent and the conditions of
516 persistence for such a system deserve further inquiry. In particular, it is a potential empirical example
517 of the model proposed by M'Gonigle *et al.* (2012), in which sexual selection, spatial variation in local
518 carrying capacity, and female mate-search costs allow partially divergent species to persist despite
519 ecological equivalence. More generally, it may provide some insight into the debated role of sexual
520 selection and sexual isolation in species divergence and coexistence. Yet it is remarkable that *J.*
521 *albifrons* and *J. praeheirsuta* coexist in the long term despite their seemingly ecological equivalence. It
522 is possible that some unknown frequency-dependent mechanism (e.g. via an action of pathogens or
523 parasites) is acting to lower the likelihood of extinction of one or the two morphs.

524

525 5-Hybridizing and non-hybridizing mixed populations

526 Why do *J. praeheirsuta* and *J. albifrons* hybridize when in contact in some populations and not
527 in others? There are two obvious differences between our studied populations from Brittany and
528 Normandy. First, where the two species were found to be non-hybridizing, they live on clearly
529 different habitats (under rocks vs on brown algae). The two habitats are located centimetres to
530 meters away, and a few individuals of each species were found on the habitat favoured by the other
531 species, meaning that *J. albifrons* and *J. praeheirsuta* meet each other frequently in these sites (this is
532 why we call them mixed, or sympatric, populations: the two habitats are not separated by a large
533 geographical distance and can be reached by dispersing individuals of either species). Yet these two

534 habitats are radically different and are bound to impose a serious barrier to gene flow between
535 species (ecological isolation). While neither species is restricted to one of these habitats in other
536 parts of their range, the availability of distinct habitats may play an important role facilitating the
537 coexistence of the two species and reducing hybridization opportunities. Appropriate seaweeds can
538 also be found in hybridizing populations from Normandy, but they may not represent a suitable
539 habitat there (e.g. because of wave exposure). Whatever the reason, there are no *Jaera* on algae in
540 the hybridizing populations reported here. One plausible hypothesis is thus that ecological
541 diversification facilitates the coexistence and divergence of *J. albifrons* and *J. praeheirsuta* wherever
542 this is possible. Another interesting idea, suggested by an anonymous reviewer, is that hybridization
543 allows the hybrid and subsequently *J. praeheirsuta* to acquire adaptations to the array of challenges
544 presented in switching from a seaweed to a rocky habitat. One avenue of research to tackle these
545 questions will be to search for pure *J. praeheirsuta* populations occupying rocky habitats (and *J.*
546 *albifrons* populations on seaweeds).

547 The second difference between hybridizing and reproductively isolated populations is their
548 geographic location. Contrary to the idea that natural hybridization is the exception and not the rule
549 in this system (Solignac, 1969a, b, 1978), we suggest that hybridizing populations may be found in a
550 large geographic area. It is therefore possible that populations from two large geographic zones (one
551 encompassing Normandy and the other one encompassing Brittany) have a distinct demographic and
552 evolutionary history (e.g. with variations in the conditions of secondary contacts between species
553 after periods of isolation). The intra-specific genetic differentiation between regions appeared to be
554 similar in *J. albifrons* and *J. praeheirsuta* ($F_{ST} = 0.14$ and 0.15 , respectively) and similar in intensity to
555 the genetic differentiation between species observed in Brittany. This is also apparent on the PCA
556 where the first axis partitions the genotypes in function of their geographic origin (Brittany on the
557 left, Normandy on the right, Fig. 5).

558 Testing these ideas will require analyzing the genetic structure of the two species over a large
559 geographical scale and surveying mixed populations for morphological or genetic signs of
560 hybridization in different habitat conditions.

561

562 **Acknowledgments**

563 This study stands on a tremendous amount of past work, notably due to Michel Solignac, who
564 passed away during the writing of this article. While the authors did not know him personally, they
565 wish to acknowledge his achievements. We also thank Franck Gentil for initiating T.B. to the *Jaera*
566 *albifrons* complex. We thank Nicolas Bierne, Pierre-Alexandre Gagnaire, Julie Jaquiéry, Didier Jollivet,
567 Thomas Lenormand, Frédérique Viard, and Glenn Yannic for discussions or comments on a previous
568 draft. We are grateful to Denis Roze for bringing back useful observations from the Isles of Scilly. We
569 thank the "Centre de Recherches en Environnement Côtier" (CREC, university of Caen) and the
570 "Groupe d'Etude des Milieux Estuariens et Littoraux de Normandie" (GEMEL-Normandie) for giving us
571 access to accommodation and lab space during field surveys in Normandy. This work benefited from
572 access to the Biogenouest genomic platform at Station Biologique de Roscoff and was funded by the
573 French Agence Nationale de la Recherche (grant ANR-13-JSV7-0001-01 to T.B.).

574

575 **Data accessibility**

576 Multi-locus genotypes at each sampling location will be made available in Dryad.

577 **REFERENCES**

- 578 Arnold, M. L. 1997. *Natural hybridization and evolution*. Oxford University Press.
- 579 Barton, N. H. & Hewitt, G. M. 1985. Analysis of Hybrid Zones. *Annu Rev Ecol Syst* **16**: 113-148.
- 580 Bierne, N., Borsa, P., Daguin, C., Jollivet, D., Viard, F., Bonhomme, F. & David, P. 2003. Introgression
581 patterns in the mosaic hybrid zone between *Mytilus edulis* and *M. galloprovincialis*. *Mol Ecol*
582 **12**: 447-461.
- 583 Bocquet, C. (1953) Recherches sur le polymorphisme naturel des Jaera Marina (Fabr.)(Isopodes
584 Asellotes): Essai de systématique évolutive. pp. 436. Centre national de la recherche
585 scientifique.
- 586 Bocquet, C. 1972. La speciation des *Jaera albifrons*. Etat présent des connaissances et des problèmes.
587 131-137.
- 588 Bocquet, C. & Solignac, M. 1969. Etude morphologique des hybrides expérimentaux entre *Jaera*
589 (*albifrons*) *albifrons* et *Jaera (albifrons) praehirsuta* (isopodes asellotes). *Archives Zool Exp*
590 *Gen* **110**: 435-452.
- 591 Butlin, R. K., Saura, M., Charrier, G., Jackson, B., Andre, C., Caballero, A., Coyne, J. A., Galindo, J.,
592 Grahame, J. W., Hollander, J., Kemppainen, P., Martinez-Fernandez, M., Panova, M.,
593 Quesada, H., Johannesson, K. & Rolan-Alvarez, E. 2014. Parallel evolution of local adaptation
594 and reproductive isolation in the face of gene flow. *Evolution* **68**: 935-949.
- 595 Coyne, J. A. & Orr, H. A. 2004. *Speciation*. Sinauer associates, Sunderland.
- 596 Culumber, Z. W., Fisher, H. S., Tobler, M., Mateos, M., Barber, P. H., Sorenson, M. D. & Rosenthal, G.
597 G. 2011. Replicated hybrid zones of Xiphophorus swordtails along an elevational gradient.
598 *Mol Ecol* **20**: 342-356.
- 599 Dray, S. & Dufour, A.-B. 2007. The ade4 package: implementing the duality diagram for ecologists. *J*
600 *Stat Softw* **22**: 1-20.
- 601 Earl, D. A. 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output
602 and implementing the Evanno method. *Conserv Genet Resour* **4**: 359-361.

603 Endler, J. A. 1977. *Geographic variation, speciation, and clines*. Princeton University Press.

604 Evanno, G., Regnaut, S. & Goudet, J. 2005. Detecting the number of clusters of individuals using the
605 software STRUCTURE: a simulation study. *Mol Ecol* **14**: 2611-20.

606 Excoffier, L. & Lischer, H. E. L. 2010. Arlequin suite ver 3.5: a new series of programs to perform
607 population genetics analyses under Linux and Windows. *Mol Ecol Resour* **10**: 564-567.

608 Excoffier, L., Smouse, P. E. & Quattro, J. M. 1992. Analysis of molecular variance inferred from metric
609 distances among DNA haplotypes: application to human mitochondrial DNA restriction data.
610 *Genetics* **131**: 479-491.

611 Gagnaire, P. A., Pavey, S. A., Normandeau, E. & Bernatchez, L. 2013. The genetic architecture of
612 reproductive isolation during speciation-with-gene-flow in lake whitefish species pairs
613 assessed by RAD sequencing. *Evolution* **67**: 2483-2497.

614 Gompert, Z., Lucas, L. K., Fordyce, J. A., Forister, M. L. & Nice, C. C. 2010. Secondary contact between
615 *Lycaeides idas* and *L. melissa* in the Rocky Mountains: extensive admixture and a patchy
616 hybrid zone. *Mol Ecol* **19**: 3171-3192.

617 Goudet, J. 2001. FSTAT, a program to estimate and test gene diversities and fixation indices (version
618 2.9.3), updated from Goudet 1995. Available from
619 <http://www.unil.ch/izea/software/fstat.html>.

620 Hardy, O. J. & Vekemans, X. 2002. SPAGeDi: a versatile computer program to analyse spatial genetic
621 structure at the individual or population levels. *Mol Ecol Notes* **2**: 618-620.

622 Harrison, R. G. & Larson, E. L. 2014. Hybridization, Introgression, and the Nature of Species
623 Boundaries. *J Hered* **105**: 795-809.

624 Harrison, R. G. & Larson, E. L. 2016. Heterogeneous genome divergence, differential introgression,
625 and the origin and structure of hybrid zones. *Mol Ecol* **25**: 2454-2466.

626 Harrison, R. G. & Rand, D. M. (1989) Mosaic hybrid zones and the nature of species boundaries. In:
627 *Speciation and its consequences*, (Otte, D. & Endler, J. A., eds.). pp. 111-133. Sinauer
628 Associates, Inc.

629 Hasselman, D. J., Argo, E. E., McBride, M. C., Bentzen, P., Schultz, T. F., Perez-Umphrey, A. A. &
630 Palkovacs, E. P. 2014. Human disturbance causes the formation of a hybrid swarm between
631 two naturally sympatric fish species. *Mol Ecol* **23**: 1137-1152.

632 Jiggins, C. D. & Mallet, J. 2000. Bimodal hybrid zones and speciation. *Trends Ecol Evol* **15**: 250-255.

633 Jones, M. 1972. Effects of salinity on the survival of the *Jaera albifrons* Leach group of species
634 (Crustacea: Isopoda). *J Exp Mar Biol Ecol* **9**: 231-237.

635 Jones, M. B. & Fordy, M. R. 1971. A Stereoscan electron microscope study of male reproductive
636 characters in the *Jaera albifrons* group of species (Crustacea : Isopoda). *Mar Biol* **10**: 265-271.

637 M'Gonigle, L. K., Mazzucco, R., Otto, S. P. & Dieckmann, U. 2012. Sexual selection enables long-term
638 coexistence despite ecological equivalence. *Nature* **484**: 506-509.

639 McKinnon, J. S. & Rundle, H. D. 2002. Speciation in nature: the threespine stickleback model systems.
640 *Trends Ecol Evol* **17**: 480-488.

641 Mifsud, D. V. (2011) The genetic basis of speciation in the *Jaera albifrons* species group of intertidal
642 isopods. Vol. PhD. pp. 201. University of Aberdeen.

643 Moore, W. S. 1977. Evaluation of Narrow Hybrid Zones in Vertebrates. *Q Rev Biol* **52**: 263-277.

644 Naylor, E. & Haahtela, I. 1966. Habitat preferences and interspersions of species within the
645 superspecies *Jaera albifrons* Leach (Crustacea: Isopoda). *J Anim Ecol* **35**: 209-216.

646 Poelstra, J. W., Vijay, N., Bossu, C. M., Lantz, H., Ryll, B., Muller, I., Baglione, V., Unneberg, P.,
647 Wikelski, M., Grabherr, M. G. & Wolf, J. B. W. 2014. The genomic landscape underlying
648 phenotypic integrity in the face of gene flow in crows. *Science* **344**: 1410-1414.

649 Powell, T. H. Q., Hood, G. R., Murphy, M. O., Heilveil, J. S., Berlocher, S. H., Nosil, P. & Feder, J. L.
650 2013. Genetic Divergence Along the Speciation Continuum: The Transition from Host Race to
651 Species in *Rhagoletis* (Diptera: Tephritidae). *Evolution* **67**: 2561-2576.

652 Pritchard, J. K., Stephens, M. & Donnelly, P. 2000. Inference of population structure using multilocus
653 genotype data. *Genetics* **155**: 945-959.

654 Pritchard, V. L. & Edmands, S. 2013. The Genomic Trajectory of Hybrid Swarms: Outcomes of
655 Repeated Crosses between Populations of *Tigriopus Californicus*. *Evolution* **67**: 774-791.

656 Ribardière, A., Broquet, T. & Daguin-Thiebaut, C. 2015. Microsatellite markers for the *Jaera albifrons*
657 species complex (marine isopods). *BMC research notes* **8**: 632.

658 Rougemont, Q., Gagher, A., Lasne, E., Cote, J., Coke, M., Besnard, A. L., Launey, S. & Evanno, G. 2015.
659 Low reproductive isolation and highly variable levels of gene flow reveal limited progress
660 towards speciation between European river and brook lampreys. *J Evolution Biol* **28**: 2248-
661 2263.

662 Rougemont, Q., Roux, C., Neuenschwander, S., Goudet, J., Launey, S. & Evanno, G. 2016.
663 Reconstructing the demographic history of divergence between European river and brook
664 lampreys using approximate Bayesian computations. *Peerj* **4**.

665 Rousset, F. 2008. genepop'007: a complete re-implementation of the genepop software for Windows
666 and Linux. *Mol Ecol Resour* **8**: 103-106.

667 Seehausen, O. (2009) Progressive levels of trait divergence along a 'speciation transect' in the lake
668 Victoria cichlid fish *Pundamilia*. In: *Speciation and patterns of diversity*, (Butlin, R., Bridle, J. &
669 Schluter, D., eds.). pp. 155-176 Ecological Reviews. Cambridge University Press.

670 Seehausen, O., vanAlphen, J. J. M. & Witte, F. 1997. Cichlid fish diversity threatened by
671 eutrophication that curbs sexual selection. *Science* **277**: 1808-1811.

672 Solignac, M. 1969a. Contiguous distributions sympatry and natural hybridization in a super-species
673 *Jaera Albifrons* Leach (Isopods Asellidae). *Cr Acad Sci D Nat* **268**: 1610-&.

674 Solignac, M. 1969b. Hybridation introgressive dans la population complexe des *Jaera albifrons* de
675 Luc-sur-Mer. *Arch Zool Exp Gen* **110**: 629.

676 Solignac, M. (1978) Nature, déterminisme et origine des mécanismes d'isolement dans le complexe
677 *Jaera albifrons* (Isopodes, Asellotes). Phd Thesis. pp. 232. Université Paris 6.

678 Solignac, M. 1981. Isolating mechanisms and modalities of speciation in the *Jaera albifrons* species
679 complex (*Crustacea, Isopoda*). *Syst Zool* **30**: 387-405.

680 Taylor, E. B., Boughman, J. W., Groenenboom, M., Sniatynski, M., Schluter, D. & Gow, J. L. 2006.
681 Speciation in reverse: morphological and genetic evidence of the collapse of a three-spined
682 stickleback (*Gasterosteus aculeatus*) species pair. *Mol Ecol* **15**: 343-355.

683 Team, R. C. 2016. R: A language and environment for statistical computing.

684 Teeter, K. C., Thibodeau, L. M., Gompert, Z., Buerkle, C. A., Nachman, M. W. & Tucker, P. K. 2010. The
685 Variable Genomic Architecture of Isolation between Hybridizing Species of House Mice.
686 *Evolution* **64**: 472-485.

687 Toews, D. P. L. & Brelsford, A. 2012. The biogeography of mitochondrial and nuclear discordance in
688 animals. *Mol Ecol* **21**: 3907-3930.

689 Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M. & Shipley, P. 2004. Micro-checker: software for
690 identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Notes* **4**: 535-
691 538.

692 Veuille, M. 1976. Biogeography of the *Jaera albifrons* superspecies (Isopoda, Asellota) on the Atlantic
693 coast of Canada. *Can J Zool* **54**: 1235-1241.

694 Westram, A. M., Panova, M., Galindo, J. & Butlin, R. K. 2016. Targeted resequencing reveals
695 geographical patterns of differentiation for loci implicated in parallel evolution. *Mol Ecol* **25**:
696 3169-3186.

697

Table 1: *Jaera albifrons* and *J. praehirsuta* sampling locations, sample sizes, and genetic diversity (observed and expected heterozygosity H_o and H_e , and F_{IS}). Individuals with an intermediate phenotype ($n=9$) from sites 7 and 8 are not included. The statistical significance of F_{IS} is indicated (*: p -value < 0.05)

Sampled site	Coordinates	<i>J. albifrons</i>				<i>J. praehirsuta</i>			
		N	H_o	H_e	F_{IS}	N	H_o	H_e	F_{IS}
BRITTANY									
1	48°40'27.6"N, 3°57'11.3"W	24	0.48	0.55	0.137*	10	0.53	0.61	0.145
2	48°40'20.9"N, 3°57'00.3"W	25	0.47	0.57	0.188*	7	0.55	0.64	0.142
3	48°39'10.8"N, 3°57'03.0"W	22	0.51	0.54	0.070	18	0.54	0.62	0.135*
4	48°39'12.3"N, 3°57'00.4"W	24	0.5	0.57	0.131*	14	0.52	0.62	0.168*
5	48°39'34.1"N, 3°56'25.7"W	25	0.48	0.55	0.115	5	0.63	0.69	0.097
6	48°39'33.5"N, 3°56'31.2"W	22	0.45	0.52	0.132*	26	0.5	0.58	0.153*
NORMANDY									
7	49°23'30.4"N, 1°02'09.6"W	20	0.49	0.52	0.059	12	0.45	0.53	0.163*
8	49°21'15.7"N, 0°47'54.2"W	9	0.42	0.48	0.141	44	0.52	0.58	0.107*
9	49°20'53.5"N, 0°41'03.2"W	15	0.43	0.47	0.094	1	-	-	-

Table 2: Distribution of the genetic variation estimated through hierarchical analyses of molecular variance in regions Brittany and Normandy (as defined in Fig. 1). We present the results obtained with and without loci Ja41 and Ja64, which show a very strong differentiation in Normandy (see F_{CT} at these two loci).

Source of variation	Brittany			Normandy		
	% of total variation	F-stat	p-value	% of total variation	F-stat	p-value ^a
23 loci						
Among sampling sites						
between species	13.2	$F_{CT} = 0.132$	$p = 0.001$	7.4	$F_{CT} = 0.074$	$(p = 0.336)$
within species	1.3	$F_{SC} = 0.015$	$p < 0.001$	1.5	$F_{SC} = 0.016$	$p = 0.003$
Within sampling sites						
among individuals	9.7	$F_{IS} = 0.114$	$p < 0.001$	8.8	$F_{IS} = 0.097$	$p < 0.001$
within individuals	75.8		$p < 0.001$	82.3		$p < 0.001$
21 loci (without Ja41 & Ja64)						
Among sampling sites						
between species	12.5	$F_{CT} = 0.125$	$p = 0.002$	1.7	$F_{CT} = 0.017$	$(p = 0.328)$
within species	1.2	$F_{SC} = 0.013$	$p < 0.001$	1.8	$F_{SC} = 0.019$	$p = 0.001$
Within sampling sites						
among individuals	10.1	$F_{IS} = 0.117$	$p < 0.001$	8.4	$F_{IS} = 0.087$	$p < 0.001$
within individuals	76.3		$p < 0.001$	88.1		$p < 0.001$
2 loci (Ja41 & Ja64)						
Among sampling sites						
between species	20.2	$F_{CT} = 0.202$	$p = 0.003$	42.8	$F_{CT} = 0.428$	$(p = 0.332)$
within species	2.5	$F_{SC} = 0.032$	$p < 0.001$	-0.7	$F_{SC} = -0.012$	$p = 0.65$
Within sampling sites						
among individuals	8.9	$F_{IS} = 0.115$	$p < 0.001$	11.6	$F_{IS} = 0.2$	$p < 0.001$
within individuals	68.4		$p < 0.001$	46.3		$p < 0.001$

a) The statistical significance of between-species variation was tested using permutations of sites between species, which is essentially powerless in region Normandy where only two sites harboring mixed populations were found. The relevant p-values (indicated in grey) are thus meaningless, and the differentiation between species was better tested in this case using non-hierarchical F-statistics within each site (see text).

Table 3: Pairwise genetic differentiation between populations in Brittany. Above diagonal: pairwise F_{ST} . Below diagonal: exact test of population differentiation p -value (in bold when significant). Values in the grey area correspond to inter-specific F_{ST} .

	<i>J. albifrons</i>						<i>J. praeirsuta</i>					
	1	2	3	4	5	6	1	2	3	4	5	6
<i>J. albifrons</i>												
1	-	0.02	0.061	0.06	0.035	0.034	0.178	0.161	0.14	0.14	0.062	0.167
2	0.003	-	0.066	0.055	0.036	0.05	0.175	0.151	0.137	0.141	0.059	0.161
3	0	0	-	0.005	0.021	0.017	0.182	0.177	0.155	0.149	0.099	0.187
4	0	0	0.190	-	0.015	0.024	0.174	0.167	0.141	0.14	0.092	0.179
5	0	0	0.003	0.014	-	0.009	0.207	0.195	0.17	0.166	0.1	0.196
6	0	0	0.008	0.001	0.116	-	0.201	0.184	0.157	0.152	0.103	0.19
<i>J. praeirsuta</i>												
1	0	0	0	0	0	0	-	-0.007	0.01	-0.005	-0.048	0.017
2	0	0	0	0	0	0	0.671	-	-0.005	-0.004	-0.068	0.008
3	0	0	0	0	0	0	0.160	0.715	-	-0.008	-0.061	0.004
4	0	0	0	0	0	0	0.737	0.657	0.919	-	-0.074	0.005
5	0.004	0.003	0	0	0	0	0.979	0.988	0.999	0.999	-	-0.075
6	0	0	0	0	0	0	0.042	0.278	0.295	0.268	1	-

Table 4: Pairwise genetic differentiation between populations in Normandy.

Above diagonal: pairwise F_{ST} . Below diagonal: exact test of population differentiation p -value (in bold when significant). Values in the grey area correspond to inter-specific F_{ST} .

	<i>J. albifrons</i>		<i>J. praehirsuta</i>	
	7	8	7	8
<i>J. albifrons</i>				
7	-	0.025	0.104	0.073
8	0.037	-	0.153	0.101
<i>J. praehirsuta</i>				
7	0	0	-	0.018
8	0	0	0.015	-

Figure legends

Figure 1: Morphological differentiation at pereopods (numbered P1-P7) between males *Jaera albifrons* (carpus of pereopods P6 and P7 extended as a lobe with a number of straight setae) and *J. praehirsuta* (pereopods P1-4 with many curved setae) as found in region Brittany (western France, drawings modified from Solignac, 1981). A few individuals with secondary sexual traits intermediate to *J. albifrons* and *J. praehirsuta* were found in region Normandy. The length of the individual (a female) represented on the picture is 4 mm. Photography credit to Guillaume Evanno & Thomas Broquet.

Figure 2: Sampling sites in two regions of Western France ("Brittany" and "Normandy"). Pie charts give the number of males showing secondary sexual traits typical of *Jaera albifrons* (in blue), *Jaera praehirsuta* (in green), and intermediate phenotypes (in red) sampled at each location. Note that these numbers reflect the relative proportions of each species at each site in Normandy, where the two species occupy the same microhabitat and cannot be distinguished in the field. By contrast, in Brittany pie charts are representative of sample sizes but not necessary of the relative density of each species (because there the two species occupy two different habitats with little overlap, see text). Mixed populations from Normandy were found following an intensive survey (geographic extent shown in yellow, details in text), revealing that the hybridizing population originally studied by Michel Solignac in 1965-1970 (Luc-sur-Mer, indicated by a star) is now extinct. The nearest non-mixed populations that we could find comprised only *Jaera albifrons* (empty circles). Additional mixed populations with some individuals showing an intermediate phenotype were further found in Yport and the Isles of Scilly (black dots, see text).

Figure 3: Locus-by-locus genetic differentiation between *J. albifrons* and *J. praehirsuta* (F_{CT} , expressed in this figure as the percentage of genetic variation due to differences between species) in Brittany (black dots) and in Normandy (white dots). Significant differentiation between $F_{CT-Brittany}$ and $F_{CT-Normandy}$ is represented by a star above the locus name. Here the loci are arranged by increasing order of $F_{CT-Normandy}$. We see that all white dots but two lie near zero (no differentiation between species in Normandy) while the two remaining loci (Ja41 and Ja64) show a strong differentiation. These two microsatellites are the only loci showing a significantly stronger differentiation between species in Normandy than in Brittany (white vs black dots). In Brittany the average level of differentiation between species is near 13% and there is also heterogeneity across loci.

Figure 4: STRUCTURE clusters (K=2) defined in Brittany (A) and Normandy (B). Because loci Ja41 and Ja64 are outliers in Normandy (see text and Fig. 3), STRUCTURE results for this region are presented with all loci (23 loci, panel B1), without the two outliers (21 loci, panel B2), and considering only Ja41 and Ja64 (2 loci, panel B3). Numbers refer to sampling sites (Figure 2). We see that individuals morphologically identified as *J. albifrons* or *J. praehirsuta* cluster into two distinct groups in Brittany (regardless of sampling location, panel A), while this will remain true in Normandy only due to the effect of two markers out of twenty-three (panels B1, B2 and B3).

Figure 5: Principal component analysis based on individual multi-locus genotypes at 23 microsatellite loci. The first axis separates individuals from Brittany (left) and Normandy (right). The second axis shows the genetic divergence between males bearing sexual traits typical of *J. albifrons* vs *J. praehirsuta*. We see that these two types of males are less genetically differentiated in region Normandy (in agreement with results from the analysis of molecular variance and STRUCTURE results from figure 4) and males with an intermediate phenotype are undifferentiated from *J. praehirsuta*.

Figure 1





Figure 2




Figure 3





Figure 4


A) BRITTANY

23 loci ($F_{CT} = 13.2\%$)




B) NORMANDY

B1) 23 loci ($F_{CT} = 7.4\%$)



B2) 21 loci ($F_{CT} = 1.7\%$)



B3) 2 loci (Ja41 & Ja64, $F_{CT} = 42.8\%$)





Figure 5



Supplementary material

Figure S1: Distribution of individuals along the coast at Ste-Honorine-des-Pertes (site 8, Normandy) during a survey realized in summer 2014. Using a precise localization system (5.5 cm horizontal accuracy), we found that male individuals bearing sexual traits typical of *J. albifrons* or *J. praeheirsuta* occupied the same microhabitats and show no particular distribution patterns (both species are intermixed with no apparent distributional gradient). Females were included in this analysis, and some of them were identified as *J. albifrons* or *J. praeheirsuta* based on the sexual characters born by their sons (raised in the lab). For clarity purpose we did not distinguish males and females in this figure. Note also that many individuals are not visible here due to the near superposition of their locations.

Figure S2: Bootstrap distribution of F_{CT} (genetic differentiation between species) in Brittany (black) and Normandy (white) at loci Ja41 (Panel A) and Ja64 (B). Here we see that 1000 resampling iterations rarely produced a situation where $F_{CT-Normandy}$ was greater or equal to $F_{CT-Brittany}$ at these two loci (p -value ≤ 0.001). Observed F_{CT} are indicated by triangles (Brittany: black, Normandy: white).

Figure S3: Bootstrap distribution of F_{CT} (genetic differentiation between species) in Brittany (black) and Normandy (white) at 23 microsatellite loci. Observed F_{CT} in each region are indicated by triangles.

Figure S4: Allelic frequencies at locus Ja41 in *Jaera albifrons* (blue) and *Jaera praeheirsuta* (green) from Brittany (panel A) and Normandy (panel B). Note that the F_{CT} values reported here were calculated in hierarchical analyses of variance without the individuals showing an intermediate phenotype (see main text).

Figure S5: Allelic frequencies at locus Ja64 in *Jaera albifrons* (blue) and *Jaera praeheirsuta* (green) from Brittany (panel A) and Normandy (panel B). Note that the F_{CT} values reported here were calculated in

hierarchical analyses of variance without the individuals showing an intermediate phenotype (see main text).


Figure S6: Patterns of isolation-by distance observed in Brittany for *Jaera albifrons* (blue circles; $y=0.0209x+0.0011$, $R^2 = 0.73$, Mantel test p -value < 0.01) and *J. prae-hirsuta* (green circles; $y=0.0065x-0.0133$, $R^2=0.11$, Mantel test p -value = 0.02).

Figure S1



Figure S2

A



B




Figure S3




Figure S3 (continued)














Figure S4





Figure S5





Figure S6

