

**First occurrence of Mosasauridae (Squamata) in the Maastrichtian (latest Cretaceous) of Alicante (Valencia Community, Eastern Spain)**

N Bardet, J F Baeza Carratala, V. Diez Diaz, A Carbonell, Manuel Garcia Avila, V Giner

► **To cite this version:**

N Bardet, J F Baeza Carratala, V. Diez Diaz, A Carbonell, Manuel Garcia Avila, et al.. First occurrence of Mosasauridae (Squamata) in the Maastrichtian (latest Cretaceous) of Alicante (Valencia Community, Eastern Spain). Estudios Geologicos-Madrid, 2013, 69 (1), pp.97-104. <10.3989/egeol.40792.169>. <hal-01537159>

**HAL Id: hal-01537159**

**<http://hal.upmc.fr/hal-01537159>**

Submitted on 12 Jun 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# First occurrence of Mosasauridae (Squamata) in the Maastrichtian (latest Cretaceous) of Alicante (Valencia Community, Eastern Spain)

*Primer hallazgo de Mosasauridae (Squamata) en el Maastrichtiense (Cretácico final) de Alicante (Comunidad de Valencia, Levante Español)*

N. Bardet<sup>1</sup>, J.F. Baeza Carratalá<sup>2</sup>, V. Díez Díaz<sup>3</sup>, Á. Carbonell<sup>4</sup>, Manuel García Ávila<sup>5</sup>, V. Giner<sup>4</sup>

## ABSTRACT

Although the mosasaurid genus *Prognathodon* is known worldwide during the latest Cretaceous (Campanian-Maastrichtian), we report here its first occurrence in the Eastern area of the Iberian Peninsula. It was previously known from coeval levels of the Basque Country. The specimen from Castalla (Alicante) corresponds to a pterygoid tooth. Though it cannot be precisely determined at the specific level, the tooth belongs to a *Prognathodon* species with 'slender' teeth such as *P. compressidens*, *P. sectorius* and *P. mosasauroides*, all known in the Campanian-Maastrichtian of Europe, or *P. kianda* from the Maastrichtian of Angola.

**Keywords:** Mosasauridae, *Prognathodon*, latest Cretaceous, Betic Cordillera, Spain.

## RESUMEN

Aunque el género de mosasaurio *Prognathodon* está conocido mundialmente durante el Cretácico final (Campaniense-Maastrichtiense), damos a conocer su primera ocurrencia en el Levante Español. Previamente ya se conocía en niveles contemporáneos del País Vasco. El espécimen de Castalla corresponde a un diente del pterigoides. Aunque no se puede identificar a nivel específico, el diente pertenece a una especie de *Prognathodon* con dientes 'delgados' como *P. compressidens*, *P. sectorius* y *P. mosasauroides*, del Campaniense-Maastrichtiense de Europa, ó *P. kianda* del Maastrichtiense de Angola.

**Palabras claves:** Mosasauridae, *Prognathodon*, Cretácico final, Cordillera Bética, España.

## Introduction

Mosasaurid remains are extremely scarce in the latest Cretaceous (Campanian-Maastrichtian) of the Iberian Peninsula and are exclusively known in two areas: the Lusitanian Basin and the Basque-Cantabrian Region (see Bardet *et al.*, 2008).

In Portugal, only few isolated remains consisting of teeth and vertebrae were described by Sauvage (1897-1898). More recently, a few mosasaurid vertebrae have been mentioned (Antunes & Broin, 1988).

In the northern Iberian Peninsula, mosasaurid specimens found in the Campanian-Maastrichtian of the Basque-Cantabrian Basin have been described in

<sup>1</sup> CNRS UMR 7207 CNRS-MNHN-UPMC, Département Histoire de la Terre, Muséum National d'Histoire Naturelle, CP 38, 8 rue Buffon, 75005 Paris, France. Email: bardet@mnhn.fr

<sup>2</sup> Departamento de Ciencias de la Tierra y del Medio Ambiente, Universidad de Alicante, Campus San Vicente del Raspeig s/n, 03690, Alicante, Spain. Email: jf.baeza@ua.es

<sup>3</sup> Departamento Estratigrafía y Paleontología, Facultad de Ciencia y Tecnología, Universidad del País Vasco, Apto 644, 48002 Bilbao, Spain. Email: daniajinn@gmail.com

<sup>4</sup> Asociación Paleontológica Alcoyana Isurus, Pintor Cabrera 61, 03802 Alcoy, Alicante, Spain. Email: asociacion@paleoisurus.com

<sup>5</sup> Asociación Paleontológica Alcarreña Nautilus, Guadalajara.

recent years by Bardet *et al.* (1993, 1997a, 1999, 2006, 2012). The material, which also mainly consists of isolated teeth and vertebrae, represents however the most diverse mosasaurid assemblage found to date in the Iberian Peninsula (Bardet *et al.*, 2008). Here, the genera *Mosasaurus*, *Prognathodon*, *Platecarpus* and *Tylosaurus* have been recognised.

*Prognathodon* has been reported as follows : 1) *Prognathodon* sp. in the Upper Campanian Vitoria Formation of Castillo-Lasarte, Álava (Bardet *et al.*, 1997a) ; 2) *Prognathodon solvayi* Dollo, 1889 and *Prognathodon* sp. in the Upper Maastrichtian of an unnamed unit (lateral equivalent of the Torme Formation) of Albaina, Condado de Treviño (Bardet *et al.*, 1999); 3) *Prognathodon* cf. *sectorius* in the Upper Campanian Vitoria Formation of Olazti (Olazagutia), Navarre (Bardet *et al.*, 2012).

Here we report on the discovery of new mosasaurid specimen, an isolated tooth, from the Late Cretaceous of near Alicante, Valencia Community.

*Institutional abbreviation*—CVAI, Colección Vertebrados Asociación Isurus.

## Geographical and geological setting

The studied specimen came from Sierra de La Argueña (Alicante, SE Spain), a calcareous terrain characterized by a predominance of marine hemipelagic sedimentation during the Late Cretaceous (Fig. 1B). The level where the specimen has been found is located in the Foia Redona locality, an old quarry for the extraction of marls.

This area belongs to the External Zones of the Betic Cordillera, which, during the Mesozoic, formed the south Iberian Paleomargin (Vera, 1988), in an area that previous authors have set in the easternmost Internal Prebetic Domain (Azéma *et al.*, 1979; García Hernández *et al.*, 1980; De Ruig, 1992, Martín-Chivelet, 1992, Chacón, 2002; Chacón & Martín-Chivelet, 2003, among others) (Fig. 1A), and included into the recently individualized Prebetic zone of the Aspe-Jijona-Alicante Sector (Arias *et al.*, 2004).

The stratigraphy of this Prebetic zone during the Cretaceous has been studied in detail by many authors (e.g.: Vera *et al.*, 1982, Martínez del Olmo *et al.*, 1982, Martín-Chivelet, 1992, Chacón, 2002; Chacón & Martín-Chivelet, 2001, 2003, 2005). Recent studies by Chacón & Martín-Chivelet (2005) divided the stratigraphical sequence of this region in different lithostratigraphical formations. Among them, in the




Figure 1—A. Geological sketch showing the studied area within the context of the Betic Cordillera in Alicante Province (modified after Estévez *et al.*, 2004). B. Situation map and stratigraphical setting of Foia Redona section into Cretaceous outcrops in the northern Alicante province.

Sierra de La Argueña, very homogeneous facies are observed; however some successive levels referred to the Maastrichtian could be characterized.

The level that has yielded the mosasaur tooth consists of marly sediments that can be assigned to the Raspay Formation (Martín-Chivelet, 1994), widely outcropping in this eastern part of the Cordillera according to Chacón & Martín-Chivelet (2005), and which is assigned to the upper Maastrichtian. In the studied stratigraphical section, it corresponds to the predominantly marly member deposited into the basin after the middle Maastrichtian discontinuity that implies a change from the underlying carbonate succession to marly hemipelagic sedimentation (Martín-Chivelet *et al.*, 2002; Chacón, 2002; Chacón




Figure 2—Foia Redona lithostratigraphical section. a. pre-Middle Maastrichtian carbonates succession in La Argueña Sierra; b. Outcrop detail showing the ferruginous crusted surface. c, d. Detailed view of olistholiths. e. Marls strata with *Prognathodon* remains. f, g. Outcrop perspective of the Raspay Fm. in Foia Redona site, showing the chaotic appearance because of the olistholiths and extractive activities.

& Martín-Chivelet, 2003, 2005). The paleoenvironmental conditions of these sediments have been interpreted as an open-marine environment with significant terrigenous content, located between the outer shelf and the upper bathyal zone (Chacón & Martín-Chivelet, 2005).

### Stratigraphical data

In the stratigraphic section of Foia Redona where the outcrop occurs (Fig. 2) marly levels are mainly represented from the Raspay Formation (Martín-Chivelet, 1994), overlying the mostly carbonated succession (corresponding to the Aspe or Carche Formation, according to the Prebetic zone in which they crop out; Chacón & Martín-Chivelet, 2005).

The base of the section is alternating greenish marls and yellowish marly/silty limestones, with abundant Fe-oxides and a typical conchoidal fracture. The levels are frequently well-bedded, with decimetric to metric-thick bedding. This bottom interval ends with a marly limestone level overlain by an encrusted ferruginous surface, with abundant bioturbation and a rich, well-preserved macrofauna in which is predominantly corals, bivalves and limonitized nucleus of gastropods and ammonoids. This surface presumably coincides with the middle Maastrichtian discontinuity that marks the transition to marly facies in the basin (Chacón & Martín-Chivelet, 2003, 2005, among others).

Overlying this level, there is a thick set of grey and greenish marly beds evolving upwards to darker marls; the underlying ferruginous surface is often found reworked in the basal tract of this levels.

The entire marly tract has a chaotic appearance. The apparent homogeneity is disrupted by the presence of numerous olistholiths. In the studied area reworking due to old mining activities must also be taken into account.

The olisthons consist of beige marly limestones, with wackstone-packstone texture; they are rounded and decimetric to metric in diameter and include small echinoderms (*Salenia* sp.), agglutinated benthic macroforaminifera, and bryozoans. This association is also found in the marls embedding the olistholiths, which also include *Isocrania* sp., *Magas* sp., bivalves, serpulids and small selachian teeth.

The first marly tract with olistholiths is followed by a second level of similar lithology, but lighter in colour, in which the studied tooth has been found,

associated with a rich fauna of echinoderms in which *Echinocorys* sp. and *Cyclaster* sp. dominate.

In these marls, interbedded greenish to yellow marly levels are present and are about 50 cm thick. The marly sediments become darker upwards, and in their upper part, the marly limestones levels are well-bedded and acquire boudinaged morphology, and lighter shades.

The Foia Redona section finishes with a level of beige marls on which lays a metric-thick distinctive strata which lithology corresponds to yellow calcarenites with basal parallel lamination.

In addition to the data arising from the recorded macrofauna, the age of the deposits has been contrasted with the several sampling (samples FR-1 to 3 in Fig. 2) that have yielded a rich microfauna of foraminifera. Samples were taken from immediately above and below the level in which the mosasaur tooth was found. Both samples preserve a similar fauna. The underlying assemblage is dominated by Heterohelicids, Contusotruncanids, the notable occurrence of *Planoglobulina acervulinoides* and occasional ostracods. The overlying assemblage includes numerous Heterohelicids, several Globotruncanid species, *Rugoglobigerina* cf. *hexacamerata*, occasional *Racemiguembelina fructicosa*, *Globigerinelloides* sp. and few ostracods. The younger recorded assemblage includes almost the same microfauna but with an increase in the presence of *R. fructicosa*. In all samples the benthic foraminifera are lesser extent.

By comparing these data with the zonation proposed by Premoli-Silva and Sliter (2002) and the precise biostratigraphical data of Chacón (2002), and Chacón and Martín-Chivelet (2003, 2005) for the Raspay Formation, the assemblages found in the Foia Redona section can be assigned to the Late (but not latest) Maastrichtian, presumably to the upper part of the *Gansserina gansseri* Zone but not reaching the uppermost part of the Maastrichtian as the index fossil of this interval *Abathomphalus mayaroensis* having not been recorded.

### Systematic palaeontology

Squamata Oppel, 1811

Mosasauridae Gervais, 1853

*Prognathodon* Dollo, 1889

*Prognathodon* sp.

**Material:** CVAI 00141, an isolated pterygoid tooth crown (Fig. 3).




Figure 3—*Prognathodon* sp., CVAI 00141, pterygoid tooth crown; Raspay Formation; Late Maastrichtian (*Gansserina gansseri* Zone?); latest Cretaceous); Castalla (Alicante), Valencia Community, Eastern Spain. A, labial view; B, lingual view; C, anterior view; D, posterior view; E, apical view. Scale = 1 cm.

**Geographical occurrence:** Font de la Carrasca Quarry, East of Castalla, northwest of Alicante Province, Community of Valencia, Eastern Spain. UTM: 30S 6992204273190 (Fig. 1).

**Stratigraphical occurrence:** Raspay Formation; Late Maastrichtian, probably in the upper part of the *Gansserina gansseri* Zone (Fig. 2).

### Description

The tooth preserves only the enamel crown which is 23.8 mm high, 16 mm long and 11.4 mm wide. In lateral view (Figs. 3A-B), the crown is a robust posteriorly recurved triangle, with a pointed apex, a convex anterior surface and a slightly concave posterior one. The crown is compressed labiolingually and the basal cross-section is teardrop shaped (Figs. 3C-E). The labial surface is slightly convex whereas the lingual one is almost flat (Figs. 3C-D). There is only a posterior marked carina slightly displaced laterally (Fig. 3C). It seems to be 'pinched' from the main shaft and bears minute serrations. The anterior surface is regularly curved without any carina. The enamel is completely smooth and has a shiny aspect.

Because of the lack of anterior carina, it could correspond either to an anterior marginal tooth or to a pterygoid one. Usually, the anterior marginal teeth have a slender appearance than the pterygoid teeth, as they are higher than long (height twice the length *versus* less than twice high than long—compared

Fig. 3 to Fig. 5 of Schulp *et al.*, 2008). The Castalla crown is thus considered here as a pterygoid tooth.

### Comparisons and systematic attribution

Mosasaurid teeth (both marginal and pterygoid ones) are generally highly diagnostic at both generic and even specific level (Russell, 1967, Lindgren & Siverson, 2002) so that even isolated teeth can be identified taxonomically.

The absence of medial striae on the tooth excludes the Castalla tooth from the Russellosaurinae (Bell, 1997; Russellosaurina of Bell & Polcyn, 2005). Among Mosasaurinae, its relatively large size and robustness, as well as the occurrence of a completely smooth enamel, support its attribution to the *Globidensini* (*sensu* Bell, 1997; Bell & Polcyn, 2005) taxon *Prognathodon*, the only genus of this clade devoid of low blunt teeth, contrary to *Globidens* and *Carinodens* (Schulp *et al.*, 2004).

The combination of the following characters, that are, a crown moderately posteromedially recurved, with subequal convex lingual and labial surfaces, a fairly well marked 'pinched' posterior carina, and a smooth shiny enamel, permits to refer the Castalla's mosasaurid to *Prognathodon* Dollo, 1889. This genus includes about ten species from the Campanian-Maastrichtian of many parts of the world, including Europe, North America, Africa, Middle-East and New Zealand (see Schulp *et al.*, 2008).

*Prognathodon* exhibits a large tooth morphology variation interval. The Castalla specimen is clearly distinguishable from the strongly faceted teeth of the type species *P. solvayi* Dollo, 1889 from the Maastrichtian of Belgium (Lingham-Soliar & Nolf, 1989), as well as from species with large robust blunt teeth ornamented by a coarse thick ‘anastomosed’ enamel, that are, *P. currii* from the Maastrichtian of Negev and Morocco (Christiansen & Bonde, 2002; Bardet *et al.*, 2005), *P. giganteus* from the Campanian-Maastrichtian of Europe, Syria and Morocco (Lingham-Soliar & Nolf, 1989; Bardet *et al.*, 1997b; Bardet *et al.*, 2000, 2010), *P. overtoni* (Williston, 1897) from the Campanian of South Dakota (Lingham-Soliar & Nolf, 1989; Schulp, 2006), *P. saturator* Dortangs *et al.*, 2002 from the Maastrichtian of The Netherlands (Dortangs *et al.*, 2002), *P. waiparaensis* Welles & Gregg, 1971 from the Maastrichtian of New Zealand, (Welles & Gregg, 1971), and a new, yet undescribed species from the Maastrichtian of Morocco (N.B., pers. obs.). Though the Castalla tooth resembles in general shape a tooth from the Maastrichtian of Normandy (Northwestern France) referred to as *Prognathodon* sp., this one is however distinctly more robust, bears an anterior carina (though slight) and a thick ‘anastomosed’ enamel (Buffetaut & Bardet, 2012).

As a whole, the Castalla tooth general appearance is more reminiscent of that of *Prognathodon* species possessing ‘slender teeth’, such as *P. compressidens* (Gaudry, 1892) from the Campanian of France (Schulp *et al.*, 2008), *P. mosasauroides* (Gaudry, 1892) from the Maastrichtian of France (Schulp *et al.*, 2008), *P. kianda* Schulp *et al.*, 2008 from the Maastrichtian of Angola (Schulp *et al.*, 2008), and *P. sectorius* (Cope, 1871) from the Maastrichtian of New-Jersey, The Netherlands and the Basque Country (Schulp *et al.*, 2008, Bardet *et al.*, 2012). As a whole, these species bear marginal teeth labiolingually compressed with completely smooth and shiny enamel. Those of *P. compressidens* are the smallest and are slender and notably posteriorly recurved. Those of *P. mosasauroides* are very large and compressed, acute, with a straight posterior surface. Those of *P. kianda* are the most slender whereas those of *P. sectorius* are the most robust.

Unfortunately, pterygoid teeth are unknown in *P. compressidens*, *P. mosasauroides* and *P. sectorius*, so that comparisons are only possible with *P. kianda*. The morphology of the largest preserved pterygoid teeth of this species fits pretty well with that of the Castalla one (compare Fig. 2 to Fig. 5 of Schulp *et al.*, 2008). However, as comparisons cannot be made

with the three species above mentioned, we cannot confidently refer the Castalla tooth to this African species so that it appears safer to refer it only to *Prognathodon* sp.

## Conclusion

The specimen from Castalla (Alicante) provides additional evidence of the potential richness of the Iberian Peninsula in mosasaurid remains, though they currently consist mainly on isolated teeth only. It also confirms once more that *Prognathodon* was a cosmopolitan predator during Campanian-Maastrichtian times. In Europe, several species of *Prognathodon* have been described, that are the Campanian *P. compressidens*, the Campanian-Maastrichtian *P. giganteus*, and the Maastrichtian *P. mosasauroides*, *P. saturator*, *P. sectorius* and *P. solvayi*. However, either the Castalla tooth does not fit in general morphology with some of them (*P. solvayi*, *P. giganteus*, *P. saturator*), or pterygoid teeth are unknown in other ones preventing direct comparison (*P. compressidens*, *P. mosasauroides*, *P. sectorius*). Only *P. kianda*, a species with ‘slender’ teeth from the Maastrichtian of Angola, is suitable for comparisons and shows possible affinities. However, due to the incompleteness of the data precluding for a specific assignment, the Castalla’s specimen is here referred to *Prognathodon* sp. *Prognathodon* was previously known in the Iberian Peninsula by remains found in the Campanian-Maastrichtian of the Basque Country referred to *P. solvayi*, *P. cf. sectorius* and *Prognathodon* sp.

## ACKNOWLEDGEMENTS

The present paper is supported by the Research Group VIGROB-167 of the University of Alicante (J.F.B.C.). The tooth has been donated by one of the authors (M.G.A.) for its study. We thank Anne Schulp (Maastricht, The Netherlands) and Mike Polcyn (Dallas, Texas, USA) for their constructive reviews that permitted to improve the manuscript.

## References

- Antunes, M.T. & Broin, F. de (1988). Le Crétacé terminal de Beira Litoral, Portugal: remarques stratigraphiques et écologiques, étude complémentaire de *Rosasia soutoi* (Chelonii, Bothremydidae). *Ciências Terra*, 9: 153-200.
- Arias, C.; Castro, J.M.; Chacón, B.; Company, M.; Crespo-Blanc, A.; Díaz de Federico, A.; Estévez, A.; Fernán-

- dez, M., García-Hernández, M.; De Gea, G. A.; López Garrido, A.C.; Martín-Algarra, A.; Martín-Chivelet, J.; Molina, J.M.; Morata, D.; Nieto, L.M.; O'Dogherty, L.; Pérez-López, A.; Puga, E.; Rey, J.; Rivas, P.; Ruiz-Ortiz, P.A.; Sandoval, J.; Sanz de Galdeano, C.; Vera, J.A. & Vilas, L. (2004). Zonas Externas Béticas. In: *Geología de España* (Vera, J.A., Ed.). Sociedad Geológica de España, I.G.M.E, 354-372.
- Azéma, J.; Foucault, A.; Fourcade, E.; García-Hernández, M.; González Donoso, J.M.; Linares, A.; Linares, D.; Lopez-Garrido, A.C.; Rivas, P. & Vera, J.A. (1979). *Las microfácies del Jurásico y Cretácico de las Zonas Externas de las Cordilleras Béticas*. Secretariado de Publicaciones de la Universidad de Granada, 172 pp.
- Bardet, N.; Corral, J.C. & Pereda Suberbiola, J. (1993). Primeros restos de reptiles marinos en el Cretácico superior de la Cuenca Vasco-Cantábrica. *Estudios del Museo de Ciencias Naturales de Álava*, 8: 27-35.
- Bardet, N.; Corral, J. C. & Pereda Suberbiola, X. (1997a). Les mosasaures (Squamata) du Crétacé supérieur du Bassin Basco-Cantabrique. *Geobios*, 20: 19-26. doi: 10.1016/S0016-6995(97)80005-8
- Bardet, N.; Barbin, V.; Laurain, M. & Janin, M.C. (1997b). Première découverte du mosasaure *Prognathodon giganteus* (Squamata) dans le Campanien (Crétacé supérieur) de Champagne, France. *Revue de Paléobiologie*, 16 (1): 225-230.
- Bardet, N.; Corral, J.C. & Pereda Suberbiola, X. (1999). Marine reptiles from the uppermost Cretaceous of the Laño quarry (Iberian Peninsula). *Estudios del Museo de Ciencias Naturales de Álava*, 14 (Número especial 1): 373-380.
- Bardet, N.; Cappetta, H.; Pereda Suberbiola, X.; Mouty, M.; Al Maleh, A.K.; Ahmad, A.M.; Khrata, O. & Gannoun, N. (2000). The marine vertebrate faunas from the Late Cretaceous phosphates of Syria. *Geological Magazine*, 137: 269-290. doi: 10.1017/S0016756800003988
- Bardet, N.; Pereda Suberbiola, X.; Iarochène, M.; Amalik, M. & Bouya, B. (2005). Durophagous Mosasauridae (Squamata) from the Upper Cretaceous Phosphates of Morocco, with the description of a new species of *Globidens*. *Netherlands Journal of Geosciences*, 84(3):167-175.
- Bardet, N.; Pereda Suberbiola, X. & Corral, J.C. (2006). A tylosaurine Mosasauridae (Squamata) from the Late Cretaceous of the Basque-Cantabrian Region. *Estudios Geológicos*, 62: 213-218. doi: 10.3989/egeol.0662121.
- Bardet, N.; Pereda Suberbiola, X. & Ruíz-Omeñaca, J.I. (2008). Mesozoic marine reptiles from the Iberian Peninsula. *Geo-Temas*, 10: 1245-1248.
- Bardet, N.; Pereda Suberbiola, X.; Jouve, S.; Bourdon, E.; Vincent, P.; Houssaye, A.; Rage, J.C.; Jalil, N.E.; Bouya, B. & Amaghaz, M. (2010). Reptilian assemblages from the latest Cretaceous – Palaeogene phosphates of Morocco: from Arambourg to present time. *Historical Biology*, 22(1-3): 186-199. doi: 10.1080/08912961003754945
- Bardet N.; Pereda Suberbiola, X.; Carmelo Corral, J.; Baceta, J.I.; Torres, J.Á.; Botantz, B. & Martin, G. (2012). A skull fragment of the mosasaurid *Prognathodon* cf. *sectorius* from the Late Cretaceous of Navarre (Basque-Cantabrian Region). *Bull. Soc. Geol. France*, 183(2): 115-119. doi: 10.2113/gssgfbull.183.2.117
- Bell, G.L. Jr. (1997). Mosasauridae. Introduction. In: *Ancient Marine Reptiles* (Callaway, J.M. & Nicholls, E.L., eds.). Academic Press, San Diego, 281-292.
- Bell, G.L. Jr. & Polcyn, M.J. (2005). *Dallasaurus turneri*, a new primitive mosasauroid from the Middle Turonian of Texas and comments on the phylogeny of Mosasauridae (Squamata). *Netherlands Journal of Geosciences*, 84(3): 177-194.
- Buffetaut, E. & Bardet, N. (2012). The mosasaurid (Squamata) *Prognathodon* in the Maastrichtian (Late Cretaceous) of the Cotentin (Normandy, north-western France). *Bull. Soc. Geol. France*, 183(2): 109-113. doi: 10.2113/gssgfbull.183.2.111
- Chacón, B. (2002) *Las sucesiones hemipelágicas del final del Cretácico e inicio del Paleógeno en el SE de la Placa Ibérica: Estratigrafía de eventos y evolución de la cuenca*. Doctoral thesis, Universidad Complutense Madrid, 440 p.
- Chacón, B. & Martín-Chivelet, J. (2001) Implicaciones tectosedimentarias de la discontinuidad estratigráfica del Maastrichtiense medio en Aspe (Prebético de Alicante). *Revista de la Sociedad Geológica de España*, 14 (1-2): 123-133.
- Chacón, B. & Martín-Chivelet, J. (2003) Discontinuidades estratigráficas regionales en las sucesiones hemipelágicas finicretácicas del Prebético (sector Jumilla-Callosa-Aspe). *Journal of Iberian Geology*, 29: 89-109.
- Chacón, B. & Martín-Chivelet, J. (2005) Subdivisión litoestratigráfica de las series hemipelágicas de edad Coniaciense-Thanetiense en el Prebético Oriental (SE de España). *Revista de la Sociedad Geológica de España*, 18 (1-2): 3-20
- Christiansen, P. & Bonde, N. (2002). A new species of gigantic mosasaur from the Late Cretaceous of Israel. *Journal of Vertebrate Paleontology*, 22: 629-644. doi: 10.1671/0272-4634(2002)022[0629:ANSOGM]2.0.CO;2
- Cope, E.D. (1871). Supplement to the "Synopsis of the extinct Batrachia and Reptilia of North America". *Proceedings of the American Philosophical Society*, 12: 41-52.
- De Ruig, M.J. (1992) *Tectono-sedimentary evolution of the Prebetic fold belt of Alicante (SE Spain)*. Doctoral thesis, Free University of Amsterdam, 207 pp.
- Dollo, L. (1889). Note sur les vertébrés récents offerts au Musée de Bruxelles par M. Alfred Lecomnier. *Bulletin de la Société belge de Géologie, de Paléontologie et d'Hydrologie*, 3: 181-182.
- Dortangs, R.W.; Schulp, A.S.; Mulder, E.W.A.; Jagt, J.W.M.; Peeters, H.H.G. & de Graaf, D.T.H. (2002). A large new mosasaur from the Upper Cretaceous of The Netherlands. *Netherlands Journal of Geosciences*, 81: 1-8.
- Estévez, A., Vera, J.A., Alfaro, P.; Andreu, J.M., Tent-Manclús, J.E. & Yébenes, A. (2004) Alicante en la Cordillera Bética. In: *Geología de Alicante* (Alfaro, P.; Andreu, J.M., Estévez, A., Tent-Manclús, J.E. & Yébenes, A., eds.), Universidad de Alicante: 39-50.



- García-Hernández, M.; López-Garrido, A.C., Rivas, P., Sanz de Galdeano, C., & Vera, J.A. (1980) Mesozoic Paleogeographic evolution of the External Zones of the Betic Cordillera. *Geologie en Mijnbouw*, 59: 155-168.
- Gaudry, A. (1892). Les Pythonomorphes de France. *Mémoires Société Géologique de France*, 10: 13 pp, 2 pl.
- Gervais, P. (1853). Observations relatives aux reptiles fossiles de France. *Comptes Rendus de l'Académie des Sciences Paris*, 36: 374-377, 470-474.
- Lingham-Soliar, T. & Nolf, D. (1989). The mosasaur *Prognathodon* (Reptilia, Mosasauridae) from the Upper Cretaceous of Belgium. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Sciences de la Terre*, 59: 137-190.
- Martín-Chivelet, J. (1992) *Las plataformas carbonatadas del Cretácico superior de la Margen Bética (Altiplano de Jumilla-Yecla, Murcia)*. Doctoral thesis, Universidad Complutense Madrid, 899 pp.
- Martín-Chivelet, J. (1994) Litoestratigrafía del Cretácico superior del Altiplano de Jumilla-Yecla (Zona Prebética). *Cuadernos de Geología Ibérica*, 18: 117-173.
- Martín-Chivelet, J.; Berástegui, X.; Rosales, I.; Vilas, L.; Vera, J.A.; Caus, E.; Gräfe, K.-U.; Segura, M.; Puig, C.; Mas, R.; Robles, S.; Floquet, M.; Quesada, S.; Ruiz-Ortiz, P.A.; Fregenal-Martínez, M.A.; Salas, R.; García, A.; Martín- Algarra, A.; Arias, C.; Meléndez, N.; Chacón, B.; Molina, J.M.; Sanz, J.L.; Castro, J.M.; García-Hernández, M.; Carenas, B.; García-Hidalgo, J.; Gil, J. & Ortega, F. (2002) Cretaceous. In: *Geology of Spain* (Gibbons, W. y Moreno, M.T., eds.). Geological Society of London, 255-292.
- Martínez del Olmo, W.; Leret, G. & Megías, A. G. (1982) El límite de la plataforma carbonatada del Cretácico Superior en la zona prebética. *Cuadernos de Geología Ibérica*, 8: 597-614.
- Oppel, M. (1811). *Die Ordnungen, Familien und Gattungen der Reptilien als Prodrom einer Naturgeschichte derselben*. Joseph Lindauer, München, 87 pp. doi: 10.5962/bhl.title.4911
- Premoli-Silva, I. & Slitter, W. V. (2002) *Practical manual of Cretaceous planktonic foraminifera* (Premoli-Silva, I. & Rettori, R., eds.). University of Perugia, Italy, 462 pp.
- Russell, D.A. (1967) Systematics and Morphology of American Mosasaurs. *Bulletin of the Peabody Museum of Natural History, Yale University*, 23: 1-241.
- Sauvage, H.E. (1897-1898). Les Vertébrés fossiles du Portugal. Contributions à l'étude des poissons et des reptiles du Jurassique et du Crétacé. *Mémoires et Communications du Service géologique du Portugal*, 1-46.
- Schulp, A.S. (2006). On Maastricht Mosasaurs. *Publicaties van het Natuurhistorisch Genootschap in Limburg*, 45(1): 1-140.
- Schulp, A.S.; Jagt, J.W.M. & Fonken, F. (2004). New material of the mosasaur *Carinodens belgicus* from the Upper Cretaceous of The Netherlands. *Journal of Vertebrate Paleontology*, 2: 744-747. doi: 10.1671/0272-4634(2004)024[0744:NMOTMC]2.0.CO;2
- Schulp, A.; Polcyn, M.; Mateus, O.; Jacobs, L. & Morais, M.L. (2008). A new species of *Prognathodon* (Squamata, Mosasauridae) from the Maastrichtian of Angola, and the affinities of the mosasaur genus *Liodon*. In: *Proceedings of the Second Mosasaur Meeting*, Fort Hays Studies, Special Issue(3): 1-12.
- Vera, J.A. (1988) Evolución de los sistemas de depósito en el margen Ibérico de la Cordillera Bética. *Revista de la Sociedad Geológica de España*, 1(3-4): 373-391.
- Vera, J.A.; García-Hernández, M.; López Garrido, A.C.; Comas, M.J.; Ruíz-Ortiz, P.A. & Martín-Algarra, A. (1982) La Cordillera Bética. In: *El Cretácico de España*. (García, A., ed.). Madrid. Universidad Complutense, 515-631.
- Welles, S.P. & Gregg D.R. (1971). Late Cretaceous marine reptiles of New Zealand. *Records of the Canterbury Museum*, 9: 1-111.
- Williston, S.W. (1897). *Brachysaurus, a new genus of mosasaurs*. *Kansas University Quarterly*, 6: 95-98.

Recibido el 6 de septiembre de 2011

Acceptado el 30 de marzo de 2012

Publicado online el 16 de julio de 2012