



Parental comprehension of the benefits/risks of first-line randomised clinical trials in children with solid tumours: a two-stage cross-sectional interview study

Hélène Chappuy, Naim Bouazza, Veronique Minard-Colin, Catherine Patte, Laurence Brugières, Judith Landman-Parker, Anne Auvrignon, Dominique Davous, Hélène Pacquement, Daniel Orbach, et al.

► To cite this version:

Hélène Chappuy, Naim Bouazza, Veronique Minard-Colin, Catherine Patte, Laurence Brugières, et al.. Parental comprehension of the benefits/risks of first-line randomised clinical trials in children with solid tumours: a two-stage cross-sectional interview study. *BMJ Open*, 2013, 3 (5), pp.e002733. 10.1136/bmjopen-2013-002733 . hal-01548392

HAL Id: hal-01548392

<https://hal.sorbonne-universite.fr/hal-01548392>

Submitted on 27 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Parental comprehension of the benefits/risks of first-line randomised clinical trials in children with solid tumours: a two-stage cross-sectional interview study

Hélène Chappuy,¹ Naim Bouazza,² Veronique Minard-Colin,³ Catherine Patte,³ Laurence Brugières,³ Judith Landman-Parker,⁴ Anne Auvrignon,⁴ Dominique Davous,⁵ Hélène Pacquement,⁶ Daniel Orbach,⁶ Jean Marc Tréluyer,⁷ François Doz⁶

To cite: Chappuy H, Bouazza N, Minard-Colin V, *et al*. Parental comprehension of the benefits/risks of first-line randomised clinical trials in children with solid tumours: a two-stage cross-sectional interview study. *BMJ Open* 2013;**3**:e002733. doi:10.1136/bmjopen-2013-002733

► Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2013-002733>).

Received 15 February 2013

Revised 27 March 2013

Accepted 15 April 2013

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

For numbered affiliations see end of article.

Correspondence to

Dr Hélène Chappuy;
helene.chappuy@nck.aphp.fr

ABSTRACT

Objectives: To analyse the parental understanding of informed consent information in first-line randomised clinical trials (RCTs) including children with malignant solid tumours and to assess parents' needs for decision-making.

Design: Observational prospective study.

Setting: 3 paediatric oncology centres in the Parisian region in France.

Participants: 53 parents were approached to participate in a RCT for their child with malignant solid tumour, over a 1-year period. 40 parents have been interviewed in our study.

Primary and secondary outcome measures:

Parental understanding of information in RCTs, parents' needs for decision-making. Parents were questioned by a psychologist, independent of the paediatric oncology teams, using a semidirected interview, 1 (M1) and 6 months (M6) after the consent was sought.

Results: 18 parents (45%) did not understand the concept of randomisation. Half of the parents could explain neither the aim of the clinical trial nor the potential benefit to their child of inclusion. 35 parents (87.5%) expressed very few specific risks related to the trial. Being mostly French-speaking ($p=0.03$) and the reading of the information sheet by the parents ($p=0.0025$) improved their understanding. The parental comprehension did not differ between M1 and M6. The principal factors underlying their decision were confidence in the medical team (39%), wish to access to the best treatment (37%) and the best quality of life (37%).

Conclusions: Despite medical explanations, parents have limited knowledge in some areas in first-line RCTs and improvements of information process are required. The risks specific to the randomised trial are underestimated by parents and the unproven nature of the treatment is not well-known or understood.

ARTICLE SUMMARY

Article focus

- How is the parental comprehension of the benefits/risks to their child with a solid tumour for the inclusion in a randomised phase 3 trial?
- Do perceived risks and benefits figure prominently in the parental decision about randomised phase 3 trial participation?

Key messages

- Most parents had understood the general aspects, by contrast, less than half the parents had correctly understood the more specific information such as the aims of the study, randomisation, the various phases of treatment and the risks specific to the randomised trial.
- Parents tend to underestimate risks specific to the randomised trial, and the unproven nature of the treatment is not well-known or understood.
- Quality of life could be a crucial element in the decision of parents to agree to the participation of their child in a randomised phase 3 trial.

Strengths and limitations of this study

- We have data that reflect the full spectrum of paediatric oncology clinical research, including children with solid or brain tumours, whereas most of the available paediatric data are related to research for newly diagnosed acute leukaemia.
- The parents were asked to participate in this study closely to having consented. Thus, the risk of recall bias was limited.

INTRODUCTION

Cancers can occur before the age of 15 years in 1 out of every 500 children and 1700 new cases are recorded each year for this age

Parental comprehension in randomised clinical trials

group in French registries (<http://sfce1.sfpediatric.com/>). Although global cure rates in paediatric cancers today achieve 80% in western countries, cancers are still the second leading cause of death in children between 1 and 15 years, after accidents. The optimisation of treatment in paediatric oncology is desirable not only to continue to improve cure rates but also to decrease late effects. In this setting, randomised clinical trials (RCTs) to test new treatment options are essential in paediatric oncology.

The legislation requires that parents give permission for a minor to enroll in a study, and it is important for researchers to increase their understanding of how parents arrive at a decision in this matter.^{1 2} The balancing of potential risks and benefits could be critical for parental decisions about allowing their children to participate in research studies and understanding how people weigh these factors is a key point. Boccia *et al*³ suggested that perceived risk and benefit figure prominently in a decision about research. The patients and their parents must, therefore, necessarily receive appropriate information about the possible risks and the uncertain nature of the benefits of participation to allow them to make truly autonomous decisions about participation in trials. They also require adequate information concerning crucial aspects of RCTs if they are to make an informed decision. However, published findings indicate that parents or patients often misunderstand the research to which they consent.^{4–7} Furthermore, most of the available paediatric data on the information process in the context of phase 3 trials in oncology are related to research for newly diagnosed acute leukaemia^{8–10}; few data are available that reflect the full spectrum of paediatric oncology clinical research, including children with solid or brain tumours.

The primary aim of this study was to analyse parental understanding of oral and written information in RCTs including children with varied solid tumours. The secondary aim was to assess the parents' statements in order to identify parental needs for decision-making.

METHODS

Sample population

This prospective study was carried out in the oncology departments of the three paediatric oncology centres in the Parisian region—Institut Curie (Paris), Institut Gustave Roussy (Villejuif) and Hôpital Armand Trousseau (Paris). This study was approved by the appropriate institutional review board (*Comité de Protection des Personnes Ile de France III Hôpital Cochin*). The parents from these three centres who had been asked to give consent for the participation of their child in a phase 3 randomised trial were included in this study, over a 1-year period. The doctors provided the parents with a copy of the information and consent forms and a summary diagram of the various phases of the research protocol. To avoid bias owing to the assessment only of patients with a positive attitude to RCTs, we also included the

parents of children who had declined to participate in the trial. The characteristics of these randomised phase 3 trial protocols are described in [table 1](#): most were trials aiming to decrease the toxicity of treatment without affecting its efficacy. With the agreement of physicians participating in the study, the physician seeking parental consent suggested for the participation in our study to the parents. Non-inclusion criteria included consent in a language other than French. Families agreeing to participate were seen at the time of a planned consultation during routine follow-up or at the time of a hospital admission in order to minimise the additional constraints (time and travel) imposed on the parents.

Instrument

The parents were seen twice by a qualified psychologist: 1 month (M1) and 6 months (M6), after they were asked to participate in a RCT. These interviews took place on site and were recorded and transcribed in their entirety. The parents were allowed to express themselves freely, in the framework of a semidirected interview, in response to standardised questions. The structure for the semidirected interviews was developed in collaboration with psychologists, the parent of a former patient representing a patients' association and some of the investigating doctors from the centres participating in this study. This structure has been validated in previous studies.^{8 11} The interview covered various aspects of informed consent (IC); we evaluated parental comprehension based on the 11 elements included in the information sheet for each protocol. The questions asked to assess parental comprehension are described in [table 2](#).

We tried to identify the elements predictive of a good understanding of the information provided at the time at which consent was sought: parental socioprofessional category (high standing/intermediate status/no profession), the principal language spoken, whether the information document was read by the parents (yes/no), personal efforts to find information (yes/no). We used the following questions to determine the reasons for which the parents had given their consent: "How difficult was it to take the decision you took concerning the participation of your child in this protocol?", "What were the principal elements underlying your decision?", "Who do you feel took the final decision?" and "What do you expect from the doctor?".

Data analysis

Each interview was independently coded by two psychologists, with the coding reviewed by a paediatrician in cases of disagreement. Comprehension was classified as 'complete', 'partial' or 'null' for each element if all, some or none of the information sheet was expressed by the parents. When considering risks, only those specific to the randomised trial were considered ([table 1](#)). To test the influence of covariates on comprehension, partial and complete comprehension were pooled for each of the 11 elements and compared with the absence

Table 1 Description of the protocols studied according to the information leaflets

n	Disease	Main objective	Individual benefit	Randomisation	Treatment duration	Specific risks	Alternative
17	Retinoblastoma	Conservative treatment, reduction of adverse effects	Conservative treatment, reduction of adverse effects	Standard chemotherapy vs new regimen	4–6 months	Auditory, renal, consciousness, pains in the limbs and jaw	Standard CT, enucleation
4	Ewing's sarcoma	R1: reduction of toxicity R2: evaluation of the efficacy	Lower renal toxicities	Standard CT vs new regimen	Not specified	Renal, decrease in sperm count/fertility++	Standard CT
7	High-risk neuroblastoma	Comparison of efficacy and toxicity of two high-dose chemotherapy regimens	Same efficacy, but fewer adverse effects	High-dose CT vs another high-dose CT	Not specified	Hepatic, thyroid, renal, auditory, endocrine, fertility	Standard CT
2	Low-grade glioma	Evaluation of the efficacy of a new chemotherapy combination	Best possible CT	Standard CT vs the new CT combination	81 weeks	Secondary induction of leukemia, haematological and infectious toxicity	Standard CT
4	Standard-risk medulloblastoma	Evaluation of the efficacy of hyperfractionated RT and reduction of toxicity	Best possible RT, lower long term toxicity	Classical RT dose vs hyperfractionation	16 months	Neuro-cognitive impairment and endocrine problems	Classical fractionation of RT
5	Localised nephroblastoma	Assessment of the equivalence of two CT regimens	Same efficacy, but fewer adverse effects	Standard CT vs new regimen	25–30 weeks	Cardiac	Standard CT
1	Standard-risk hepatoblastoma	Reduction of toxicity	Same efficacy, but fewer adverse effects	Standard CT vs new regimen	3–5 months	Cardiac	Standard CT

n, number of children included in this study; CT, chemotherapy; RT, radiotherapy.

Parental comprehension in randomised clinical trials

Table 2 Questions asked during the interview addressing the level of understanding

No.	Concept	Question
1	Participation in a research protocol	'Is your child being treated as part of a research protocol?'
2	Aim of the protocol	'What is the aim of this protocol?'
3	Course of the protocol	'What is planned for your child in the framework of this protocol?'
4	Principle of randomisation	'If you gave consent for a protocol in which two different treatments might be given, do you know how the treatment given to your child was chosen? If yes, how?'
5	Individual benefit	'What benefits do you expect your child to gain from participation in this protocol?'
6	Collective benefit	'Could you describe the possible benefits to other children of the participation of your child in this protocol?'
7	Risks	'What are the possible risks to your child of participating in this protocol?'
8	Alternatives	'If you had not consented to the participation of your child in this protocol, what care would your child have received?'
9	Voluntary nature of participation	'Was the participation of your child in this protocol voluntary?'
10	Duration of participation	'How long were you told that the participation of your child in this protocol would last?'
11	Freedom to withdraw from the project at any time	'Could you change your mind once the study had begun?'

of comprehension. We evaluated the influence of the covariates on the percentage of the elements understood by the parents. A mixed effects model taking into account the subject and items as crossed random effects was used to assess the stability of parental comprehension between the two interviews at M1 and M6. The data collected were put into an Access database. The results are expressed as percentages, means and standard deviations, medians and ranges. χ^2 tests were carried out to compare qualitative values, t tests were used to compare quantitative values (normality assumption was checked by quantile–quantile plots and Shapiro-Wilks' test) and logistic regression analyses were carried out to assess the influence of covariates. Agreement between the two psychologists was assessed with a weighted κ reliability test for ordered data.

RESULTS

During our study period, 53 families were approached to participate in a RCT. Forty families have been interviewed in our study. Our recruitment rate was 75%. Thirteen families were non-interviewed for several reasons: the delay of 1 month was exceeded or the physicians have forgotten to propose our study.

We carried out 40 first interviews, including four with parents who declined participation in a randomised phase 3 trial (1 child with medulloblastoma, 1 with hepatoblastoma and 2 with retinoblastoma). We interviewed 37 mothers and 20 fathers (17 couples, 3 men attending alone and 20 women alone). We carried out 32 s interviews: four of the original interviewees (or couples) declined a second interview and four others were lost to follow-up for this study (management continued at another center). We interviewed 25 mothers and 14 fathers (7 couples, 7 men attending alone and 18 women alone). We considered the responses of each

of the couples as if they were individuals (ie, below, 'parent' refers to the mother, father or couple interviewed). Regarding the values of weighted κ obtained, 0.84 (95% CI 0.72 to 0.90) and 0.93 (95% CI 0.85 to 0.96) for M1 and M6, respectively, there was a high degree of agreement between the two psychologists about the coding of each interview. The characteristics of the study group are summarised in [table 3](#).

In 64% of the cases, information about the protocol had been provided by a single doctor, to both parents simultaneously in 91% of these cases. Twenty-seven parents (68%) had reread the information document. When asked if they had looked for information themselves, 13 parents (32.5%) replied in the negative, while others (67.5%) having mostly (24/40) searched for information about the disease rather than about the randomised clinical trial (7/40). Internet was the principal source of information for these parents.

Parental comprehension

At the first interview, six parents (15%) confused consent for care with consent for research. Each of these cases of confusion concerned a different protocol. Thirteen of the parents (34.2%) did not know that randomisation had occurred and five (13.2%) knew that randomisation existed but could not explain it clearly. When asked about the risks specific to the randomised trial, four parents (10%) were unable to cite a single risk, 35 parents (87.5%) were able to cite less than half the risks and one parent (2.5%) was able to cite more than half the risks.

The results of the second interviews were compared with those of the first interview: the level of parental comprehension did not differ markedly between M1 and M6 ($p=0.5$). [Figure 1](#) summarises the comprehension of the 11 elements described in the methods for the two interviews.

Parental comprehension in randomised clinical trials

Table 3 Characteristics of participating parents

Characteristic	Study group
Relationship to child (n, %)	
Father	20 (35)
Mother	37 (65)
Marital status (n, %)	
Married/living with partner	37 (92)
Divorced/separated	3 (8)
Parents' profession (n, %)	
High standing	11 (19)
Intermediate status	37 (65)
No profession	9 (16)
Mostly French-speaking (n, %)	
Yes	50 (86)
No	7 (14)
No. of children (n, %)	
1	11 (27.5)
2–3	21 (52.5)
4+	8 (20)
Patients' age (years)	
Mean (SD)	4.23 (4.5)
Median	1.93
Range	5 months–15 years
First interview (n=40, 17 couples)	
Duration (minutes, mean (SD)/median, range)	51(17.8)/50.0, 20–90
Time since inclusion (days, mean (SD)/median, range)	34(6.3)/31.5, 23–50
Second interview (n=32, 7 couples)	
Duration (minutes, mean (SD)/median, range)	34(13.2)/35.0, 15–60
Time since inclusion (months, mean (SD)/median, range)	8.4(2.8)/8.0, 5–16

As far as the benefit to their children was concerned, parents giving a different response from that indicated in the information document (first interview, 35% of parents/second interview, 11.1% of parents) expressed other aspects of the benefits anticipated for their child. Some felt there was a benefit in terms of quality of life (less time spent in hospital, fewer displacements), decreasing the amount of time for which their child would not attend school

I think I would have had to stay here all day and he would have had to have two sessions, but he wanted to go to school and that's why he preferred to do it in one go, because the old treatment takes three days and this one only takes one day, so for that reason alone, it's better for him in my opinion.


Other parents had hopes of beneficial effects in terms of their child being cured

Anyway, for him, if it's beneficial he's cured... and that's the most important thing.

Some parents expressed notions of altruism

Their aim and our aim are the same, because their goal is to progress in their studies, to save children..., C. has benefited from what went before and so he should help those who will come after him too.

Finally, some parents agreed to participate because they considered the research protocol to be providing access to better follow-up than they would have obtained with the standard treatment

Figure 1 Percentage of parents who understood each of the 11 items at M1 and M6.

Parental comprehension in randomised clinical trials

We were told, relatively clearly, that children who followed research protocols... I'm not saying that the others are neglected... but they are possibly followed a bit more closely than children given the classic treatment, because of the phenomenon of data collection for research.

If we look more closely at the responses of the four parents who refused to allow the participation of their children in a research protocol, they had no notion (0/4) of an expected benefit for their child (*Yes but I don't see why it's beneficial for him, because in some ways it's still being evaluated*).

Fluency in French ($p=0.03$) and reading the information notice ($p=0.0025$) were the only covariates tested that significantly increased parental comprehension (table 4).

Reasons for participation and physician's influence on the decision to participate

When asked how they felt about taking this decision, 24 parents (60%) said that they felt it was 'a logical decision' and 10 parents (25%) said that they had found it difficult. The others said that they were not really 'ready' to take a decision. In response to the question about the elements guiding the decision, the most frequently given answers were confidence in the medical team ($n=15$, 39%), access to the best possible treatment for the child ($n=14$, 37%) and improvements in quality of life ($n=14$, 37%). Most of the parents ($n=34$, 85%) said that they had taken the decision together, whereas the others had shared the task of decision-making with the doctor ($n=1$, 2.5%) or with the child ($n=5$, 12.5%). The parents expected the following from the doctor: communication (17/22, 77%), sincerity (11/22, 50%), availability (11/22, 50%), humanity (7/22, 32%) and competence (7/22, 32%).

DISCUSSION

Ideally, parents giving consent for inclusion in a randomised clinical trial should have understood all the

elements required by law to protect the participant. In reality, parental comprehension is highly variable and depends on the elements considered. Most parents had understood the general aspects such as the potential benefit to other children, the nature of the study (a clinical trial), the notion of voluntary participation, the duration of participation, the alternatives opened to them and the possibility of withdrawal at any time. By contrast, less than half the parents had correctly understood the more specific information such as the aims of the study, randomisation, the various phases of treatment and the risks specific to the randomised trial. These results are similar to published findings.^{4 5 8 12} For example, for randomisation in our study, some of the parents had views about the best 'arm' for their child: some preferred the standard treatment, which they considered to be less risky,¹³ whereas others were disappointed that their child was in the standard treatment arm because they could not see the point of having consented to participation in the study.¹⁴ We wished to focus more specifically on parental comprehension of the potential (or expected) benefits to their child of inclusion in a randomised phase 3 research protocol. Indeed, less than one-third of the parents were able to describe specifically the potential benefits to their child, as stated in the information document. In a recent study, Tait *et al*¹⁵ showed, through the use of scenarios, that only a one-quarter of the parents had fully understood the benefits to their child, and comprehension was even lower for studies of treatments with efficacy levels similar to those of the standard treatment, but lower expected toxicity. In our study, the parents expressed other expectations such as benefits in terms of quality of life (less time spent in hospital, fewer displacements), resulting in fewer hours of absence from school. This aspect of the perception of the benefits anticipated by parents has been identified before in phase I trials in adults.^{16 17} Other parents had hopes for the cure of their child, not differentiating between the trial and standard management, as described by Caldwell *et al*¹⁸ Finally, some parents agreed to the participation of their child in a phase three randomised trial because they considered that it would provide access to better follow-up than standard treatment as this has also been reported in non-oncological protocols.¹⁹⁻²¹ In our study, reading the information document improved parental comprehension. Unfortunately, some of these documents are difficult to read,^{22 23} so it is important for the investigating doctors to pay particular attention when writing such documents, to ensure that they are clear, as short as possible and understandable by all. O'Lonergan *et al*²⁴ recently reported a novel approach in which visual and audio media (multimedia) were used, rather than a paper-based written text document, to make the information easier to comprehend. However, this was a hypothetical research study involving simple, low-risk procedures and may therefore not be representative of actual research studies.

Table 4 Factors predictive of sufficient understanding of the information

Covariates	Per cent items understood	p
Parents' profession		
High standing	88.7 (SD 13.5)	0.09
Intermediate status	86.0 (SD 15.8)	
No profession	72.5 (SD 27.6)	
Mostly French-speaking		
Yes	84.8 (SD 18.7)	0.03*
No	61.9 (SD 29.6)	
Parents read the information sheet		
Yes	88.7 (SD 14.8)	0.0025*
No	68.5 (SD 24.6)	
Parents sought additional informations		
Yes	84.5 (SD 18.1)	0.29
No	77.2 (SD 25.1)	

As shown above, some of the elements of the research protocol were well understood, but not by all the parents. So, how can the parents decide? Although not yet demonstrated in paediatrics to our knowledge, quality of life could be a crucial element in the decision of parents to agree to or refuse the participation of their child in a randomised phase 3 trial. The smaller amount of time spent in hospital and the smaller number of journeys for a research protocol than for standard treatment was an element that appeared frequently in the decision-making process. Recent studies have reported that logistic constraints are a major element in the decisions of adult patients,^{25 26} sometimes even outweighing benefits. In our study, the parents took into account the benefits and constraints of inclusion to a much greater extent than the risks specific to the research protocol, as indicated by their poor 'recall' of the risks. However, there may be several reasons for this recall failure. For example, having given their consent for participation, the parents may minimise the impact of potential risks, focusing instead on the possible benefits of participation. Other parents may try to protect themselves psychologically by blocking out the largest risks. The parents may also feel that their child's illness poses a greater and more immediate threat, in terms of morbidity and mortality, than the risks inherent to a clinical trial. These and the other factors affect people's perceptions of risk, potentially accounting for the poor recall of risks specific to the randomised trial.²⁷ Overall, even if the decision was difficult for some of the parents, more than half felt that it was logical. The decision seemed to be linked to the confidence of the parents in the investigating doctor and their relationship with that doctor.²⁸ This confidence overcame the imbalance in knowledge between the parents and the doctor.^{8 14} By contrast, a lack of confidence in the doctor may be associated with the decision being more difficult for the parents.¹¹ In previous studies,^{29–31} the way in which the investigating doctor presented the trial to the patient was found to affect the likelihood of participation. In our study, the parents recognised that the doctor played an important role in their decision, but they did never feel obliged to sign the consent form.

What may appear to the investigating doctor to be a poor understanding of the research protocol may actually correspond to the parents' construction of the situation to render it acceptable, thus allowing them to fulfill their role as protectors of their children. As suggested by Shilling,²⁸ given that it is difficult to accept that there is uncertainty about the effectiveness of trial treatments and assumptions that the experimental treatment will necessarily be superior, these responses indicate that the parents seek hope and certainty in this uncertain situation in which they feel threatened. Parents need to give their decision some meaning, by seeing it as being in the best interests of their child.

In conclusion, most parents understand most of the elements of IC well. However, knowledge is limited in

some areas and improvements of information process are required. Parents tend to underestimate risks specific to the randomised trial and the unproven nature of the treatment is not well known or understood. Investigators should also systematically ask the parents to reformulate the information they have been given to verify that they have understood; this would also encourage active parental participation in a two-way information exchange process.

Author affiliations

¹Pediatric Emergency Department, Hôpital Necker Enfants Malades, Université Paris Descartes, Paris, France

²Clinical Research Unit Paris centre, APHP, Paris, France

³Department of Pediatric and Adolescent Oncology, Institut Gustave Roussy, Paris, France

⁴Département of Pediatric Oncology, Hôpital Armand Trousseau, Paris, France

⁵Association Apprivoiser l'absence, Cent pour Sang la Vie, Paris, France

⁶Department of Pediatric Oncology, Institut Curie, Paris, France

⁷Department of Pharmacology, APHP, Paris, France

Acknowledgements We thank the parents who agreed to participate in this study, making this work possible. We thank all the investigating doctors from the Société Française des Cancres de l'Enfant (SFCE), the psychologists (Amélie de Haut de Sigy and Anne Charlotte Foubert) and also the clinical research assistants from all centres.

Contributors HC, DD and FD conceived this study and HC obtained research funding. JMT, NB and HC was in charge of all the data analyses. HC took the lead in reviewing the literature and crosschecking references. HC and JMT drafted major portions of the initial manuscript, whereas DD and FD helped refined it. The other authors CP, JLP, AA and HP were investigators. HC, JMT, NB, DD, FD, VMC, LB and DO have seen and approved the submission of this version of the manuscript and takes full responsibility for the manuscript.

Funding This work received funding from a clinical research initiation contract (DRRC APHP).

Competing interests None.

Ethics approval Institutional Review Board (Comité de Protection des Personnes Ile de France III Hôpital Cochin).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

REFERENCES

- Loi n°2004-806 du 9 août 2004 sur la recherche biomédicale. Code de la Santé Publique livre 1 titre 2. <http://www.legifrance.fr>
- Directive 2001/20/EC of the European parliament and of the council of 4 april 2001 on the approximation of the laws, regulations and administrative provisions of the member states relating to the implementation of good clinical practice in the conduct of clinical trials on medicinal products for human use. <http://europa.eu.int>
- Boccia ML, Campbell FA, Goldman BD, *et al.* Differential recall of consent information and parental decisions about enrolling children in research studies. *J Gen Psychol* 2009;136:91–108.
- Kodish E, Eder M, Noll RB, *et al.* Communication of randomization in childhood leukemia trials. *JAMA* 2004;291:470–5.
- Kupst MJ, Patenaude AF, Walco GA, *et al.* Clinical trials in pediatric cancer: parental perspectives on informed consent. *J Pediatr Hematol Oncol* 2003;25:787–90.
- Falagas ME, Korbila IP, Giannopoulou KP, *et al.* Informed consent: how much and what do patients understand? *Am J Surg* 2009;198:420–35.
- Bergemar M, Molin C, Wilking N, *et al.* Knowledge and understanding among cancer patients consenting to participate in clinical trials. *Eur J Cancer* 2008;44:2627–33.
- Chappuy H, Baruchel A, Leverger G, *et al.* Parental comprehension and satisfaction in informed consent in paediatric clinical trials: a prospective study on childhood leukaemia. *Arch Dis Child* 2010;95:800–4.

Parental comprehension in randomised clinical trials

9. Miller VA, Drotar D, Burant C, *et al*. Clinician-parent communication during informed consent for pediatric leukemia trials. *J Pediatr Psychol* 2005;30:219–29.
10. Greenley RN, Drotar D, Zyzanski SJ, *et al*. Stability of parental understanding of random assignment in childhood leukemia trials: an empirical examination of informed consent. *J Clin Oncol* 2006;24:891–7.
11. Chappuy H, Doz F, Blanche S, *et al*. Parental consent in paediatric clinical research. *Arch Dis Child* 2006;91:112–16.
12. Wiley FM, Ruccione K, Moore IM, *et al*. Parents' perceptions of randomization in pediatric clinical trials. *Children Cancer Group. Cancer Pract* 1999;7:248–56.
13. Snowdon C, Garcia J, Elbourne D. Making sense of randomization; responses of parents of critically ill babies to random allocation of treatment in a clinical trial. *Soc Sci Med* 1997;45:1337–55.
14. Eiser C, Davies H, Jenney M, *et al*. Mothers' attitudes to the randomized controlled trial (RCT): the case of acute lymphoblastic leukaemia (ALL) in children. *Child Care Health Dev* 2005;31:517–23.
15. Tait AR, Zikmund-Fisher BJ, Fagerlin A, *et al*. Effect of various risk/benefit trade-offs on parents' understanding of a pediatric research study. *Pediatrics* 2010;125:e1475–82.
16. Weinfurt KP, Castel LD, Li Y, *et al*. The correlation between patient characteristics and expectations of benefit from Phase I clinical trials. *Cancer* 2003;98:166–75.
17. Cheng JD, Hitt J, Koczwara B, *et al*. Impact of quality of life on patient expectations regarding phase I clinical trials. *J Clin Oncol* 2000;18:421–8. Erratum in: *J Clin Oncol* 2000;18:1807.
18. Caldwell PH, Butow PN, Craig JC. Parents' attitudes to children's participation in randomized controlled trials. *J Pediatr* 2003;142:554–9.
19. Pletsch PK, Stevens PE. Inclusion of children in clinical research: lessons learned from mothers of diabetic children. *Clin Nurs Res* 2001;10:140–62.
20. Gammelgaard A, Knudsen LE, Bisgaard H. Perceptions of parents on the participation of their infants in clinical research. *Arch Dis Child* 2006;91:977–80.
21. van Stuijvenberg M, Suur MH, de Vos S, *et al*. Informed consent, parental awareness, and reasons for participating in a randomised controlled study. *Arch Dis Child* 1998;79:120–5.
22. Ménoni V, Lucas N, Leforestier JF, *et al*. Readability of the written study information in pediatric research in France. *PLoS ONE* 2011;6:e18484.
23. Ménoni V, Lucas N, Leforestier JF, *et al*. The readability of information and consent forms in clinical research in France. *PLoS ONE* 2010;5:e10576.
24. O'Lonergan TA, Forster-Harwood JE. Novel approach to parental permission and child assent for research: improving comprehension. *Pediatrics* 2011;127:917–24.
25. Avis NE, Smith KW, Link CL, *et al*. Factors associated with participation in breast cancer treatment clinical trials. *J Clin Oncol* 2006;24:1860–7.
26. Cox K, McGarry J. Why patients don't take part in cancer clinical trials: An overview of the literature. *Eur J Cancer Care* 2003;12:114–22.
27. Lloyd AJ. The extent of patients' understanding of the risk of treatments. *Qual Health Care* 2001;10(Suppl 1):i14–18.
28. Shilling V, Young B. How do parents experience being asked to enter a child in a randomised controlled trial? *BMC Med Ethics* 2009;10:1.
29. Albrecht TL, Ruckdeschel JC, Riddle DL, *et al*. Communication and consumer decision making about cancer clinical trials. *Patient Educ Couns* 2003;50:39–42.
30. Jenkins V, Fallowfield L, Solis-Trapala I, *et al*. Discussing randomized clinical trials of cancer therapy: Evaluation of a Cancer Research UK training programme. *BMJ* 2005;330:400.
31. Harth SC, Thong YH. Parental perceptions and attitudes about informed consent in clinical research involving children. *Soc Sci Med* 1995;41:1647–51.

Parental comprehension of the benefits/risks of first-line randomised clinical trials in children with solid tumours: a two-stage cross-sectional interview study

Hélène Chappuy, Naim Bouazza, Veronique Minard-Colin, Catherine Patte, Laurence Brugières, Judith Landman-Parker, Anne Auvrignon, Dominique Davous, Hélène Pacquement, Daniel Orbach, Jean Marc Tréluyer and François Doz

BMJ Open 2013 3:
doi: 10.1136/bmjopen-2013-002733

Updated information and services can be found at:
<http://bmjopen.bmj.com/content/3/5/e002733>

These include:

References

This article cites 29 articles, 12 of which you can access for free at:
<http://bmjopen.bmj.com/content/3/5/e002733#BIBL>

Open Access

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license. See: <http://creativecommons.org/licenses/by-nc/3.0/> and <http://creativecommons.org/licenses/by-nc/3.0/legalcode>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Ethics](#) (80)
[Oncology](#) (404)
[Paediatrics](#) (609)
[Research methods](#) (590)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>