

Primary osteomyelitis caused by an NDM-1-producing *K. pneumoniae* strain of the highly virulent sequence type 23

Fabrice Compain, Alexis Vandenberghe, Marie Gominet, Nathalie Genel, David Lebeaux, Astrid Ramahefasolo, Isabelle Podglajen, Dominique Decré

► **To cite this version:**

Fabrice Compain, Alexis Vandenberghe, Marie Gominet, Nathalie Genel, David Lebeaux, et al.. Primary osteomyelitis caused by an NDM-1-producing *K. pneumoniae* strain of the highly virulent sequence type 23. *Emerging microbes*

infections, Nature Publishing Group, 2017, 6, pp.e57. <10.1038/emi.2017.43>. <hal-01549095>

HAL Id: hal-01549095

<https://hal.sorbonne-universite.fr/hal-01549095>

Submitted on 28 Jun 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



LETTER TO THE EDITOR

Primary osteomyelitis caused by an NDM-1-producing *K. pneumoniae* strain of the highly virulent sequence type 23

Fabrice Compain^{1,2,3}, Alexis Vandenberghe⁴, Marie Gominet¹, Nathalie Genel⁴, David Lebeaux^{1,2,3}, Astrid Ramahefasolo¹, Isabelle Podglajen^{1,2,3,*} and Dominique Decré^{3,4,5,*}

Emerging Microbes & Infections (2017) **6**, e57; doi:10.1038/emi.2017.43; published online 21 June 2017

Dear Editor,

Hypervirulent variant of *Klebsiella pneumoniae* (hvKP) is characterized by its hypermucoviscosity and its capacity to cause invasive community-acquired infections in previously healthy young individuals.¹ The most common clinical presentation is a pyogenic liver abscess with metastatic spread, while various other invasive infections have been reported.^{1,2} Strains of hvKP belong predominantly to the capsular serotypes K1 and K2² and their genetic background seems to differ from that of multidrug-resistant (MDR) isolates.³ The most commonly encountered sequence types (STs) among hvKP include ST23 (associated with serotype K1), ST25, ST65, ST86, ST375 and the recently emerging ST380.^{4,5}

Most hvKP isolated to date are highly sensitive to nearly all antibiotics except ampicillin. However, MDR isolates belonging to virulent clones have been described recently and extended-spectrum β -lactamase (ESBL) producers were reported among hvKP.⁶ Strains belonging to highly virulent clone ST23 and producing the KPC-2 carbapenemase are rare but have been observed on at least three continents.^{7–9}

We describe here the first NDM-1-producing *K. pneumoniae* isolate of capsular serotype K1 belonging to the highly virulent ST23 clone. It caused primary osteomyelitis in a young diabetic patient from Saudi Arabia. The rise of multidrug resistance in a highly virulent strain is a cause of great concern.

A 30-year-old man with unremarkable medical history except type I diabetes mellitus was transferred to our hospital for maxillofacial surgery subsequent to a road accident 2 months earlier and emergency facial surgery in Saudi Arabia. Surgery at our institution consisted in the ablation of all osteosynthesis material as well as osteotomy of the zygomatic arch, maxilla and mandible. Abundant hvKP resistant to all carbapenems was found in all surgical samples and osteomyelitis was diagnosed. Whole-body CT scanning revealed no other foci of infection. The patient was treated with parenteral meropenem,

amikacin and fosfomycin for 15 days, followed by ciprofloxacin and sulfamethoxazole/trimethoprim for a total of three months and was discharged in good condition.

Four unique carbapenem-resistant *K. pneumoniae* isolates were obtained from the patient. One isolate was the sole organism cultured from all perioperative samples, while three were found only in systematic rectal swab screening sample obtained at admission (Table 1). The four isolates were high-level resistant to all β -lactams including carbapenems. The isolate responsible for osteomyelitis (KP43-45) remained susceptible to aminoglycosides, ciprofloxacin, sulfamethoxazole/trimethoprim, tigecycline and fosfomycin, while the other isolates showed various associated resistance profiles (Table 1).

PCR and sequencing revealed the presence in all isolates of the carbapenemase *bla*_{NDM-1} gene, associated with the carbapenemase *bla*_{OXA-48} gene in three of them, that is, KP43-53, KP43-54 and KP43-55 (Table 1). ESBL production was detected in all four isolates. Plasmid analysis was performed using the relaxase gene typing method as previously described.¹⁰ KP43-45 (harboring only *bla*_{NDM-1}) contained a plasmid of the FIIk incompatibility (Inc) group, while plasmids belonging to IncFIIk and IncL/M were detected in isolates carrying both *bla*_{NDM-1} and *bla*_{OXA-48}.

Multilocus sequence typing (<http://bigsd.web.pasteur.fr/>) revealed two distinct allelic profiles, one for KP43-45, KP43-53 and KP43-54 belonging to ST23 and one for KP43-55 belonging to ST437. This was in concordance with molecular fingerprints obtained using random amplified polymorphic DNA and repetitive element palindromic sequence-based PCR (data not shown).

The three hypermucoviscous isolates of capsular serotype K1 belonging to the ST23 clone were string-test positive¹ and carried the *rmpA* (regulator of mucoid phenotype A), *entB* (siderophore, enterobactin), *ybtS* (siderophore, yersiniabactin), *kfu* (siderophore), *iutA* (siderophore, aerobactin), *mrkD* (type 3 fimbrial adhesin) and *allS* (allantoin metabolism) virulence genes (Table 1). KP43-55

¹Microbiology, AP-HP, Hôpital Européen Georges Pompidou, Paris 75015, France; ²Université Paris Descartes, Sorbonne Paris Cité, Paris 75006, France; ³Institut National de la Santé et de la Recherche Médicale, Paris 75013, France; ⁴UPMC Univ Paris 06 INSERM, CIMI-Paris, UMR 1135, ERL, CNRS 8544, Paris 75013, France and ⁵Microbiology, AP-HP, Hôpital Saint Antoine, Paris 75011, France

*These authors contributed equally to this work.

Correspondence: I Podglajen; D Decré

E-mail: isabelle.podglajen@aphp.fr; dominique.decre@aphp.fr

Received 15 January 2017; accepted 16 April 2017

Table 1 Resistance profile, virulence properties and genetic background of the four *K. pneumoniae* isolates analyzed

Variable	KP 43-45	KP 43-53	KP 43-54	KP 43-55
Isolate source	Frontal sinus mucocele and sequestra; rectal swab	Rectal swab	Rectal swab	Rectal swab
<i>Antibiotic and disk diffusion zone diameter (mm)</i>				
AMP	6 (R)	6 (R)	6 (R)	6 (R)
AMC	6 (R)	6 (R)	6 (R)	9 (R)
TZP	6 (R)	6 (R)	6 (R)	11 (R)
CTX	6 (R)	6 (R)	6 (R)	6 (R)
CAZ	6 (R)	6 (R)	6 (R)	6 (R)
ATM	6 (R)	14 (R)	6 (R)	10 (R)
FEP	6 (R)	12 (R)	6 (R)	11 (R)
IPM	16 (I) ^a	17 (I)	18 (I)	16 (I)
MEM	15 (R) ^a	15 (R)	16 (I)	15 (R)
AN	22 (S)	9 (R)	10 (R)	9 (R)
GE	24 (S)	27 (S)	30 (S)	13 (R)
CIP	22 (S) ^a	21 (I)	21 (I)	6 (R)
SXT	20 (S)	23 (S)	27 (S)	6 (R)
FOS	20 (S)	22 (S)	21 (S)	24 (S)
Carbapenemase content	NDM-1	NDM-1, OXA-48	NDM-1, OXA-48	NDM-1, OXA-48
Plasmid Inc type	FIik	FIik, L/M	FIik, L/M	FIik, L/M
MLST	ST23	ST23	ST23	ST437
Rep-PCR or RAPD profile	A	A	A	B
<i>Virulence properties</i>				
String-test	Positive	Positive	Positive	Negative
Capsular polysaccharide	K1	K1	K1	non-K1/K2
Virulence genes	<i>magA, rmpA, ybtS, kfu, iutA, allS, entB, mrkD</i>	<i>magA, rmpA, ybtS, kfu, iutA, allS, entB, mrkD</i>	<i>magA, rmpA, ybtS, kfu, iutA, allS, entB, mrkD</i>	<i>ybtS, entB, mrkD</i>

Abbreviations: ampicillin, AMP; co-amoxiclav, AMC; piperacillin-tazobactam, TZP; cefotaxime, CTX; ceftazidime, CAZ; aztreonam, ATM; cefepime, FEP; imipenem, IPM; meropenem, MEM; amikacin, AN; gentamicin, GE; ciprofloxacin, CIP; sulfamethoxazole/trimethoprim, SXT; fosfomycin, FOS; tigecycline, TGC; susceptible, S; intermediate, I; resistant, R, according to EUCAST breakpoints version 5.0 (www.eucast.org); multilocus sequence typing, MLST.

^aFor the clinical isolate, MIC of ertapenem was of 32 µg/mL, IPM 8 µg/mL, MEM 12 µg/mL, CIP 0.5 µg/mL and TGC 0.75 µg/mL. Minimal inhibitory concentrations were determined with the E-test (bioMérieux).

(ST437) was string-test negative, non-K1/K2 and tested positive only for the *entB*, *ybtS* and *mrkD* virulence genes.

Hypervirulent variant *K. pneumoniae* is a globally emerging pathogen that was first described in Taiwan and has now spread to other regions in Southeast Asia, Europe and the Americas.^{2,5} Invasive syndromes due to hvKP have been reported to occur in the Middle-East. Our patient was repatriated from Saudi Arabia, a country where a case of liver abscess caused by a serotype K1 *K. pneumoniae* isolate was recently described.¹¹ The novel *bla*_{NDM-1}-producing isolate described here may well have been acquired in that country.

K. pneumoniae ST23 strains belong to a virulent clone that is highly prevalent in *K. pneumoniae* liver abscesses.^{2,4} Most ST23 strains are hypermucoviscous, which seems to correlate with the presence of the *magA* and *rmpA* genes.¹ The ST23 strains described in our study also carried seven virulence genes often found in this clone.^{3,4} A non-hvKP isolate of ST437 was found in a rectal swab sample. ST437 is part of clonal complex 11, a high-risk MDR complex.¹² A horizontal transfer of resistance genes from the ST437 isolate to the ST23 isolates is highly probable; however, our mating experiments to substantiate this were not successful.

K. pneumoniae ST23 isolates have been found to be susceptible to most drugs including third-generation cephalosporins (3GC), fluoroquinolones and aminoglycosides. Although some studies have described ESBL-producing isolates resistant to 3GC or carbapenems,⁶ reports of carbapenemase-producing ST23 isolates

remain rare. KPC-2-producing ST23 isolates were described in China and Poland in 2010 and 2011, respectively, but without any clinical or microbiological supplementary information.^{8,9} A hypermucoviscous KPC-2-producing *K. pneumoniae* belonging to ST23 was recovered in a respiratory sample of a patient from Argentina.⁷ Although carbapenemase-producing Gram-negative bacteria have largely spread in the Arabian Peninsula, no carbapenemase-producing *K. pneumoniae* belonging to ST23 have been described before in that region.¹³

In this study, we describe the first hvKP isolate belonging to ST23 and producing the NDM-1 carbapenemase. This isolate had all the hvKP characteristics described by Shon *et al.*,¹ as it was responsible for an invasive infection in a previously healthy young patient and displayed particular microbiological features such as hypermucoviscosity and the presence of virulence factors. The only underlying disease that could be identified in our patient was type I diabetes mellitus, which has been suggested in some studies to be a potent risk factor for hvKP infection.^{1,2} An interesting feature was the intestinal-tract carriage of three unique hypermucoviscous, carbapenem-resistant *K. pneumoniae* isolates, including that responsible for osteomyelitis. Asymptomatic carriage of hvKP isolates has been suggested to be a major risk factor for infection in Asian patients.¹⁴ Therefore colonization with hvKP ST23 isolates harboring both NDM-1 and OXA-48 carbapenemases with associated resistance to aminoglycosides and ciprofloxacin in our patient should cause concern. Moreover, plasmid typing identified IncFIik and IncL/M replicons, which are part of

conjugative plasmids involved in horizontal transfer of carbapenemase genes.¹⁵ IncL/M plasmids have been previously associated with the dissemination of the *bla*_{OXA-48} gene.¹⁵

Acquisition of carbapenemase genes by the hvKP ST23 strains is worrisome and should be closely monitored. This can be done easily with the string test, MLST and identification of virulence factors. The acquisition of *bla*_{NDM-1} might be one step leading to treatment failure in patients infected with hypervirulent *K. pneumoniae*.

- 1 Shon AS, Bajwa RPS, Russo TA. Hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*: a new and dangerous breed. *Virulence* 2013; **4**: 107–118.
- 2 Siu LK, Yeh K-M, Lin J-C *et al*. *Klebsiella pneumoniae* liver abscess: a new invasive syndrome. *Lancet Infect Dis* 2012; **12**: 881–887.
- 3 Bialek-Davenet S, Criscuolo A, Ailloud F *et al*. Genomic definition of hypervirulent and multidrug-resistant *Klebsiella pneumoniae* clonal groups. *Emerg Infect Dis* 2014; **20**: 1812–1820.
- 4 Brisse S, Fevre C, Passet V *et al*. Virulent clones of *Klebsiella pneumoniae*: identification and evolutionary scenario based on genomic and phenotypic characterization. *PLoS ONE* 2009; **4**: e4982.
- 5 Decré D, Verdet C, Emirian A *et al*. Emerging severe and fatal infections due to *Klebsiella pneumoniae* in two university hospitals in France. *J Clin Microbiol* 2011; **49**: 3012–3014.
- 6 Li W, Sun G, Yu Y *et al*. Increasing occurrence of antimicrobial-resistant hypervirulent (hypermucoviscous) *Klebsiella pneumoniae* isolates in China. *Clin Infect Dis* 2014; **58**: 225–232.
- 7 Cejas D, Fernández Canigia L, Rincón Cruz G *et al*. First isolate of KPC-2-producing *Klebsiella pneumoniae* sequence type 23 from the Americas. *J Clin Microbiol* 2014; **52**: 3483–3485.

- 8 Qi Y, Wei Z, Ji S *et al*. ST11, the dominant clone of KPC-producing *Klebsiella pneumoniae* in China. *J Antimicrob Chemother* 2011; **66**: 307–312.
- 9 Baraniak A, Grabowska A, Izdebski R *et al*. Molecular characteristics of KPC-producing Enterobacteriaceae at the early stage of their dissemination in Poland, 2008–2009. *Antimicrob Agents Chemother* 2011; **55**: 5493–5499.
- 10 Compain F, Poisson A, Le Hello S *et al*. Targeting relaxase genes for classification of the predominant plasmids in Enterobacteriaceae. *Int J Med Microbiol* 2014; **304**: 236–242.
- 11 Enani MA, El-Khizzi NA. Community acquired *Klebsiella pneumoniae*, K1 serotype. Invasive liver abscess with bacteremia and endophthalmitis. *Saudi Med J* 2012; **33**: 782–786.
- 12 Oliveira S, Moura RA, Silva KC *et al*. Isolation of KPC-2-producing *Klebsiella pneumoniae* strains belonging to the high-risk multiresistant clonal complex 11 (ST437 and ST340) in urban rivers. *J Antimicrob Chemother* 2014; **69**: 849–852.
- 13 Zowawi HM, Balkhy HH, Walsh TR *et al*. β -lactamase production in key Gram-negative pathogen isolates from the Arabian peninsula. *Clin Microbiol Rev* 2013; **26**: 361–380.
- 14 Fung C-P, Lin Y-T, Lin J-C *et al*. *Klebsiella pneumoniae* in gastrointestinal tract and pyogenic liver abscess. *Emerg Infect Dis* 2012; **18**: 1322–1325.
- 15 Carattoli A. Plasmids and the spread of resistance. *Int J Med Microbiol* 2013; **303**: 298–304.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017