



HAL
open science

Blood lead levels and risk factors for lead exposure among pregnant women in western French Guiana: the role of manioc consumption

Diane Rimbaud, Marion Restrepo, Anne Louison, Rachida Boukhari, Vanessa Ardillon, Gabriel Carles, Véronique Lambert, Anne Jolivet

► To cite this version:

Diane Rimbaud, Marion Restrepo, Anne Louison, Rachida Boukhari, Vanessa Ardillon, et al.. Blood lead levels and risk factors for lead exposure among pregnant women in western French Guiana: the role of manioc consumption. *Journal of Toxicology and Environmental Health, Part A: Current Issues*, 2017, pp.1 - 12. 10.1080/15287394.2017.1331490 . hal-01552928

HAL Id: hal-01552928

<https://hal.sorbonne-universite.fr/hal-01552928>

Submitted on 3 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1
2
3 Blood lead levels and risk factors for lead exposure among pregnant women in western
4
5 French Guiana: the role of manioc consumption
6
7

8
9 Diane Rimbaud^a, Marion Restrepo^a, Anne Louison^b, Rachida Boukhari^c, Vanessa Ardillon^d,
10
11 Gabriel Carles^e, Véronique Lambert^e, Anne Jolivet^{a,f,*}
12
13

14
15 ^a Department of Public Health, Centre Hospitalier de l'Ouest Guyanais, Saint-Laurent du
16
17 Maroni, French Guiana
18

19 ^b Réseau Périnatal-Guyane, Cayenne, French Guiana
20

21 ^c Department of Clinical Biology, Centre Hospitalier de l'Ouest Guyanais, Saint-Laurent du
22
23 Maroni, French Guiana
24

25 ^d CIRE Antilles-Guyane, Cayenne, French Guiana
26

27 ^e Department of Obstetrics & Gynecology, Centre Hospitalier de l'Ouest Guyanais, Saint-
28
29 Laurent du Maroni, French Guiana
30

31 ^f Sorbonne Universités, UPMC Univ Paris 06, INSERM, Institut Pierre Louis d'épidémiologie
32
33 et de Santé Publique (IPLESP UMRS 1136), Department of social epidemiology, F75012,
34
35 Paris, France
36
37

38
39 **RUNNING TITLE** : Lead poisoning among pregnant women in western French Guiana
40
41

42
43 **Corresponding author:**
44

45 Anne JOLIVET, MD, PhD
46

47 Department of Public Health
48

49 Centre Hospitalier de l'Ouest Guyanais
50

51 16 Boulevard du Général de Gaulle - BP 245
52

53 97 393 Saint Laurent du Maroni, French Guiana
54

55 E-mail : a.jolivet@ch-ouestguyane.fr
56

57 Tel : (00 594) 5 94 34 89 12
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Concerns about lead (Pb) poisoning in French Guiana first arose in 2011, after the discovery of excessively high levels of the metal among children in a small neighborhood without any clear source of Pb. Since 2012, blood lead levels (BLL) measurement has been proposed to all pregnant women in western French Guiana. The aim of this study was to determine BLL in pregnant women in this region and identify factors associated with elevated BLL. An observational study of a consecutive sample of women who delivered in the maternity ward of the hospital was conducted. Risk factors were investigated using a questionnaire administered post-delivery by midwives (N = 531). Approximately 25 and 5% of women displayed BLL of $\geq 50 \mu\text{g/L}$ and $\geq 100 \mu\text{g/L}$, respectively. The geometric mean was $32.6 \mu\text{g/L}$. Factors that were significantly associated with an elevated BLL after modeling (multivariate linear regression) included: place of residence along the Maroni river, low level of education, daily consumption of manioc derivatives, weekly and daily consumption or personal preparation of manioc flour during pregnancy and weekly consumption of wild game. This study provides insight into the regional and social disparities in BLL in French Guiana and on potential sources of exposure. Evidence indicates that foods that are primarily produced and consumed in the Guiana Shield significantly affect BLL. Taken together with existing data, our results demonstrate that specific actions in terms of prevention, screening and care are required to be adapted and put into place in order to reduce exposure.

Keywords: Lead poisoning; Epidemiological Study; French Guiana; Manioc

Introduction

Lead (Pb) is a non-essential metal that is highly toxic to several physiological systems in humans, including the nervous, renal, cardiovascular, reproductive, immune, and hematologic systems (Counter et al., 2015; García-Lestón et al., 2012; Pollack et al., 2015). The most vulnerable populations are children and pregnant women (Buchanan et al., 2011; Lamadrid-Figueroa et al., 2006), and there are no apparent established thresholds for levels at which a number of critical health effects occur (NTP, 2012). Low intelligence quotients (IQ) were reported in children with blood lead levels (BLL) of less than 100 µg/L (Bellinger, 2008; Lanphear et al., 2005); however it should be noted that these findings are contradictory. BLL vary over the course of pregnancy, following a U-shaped curve with the lowest point between 12 and 20 weeks after the last menstrual period. In the second trimester of pregnancy, low BLL can be explained by hemodilution, whereas in the third trimester, higher BLL may occur due to the mobilization of calcium and Pb from maternal bone to mineralize the fetal skeleton (Miranda et al., 2010; Rothenberg et al., 1994). Lead readily crosses the placental barrier (Lamadrid-Figueroa et al., 2006; Osman et al., 2000; Pollack et al., 2015), and its primary effects on pregnancy progression include: the risk of gestational hypertension and/or pre-eclampsia (Kennedy et al., 2012), an increased risk of spontaneous abortion (Borja-Aburto et al., 1999), danger of premature labor, prematurity, and low birth weight and height (Taylor et al., 2015; Zhu et al., 2010). Finally, Pb was found to exert a detrimental effect on fetal development, and especially neurocognitive function, *in utero* (Jedrychowski et al., 2008; Hu et al., 2006; Schnaas et al., 2006), making pregnant women a high-risk group.

Lead is responsible for a significant disease burden in developed countries (Bellinger, 2012; Valent et al., 2004) but the largest disease burden is seen in developing countries, including Latin America, where this metal continues to pose a significant public health risk and entails high costs (Attina and Trasande, 2013). Legislative measures have been gradually introduced to reduce exposure by removing Pb from paint, food cans, water pipes

1
2
3 and petrol (Tagne-Fotso et al., 2016), and population-wide BLL have fallen quickly in the
4
5 USA (Tsoi et al., 2016) and Europe (WHO, 2009) in response to these measures.
6

7 French Guiana is a French overseas department situated in northeastern South
8
9 America, between Brazil to the east and south and Suriname to the west. It is a large country
10
11 of approximately 84,000 km², and with a population of 246,500 in 2013. St-Laurent-du-
12
13 Maroni is the second largest city, separated from Suriname by the natural border created by
14
15 the Maroni River. The population of the city is growing rapidly (+4.3% per year), and
16
17 characterized by its youth (52.5% of inhabitants were younger than 20 in 2013) and melting-
18
19 pot nature (due to multiple waves of immigration). Its economy is characterized by weak
20
21 development, a high level of unemployment (47.8% in 2013) and persistence of traditional
22
23 ways of life such as hunting, fishing, and slash-and-burn farming. The Western French
24
25 Guiana Hospital Center (Centre Hospitalier de l'Ouest Guyanais, CHOG) is the sole
26
27 secondary healthcare provider for a large area and treats patients from a region that spans
28
29 both sides of the border (Figure 1), which was estimated to be home to approximately
30
31 100,000 individuals in 2013.
32

33 The problem with Pb poisoning in west French Guiana arose in 2011, after discovery
34
35 of excessive metal poisoning in a 3-year-old girl (BLL = 1724 µg/L) (Rorive et al., 2015). Up
36
37 until that point, no apparent studies had been performed in French Guiana, as Pb poisoning
38
39 did not seem to be a public health concern. The results from an investigation conducted by
40
41 CIRE Antilles-Guyane after this case came to light showed that 93% (13/14) of children
42
43 under 7 year of age living in the vicinity exhibited a BLL > 100 µg/L. Environmental
44
45 investigations suggested multifactorial exposure that includes a nutritional element, primarily
46
47 from derivatives of bitter manioc (Barbosa Jr. et al., 2009; Carneiro et al., 2013; Freire et al.,
48
49 2014). Given the lack of knowledge regarding Pb exposure in French Guiana, systematic
50
51 assessment of BLL in pregnant women was implemented in 2012 in western French Guiana.
52
53 BLL assessment is performed by health professionals ideally during the prenatal visit in the
54
55 4th month of pregnancy and is repeated during the 3rd trimester if the first BLL value is > 50
56
57 µg/l. BLL is determined at delivery if not assessed previously. Thus, the objectives of this
58
59
60

1
2
3 study were to (1) determine Pb levels in pregnant women in western French Guiana
4
5 (population-wide BLL) and (2) identify factors associated with elevated BLL.
6
7

8 9 **Methods and Materials**

10 11 12 *Study population*

13
14
15 A descriptive, cross-sectional, single-center anonymous study was conducted over a
16
17 period of 12 consecutive weeks from September to November 2013. The study population
18
19 included women who delivered in the CHOG maternity ward, whose BLL was measured at
20
21 least once during the course of their pregnancy, and who agreed to participate or had
22
23 parental consent (for mothers under the age of 18). Patient information, patient consent and
24
25 data were collected by midwives on the maternity service who were trained for the study.
26
27

28 29 *Data collection*

30
31 The following information was collected using a questionnaire translated into local
32
33 languages: sociodemographic characteristics (age, place of birth, native language, primary
34
35 place of residence), socioeconomic characteristics (level of education, professional status,
36
37 legal status, perceived financial situation), nutrition throughout pregnancy (frequency with
38
39 which certain foods identified as potential risks were consumed, such as bitter manioc
40
41 derivatives: couac (manioc flour), cassava (manioc pancakes), crabio (detoxified cooking
42
43 liquid used as a sauce), tapioca (bitter manioc starch), wassaï (fruit of the wassaï palm tree),
44
45 pemba (local clay), wild game (mammals such as Capybaras, Cuniculus paca, and javelinas,
46
47 and local birds such as pigeons, curassows, etc.)), lifestyle factors' (tobacco use, and alcohol
48
49 consumption), the source of drinking water and preparing couac during pregnancy. The
50
51 maternal BLL were recorded from medical records after all other information was collected,
52
53 to avoid biasing the collection of patient information. We also recorded the gestational age
54
55 (as calculated from the date of last menses or corrected by the first trimester ultrasound) at
56
57 which the BLL was assessed.
58
59
60

Whole blood collection and determination of Pb levels

Blood samples were collected by a health professional using standard procedures. Blood was collected and stored in purple-top vacutainers (Becton-Dickinson) with EDTA. Vacutainers were frozen and stored at -20°C in the CHOG until shipping. Pb analyses were carried out in a central laboratory (Cerba, Paris) and measured by atomic absorption spectrometry. The limit of detection (LOD) and the limit of quantification (LOQ) were 10 µg/L. All specimens with a Pb concentration greater than 200 µg/L were reanalyzed for confirmation. In cases where the BLL was below the LOQ, values were rounded up to $LOQ/\sqrt{2}$ (Hornung, 1990), given that only a small percentage of the data was censored (<5%) and BLL were distributed normally when this imputation was applied.

Statistical analyses

The rate of participation was calculated based on total number of deliveries that took place during the study period. When the BLL was determined at multiple points during the pregnancy, the highest recorded level was employed. $BLL \geq 50 \mu\text{g/L}$ and $\geq 100 \mu\text{g/L}$ were evaluated with a 95% confidence interval (CI 95%). The Pb variable was log-transformed in order to obtain a variable with a normal distribution (skewness-kurtosis test > 0.05). The log of BLL were compared according to the women's characteristics using one-way analysis of variance (ANOVA) tests. Multivariate analyses (generalized linear model) were then performed. The explanatory variable was the log of the maternal BLL during pregnancy. First, all explanatory variables were tested using binary models (model 1). Secondly, socioeconomic and sociodemographic variables that were significant using a threshold value of 20% were introduced into a multivariate model and progressively eliminated to obtain an "adjusted model" that contained only significant variables (using a threshold value of 5%). Age was used as the primary adjustment variable. Thirdly, nutritional and occupational factors were tested one by one using the adjusted model. Finally, nutrition and occupational factors (that were significant using a threshold value of 20%) were introduced into a

1
2
3 multivariate model and progressively eliminated according to a descending procedure to
4 obtain a “final model” containing only the significant variables (using a threshold value of
5 5%). We calculated the adjusted R^2 and tested the validity of the final model (normality of
6 residuals). The analyses were performed using STATA® v13.1 software.
7
8

9
10
11 The study was declared to the CNIL and was approved by the Institutional Review Board
12 (IRB00003888) of the French National Institute of Health and Medical Research (INSERM).
13
14

15 16 17 **Results** 18

19
20
21 During the study period, 613 women delivered, 554 were eligible for inclusion (59 did
22 not have BLL assessed during pregnancy), and 531 were included (5 declined to participate,
23 8 left the maternity ward before inclusion, and 10 did not participate for other reasons). The
24 overall participation rate was 86.6%. The population is described in Table 1. The average
25 age was 27 years. Approximately 40% of women were born in French Guiana and 51.7% in
26 Suriname, 81.2% of the women were of bushi-nenge origin, 62.9% lived in St. Laurent du
27 Maroni, 17.2% resided along the Maroni River and 7% reported Suriname as their primary
28 residence. Approximately 25% of the population did not attend school. The geometric mean
29 (GM) of BLL was 32.6 $\mu\text{g/L}$ (max: 259 $\mu\text{g/L}$) where 25.8% (CI 95 % = [22.1-29.5]) and 5.1%
30 (CI 95 % = [3.2-7.0]) of the women had BLL $\geq 50 \mu\text{g/L}$ and $\geq 100 \mu\text{g/L}$, respectively. Only
31 4.1% of BLL were under the LOQ.
32
33
34
35
36
37
38
39
40
41
42

43 The comparison between GM of BLL according to sociodemographic and
44 socioeconomic characteristics is presented in Table 1. BLL were significantly higher in:
45 women born in Suriname whose first language was nengue tongo, who resided along the
46 Maroni River (Apatou or Grand Santi), with a low level of education (an inverse correlation
47 between BLL and level of education was observed), had irregular employment (“jobbing”) or
48 undocumented status, and did not have enough money to live on. The highest BLL were
49 noted for women living in Apatou and Grand Santi (49.4 $\mu\text{g/L}$ and 59.8 $\mu\text{g/L}$, respectively).
50 Women who had no schooling displayed BLL of 45.1 $\mu\text{g/L}$. BLL were mostly recorded from
51
52
53
54
55
56
57
58
59
60

1
2
3 week 20 to delivery (62%), then from week 12 to week 20 (32%). BLL collected from week 12
4
5 to week 20 were significantly lower.

6
7 The comparison of BLL according to nutritional habits during pregnancy is depicted in
8
9 Table 2. BLL were significantly higher in women who regularly (daily or weekly) consumed
10
11 manioc, all manioc derivatives combined (specifically couac, domi and cassava), wild game
12
13 and fish. A positive correlation was found between BLL and frequency with which manioc
14
15 derivatives were consumed. Notably, 73.9% of the population regularly consumed manioc
16
17 derivatives, and 61.4% regularly consumed couac. The BLL was higher when couac was
18
19 prepared personally (56 µg/L) and when woman helped prepare couac during her pregnancy
20
21 (49.6 µg/L) or indicated that rainwater was consumed (40.5 µg/L) (Table 3). Drinking bottled
22
23 water was associated with lower BLL (21.6 µg/L). No correlation was observed with alcohol
24
25 and tobacco consumption during pregnancy.

26
27 The results from the multivariate analysis are presented in Table 4. Among the
28
29 socioeconomic and sociodemographic variables, only native language, place of residence,
30
31 level of education and gestational age at the time BLL was assessed, remained significantly
32
33 associated with an elevated BLL in the multivariate model after the descending procedure
34
35 (data not shown). These four variables, as well as age, were included in the adjusted model.
36
37 In the final model (model 3), the variables associated with a risk of elevated BLL were:
38
39 speaking nengue tongo as a first language, place of residence (Grand Santi), a low level of
40
41 education (primary school only or no schooling), consumption of couac (weekly or daily),
42
43 participating in couac preparation during pregnancy and weekly consumption of wild game.
44
45 Drinking rainwater no longer appeared to be a risk factor for elevated BLL in this final model.
46
47 The adjusted R^2 of the final model was 0.31, and the graphic distribution of residuals was
48
49 normal. The “consumption of manioc derivatives” variable was not included in the final model
50
51 to avoid the risk of overadjustment. However, when it was substituted for the “couac
52
53 consumption” variable in the final model, daily consumption of manioc derivatives was
54
55 significantly associated with an elevated BLL. Finally, 90% of women who participated in this
56
57 study had at least one of the risk factors identified in the final model.
58
59
60

Discussion

In our study, the GM of BLL in women was 32.6 µg/L, while 25.8% had a BLL above the threshold for intervention in minors in France (50 µg/L since 2015) (HCSP, 2014). The average BLL observed is approximately 4-fold higher than that found in metropolitan France in 2011 (study of 1968 women delivering in 211 maternity wards: BLL = 8.30 µg/L) (Guldner et al. 2014), and 5-fold higher than that found in USA between 2003 and 2008 (NHANES Study, BLL = 6.4 µg/L) (Jones et al., 2010). However, the BLL observed in this study did not reach the levels noted in Brazil, where average BLL between 57 and 127 µg/L were found in pregnant women or women of reproductive age (Barbosa Jr. et al., 2009; Carneiro et al., 2013; Zentner and Rondó, 2004).

The first set of risk factors identified in our study was associated with belonging to a disadvantaged socioeconomic class. These risk factors were identified in many studies (Barbosa Jr. et al., 2009; Carneiro et al., 2013; Hertz-Picciotto et al., 2000; Jones et al., 2010; McKelvey et al., 2007). There are many potential explanations for the link between a precarious socioeconomic status and Pb poisoning: nutritional deficiencies such as iron and calcium promote absorption of metal, or environmental sources (Tagne-Fotso et al., 2016). Pregnant women in western French Guiana are particularly likely to have anemia (70%) and iron deficiency (Louison-Ferté et al., 2014). In addition, the soil in French Guiana is particularly poor in iron and calcium, which might promote the persistence of deficiencies in populations whose food intake is primarily based on farming (ANSES, 2015).

Independent of socioeconomic characteristics, elevated BLL were identified in women who consumed manioc derivatives in a dose-dependent manner. Analyses performed by the French authorities in 2011 found high Pb content in some manioc and couac samples. An analytical study of Pb in manioc and its derivatives was conducted by the French-Guiana Regional Agency of Health in 2012-2013 by sampling communities, then farmers (ANSES, 2015). In total, 86 manioc samples and 50 couac samples were collected from homemade or

1
2
3 familial sources of manioc treatment or preparation (industrial production does not exist in
4 French Guiana). The average metal content based on “fresh weight” was 0.06 mg/kg in
5 manioc roots and 0.19 mg/kg in couac. According to this investigation, 24% of couac
6 samples and 14% of root samples exceeded the limit permitted for sale (established by the
7 European Regulation CE 1881/2006 as 0.1 mg/kg of fresh manioc). These data are a cause
8 for concern, because couac is a staple food in French Guiana, as it is in Brazil (mainly in the
9 northern and northeastern regions) and Africa (Adayemi et al., 2016). In our study, 21.5% of
10 women consumed daily couac, and 46% consumed couac at least twice a week. Several
11 different hypotheses may account for this contamination, including initial contamination of the
12 manioc roots from soil or Pb transfer during manioc processing: shredding, grinding and
13 cooking on hotplates. There is no ongoing or historic industrial or mining activity (outside of
14 gold mining) that might explain contamination of soil or waterways. An expert assessment
15 conducted in French Guiana (by the Bureau de Recherches Géologiques et Minières de
16 Guyane in 2013) provided little information. This study did not measure concentrations in
17 soil, only in waterway sediments. The observed Pb levels were relatively comparable to
18 those that occur naturally in large rivers (Horowitz, 1999), although elevated levels were
19 observed in certain areas of French Guiana. No data are available for the Grand-Santi and
20 Apatou regions. Finally, these data do not exclude the possibility of soil contamination in
21 French Guiana, and more analyses are needed (ANSES, 2015). Traditional agricultural
22 practices based on slashing and burning may be at the origin of zones with the highest
23 concentrations of metal. Contamination of manioc flour by Pb transfer from hotplates during
24 cooking was proposed by Barbosa Jr. et al. (2009), who found significantly higher levels of
25 metal in manioc flour compared to manioc root (flour: 0.19 ± 0.1 mg/kg, manioc: $0.017 \pm$
26 0.016 mg/kg). This finding was also reported in a similar analysis performed later in another
27 village (Barbosa Jr. et al., 2009). In French Guiana, a significant degree of metal enrichment
28 was observed during shredding and grinding of manioc roots due to the use of homemade
29 tools, but not during the cooking stage (ANSES, 2015). This observation, which was only
30 conducted at two sites, does not allow us to draw any formal conclusions. In addition, in our
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 study, women who participated in preparing couac during their pregnancies were found to
4 exhibit higher Pb levels. Data suggest that Pb poisoning might also occur by inhaling metal
5 particles while preparing couac.
6
7

8
9 The other risk factors identified in our investigation were weekly consumption of wild
10 game, which is well-known (Tagne-Fotso et al., 2016), however, this was not apparent in this
11 study. We did not find any link between geophagy (consuming pemba, local clay) and BLL,
12 even though this has been identified as a source of lead poisoning in other contexts
13 (Bakhireva et al., 2013; Thihalolipavan et al., 2013). In contrast, Lambert et al. (2010) noted
14 elevated Pb in pemba itself in French Guiana. It is possible that this item was underreported
15 due to reluctance to admit to consuming pemba to a healthcare provider. It is also possible
16 that pemba sequesters Pb (due to its chelating ability), thereby rendering it less bioavailable
17 (ANSES, 2015). Drinking rainwater appeared to be linked to metal exposure, even though
18 the association was not significant. In French Guiana, house roofs are typically composed of
19 sheets of steel, and thus it is possible that rainwater becomes contaminated due to
20 degradation of the metal. Environmental investigations would be required to confirm this link.
21
22 Cooking utensils, particularly those made of ceramic, tin and crystal, are also known as
23 sources of Pb exposure (Lynch et al., 2008). However, these utensils were not found in
24 environmental field studies. Lead poisoning due to the use of cooking utensils of unregulated
25 quality remains a possibility in French Guiana, as Pb may be released during cooking,
26 especially of acidic foods (ANSES, 2015). This information was not collected during our
27 study, nor was information collected regarding traditional remedies or cosmetics. Waste
28 treatment is another issue in French Guiana, particularly in western French Guiana, where all
29 waste is discarded in landfills without being recycled or incinerated. In addition, there are no
30 means of recycling hazardous waste such as car batteries, and illegal waste recovery may
31 occur via Suriname. This could be the origin of localized soil contamination. Sucking on a
32 piece of metal containing a high concentration of Pb is another possible source of
33 contamination, particularly in children (Meyer et al., 2008). This source of contamination
34 could explain the Pb poisoning of the 3-year-old girl in Mana, where car frames containing
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 batteries were accessible to children (ANSES, 2015). However, this possibility is essentially
4 limited to the coastal areas, as there are still few cars in the Maroni river villages, which can
5 only be accessed from the river.
6
7

8
9 There are a number of strengths of our study: a high level of participation (86.6%), a
10 relatively large number of women included in the study (N = 531) and the fact that data
11 collection was performed in local languages. There are several limitations that need to be
12 noted. First, this was an observational study, and BLL were measured at different points
13 during pregnancy. We did observe a U-shaped curve, with the lowest point between 12 and
14 20 weeks of pregnancy, as reported in the literature (Hertz-Picciotto et al., 2000; Miranda et
15 al., 2010). However, this variable did not modify the link we observed between nutritional
16 factors and BLL. Secondly, 59 women (9.6% of the deliveries that occurred during the study
17 period) were not included because their BLL were not collected during pregnancy. We did
18 not collect any information about these women, but it is likely that they received little or no
19 follow-up, and that the BLL collection was omitted at delivery. Many women arrive at the
20 CHOG in labor without any previous follow-up. According to the regional delivery register,
21 this was true for 9.2% of the woman who delivered at the CHOG in 2013, versus only 0.6% in
22 our sample. This may have led to a recruitment bias, as these women often live in precarious
23 situations. Thirdly, is the choice of using maximal BLL, which may have led to a slight
24 overestimation of metal levels. However, this overestimate does not appear to have exerted
25 a substantial effect, as the GM of the first BLL was 32 µg/L. The fourth limitation is linked to
26 the fact that BLL during pregnancy is not solely due to contemporary exposure, but is also
27 associated with remobilization of metal that has accumulated in the bones in the past
28 (Miranda et al., 2010).
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49
50
51 Since 2012, systematic screening has been provided for pregnant women in western
52 French Guiana. Based on the results from our study, this screening needs to be continued as
53 90% of women displayed at least one risk factor and 25% of women exhibited a BLL above
54 the threshold for intervention. Consequently women may benefit from health and dietary
55
56
57
58
59
60

1
2
3 advice, as well as iron and calcium supplementation. The impact of this metal exposure on
4
5 cognitive development in children is problematic, even more so when combined with other
6
7 heavy metal poisoning. Indeed, mercury (Hg) poisoning has been recognized in French
8
9 Guiana for about 20 years, and associated with consumption of river fish and panning for
10
11 gold (Fréry et al., 2001). Mercury poisoning primarily affects the residents of Haut-Maroni.
12

13 14 15 **Conclusion**

16
17
18
19 This is the first study conducted in pregnant women in western French Guiana to
20
21 provide evidence to raise public health problem. This study has shed light on regional and
22
23 social disparities in BLL in western French Guiana and potential sources of contamination.
24
25 Our study identified foods that are widely prepared and consumed by the local population as
26
27 possible metal sources. Given the existing data, recommendations and specific actions in
28
29 terms of prevention, screening and care for pregnant women and women of reproductive age
30
31 need to be adapted and put into place, as well as cooperation with bordering countries that
32
33 are facing similar health issues.
34
35
36
37
38

39 **Acknowledgements**

40
41
42
43 The authors would like to thank the Agence-Régionale de Santé (ARS) de Guyane,
44
45 who helped fund this study, the CHOG maternity unit and the midwives who participated in
46
47 the study (Sarah Belkacem, Mathilde Firmin, Caroline Hochart, Rainer Kramer, Amélie
48
49 Lehouillier, Bénédicte Mence, Marlène Rémy, Marion Restrepo, Pauline Stocq), the Périnat
50
51 Guyane network (Mme Anne Louison and Dr Anne Favre), the CICEC Antilles-Guyane (Mme
52
53 Stéphanie Rogier), and Dr. Emily Crow who assisted with translation of the manuscript. The
54
55 ARS did not participate in the writing of the article.
56
57
58
59
60

References

- Adeyemi, J.A., Adedire, C.O., Paulelli, A.C., da Cunha Martins Jr., A., Ileke, K.D., Barbosa Jr., F., 2016. Levels and daily intake of lead (Pb) and six essential elements in gari samples from Ondo State, Southwest Nigeria: A potential risk factor of health status. *J. Food Compos. Anal.* 45: 34–38.
- ANSES (French Agency for Food, Environmental and Occupational Health Safety), 2015. AVIS de l'Anses relatif à une demande d'appui scientifique et technique concernant le signalement d'une contamination au plomb de tubercules de manioc et des produits dérivés consommés en Guyane, Paris, France. <https://www.anses.fr/fr/content/avis-de-l%E2%80%99anses-relatif-%C3%A0-une-demande-dappui-scientifique-et-technique-concernant-le> (accessed 11.23.15).
- Attina, T.M., Trasande, L., 2013. Economic costs of childhood lead exposure in low- and middle-income countries. *Environ. Health Perspect.* 121: 1097–1102.
- Bakhireva, L.N., Rowland, A.S., Young, B.N., Cano, S., Phelan, S.T., Artyushkova, K., Rayburn, W.F., Lewis, J., 2013. Sources of potential lead exposure among pregnant women in New Mexico. *Matern. Child Health J.* 17: 172–179.
- Barbosa Jr., F., Fillion, M., Lemire, M., Sousa Passos, C.J., Lisboa Rodrigues, J., Philibert, A., Guimarães, J.-R., Mergler, D., 2009. Elevated blood lead levels in a riverside population in the Brazilian Amazon. *Environ. Res.* 109, 594–599.
- Bellinger, D.C., 2012. Comparing the population neurodevelopmental burdens associated with children's exposures to environmental chemicals and other risk factors. *Neurotoxicology* 33: 641–643.
- Bellinger, D.C., 2008. Very low lead exposures and children's neurodevelopment. *Curr. Opin. Pediatr.* 20: 172–177.

- 1
2
3 Borja-Aburto, V.H., Hertz-Picciotto, I., Rojas Lopez, M., Farias, P., Rios, C., Blanco, J., 1999. Blood lead
4
5 levels measured prospectively and risk of spontaneous abortion. *Am. J. Epidemiol.* 150:
6
7 590–597.
8
- 9 Buchanan, L.H., Counter, S.A., Ortega, F., 2011. Environmental lead exposure and otoacoustic
10
11 emissions in Andean children. *J Toxicol Environ Health A* 74: 1280-1293.
12
13
- 14 Carneiro, M.F.H., Evangelista, F.S. de B., Barbosa, F., Jr, 2013. Manioc flour consumption as a risk
15
16 factor for lead poisoning in the Brazilian Amazon. *J. Toxicol. Environ. Health A* 76: 206–216.
17
18
- 19 Counter, S.A. , Buchanan, L.H., Ortega, F., 2015. Blood levels in Andean infants and young children in
20
21 Ecuador: An international comparison. *J. Toxicol. Environ. Health A* 78: 778-787.
22
23
- 24 Fewtrell, L.J., Prüss-Ustün, A., Landrigan, P., Ayuso-Mateos, J.L., 2004. Estimating the global burden
25
26 of disease of mild mental retardation and cardiovascular diseases from environmental lead
27
28 exposure. *Environ. Res.* 94: 120–133.
29
- 30 Freire, C., Koifman, R.J., Fujimoto, D., de Oliveira Souza, V.C., Barbosa, F., Koifman, S., 2014.
31
32 Reference values of lead in blood and related factors among blood donors in the Western
33
34 Amazon, Brazil. *J. Toxicol. Environ. Health A* 77, 426–440.
35
36
- 37 Fréry, N., Maury-Brachet, R., Maillot, E., Deheeger, M., de Mérona, B., Boudou, A., 2001. Gold-mining
38
39 activities and mercury contamination of native Amerindian communities in French Guiana:
40
41 key role of fish in dietary uptake. *Environ. Health Perspect.* 109: 449–456.
42
43
- 44 Garcia-Leston, J., Roma-Torres, J., Mayan O., Schroecksadel, S., Fuchs, D., Moreira, A.O., Pásaro, E.,
45
46 Méndez, J., Teixeira, J.P., Laffon, B., 2012. Assessement of immunotoxicity parameters in
47
48 individuals occupationally exposed to lead. *J. Toxicol. Environ. Health A* 75: 807-818..
49
- 50 Guldner, L., Saoudi, A., Deremeaux, C., Pécheux, M., Lefranc, A, 2014. Describing exposures to
51
52 environmental contaminants in mothers of newborns in France, 2011 : First results
53
54 obtained in the framework of the French biomonitoring program. Poster presented at: ISEE-
55
56 EUROPE 2014. Young Researchers Conference on Environmental Epidemiology; 2014 Oct
57
58
59
60

1
2
3 20-21; Barcelona, Spain.
4
5 http://invs.santepubliquefrance.fr/content/download/98210/354381/version/2/file/Describing_exposures_environmental_contaminants_mothers_newborns_France_2011.pdf
6
7
8

9
10 HCSP (French High Council for Public Health), 2015. Détermination de nouveaux objectifs de gestion
11 des expositions au plomb: Synthèse et recommandations, Paris, France.
12
13 <http://www.hcsp.fr/explore.cgi/avisrapportsdomaine?clefr=445>.
14
15

16 Hertz-Picciotto, I., Schramm, M., Watt-Morse, M., Chantala, K., Anderson, J., Osterloh, J., 2000.
17
18 Patterns and determinants of blood lead during pregnancy. *Am. J. Epidemiol.* 152: 829–837.
19

20 Horowitz, H.J., Meybeck, M., Idlafkih, Z., Biger, E., 1999. Variations in trace element geochemistry in
21 the Seine River Basin based on floodplain deposits and bed sediments. *Hydrol. Process* 13:
22 1329-1340.
23
24
25
26

27
28 Hu, H., Téllez-Rojo, M.M., Bellinger, D., Smith, D., Ettinger, A.S., Lamadrid-Figueroa, H., Schwartz, J.,
29
30 Schnaas, L., Mercado-García, A., Hernández-Avila, M., 2006. Fetal lead exposure at each
31 stage of pregnancy as a predictor of infant mental development. *Environ. Health Persp.* 114:
32 1730–1735.
33
34
35
36

37 Jedrychowski, W., Perera, F., Jankowski, J., Rauh, V., Flak, E., Caldwell, K.L., Jones, R.L., Pac, A.,
38
39 Lisowska-Miszczuk, I., 2008. Prenatal low-level lead exposure and developmental delay of
40 infants at age 6 months (Krakow inner city study). *Int. J. Hyg. Environ. Health* 211: 345–351.
41
42

43 Jones, L., Parker, J.D., Mendola, P., 2010. Blood lead and mercury levels in pregnant women in the
44 United States, 2003-2008. NCHS Data Brief 1–8.
45
46

47 Kennedy, D.A., Woodland, C., Koren, G., 2012. Lead exposure, gestational hypertension and pre-
48 eclampsia: A systematic review of cause and effect.
49
50 *J. Obstet. Gynaecol.* 32: 512–517
51
52

53 Laborde, A., Tomasina, F., Bianchi, F., Bruné, M.-N., Buka, I., Comba, P., Corra, L., Cori, L., Duffert,
54
55 C.M., Harari, R., Iavarone, I., McDiarmid, M.A., Gray, K.A., Sly, P.D., Soares, A., Suk, W.A.,
56
57
58
59
60

- 1
2
3 Landrigan, P.J., 2015. Children's health in Latin America: The influence of environmental
4 exposures. *Environ. Health Perspect.* 123: 201–209.
- 5
6
7 Lamadrid-Figueroa, H., Téllez-Rojo, M.M., Hernández-Cadena, L., Mercado-García, A., Smith, D.,
8
9 Solano-González, M., Hernández-Avila, M., Hu, H., 2006. *J. Toxicol. Environ. Health A* 69:
10
11 1781-1796.
- 12
13
14 Lambert, V., Boukhari, R., Nacher, M., Goullé, J.-P., Roudier, E., Elguindi, W., Laquerrière, A., Carles,
15
16 G., 2010. Plasma and urinary aluminum concentrations in severely anemic geophagous
17
18 pregnant women in the Bas Maroni region of French Guiana: A case-control study.
19
20
21 *Am. J. Trop. Med. Hyg.* 83: 1100–1105.
- 22
23 Lanphear, B.P., Hornung, R., Khoury, J., Yolton, K., Baghurst, P., Bellinger, D.C., Canfield, R.L., Dietrich,
24
25 K.N., Bornschein, R., Greene, T., Rothenberg, S.J., Needleman, H.L., Schnaas, L., Wasserman,
26
27 G., Graziano, J., Roberts, R., 2005. Low-level environmental lead exposure and children's
28
29 intellectual function: An international pooled analysis. *Environ. Health Perspect.* 113: 894–
30
31 899.
- 32
33
34 Louison-Ferté, A., Jolivet, A., Lambert, V., Bosquillon, L., Carles, G., 2014. Lutte contre l'anémie de la
35
36 femme enceinte dans l'Ouest guyanais : diagnostic et mise en oeuvre d'actions par le
37
38 réseau Périnatal Guyane autour d'une évaluation des pratiques professionnelles. *Rev. Med*
39
40
41 *Perinat* 6: 116–121.
- 42
43 Lynch, R., Elledge, B., Peters, C., 2008. An assessment of lead leachability from lead-glazed ceramic
44
45 cooking vessels. *J. Environ. Health* 70: 36–40, 53.
- 46
47 McKelvey, W., Gwynn, R.C., Jeffery, N., Kass, D., Thorpe, L.E., Garg, R.K., Palmer, C.D., Parsons, P.J.,
48
49 2007. A biomonitoring study of lead, cadmium, and mercury in the blood of New York City
50
51 adults. *Environ. Health Perspect.* 115: 1435–1441.
- 52
53 Meyer, P.A., Brown, M.J., Falk, H., 2008. Global approach to reducing lead exposure and poisoning.
54
55
56 *Mutat. Res.* 659, 166–175.
- 57
58
59
60

- 1
2
3 Miranda, M.L., Edwards, S.E., Swamy, G.K., Paul, C.J., Neelon, B., 2010. Bloodlead levels among
4 pregnant women: Historical versus contemporaneous exposures.
5
6 *Int. J. Environ. Res. Public. Health* 7: 1508–1519.
7
8
9 NTP (National Toxicology Program), 2012. NTP monograph on health effects of low-level lead. NTP
10 Monogr. xiii, xv-148.
11
12
13 Osman, K., Akesson, A., Berglund, M., Bremme, K., Schütz, A., Ask, K., Vahter, M., 2000. Toxic and
14 essential elements in placentas of Swedish women. *Clin. Biochem.* 33: 131–138.
15
16
17 Pollack, A.Z., Mumford, S.L., Mendola, P., Perkins, N.J., Rotman, Y., Wactawski-Wende, J.,
18 Schisterman, E.F., 2015. Kidney biomarkers associated with blood lead, mercury and
19 cadmium in premenopausal women: A prospective cohort study. *J. Toxicol. Environ. Health*
20 *A* 78: 119-131.
21
22
23
24
25
26
27 Rorive, S., Boukhari, R., Harrois, D., 2015. Saturnisme: Un premier cas en Guyane découvert sur le
28 frottis sanguin. *Rev. Francoph. Lab.* 2015: 79–80.
29
30
31
32 Rothenberg, S.J., Karchmer, S., Schnaas, L., Perroni, E., Zea, F., Fernández Alba, J., 1994. Changes in
33 serial blood lead levels during pregnancy. *Environ. Health Perspect.* 102: 876–880.
34
35
36 Schnaas, L., Rothenberg, S.J., Flores, M.-F., Martinez, S., Hernandez, C., Osorio, E., Velasco, S.R.,
37 Perroni, E., 2006. Reduced intellectual development in children with prenatal lead
38 exposure. *Environ. Health Perspect.* 114: , 791–797.
39
40
41
42 Suk, W.A., Ahanchian, H., Asante, K.A., Carpenter, D.O., Diaz-Barriga, F., Ha, E.-H., Huo, X., King, M.,
43 Ruchirawat, M., da Silva, E.R., Sly, L., Sly, P.D., Stein, R.T., van den Berg, M., Zar, H.,
44 Landrigan, P.J., 2016. Environmental pollution: An under-recognized threat to children’s
45 health, especially in low- and middle-income countries. *Environ. Health Perspect.* 124: A41-
46 A45.
47
48
49
50
51
52
53 Tagne-Fotso, R., Leroyer, A., Howsam, M., Dehon, B., Richeval, C., Members of Health Examination
54 Centres of Nord-Pas-de-Calais region network, Nisse, C., 2016. Current sources of lead
55 exposure and their relative contributions to the blood levels in the general adult population
56
57
58
59
60

1
2
3 of Northern France: The IMEPOGE Study, 2008-2010. *J. Toxicol. Environ. Health A* 79: 245-
4
5 265.

6
7
8 Taylor, C., Golding, J., Emond, A., 2015. Adverse effects of maternal lead levels on birth outcomes in
9
10 the ALSPAC study: A prospective birth cohort study. *Br J Obstet Gynaecol* 122: 322–328.

11
12 Thihalolipavan, S., Candalla, B.M., Ehrlich, J., 2013. Examining pica in NYC pregnant women with
13
14 elevated blood lead levels. *Matern. Child Health J.* 17: 49–55.

15
16 Tsoi, M.-F., Cheung, C.-L., Cheung, T.T., Cheung, B.M.Y., 2016. Continual decrease in blood lead level
17
18 in Americans: United States National Health Nutrition and Examination Survey 1999-2014.
19
20 *Am. J. Med.* 129: 1213-1218.

21
22 Valent, F., Little, D., Bertollini, R., Nemer, L.E., Barbone, F., Tamburlini, G., 2004. Burden of disease
23
24 attributable to selected environmental factors and injury among children and adolescents
25
26 in Europe. *Lancet* 363: 2032–2039.

27
28
29 WHO (World Health Organization) Europe, 2009. Levels of lead in children's blood. European
30
31 Environnement and Health Information System. Fact sheet No 4.5,
32
33 [http://www.euro.who.int/__data/assets/pdf_file/0003/97050/4.5.-Levels-of-lead-in-](http://www.euro.who.int/__data/assets/pdf_file/0003/97050/4.5.-Levels-of-lead-in-childrens-blood-EDITING_layouted.pdf)
34
35 [childrens-blood-EDITING_layouted.pdf](http://www.euro.who.int/__data/assets/pdf_file/0003/97050/4.5.-Levels-of-lead-in-childrens-blood-EDITING_layouted.pdf). n.d.

36
37
38 Zentner, L.E.A., Rondó, P.H.C., 2004. Lead contamination among pregnant Brazilian women living
39
40 near a lead smelter. *Int. J. Gynaecol. Obstet. Off. Organ Int. Fed. Gynaecol. Obstet.* 87: 147–
41
42 148.

43
44
45 Zhu, M., Fitzgerald, E.F., Gelberg, K.H., Lin, S., Druschel, C.M., 2010. Maternal low-level lead exposure
46
47 and fetal growth. *Environ. Health Persp.* 118: 1471–1475.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Area served by the St-Laurent-du-Maroni Hospital



Table 1: Participants' characteristics and BLL ($\mu\text{g/L}$).

		n	%	GM*	P-value
Age	<18	37	7.0	32.7	0.42
	18-24	198	37.3	31.2	
	25-35	219	41.2	32.7	
	>35	77	14.5	36.3	
Place of birth	French Guiana	214	40.4	32.2	$p < 10^{-3}$
	Surinam	274	51.7	35.9	
	Brazil	17	3.2	25.6	
	Other	25	4.7	15.1	
Native language	French	33	6.3	17.2	$p < 10^{-3}$
	Nenge tongo	427	81.2	37.3	
	Amerindian language	7	1.3	19.4	
	Portuguese	20	3.8	23.8	
	Other	39	7.4	18.1	
Primary place of residence	St Laurent	333	62.9	30.0	$p < 10^{-3}$
	Mana/Awala	54	10.2	25.6	
	Apatou	49	9.3	48.7	
	Grand Santi	42	7.9	59.8	
	Surinam	37	7.0	28.9	
	Other	14	2.7	28.0	
Level of education	No schooling	124	23.6	45.1	$p < 10^{-3}$
	Primary	60	11.4	41.8	
	Secondary	146	27.8	31.1	
	High school	160	30.4	27.5	
	University	36	6.8	18.1	
Employment status before pregnancy	Student	102	19.4	30.1	$p < 10^{-3}$
	Employed	85	16.1	23.9	
	Unemployed	21	4.0	31.4	
	Irregular employment	49	9.3	40.1	
	Homemaker	270	51.2	35.7	
Legal status	French citizen or foreign citizen with 10-year French territory residence card	238	48.6	28.9	0.0007
	Temporary documented migrant (1-year stay document)	77	15.7	31.3	
	Undocumented migrant	175	35.7	37.0	
Financial means	Not enough	114	21.6	40.9	$p < 10^{-3}$
	Just enough	269	50.9	33.0	
	Adequate	145	27.5	27.0	
Gestational age at BLL collection	<12 weeks	34	6.4	29.3	0.006
	12-20 weeks	169	31.8	26.7	
	>20 weeks	328	61.8	36.6	

* GM: geometric mean of BLL ($\mu\text{g/L}$)

Table 2: BLL ($\mu\text{g/L}$) according to nutritional habits and lifestyle factors during pregnancy.

		n	%	GM**	p
Manioc	Never	171	32.6	30.6	$p < 10^{-3}$
	Monthly	140	26.7	29.0	
	Weekly	157	29.9	35.1	
	Daily	57	10.8	44.2	
All derivatives	Never	33	6.2	19.2	$p < 10^{-3}$
	Monthly	81	15.3	24.8	
	Weekly	250	47.4	33.3	
	Daily	164	31.1	40.1	
Couac*	Never	93	17.6	26.1	$p < 10^{-3}$
	Monthly	111	21.0	26.3	
	Weekly	210	39.8	35.8	
	Daily	114	21.6	40.3	
Domi*	Never	127	24.4	25.3	$p < 10^{-3}$
	Monthly	133	25.5	30.1	
	Weekly	185	35.5	37.6	
	Daily	76	14.6	42.5	
Cassava*	Never	210	40.1	29.7	0.005
	Monthly	160	30.5	31.2	
	Weekly	118	22.5	36.7	
	Daily	36	6.9	41.7	
Crabio*	Never	376	71.9	33.4	0.23
	Monthly	126	24.1	30.2	
	Weekly	11	2.1	42.8	
	Daily	10	1.9	28.8	
Tapioca*	Never	359	68.9	33.6	0.03
	Monthly	127	24.4	29.6	
	Weekly	30	5.8	30.9	
	Daily	5	0.9	65.8	
Wild game	Never	210	40.4	28.9	$p < 10^{-3}$
	Monthly	205	39.4	32.7	
	Weekly	85	16.4	43.9	
	Daily	20	3.8	37.9	
Fish	Never	25	4.8	30.0	0.001
	Monthly	86	16.3	27.2	
	Weekly	291	55.2	32.1	
	Daily	125	23.7	38.9	
Wassaï*	Never	103	19.6	27.2	0.02
	Monthly	130	24.7	33.9	
	Weekly	205	39.0	33.6	
	Daily	88	16.7	35.5	
Pemba*	Never	286	54.8	31.9	0.27
	Monthly	112	21.5	33.5	
	Weekly	64	12.2	31.8	
	Daily	60	11.5	38.3	
Tobacco	Yes	18	3.4	25.9	0.14
	No	513	96.6	32.9	
Alcohol	Yes	180	33.9	34.1	0.27
	No	351	66.1	31.9	

*Couac (manioc flour), Cassava (manioc pancakes), Crabio (detoxified cooking liquid used as a sauce), Tapioca (bitter manioc starch), Wassaï (fruit of the wassaï palm tree), Pemba (local clay)

** GM: geometric mean of BLL ($\mu\text{g/L}$)

Table 3: BLL ($\mu\text{g/L}$) according to couac preparation and water consumption during pregnancy.

			n	%	GM**	p-value
Source of the couac		Personal	69	16.6	55.7	$p < 10^{-3}$
		Other*	346	83.4	31.7	
Personally prepared couac		No	313	60.8	27.2	$p < 10^{-3}$
		Yes	202	39.2	44.8	
Helped prepare couac		No	415	81.4	30.3	$p < 10^{-3}$
		Yes	95	18.6	49.3	
Water consumption	Public supply	No	235	44.3	35.1	0.03
		Yes	296	55.7	30.8	
	Bottled water	No	463	87.2	34.9	$p < 10^{-3}$
		Yes	68	12.8	20.7	
	Rain water	No	386	72.7	30.1	$p < 10^{-3}$
		Yes	145	27.3	40.4	
	Well water	No	498	93.8	32.4	0.33
		Yes	33	6.2	36.4	
	River water	No	498	93.8	32.3	0.15
		Yes	33	6.2	38.5	

* market, grocery store, family/friends, from the producer.

** GM: geometric mean of BLL ($\mu\text{g/L}$)

Table 4: Linear regression model of the logarithm of the BLL according to socioeconomic characteristics and nutritional factors (regression β coefficients and CI 95%).

		Model 1 (binary)	Model 2: Model 1 + Adjusted variables*	Model 3: Final model
Age	<25 years	0		0
	[25-35 years]	0.04 [-0.08 ; 0.16]		0.02 [-0.10 ; 0.13]
	>35 years	0.14 [-0.03 ; 0.32]		0.09 [-0.06 ; 0.25]
Native language	French and other	0		0
	Nenge tongo	0.68 [0.54 ; 0.81]		0.40 [0.25 ; 0.55]
Place of residence	SLM	0		0
	Mana/Awala	-0.16 [-0.34 ; 0.02]		0.02 [-0.15 ; 0.20]
	Apatou	0.48 [0.29 ; 0.67]		0.16 [-0.02 ; 0.34]
	Grand-Santi	0.69 [0.49 ; 0.89]		0.28 [0.07 ; 0.48]
	Other	-0.05 [-0.23 ; 0.14]		-0.10 [-0.27 ; 0.08]
Education level	High School	0		0
	University	-0.42 [-0.65 ; -0.19]		-0.25 [-0.47 ; -0.03]
	Secondary	0.12 [-0.02 ; 0.26]		-0.04 [-0.17 ; 0.10]
	Primary	0.42 [0.23 ; 0.60]		0.19 [0.01 ; 0.37]
	None	0.49 [0.35 ; 0.64]		0.20 [0.05 ; 0.35]
Gestational age	12-20 weeks	0		0
	<12 weeks	0.09 [-0.15 ; 0.33]		0.21 [-0.002 ; 0.42]
	>20 weeks	0.32 [0.19 ; 0.44]		0.28 [0.17 ; 0.39]
Consumption of Manioc	Never or monthly	0	0	
	Weekly	0.16 [0.03 ; 0.28]	0.06 [-0.06 ; 0.17]	
	Daily	0.37 [0.19 ; 0.55]	0.14 [-0.03 ; 0.31]	
Consumption of manioc derivatives	Never or monthly	0	0	
	Weekly	0.34 [0.21 ; 0.48]	0.17 [0.04 ; 0.30]	
	Daily	0.52 [0.38 ; 0.67]	0.24 [0.10 ; 0.39]	
Consumption of Couac	Never or monthly	0	0	0
	Weekly	0.29 [0.17 ; 0.41]	0.15 [0.04 ; 0.26]	0.15 [0.03 ; 0.26]
	Daily	0.40 [0.26 ; 0.55]	0.19 [0.05 ; 0.33]	0.17 [0.02 ; 0.31]
Participation in the Preparation of Couac	No	0	0	0
	Yes	0.48 [0.34 ; 0.62]	0.24 [0.10 ; 0.38]	0.20 [0.05 ; 0.34]
Consumption of Fish	Never or monthly	0	0	
	Weekly	0.14 [-0.003 ; 0.28]	0.10 [-0.03 ; 0.22]	
	Daily	0.32 [0.16 ; 0.49]	0.15 [-0.005 ; 0.30]	
Consumption of Wassai	Never or monthly	0	0	
	Weekly	0.07 [-0.05 ; 0.19]	-0.04 [-0.15 ; 0.07]	
	Daily	0.12 [-0.04 ; 0.28]	-0.01 [-0.16 ; 0.14]	
Consumption of Game	Never or monthly	0	0	0
	Weekly	0.35 [0.20 ; 0.50]	0.19 [0.05 ; 0.33]	0.17 [0.03 ; 0.30]
	Daily	0.19 [-0.09 ; 0.48]	0.01 [-0.26 ; 0.27]	0.00 [-0.26 ; 0.26]
Consumption of Pemba	Never or monthly	0	0	
	Weekly	-0.04 [-0.20 ; 0.13]	-0.05 [-0.21 ; 0.10]	
	Daily	0.15 [-0.02 ; 0.33]	0.05 [-0.11 ; 0.21]	
Rain water	No	0	0	
	Yes	0.28 [0.15 ; 0.40]	0.10 [-0.01 ; 0.22]	

* The adjusted variables used are age, level of education, native language, place of residence and gestational age
Significant β coefficients are indicated in bold