



HAL
open science

Phospholipases D ζ 1 and D ζ 2 have distinct roles in growth and antioxidant systems in *Arabidopsis thaliana* responding to salt stress

Alhem Ben Othman, Hasna Ellouzi, Séverine Planchais, Delphine de Vos, Bualuang Faiyue, Pierre Carol, Chedly Abdelly, Arnould Savouré

► To cite this version:

Alhem Ben Othman, Hasna Ellouzi, Séverine Planchais, Delphine de Vos, Bualuang Faiyue, et al.. Phospholipases D ζ 1 and D ζ 2 have distinct roles in growth and antioxidant systems in *Arabidopsis thaliana* responding to salt stress. *Planta*, 2017, 246 (4), pp.721-735. 10.1007/s00425-017-2728-2. hal-01557388

HAL Id: hal-01557388

<https://hal.sorbonne-universite.fr/hal-01557388>

Submitted on 6 Jul 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - ShareAlike 4.0 International License

1 Phospholipases D ζ 1 and D ζ 2 have distinct roles in growth and antioxidant systems in *Arabidopsis*
2 *thaliana* responding to salt stress

3
4 Ahlem Ben Othman^{1,2}, Hasna Ellouzi^{2,*}, Séverine Planchais^{1,*}, Delphine De Vos^{1,3}, Bualuang
5 Faiyue^{1,4}, Pierre Carol¹, Chedly Abdelly² and Arnould Savourel^{1,‡}

6
7 1Sorbonne Universités, UPMC Univ Paris 06, iEES, UMR 7618, UPMC Paris06 - Sorbonne
8 (UPEC, UPMC, CNRS, IRD, INRA, Paris Diderot), case 237, 4 place Jussieu, F-75252 Paris cedex
9 05, France

10 2 Laboratoire des Plantes Extrêmophiles, Centre de Biotechnologie de Borj-Cedria (CBBC), BP
11 901, Hammam-Lif 2050, Tunisia

12 3 Present address: Institut Jean-Pierre Bourgin, UMR 1318 INRA-AgroParisTech, Centre INRA
13 Versailles, 78026 Versailles Cedex, France.

14 4 Present address: Department of Biology, Mahidol Wittayanusorn School, Salaya, Phuttamonthon,
15 Nakhon Pathom 73170, Thailand

16
17 * These authors contributed equally to this work

18
19 ‡ Corresponding author

20 Telephone: +33 1 44 27 26 72

21 Fax : +33 1 44 27 35 16

22 Email: arnould.savourel@upmc.fr

23

24

25

26 Short running title

27 Roles of phospholipases D ζ during salt stress in *Arabidopsis*

28

29

30

31

32

33

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67

Main conclusion

Phospholipases D ζ play different roles in Arabidopsis salt tolerance affecting the regulation of ion transport and antioxidant responses.

Abstract

Lipid signalling mediated by phospholipase D (PLD) plays essential roles in plant growth including stress and hormonal responses. Here we show that *PLD ζ 1* and *PLD ζ 2* have distinct effects on Arabidopsis responses to salinity. A transcriptome analysis of a double *pld ζ 1pld ζ 2* mutant revealed a cluster of genes involved in abiotic and biotic stresses, such as the high salt stress responsive genes *DDF1* and *RD29A*. Another cluster of genes with a common expression pattern included ROS detoxification genes involved in electron transport and biotic and abiotic stress responses. Total SOD activity was induced early in the shoots and roots of all *pld ζ* mutants exposed to mild or severe salinity with the highest SOD activity measured in *pld ζ 2* at 14 days. Lipid peroxidation in shoots and roots was higher in the *pld ζ 1* mutant upon salt treatment and *pld ζ 1* accumulated H₂O₂ earlier than other genotypes in response to salt. Salinity caused less deleterious effects on K⁺ accumulation in shoots and roots of the *pld ζ 2* mutant than of wild type, causing only a slight variation in Na⁺/K⁺ ratio. Relative growth rates of wild-type plants, *pld ζ 1*, *pld ζ 2* and *pld ζ 1pld ζ 2* mutants were similar in control conditions but strongly affected by salt in WT and *pld ζ 1*. The efficiency of photosystem II, estimated by measuring the ratio of chlorophyll fluorescence (Fv/Fm ratio), was strongly decreased in *pld ζ 1* under salt stress. In conclusion, *PLD ζ 2* plays a key role in determining Arabidopsis sensitivity to salt stress allowing ion transport and antioxidant responses to be finely regulated.

Keywords: Ion relations; Phospholipase D; PLD ζ ; Reactive oxygen species; Salt stress; Transcriptome.

68
69
70

Introduction

71
72
73
74
75
76
77

Salinity is a major environmental constraint on the growth, productivity and diversity of plants. The effect of salt stress on plants depends on the salt concentration, the duration of exposure and the plant genotype (Ellouzi et al. 2011). The presence of salt in the environment induces water deficit in plants because the external water potential is lowered, while ion toxicity and nutritional alterations disturb ion transport systems (Munns and Tester 2008; Julkowska and Testerink 2015). Salt stress also causes membrane damage, alters levels of growth regulators, inhibits some enzymes, and disrupts photosynthesis, and may thus lead to plant death.

78
79
80
81
82
83
84
85
86
87
88
89
90
91

One of the known plant responses to salt stress is ROS production (for review see Miller et al. 2010; Ben Rejeb et al. 2014). Plant cells need to regulate ROS production as excess ROS is potentially harmful to nucleic acids, proteins and lipids, and may therefore lead to cell injury and death (Gill and Tuteja 2010). ROS produced through NADPH oxidase activity was shown to be mediated by phospholipid signalling (Zhang et al. 2009). The second messenger phosphatidic acid (PA) is a phospholipid which targets specific proteins to bring about cellular and physiological changes that allow plants to adapt to abiotic stresses (for review see Hong et al. 2010; Hou et al. 2016). PA is formed when phospholipase D (PLD) hydrolyses structural phospholipids at the terminal phosphoesteric bond with release of the hydrophilic head group. In plants, phospholipase D (PLD) is predominant among the phospholipase families (for review see Li et al. 2009; McLoughlin and Testerink, 2013). PLD activity increases rapidly in response to various environmental stresses such as cold, drought, and salinity (Vergnolle et al. 2005; Bargmann et al. 2009; Hong et al. 2010). Proline accumulation, another common physiological response to stress, was shown to be negatively regulated by PLD activity in *Arabidopsis thaliana* (Thiery et al. 2004).

92
93
94
95
96
97
98
99
100
101
102

In *A. thaliana*, the PLD family includes 12 members that are classified into six types, PLD α (3 isoforms), - β (2 isoforms), - γ (3 isoforms), - δ , - ϵ , and - ζ (2 isoforms), according to their sequences and enzymatic properties (Bargmann and Munnik 2006). Biochemical studies have revealed that the phospholipid-hydrolysing activities of PLD are either calcium-dependent through a C2 domain (C2-PLD) or calcium-independent having pleckstrin homology (PH) and phox homology (PX) domains (PXPH-PLD) (Meijer and Munnik 2003). Several PLD isoforms have been functionally characterized. The two PLD ζ s (*PLD ζ 1* and *PLD ζ 2*) are structurally different from other PLDs (for review see Li et al. 2009). PLD ζ 1 and PLD ζ 2 do not have C2 domains and do not require Ca²⁺ for enzymatic activity (Qin and Wang 2002), but they do have PH and PX domains (Qin and Wang 2002). *PLD ζ 1* gene expression is five-fold greater in roots than in leaves and the gene function is required for root hair morphogenesis (Li et al. 2006). The *PLD ζ 2* gene is mainly expressed in roots

103 (Li et al. 2006). *PLDζ2* gene expression is triggered by exogenous auxin (Li and Xue 2007) and
104 phosphate (Li et al. 2006). *PLDζ1* and *PLDζ2* were shown to be involved in the process by which
105 root architecture adapts to the lack of phosphate (Li et al. 2006). *PLDζ2* is also involved in vesicle
106 trafficking and auxin transport (Li and Xue 2007). The *pldζ2* mutant has a reduced halotropic
107 response, i.e. the capacity to change the direction of root growth to avoid salt, due to impaired
108 vesicle trafficking (Galvan-Ampudia et al. 2013).

109 In this study, the roles of *PLDζ1* and *PLDζ2* were investigated in Arabidopsis responding to salt
110 stress. The global transcriptome of the *pldζ1pldζ2* double mutant treated with salt was analysed to
111 identify candidate genes regulated by these PLDs. Changes in growth, ion balance, ROS content,
112 antioxidants, and lipid peroxidation were compared in wild type, *pldζ1*, *pldζ2*, and *pldζ1pldζ2* after
113 short and long periods of salt treatment. *pldζ1* and *pldζ2* mutants responded differently to salt stress
114 suggesting they have distinct physiological roles in Arabidopsis.

115

116

117 **Materials and methods**

118 Plant materials and growth conditions

119 *Arabidopsis thaliana* (L.) Heynh ecotype Columbia-0 (Col-0) was used as the wild type in this
120 study and was obtained from Nottingham Arabidopsis Stock Centre, Loughborough, United
121 Kingdom. Homozygous T-DNA insertion mutant lines *pldζ1* (SALK_083090) and *pldζ2A*
122 (SALK_094369) were obtained from the Salk Institute, La Jolla, USA (Alonso and Stepanova
123 2003) and have been described by Li et al. (2006). Double homozygous mutant *pldζ1pldζ2* plants
124 were isolated from the F2 progeny of crosses between *pldζ1* and *pldζ2*. The presence and
125 homozygosity of the T-DNA alleles were checked by triplex PCR in all experimental lines using a
126 primer specific to the left border of the T-DNA and two gene-specific primers flanking the insertion
127 site (Suppl. Fig. S1 and Suppl. Table S1).

128 Surface sterilized seeds of wild-type and mutant lines were sown onto grids placed on Petri dishes
129 containing 0.5 × Murashige and Skoog (MS; Murashige and Skoog 1962) solid medium (0.8 %
130 agar) according to Parre et al. (2007). Seeds were placed in a cold room at 4 °C for 1 day to break
131 dormancy and then transferred to a growth chamber at 22 °C with continuous light (90 μmol
132 photons m⁻² s⁻¹). Twelve-day-old Arabidopsis seedlings were exposed to 200 mM NaCl for up to 24
133 h.

134 For physiological experiments, wild type, *pldζ1*, *pldζ2* and *pldζ1pldζ2* seeds were germinated in
135 plastic pots (70 mL) filled with inert sand and watered daily with distilled water for one week.
136 Seedlings of each genotype were then irrigated with Hewitt nutrient solution (Hewitt 1966) for 3
137 weeks. The experiments were performed in a glasshouse under controlled conditions, photoperiod
138 of 16h/8h (day/night), at 20-25 °C and 65-75 % relative humidity. Four-week-old plants were
139 treated with 75 mM or 150 mM NaCl and collected after 3 h, 24 h and 72 h for short exposure to
140 salt stress and after 7 and 14 days for long exposure to salinity. Non-stressed plants grown without
141 added salt were collected and used as controls.

142

143 Transcriptome analysis of the *pldζ1pldζ2* double mutant

144 Analysis of the *pldζ1pldζ2* transcriptome was conducted at the Unité de Recherche en Génomique
145 Végétale (Evry, France) using the Complete Arabidopsis Transcriptome MicroArray (CATMA)
146 containing 24576 gene-specific tags from Arabidopsis (Crowe et al. 2003). Six biological replicates
147 and 14 dye swaps (technical replicates) were made. Twelve-day-old seedlings treated for 3 h with
148 either 200 mM NaCl or 400 mM mannitol for stress conditions or with 0.5 × MS for control
149 conditions were rapidly washed with water and immediately frozen in liquid nitrogen and stored at

150 –80 °C. Total RNA was isolated from seedlings by the guanidinium thiocyanate-CsCl purification
151 method (Sambrook et al. 1989). Labelling of cRNAs with Cy3-dUTP or Cy5-dUTP (Perkin-Elmer-
152 NEN Life Science Products, Courtaboeuf, France), hybridization to slides, and scanning were
153 performed as described in Lurin et al. (2004). Methods and data were deposited in CATMA
154 database RS06-01_PLD ([http://urgv.evry.inra.fr/cgi-bin/projects/CATdb/consult_expce.pl?](http://urgv.evry.inra.fr/cgi-bin/projects/CATdb/consult_expce.pl?experiment_id=113)
155 [experiment_id=113](http://urgv.evry.inra.fr/cgi-bin/projects/CATdb/consult_expce.pl?experiment_id=113)) and GEO (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE9459>).
156 Statistical analysis of transcriptome data was done as in Planchais et al. (2014).

157
158 Reverse transcription PCR and quantitative PCR analysis of gene expression
159 Total RNA was extracted from 100 mg of homogenized tissue from different genotypes using the
160 RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions. RNA
161 was quantified by measuring the absorbance at 260 nm using a Nanovue spectrophotometer
162 (Nanodrop Spectrophotometer ND 1000, Wilmington, DE, USA). RNA was incubated with DNase
163 I (Sigma-Aldrich, Saint-Louis, MO, USA) to eliminate genomic DNA. For reverse transcription,
164 first-strand cDNA was synthesized from 1.5 µg of total RNA using the RevertAid reverse
165 transcriptase kit (Thermo Fisher Scientific Inc., Waltham, MA, USA). Complementary DNAs were
166 amplified using DreamTaq Green polymerase (Thermo Fisher Scientific Inc.) and gene-specific
167 primers (Supplementary Table S1). The *APT1* gene (At1g27450) was used as a positive control for
168 quantifying relative amounts of cDNA. Amplified PCR fragments were separated on 2 % (w/v)
169 agarose gels and visualized by staining with ethidium bromide and analysing the image with
170 GelDoc (Bio-Rad, Hercules, USA). For real-time PCR, 5 µL of diluted cDNA was used with 10 µL
171 of SYBR®GreenqPCR Master Mix (Thermo Fisher Scientific Inc.) and gene-specific primers in an
172 Eppendorf Master cycler® (Eppendorf France SAS, Montesson, France). Critical thresholds (Ct)
173 were determined by using the Eppendorf Master cycler® realplex software and ratios were
174 calculated by using the method described by Pfaffl (2001). For each gene, a standard curve made
175 with dilutions of cDNA pools was used to calculate reaction efficiencies, and the levels of gene
176 transcript were normalized and expressed relative to the amounts observed under control conditions
177 with *APT1* (At1g27450) as housekeeping gene. All semi-quantitative RT-PCR and real-time PCR
178 experiments were carried out with three biological replicates.

179
180 Measurements of plant biomass, maximum efficiency of PSII photochemistry, and leaf water status
181 Fresh weight (FW) and dry weight (DW) of rosettes and roots were measured at each experimental
182 time point to evaluate leaf and root growth. The accumulation of biomass was estimated by
183 determining the relative growth rate (RGR), which is a measure of biomass production relative to

184 treatment duration and/or initial plant size. RGR was calculated as $\Delta \ln(DW)/\Delta t$, where DW is the dry
185 weight, \ln is the natural logarithm and Δ represents the difference between final and initial values for a given
186 time interval Δt (Hunt, 1990). The maximum efficiency of PSII photochemistry (F_v/F_m) was
187 monitored throughout the experiment using a portable mini-PAM fluorometer (ADC BioScientific
188 LC pro System Serial, Hoddesdon, UK).

189 The relative leaf water content (RWC) was calculated as $100 \times (FW - DW) / (TW - DW)$, where TW was
190 the turgid weight of leaves saturated for 24 h in deionized water at 4 °C in darkness and DW was
191 obtained after oven-drying the leaves at 60 °C for 72 h according to Ellouzi et al. (2013).

192

193 Measurement of ion content

194 Desiccated leaf and root tissues were ground to a fine powder and then broken down with
195 concentrated 0.5 % HNO_3 according to Deal (1954). Na^+ and K^+ contents were determined using
196 flame emission photometry (Corning, Tewksbury, MA, USA). In emission photometry the soluble
197 mineral component is injected into an air-propane flame. Following thermal excitation a
198 characteristic spectrum of lines is emitted that are selected by monochromator filters. Behind each
199 filter a photoreceptor cell detects the intensity of the emitted light which is proportional to the
200 amount of the emitting element contained in the vaporized solution.

201

202 Histochemical detection of hydrogen peroxide

203 The localization of hydrogen peroxide was determined according to Ben Rejeb et al. (2015). Fresh
204 leaf and root samples were collected from each mutant and infiltrated with a solution containing 0.5
205 $mg\ ml^{-1}$ diaminobenzidine (DAB) and 10 mM Mes buffer pH 5.8. Samples were then kept at room
206 temperature until the brown residue generated by H_2O_2 -DAB polymerization developed. Samples
207 were incubated for 60 min in boiling ethanol before being photographed.

208

209 Measurement of hydrogen peroxide concentration

210 Fresh leaf and root samples were ground in 0.2 % trichloroacetic acid (TCA) in a mortar and pestle
211 chilled on ice. The homogenate was centrifuged at 15000 g for 20 min at 4 °C. The supernatant was
212 mixed with 10 mM sodium phosphate buffer pH 7 and 1 M KI. The H_2O_2 content was determined
213 by comparing the absorbance of the sample at 390 nm with a standard calibration curve, expressing
214 values in $\mu mol\ H_2O_2\ g^{-1}\ DW$.

215

216 Membrane lipid peroxidation assays

217 Levels of lipid peroxidation were assessed by measuring the amount of malondialdehyde (MDA) in
218 tissue. Fresh leaf and root samples were homogenized in 10 % TCA. The homogenate was
219 centrifuged at 15000 g for 20 min at 4 °C. The supernatant was collected and mixed with 0.5 %
220 thiobarbituric acid in 20 % TCA. Samples were heated at 95 °C for 25 min in a water bath, and then
221 cooled on ice. The samples were centrifuged at 10000 g for 10 min and the absorbance of solutions
222 at 532 and 600 nm was recorded. The MDA level was calculated using the extinction coefficient for
223 MDA [$\epsilon = 155 \mu\text{M cm}^{-1}$] expressed in nmol MDA g⁻¹ DW.

224

225 Protein extraction and antioxidant enzyme assays

226 Leaf and root samples of each genotype were homogenized in a chilled mortar containing 10 %
227 (w/w) poly-vinyl-polypyrrolidone in 50 mM potassium phosphate buffer pH 7, 0.1 mM
228 ethylenediaminetetraacetic acid (EDTA) and 1 mM dithiothreitol. Homogenates were then
229 centrifuged at 10000 g for 15 min at 4 °C. The supernatants were collected and stored at -20 °C for
230 protein and enzyme assays. The soluble protein content of leaves and roots was estimated according
231 to Bradford (1976) with bovine serum albumin as the standard. Superoxide dismutase (SOD,
232 EC.1.15.1.1) activity was assayed based on the inhibition of nitro blue tetrazolium (NBT) reduction
233 (Beyer and Fridovich 1987). The assay was conducted in 50 mM potassium phosphate pH 7.8
234 containing 50 μL of enzyme extract, 2.25 mM NBT, 13 mM methionine, 2 mM riboflavin and 1
235 mM EDTA. The reaction was started by illuminating the sample for 15 min and stopped by
236 switching the light off. The amount of blue formazan formed was determined by measuring the
237 absorbance at 560 nm. One unit of SOD activity (U) was defined as the amount of enzyme that
238 caused 50 % inhibition of NBT reduction.

239

240 Statistical analysis

241 Differences between measurements and between genotypes at different times were evaluated by
242 analysis of variance (one-way ANOVA) using SPSS (Chicago, IL, USA). Differences were
243 considered as statistically significant when $P \leq 0.05$.

244

245

246

247

248

249

250

251 **Results**

252 Analysis of the *pldζ1pldζ2* double mutant transcriptome

253 If Arabidopsis PLD ζ act in lipid signalling in response to salt stress then a *pldζ1pldζ2* double
254 mutant, lacking both PLD ζ and PLD ζ 2, would be expected to differ from WT in the way it responds
255 to salt stress. We first investigated gene expression in WT after 3 h exposure to salt or mannitol.
256 **This was to test whether the ionic stress component of the response could be distinguished from the**
257 **osmotic stress component.** A large common set of genes was induced (560 genes) or repressed (221
258 genes) by the two individual treatments, which was attributed to the similar effects of osmotic stress
259 on the plants (Supplementary Fig. S2). However a smaller subset of genes was specifically induced
260 (170 genes) or repressed (82 genes) only by salt. For example, the known salt-stress responsive
261 gene *DDF1* was specifically upregulated in response to NaCl in WT but was not affected by
262 hyperosmotic stress **caused by** mannitol. Global transcriptome analysis of the *pldζ1pldζ2* double
263 mutant also revealed common sets of genes that were triggered or repressed by both NaCl and
264 mannitol, but also differential expression of subsets of genes in a salt-specific and mannitol-specific
265 manner (data not shown). In further experiments we were thus able to distinguish between ionic and
266 osmotic stress effects, focusing on the former. As some genes were differentially and specifically
267 regulated by NaCl, **the perception and transduction of the ionic component of the salt stress signal**
268 **must have occurred within the 3 h of the stress treatment.**

269 We then compared differences in the NaCl stress transcriptomes of *pldζ1pldζ2* and WT with the
270 WT NaCl stress response. In WT genes were either induced (642 genes) or repressed (241 genes)
271 after 3 h salt stress (Fig. 1). Twenty of the salt-induced genes in WT were found to be expressed at
272 lower levels in *pldζ1pldζ2* after 3 h salt treatment than in WT in the same conditions (intersection in
273 Fig. 1a). This cluster of genes induced in WT, but less strongly induced in *pldζ1pldζ2* was defined
274 as cluster 1 (Fig. 1 and Suppl. Fig. S3). Analysis of gene ontology using BAR Classification
275 SuperViewer (<http://bar.utoronto.ca>) showed an over-representation of genes involved in abiotic
276 and biotic stresses in cluster 1 (Suppl. Fig. S4). It was noted that the salt-stress responsive genes
277 *DDF1* and *RD29A* (Magome et al. 2008) were present in cluster 1. We investigated the expression
278 of these two genes in WT, the *pldζ1* and *pldζ2* single mutants and the *pldζ1pldζ2* double mutant
279 using RT-PCR and quantitative PCR (Fig. 2a, c). The RT-PCR analysis confirmed that *DDF1*
280 expression was induced by salinity in WT plants (Fig. 2a). *DDF1* gene expression was induced in
281 *pldζ1pldζ2* but not as strongly (4.9-fold) as in WT (6-fold) confirming the transcriptome data (Fig.
282 2c). In comparison, *DDF1* expression was induced 9.7-fold in *pldζ2* and 7.8-fold in *pldζ1* (Fig. 2a,
283 c). Similarly, *RD29A* expression was strongly induced by salt treatment in WT (Fig. 2a, c) but less

284 so in *pldζ1pldζ2* as expected from the transcriptomic data. Of the four genotypes tested *pldζ2* again
285 showed the highest level of salt-induction with a 10.4-fold increase in *RD29A* expression.

286 Another cluster of genes, cluster 2, was identified by comparing the differences in the *pldζ1pldζ2*
287 and WT transcriptomes between control and salt stress conditions. In this cluster 23 genes were
288 deregulated in the double mutant in both conditions, 8 upregulated and 15 downregulated (Fig. 1 b,
289 Suppl. Fig. S5). Cluster 2 gene functions mainly related to electron transport and biotic and abiotic
290 stresses (Suppl. Fig. S4). The ROS detoxification genes *SOD1* (At1g08830) and *CCS1* (*Copper*
291 *Chaperone for SOD1*, At1g12520) were overexpressed in *pldζ1pldζ2* compared to WT after salt
292 treatment and in control conditions (Suppl. Fig. S5).

293

294 ROS detoxification was impaired in *pldζ* mutants

295 We verified whether *SOD1* expression was altered in *pldζ* single and double mutants by RT-PCR,
296 and compared it to expression of two other ROS detoxifying genes, ascorbate peroxidase 1 (*APX1*)
297 and catalase (*CAT2*), which are commonly used as oxidative burst markers (Fig. 2b). *APX1*
298 expression was slightly higher in *pldζ2* than in WT and *pldζ1*, but was lower in *pldζ1pldζ2*. *SOD1*
299 gene expression was upregulated by salt treatment in both *pldζ* single mutants but not in WT.
300 Expression of *CAT2* was highest in *pldζ2* compared to the other genotypes even in control
301 conditions (Fig. 2).

302 To test whether the lack of both or either PLD ζ affected the ROS detoxification activity of SOD,
303 SOD activity was monitored for 14 days in the shoots and roots of WT and *pldζ* mutants exposed to
304 moderate (75 mM NaCl) or severe salinity (150 mM NaCl) (Fig. 3). In leaves treated with 75 mM
305 NaCl SOD activity increased transiently in the first 72 h up to 316 % in WT, 169 % in *pldζ1* and
306 108 % in *pldζ1pldζ2*, declining less rapidly then remaining constant until the end of the time course.
307 In roots, the kinetics of SOD activity were markedly different in *pldζ1pldζ2* in comparison to other
308 genotypes as the sharp transient increase occurred within the first three days of mild salt treatment
309 (Fig. 3). Under the same salt conditions, no significant variations were detected in stressed roots of
310 WT and *pldζ1* mutants. By contrast SOD activity in both the shoots and roots of *pldζ2* treated with
311 75 mM NaCl increased linearly to reach a plateau that remained higher than levels in the other
312 genotypes at the end of the 14-day experiment. Upon severe NaCl stress, SOD activity in leaves of
313 *pldζ2* and *pldζ1pldζ2* mutants showed a distinct profile with a transient maximum activity at 3 h,
314 remaining higher than that of WT and *pldζ1* mutants until the end of the salt treatment (Fig. 3). No
315 substantial changes in SOD activity were observed in shoots of WT and *pldζ1* mutants during stress
316 from 150 mM NaCl. SOD activities increased distinctly in the roots of all genotypes except *pldζ1*
317 under severe salt stress. Although there was a transient maximal peak in SOD activity in WT during

318 the first 72 h of exposure to 150 mM NaCl, the highest SOD activities were measured in *pldζ2* and
319 *pldζ1pldζ2* mutants after 14 days.

320
321 *pldζ* mutants differentially accumulate ROS in response to salt stress
322 The presence of H₂O₂ is an indicator of oxidative stress. The differential accumulation of H₂O₂ in
323 leaves and roots over the duration of the stress was evaluated in the four genotypes using DAB,
324 which stains tissues brown in the presence of H₂O₂ (Fig. 4). Leaves and roots grown in the absence
325 of salt did not stain brown with DAB. After 3 h salt treatment, whole leaves and roots of WT and
326 *pldζ1pldζ2* and *pldζ1* were stained brown with DAB (Fig. 4). Although the assay is qualitative, the
327 intensity of the staining consistently increased over the duration of the experiment. Roots and
328 shoots of *pldζ2* mutant were the most faintly stained, indicating these tissues had the lowest
329 amounts of H₂O₂.

330 The differences observed in DAB staining in *pldζ* mutants led us to quantify H₂O₂ content. H₂O₂
331 measurements corresponded with the DAB staining. In the absence of salt stress, H₂O₂ did not
332 accumulate above basal levels in any of the four genotypes. Under mild salt stress, H₂O₂ content
333 increased steadily in shoots and roots in all four genotypes, accumulating faster during the first 24 h
334 (Fig. 5). H₂O₂ accumulated at a similar rate in WT and *pldζ1*, but *pldζ2* and *pldζ1pldζ2* accumulated
335 less than half of the amount measured in WT (Fig. 5). Under severe salt stress, *pldζ2* leaves and
336 roots had the lowest levels of H₂O₂ compared to the other *pldζ* genotypes.

337 Oxidative stress can lead to damage of structural macromolecules including lipids. Lipid
338 peroxidation in leaves and roots was estimated by quantifying malondialdehyde (MDA), which
339 forms when ROS attack lipids. The accumulation of MDA greatly increased in both leaves and
340 roots growing under salt stress (Fig. 6). However the genotypes accumulated MDA differently. The
341 highest amounts of MDA were found in leaves and roots of salt-stressed *pldζ1*, while *pldζ2*
342 accumulated the least MDA in leaves and roots in response to both moderate and severe salt stresses
343 (Fig. 6).

344
345 Ion distribution was altered in *pldζ* mutants responding to salt stress
346 The distribution of ions between leaves and roots of WT and the *pldζ* mutants exposed to 14 days of
347 either moderate or severe salinity is shown in Table 1. Changes in Na⁺ and Cl⁻ content followed the
348 same profile over the 14 days in all genotypes and under both salt treatments, increasing linearly
349 (data not shown) and reaching high levels by the end of the experiment. These final levels were
350 higher in roots than in leaves (Table 1). There were some differences between genotypes. *pldζ1*
351 mutants stressed by 150 mM NaCl had the most Na⁺ in leaves (7.3 mmol g⁻¹ DW) and roots (8.4

352 mmol g⁻¹ DW) (Table 1). This increase was concomitant with the highest Cl⁻ accumulation, which
353 was greater in roots (5.8 mmol g⁻¹DW) than in leaves (3.9 mmol g⁻¹ DW) (Table 1). However, Na⁺
354 content increased more than Cl⁻ content within leaves and roots upon salt stress. When grown in
355 150 mM NaCl, *pldζ2* mutants contained the least Na⁺ (3 mmol g⁻¹ DW in leaves and 3.9 mmol g⁻¹
356 DW in roots) (Table 1) and the least Cl⁻ (2.1 mmol g⁻¹ DW in leaves and 3.1 mmol g⁻¹ DW in roots).
357 Na⁺ accumulation concomitant with K⁺ loss was higher in roots than in leaves in all genotypes
358 under salt stress (Table 1). Salt-stressed *pldζ1* mutants showed the most drastic decrease in K⁺
359 content with 6-fold and 22-fold decreases in leaves and roots, respectively, when treated with 150
360 mM NaCl (Table 1). By contrast, only a slight variation in Na⁺/K⁺ ratio was observed in *pldζ2*
361 mutant. Salinity effects on K⁺ accumulation were less severe in shoots (2-fold decrease) and roots
362 (7-fold decrease) in *pldζ2* than in other genotypes.

363
364 Relative growth rate, relative water content and efficiency of chlorophyll fluorescence in *pldζ*
365 mutants during salt stress

366 Relative growth rate (RGR) of WT, *pldζ1*, *pldζ2* and *pldζ1pldζ2* had decreased after 14 days of
367 exposure to NaCl (Fig. 7). RGRs of *pldζ1pldζ2* and *pldζ2* mutants were less affected by salt than
368 those of *pldζ1* and WT, which were severely inhibited by even mild salt stress. By measuring the
369 change in relative water content (RWC) over time, we found that salt stress induced dehydration in
370 leaves of all genotypes (Fig. 8). *pldζ1* mutants became the most dehydrated with RWC decreases of
371 50 % in 75 mM NaCl and 66 % in 150 mM NaCl compared to the same genotype at the same
372 developmental stage in control conditions (Fig. 8). The *pldζ1pldζ2* leaves only lost 40 % of RWC
373 compared to plants grown in control conditions.

374 The *F_v/F_m* ratio is a convenient measure of photosynthesis efficiency at the photosystem II level
375 (PSII). The *F_v/F_m* ratios for all genotypes were constant in control conditions but decreased under
376 salt stress (Fig. 8). After 14 days of salt treatment, the *F_v/F_m* ratios of *pldζ1* plants had decreased
377 by as much as 64 % in 150 mM NaCl. By contrast *pldζ1pldζ2* plants maintained higher *F_v/F_m* ratio
378 values, which were 1.16 and 1.32-fold higher than those measured in WT, respectively under mild
379 and severe stress, suggesting PSII functions better under stress in the absence of PLDζ (Fig. 8).

380

381

382 Discussion

383 Recent studies have provided valuable insights into the molecular and cellular mechanisms by
384 which plants respond to and tolerate salinity stress (Deinlein et al. 2014; Julkowska and Testerink
385 2015; Slama et al. 2015). Although *Arabidopsis thaliana* is considered to be a glycophyte, it has

386 been widely used as a genetic model to investigate salt signalling mechanisms. Salt treatment
387 significantly inhibits growth of *A. thaliana* with intracellular Na⁺ concentration increasing at the
388 expense of K⁺ (Ghars et al. 2008; Ellouzi et al. 2011). The replacement of K⁺ by Na⁺ affects the
389 cell's Na⁺-sensitive enzymes, including components of the photosynthetic machinery, that
390 determine plant growth and yield. Tolerance to salt stress depends on complex signalling networks
391 enabling plants to respond rapidly and efficiently to this constraint (Zhu 2002). Many signal
392 transduction pathways have been shown to be stimulated in response to high salinity (Bragmann et
393 al. 2009; Julkowska and Testerink 2015). Lipid mediators are key components in the signalling
394 network of plant stress adaptation, including adaptation to salinity (Julkowska and Testerink 2015).
395 As an enzyme responsible for producing PA, we focused on the roles of the two *A. thaliana* PLD ζ
396 in salt tolerance.

397 To identify changes in gene expression affected by PLD ζ activity, the transcriptome of a double
398 *pld ζ 1pld ζ 2* mutant was compared to the WT transcriptome. In gene cluster 1, the stress-responsive
399 genes *DDF1* and *RD29A* (Thiery et al. 2004; Magome et al. 2004; Magome et al. 2008) were
400 strongly induced by severe salt stress (200 mM NaCl) within 3 h in the WT but were less strongly
401 induced in *pld ζ 1pld ζ 2* under the same conditions. Transgenic Arabidopsis lines overexpressing
402 *DDF1* were shown to have increased tolerance to high salt stress (170 mM NaCl) by repressing
403 plant growth through the induction of genes coding gibberellin (GA) deactivating enzymes
404 (Magome et al. 2004; Magome et al. 2008). We found that the *pld ζ 2* single mutant showed higher
405 expression of *DDF1* and *RD29A* under salt stress compared with the WT and other *pld ζ mutants*
406 (Fig. 2). The expression of auxin-responsive genes such as *IAA5*, *IAA19*, and *GH3-3* was previously
407 found to be reduced in *pld ζ 2* mutants of Arabidopsis in response to external IAA (Li and Xue
408 2007). Therefore, *PLD ζ 2* may be specifically involved in the regulation of gene expression in stress
409 responses and hormonal signalling for growth. Controlled growth reduction may be an effective
410 strategy to save energy and minimize ROS accumulation while facing the deleterious impact of salt
411 stress (Rangani et al. 2016).

412 Transcriptomic analysis revealed a second gene cluster that included ROS detoxification genes such
413 as *SOD1* and *CCS1*, which were overexpressed in *pld ζ 1pld ζ 2* mutants (Suppl. Fig. S5). It is known
414 that ROS act as second messengers in intracellular signalling cascades to trigger plant tolerance to
415 various abiotic and biotic stresses (Ben Rejeb et al. 2014; Mittler, 2016). When *Cakile maritima*, a
416 salt-tolerant plant, is treated with salt, there is a transient and rapid increase in SOD activity and
417 H₂O₂ content within 4 h. The SOD activity measured in *C. maritima* was eight-fold higher than was
418 found in Arabidopsis (Ellouzi et al. 2013). In the present study, the highest *SOD1*, *CAT2* and *APX*
419 gene expression was observed in *pld ζ 2* mutant under salt stress (Fig. 2). High expression of *SOD*

420 under salt stress corresponded with high SOD activity and low H₂O₂ levels in leaves and roots of
421 *pldζ2* mutants, indicating that *PLDζ2* participates in the regulation of ROS generation upon salt
422 stress in Arabidopsis. Several reports show that different PLD family enzymes function in concert
423 with ROS to mediate tolerance responses to various abiotic stresses (Hong et al. 2010, Singh et al.
424 2012). For example, mutation in *PLDα1* leads to PA deficiency, reduced plasma membrane
425 NADPH oxidase activity, and less ROS in stomata guard cells in response to abscisic acid (Zhang et
426 al. 2009). As a consequence, *plda1* mutants can not fully close stomata leading to increased water
427 loss (Zhang et al. 2009). We found that *pldζ1* mutants accumulated H₂O₂ faster in leaves and roots
428 than *pldζ2* mutants do in response to salt stress. Moreover, *pldζ1* plants displayed the lowest SOD
429 activity which, associated with the highest levels of H₂O₂, implies that *pldζ1* is more sensitive to
430 salt stress than WT and the other *pldζ* mutants. Interestingly other *pld* mutants such as *plda1*, *plda3*
431 and *pldδ* have also been shown to be hypersensitive to salt treatment (Hong et al. 2008; Bargmann
432 et al. 2009).

433 Salinity had several striking effects on physiological indexes with distinctions becoming clear
434 between WT and *pldζ1* versus *pldζ2* and *pldζ1pldζ2*. The *pldζ1* mutant showed the highest Na⁺
435 content in leaves and roots compared with the other genotypes upon salt stress, leading to a drastic
436 decrease in K⁺ content in roots and shoots (Table 1). The loss of K⁺ could be explained by
437 downregulation of the expression of genes involved in K⁺ transport like the HAK5 transporter gene.
438 HAK5 is expressed under the control of *DDF2* gene, a transcription factor which is homologous to
439 *DDF1* (Hong et al. 2013). The massive K efflux could also be mediated by the opening of outward-
440 rectifying depolarisation-activated (GORK) channels, an outward K⁺ channel (for review see
441 Anshütz et al., 2014). Phospholipase Dζ could be involved in the regulation of ion channels by
442 modulating their lipid environment. For example in yeast cells, many ion channels have been shown
443 to be located in plasma membrane microdomains called lipid rafts with channel activities dependent
444 on the lipid raft composition (for review see Mollinedo 2012).

445 RGR measurements showed that WT and *pldζ1* have only a limited capacity to withstand the
446 presence of salt. Sensitivity of WT and *pldζ1* to salt stress was associated with a decline in the leaf
447 water content, increase in Na⁺ and Cl⁻ preferentially accumulated in roots, early induction of
448 oxidative stress (excess H₂O₂ and MDA), and inhibition of photosynthesis from the toxicity of the
449 salt ions (low *Fv/Fm* ratio). For most markers *pldζ1* was more sensitive than WT, indicating that
450 *PLDζ1* is required in the regulation of growth in normal conditions. Interestingly and as described
451 by Ohashi et al. (2003) and Chen et al. (2013), this PLD isoform is greatly involved in the root
452 development and growth and the lack of this PLD may lead to enhanced salt stress sensitivity
453 (Wang, 2005).

454 Conversely, *pldζ2* mutants tolerated salt stress better than WT. This response may be related to its
455 ability to minimize redistribution of Na⁺ and Cl⁻ to the roots, resulting in the lowest Na⁺/K⁺ ratio
456 and consequently better retention of K⁺ in these organs. *pldζ2* grew less than *pldζ1 pldζ2* double
457 mutant, which produced more biomass in both roots and leaves. We can consider growth reduction
458 in *pldζ2* to be a strategy to maintain performance under salinity by first controlling ion homeostasis.
459 The activation of the Salt Overly Sensitive (SOS) signalling pathway is a key mechanism for Na⁺
460 exclusion in roots (Zhu, 2000). PLDs were shown to interact strongly with SOS and plasma
461 membrane transporters under salt stress (Yu et al. 2010). Notably, the activation of PLD in salt
462 stressed tobacco elevated the PA level which is a direct stimulator of SOS1, a central regulator of
463 ion homeostasis (Gardiner et al. 2001). Other studies suggested that there is a functional connection
464 between the activation of *PLDζ2* and SOS3, which is more prominent in the roots and has a crucial
465 role in Na⁺/K⁺ dynamics (Muzi et al. 2016). Much less is known about the cross-talk between PLD
466 and SOS signalling pathways under salt stress. However, the above findings will guide future
467 research to elucidate the mechanism of the salt tolerance in *pldζ2*.

468 It was clear from our results that the *pldζ2* mutant was the most tolerant of salt stress as evidenced
469 by the lowest Na⁺/K⁺ ratio. It has long been known that salinity stress triggers a dramatic increase in
470 ROS accumulation in plant tissues. Interestingly, our results showed that salt-stressed *pldζ2* plants
471 also displayed the lowest levels of H₂O₂ and MDA detected in both leaves and roots. In addition,
472 mild and severe salt stresses triggered higher SOD activity in *pldζ2* than in the other genotypes
473 which could explain why H₂O₂ and MDA levels were lower in both leaves and roots. The oxidative
474 burst indicators SOD, H₂O₂ and MDA indicate the presence of an efficient antioxidant defence in
475 *pldζ2* mutant. The lack of *PLDζ2* may therefore allow a better performance of the plants facing
476 moderate and severe salinity due to a primed antioxidant defence system.

477 In conclusion, our study suggests that in Arabidopsis *PLDζ* genes encode isoforms that have distinct
478 roles in plant growth particularly under salt stress. The *pldζ2* KO mutant was more salt-tolerant than
479 the WT as growth, water status, ion homeostasis and antioxidant defence systems were all better
480 adjusted to withstanding salinity. Possibly *pldζ* are differently regulated in halophytes enabling
481 them to tolerate salinity. Looking for natural variants or mutations in *PLDζ2* gene might be a way to
482 improve salt tolerance in glycophytic crops.

483

484

485 **Author contribution statement**

486 AS and CA conceived and designed the research. Data collection, analysis and interpretation were
487 performed by AO, HE, SP and BF. SP conducted the transcriptomic analysis. DDV was involved in

488 the molecular analysis of *pldζ* mutant and in obtaining the double *pldζ1pldζ2* mutant. PC
489 contributed to study conception and design. AO, HE, SP wrote the manuscript. AS and CA
490 supervised manuscript preparation and correction. All authors read and approved the manuscript.

491

492 **Acknowledgments**

493 This work was supported by the UPMC (France) and the CBBC (Tunisia) and also supported by the
494 Tunisian-French UTIQUE network (n°13G0929). We thank Frederique Bitton and Jean-Pierre
495 Renou from URG-INRA, Evry (France) for help producing the microarray data. Bualuang Faiyue
496 thanks the Junior Research Fellowship Program supported by the French Embassy in Thailand.

497

498

499

500 **References**

501 Alonso JM, Stepanova AN (2003) T-DNA mutagenesis in *Arabidopsis*. *Methods Mol Biol Clifton*
502 *NJ* 236:177–188. doi: 10.1385/1-59259-413-1:177

503 Anshütz U, Becker D, Shabala S (2014) Going beyond nutrition: regulation of potassium
504 homeostasis as a common denominator of plant adaptive responses to environment. *J Plant*
505 *Physiol* 171:670-687.

506 Bargmann BO, Laxalt AM, Riet BT, Testerink C, Merquiol E, Mosblech A, Leon-Reyes A, Pieterse
507 CM, Haring MA, Heilmann I, Bartels D, Munnik T (2009) Reassessing the role of phospholipase D
508 in the *Arabidopsis* wounding response. *Plant Cell Environ* 32 743-757. doi: 10.1111/j.1365-
509 3040.2009.01962.x

510 Bargmann BO, Munnik T (2006) The role of phospholipase D in plant stress responses. *Curr Opin*
511 *Plant Biol* 9:515–22. doi: 10.1016/j.pbi.2006.07.011

512 Ben Rejeb K, Abdelly C, Savouré A (2014) How reactive oxygen species and proline face stress
513 together. *Plant Physiol Biochem* 80:278–284. doi: 10.1016/j.plaphy.2014.04.007

514 Ben Rejeb K, Lefebvre-De Vos D, Le Disquet I, Lepince A-S, Bordenave M, Maldiney R, Jdey A,
515 Abdelly C, Savouré A (2015) Hydrogen peroxide produced by NADPH oxidases increases proline
516 accumulation during salt or mannitol stress in *Arabidopsis thaliana*. *New Phytol* 208:1138–1148.
517 doi: 10.1111/nph.13550

518 Beyer WF, Fridovich I (1987) Assaying for superoxide dismutase activity: Some large
519 consequences of minor changes in conditions. *Anal Biochem* 161:559–566. doi: 10.1016/0003-
520 2697(87)90489-1

521 Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of
522 protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–54.

523 Bargmann BO, Laxalt AM, Riet BT, van Schooten B, Merquiol E, Testerink C, Haring MA, Bartels
524 D, Munnik T (2009) Multiple PLDs required for high salinity and water deficit tolerance in
525 plants. *Plant Cell Physiol* 50:78-89. doi: 10.1093/pcp/pcn173

526 Chen MX, Lung SC, Du ZY, Chye ML (2013) Engineering plants to tolerate abiotic stresses.
527 *Biocatal Agric Biotechnol* 3:81-87. doi: 10.1016/j.bcab.2013.09.010

528 Crowe ML, Serizet C, Thareau V, Aubourg S, Rouzé P, Hilson P, Beynon J, Weisbeek P, van
529 Hummelen P, Reymond P, Paz-Ares J, Niefeld W, Trick M (2003) CATMA: a complete
530 Arabidopsis GST database. *Nucleic Acids Res* 31:156–158.

531 Deal SB (1954) Flame photometric determination of sodium and potassium. *Anal Chem* 26:598–
532 599. doi: 10.1021/ac60087a063

533 Deinlein U, Stephan AB, Horie T, Luo W, Xu G, Schroeder JI (2014) Plant salt-tolerance
534 mechanisms. *Trends Plant Sci* 19:371–379. doi: 10.1016/j.tplants.2014.02.001

535 Ellouzi H, Ben Hamed K, Cela J, Munné-Bosch S, Abdelly C (2011) Early effects of salt stress on
536 the physiological and oxidative status of *Cakile maritima* (halophyte) and *Arabidopsis thaliana*
537 (glycophyte). *Physiol Plant* 142:128–143. doi: 10.1111/j.1399-3054.2011.01450.x

538 Ellouzi H, Ben Hamed K, Asensi-Fabado MA, Müller M, Abdelly C, Munné-Bosch S (2013)
539 Drought and cadmium may be as effective as salinity in conferring subsequent salt stress tolerance
540 in *Cakile maritima*. *Planta* 237:1311–1323. doi: 10.1007/s00425-013-1847-7

541 Galvan-Ampudia CS, Julkowska MM, Darwish E, Gandullo J, Korver RA, Brunoud G, Haring MA,
542 Munnik T, Vernoux T, Testerink C (2013) Halotropism is a response of plant roots to avoid a saline
543 environment. *Curr Biol* 23:2044–2050. doi: 10.1016/j.cub.2013.08.042

544 Gardiner JC, Harper JD, Weerakoon ND, Collings DA, Ritchie S, Gilroy S, Cyr RJ, Marc J (2001)
545 A 90-kD phospholipase D from tobacco binds to microtubules and the plasma membrane. *Plant Cell*
546 13: 2143–2158.

547 Ghars MA, Parre E, Debez A, Bordenave M, Richard L, Leport L, Bouchereau A, Savoure A,
548 Abdelly C (2008) Comparative salt tolerance analysis between *Arabidopsis thaliana* and
549 *Thellungiella halophila*, with special emphasis on K⁺/Na⁺ selectivity and proline accumulation. *J*
550 *Plant Physiol* 165:588–599. doi: 10.1016/j.jplph.2007.05.014

551 Gill SS, Tuteja N (2010) Reactive oxygen species and antioxidant machinery in abiotic stress
552 tolerance in crop plants. *Plant Physiol Biochem* 48:909–930. doi: 10.1016/j.plaphy.2010.08.016

553 Hewitt E (1966) The composition of the nutrient solution. *Sand Water Cult Methods Used Study*
554 *Plant Nutr* 187–246.

555 Hong J-P, Takeshi Y, Kondou Y, Schachtman DP, Matsui M, Shin R (2013) Identification and
556 characterization of transcription factors regulating Arabidopsis *HAK5*. *Plant Cell Physiol* 54:1478–
557 1490. doi: 10.1093/pcp/pct094

558 Hong Y, Pan X, Welti R, Wang X (2008) Phospholipase D α 3 is involved in the hyperosmotic
559 response in Arabidopsis. *Plant Cell* 20:803–816. doi: 10.1105/tpc.107.056390

560 Hong Y, Zhang W, Wang X (2010) Phospholipase D and phosphatidic acid signalling in plant
561 response to drought and salinity. *Plant Cell Env* 33:627–635.

562 Hou Q, Ufer G, Bartels D (2016) Lipid signalling in plant responses to abiotic stress. *Plant Cell*
563 *Environ* 39:1029–1048. doi: 10.1111/pce.12666

564 Hunt R (1990) *Basic Growth analysis: Plant growth analysis for beginners*. Unwin Hyman. London.

565 Julkowska MM, Testerink C (2015) Tuning plant signaling and growth to survive salt. *Trends Plant*
566 *Sci* 20:586–594. doi: 10.1016/j.tplants.2015.06.008

567 Li G, Xue H-W (2007) Arabidopsis PLDzeta2 regulates vesicle trafficking and is required for auxin
568 response. *Plant Cell* 19:281–295. doi: 10.1105/tpc.106.041426

569 Li M, Hong Y, Wang X (2009) Phospholipase D- and phosphatidic acid-mediated signaling in
570 plants. *Biochim Biophys Acta - Mol Cell Biol Lipids* 1791:927–935. doi:
571 10.1016/j.bbalip.2009.02.017

572 Li M, Qin C, Welti R, Wang X (2006) Double knockouts of phospholipases D ζ 1 and D ζ 2 in
573 Arabidopsis affect root elongation during phosphate-limited growth but do not affect root hair
574 patterning. *Plant Physiol* 140:761–770. doi: 10.1104/pp.105.070995

575 Lurin C, Andrés C, Aubourg S, Bellaoui M, Bitton F, Bruyère C, Caboche M, Debast C, Gualberto
576 J, Hoffmann B, Lecharny A, Ret ML, Martin-Magniette M-L, Mireau H et al (2004) Genome-wide
577 analysis of Arabidopsis pentatricopeptide repeat proteins reveals their essential role in organelle
578 biogenesis. *Plant Cell* 16:2089–2103. doi: 10.1105/tpc.104.022236

579 Magome H, Yamaguchi S, Hanada A, Kamiya Y, Oda K (2004) *dwarf and delayed-flowering 1*, a
580 novel Arabidopsis mutant deficient in gibberellin biosynthesis because of overexpression of a
581 putative AP2 transcription factor. *Plant J* 37:720-729.

582 Magome H, Yamaguchi S, Hanada A, Kamiya Y, Oda K (2008) The DDF1 transcriptional activator
583 upregulates expression of a gibberellin-deactivating gene, *GA2ox7*, under high-salinity stress in
584 Arabidopsis. *Plant J* 56:613–626. doi: 10.1111/j.1365-313X.2008.03627.x

585 McLoughlin, F, Testerink C (2013) Phosphatidic acid, a versatile water-stress signal in roots. *Front*
586 *Plant Sci* 4: 525. doi: 10.3389/fpls.2013.00525

587 Meijer HJ, Munnik T (2003) Phospholipid-based signaling in plants. *Annu Rev Plant Biol* 54:265–
588 306.

589 Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R (2010) Reactive oxygen species homeostasis and
590 signalling during drought and salinity stresses. *Plant Cell Environ* 33:453–467. doi: 10.1111/j.1365-
591 3040.2009.02041.x

592 Mittler R (2016) ROS are good. *Trends Plant Sci* 22:11-19 doi: 10.1016/j.tplants.2016.08.002

593 Mollinedo F (2012) Lipid raft involvement in yeast cell growth and death. *Front Oncol* 2, 140.

594 Munns R, Tester M (2008) Mechanisms of salinity tolerance. *Annu Rev Plant Biol* 59:651–681. doi:
595 10.1146/Annurev.Arplant.59.032607.092911

596 Murashige T, Skoog F (1962) A revised medium for rapid growth and bio assays with tobacco
597 tissue culture. *Physiol Plant* 15:473–497.

598 Muzi C, Camoni L, Visconti S, Aducci P (2016) Cold stress affects HC- ATPase and phospholipase
599 D activity in Arabidopsis. *Plant Physiol. Biochem.*108:328–336.

600 Ohashi Y, Oka A, Rodrigues-Pousada R, Possenti M, Ruberti I, Morelli G, Aoyama T (2003)
601 Modulation of phospholipid signaling by GLABRA2 in root-hair pattern formation. *Science*.
602 300:1427-30.

603 Parre E, Ghars MA, Leprince AS, Thiery L, Lefebvre D, Bordenave M, Richard L, Mazars C,
604 Abdelly C, Saviouré A (2007) Calcium signaling via phospholipase C is essential for proline
605 accumulation upon ionic but not nonionic hyperosmotic stresses in Arabidopsis. *Plant Physiol*
606 144:503–12. doi: 10.1104/pp.106.095281

607 Pfaffl MW (2001) A new mathematical model for relative quantification in real-time RT–PCR.
608 *Nucleic Acids Res* 29:e45. doi: 10.1093/nar/29.9.e45

609 Planchais S, Cabassa C, Toka I, Justin A-M, Renou J-P, Saviouré A, Carol P (2014) BASIC AMINO
610 ACID CARRIER 2 gene expression modulates arginine and urea content and stress recovery in
611 Arabidopsis leaves. *Front Plant Sci* 5:330. doi: 10.3389/fpls.2014.00330

612 Qin C, Wang X (2002) The Arabidopsis phospholipase D family. Characterization of a calcium-
613 independent and phosphatidylcholine-selective PLDzeta1 with distinct regulatory domains. *Plant*
614 *Physiol* 128:1057–68.

615 Rangani J, Parida AK, Panda A, Kumari A (2016) Coordinated changes in antioxidative enzymes
616 protect the photosynthetic machinery from salinity induced oxidative damage and confer salt
617 tolerance in an extreme halophyte *Salvadora persica* L. *Front Plant Sci* 7:50. doi:
618 10.3389/fpls.2016.00050

619 Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning*. Cold Spring Harbor Laboratory
620 Press New York

621 Singh A, Pandey A, Baran wal V, Kapoor S, Pandey GK (2012) Comprehensive expression analysis
622 of rice phospholipase D gene family during abiotic stress and development. *Plant Signal Behav* 7:
623 847-855.

624 Slama I, Abdelly C, Bouchereau A, Flowers T, Savouré A (2015) Diversity, distribution and roles
625 of osmoprotective compounds accumulated in halophytes under abiotic stress. *Ann Bot* 115:433–
626 447. doi: 10.1093/aob/mcu239

627 Thiery L, Leprince AS, Lefebvre D, Ghars MA, Debarbieux E, Savouré A (2004) Phospholipase D
628 is a negative regulator of proline biosynthesis in *Arabidopsis thaliana*. *J Biol Chem* 279:14812–8.

629 Vergnolle C, Vaultier M-N, Taconnat L, Renou J-P, Kader J-C, Zachowski A, Ruelland E (2005)
630 The cold-induced early activation of phospholipase C and D pathways determines the response of
631 two distinct clusters of genes in *Arabidopsis* cell suspensions. *Plant Physiol* 139:1217–1233. doi:
632 10.1104/pp.105.068171

633 Wang X (2005) Regulatory functions of phospholipase D and phosphatidic acid in plant growth,
634 development, and stress responses. *Plant Physiol* 139:566–573.

635 Yu L, Nie J, Cao C, Jin Y, Yan M, Wang F, Liu J, Xiao Y, Liang Y, Zhang W (2010) Phosphatidic
636 acid mediates salt stress response by regulation of MPK6 in *Arabidopsis thaliana*. *New Phytol*
637 188:762–773.

638 Zhang Y, Zhu H, Zhang Q, Li M, Yan M, Wang R, Wang L, Welti R, Zhang W, Wang X (2009)
639 Phospholipase D α 1 and phosphatidic acid regulate NADPH oxidase activity and production of
640 reactive oxygen species in ABA-mediated stomatal closure in *Arabidopsis*. *Plant Cell* 21:2357–
641 2377. doi: 10.1105/tpc.108.062992

642 Zhu JK (2002). Salt and drought stress signal transduction in plants. *Annu. Rev. Plant. Biol.* 53:
643 247–273.

644 Zhu JK (2000) Genetic analysis of plant salt tolerance using *Arabidopsis*. *Plant Physiol* 124:941-8.
645

646


647 Table 1. Ion content (mmol g⁻¹ DW) of leaves and roots of WT and *pldζ1*, *pldζ2* and *pldζ1pldζ2*
 648 mutants after 14 days of mild or severe salt stress. Data are means of three replicates ± S.E. Means
 649 indicated by the same superscript letters are not significantly different at P ≤ 0.05 in one-way
 650 ANOVA.

651
 652


Leaves		Genotype	WT	<i>pldζ1</i>	<i>pldζ2</i>	<i>pldζ1pldζ2</i>
NaCl (mM)	Ion content					
0	Na ⁺		0.14 ± 0.01 ^a	0.14 ± 0.00 ^{ab}	0.14 ± 0.01 ^{ab}	0.15 ± 0.00 ^b
75	Na ⁺		2.36 ± 1.09 ^b	3.30 ± 0.91 ^d	1.33 ± 0.19 ^a	2.89 ± 0.87 ^c
150	Na ⁺		4.50 ± 1.06 ^c	7.31 ± 1.06 ^d	3.00 ± 1.10 ^a	3.92 ± 1.08 ^b
0	K ⁺		2.79 ± 0.24 ^b	1.86 ± 0.02 ^a	2.32 ± 0.21 ^{ab}	2.07 ± 0.17 ^a
75	K ⁺		1.33 ± 0.15 ^b	0.76 ± 0.03 ^a	1.30 ± 0.27 ^b	0.65 ± 0.01 ^a
150	K ⁺		0.77 ± 0.05 ^b	0.31 ± 0.00 ^a	1.12 ± 0.22 ^b	0.28 ± 0.02 ^a
0	Cl ⁻		0.03 ± 0.00 ^a	0.03 ± 0.00 ^a	0.09 ± 0.01 ^b	0.02 ± 0.00 ^a
75	Cl ⁻		2.37 ± 0.55 ^c	2.14 ± 0.32 ^a	1.66 ± 0.32 ^b	1.70 ± 0.09 ^a
150	Cl ⁻		3.08 ± 0.85 ^b	3.88 ± 0.66 ^d	2.09 ± 0.53 ^a	3.23 ± 1.021 ^c
0	Na ⁺ /K ⁺		0.04 ± 0.00 ^a	0.07 ± 0.00 ^c	0.06 ± 0.00 ^b	0.07 ± 0.00 ^c
75	Na ⁺ /K ⁺		1.80 ± 0.06 ^b	4.38 ± 0.19 ^c	1.05 ± 0.11 ^a	4.56 ± 0.30 ^c
150	Na ⁺ /K ⁺		5.88 ± 0.44 ^b	23.88 ± 0.22 ^d	2.88 ± 0.51 ^a	14.30 ± 1.40 ^c
Roots						
0	Na ⁺		0.12 ± 0.00 ^a	0.11 ± 0.01 ^a	0.22 ± 0.01 ^b	0.13 ± 0.00 ^a
75	Na ⁺		3.73 ± 1.11 ^b	3.72 ± 1.19 ^b	2.91 ± 0.66 ^a	2.89 ± 0.97 ^a
150	Na ⁺		6.43 ± 1.31 ^b	8.42 ± 1.16 ^c	3.92 ± 0.91 ^a	5.88 ± 1.08 ^b
0	K ⁺		1.14 ± 0.21 ^b	0.65 ± 0.24 ^a	0.98 ± 0.07 ^{ab}	0.94 ± 0.02 ^{ab}
75	K ⁺		0.40 ± 0.01 ^c	0.10 ± 0.00 ^a	0.52 ± 0.02 ^d	0.28 ± 0.00 ^b
150	K ⁺		0.09 ± 0.01 ^b	0.04 ± 0.00 ^a	0.15 ± 0.00 ^c	0.08 ± 0.00 ^b
0	Cl ⁻		0.04 ± 0.00 ^a	0.03 ± 0.00 ^a	0.06 ± 0.01 ^b	0.03 ± 0.00 ^a
75	Cl ⁻		4.10 ± 0.91 ^c	4.08 ± 0.90 ^c	1.91 ± 0.36 ^a	3.50 ± 0.33 ^b
150	Cl ⁻		5.08 ± 1.05 ^c	5.72 ± 0.28 ^d	3.00 ± 0.83 ^a	4.00 ± 0.83 ^b
0	Na ⁺ /K ⁺		0.11 ± 0.00 ^a	0.21 ± 0.06 ^{ab}	0.23 ± 0.01 ^b	0.13 ± 0.01 ^{ab}
75	Na ⁺ /K ⁺		9.61 ± 0.52 ^b	35.14 ± 2.75 ^c	5.54 ± 0.13 ^a	10.22 ± 1.05 ^b
150	Na ⁺ /K ⁺		70.67 ± 2.81 ^b	246.95 ± 40.1 ^d	55.41 ± 20.0 ^a	72.91 ± 1.30 ^c

653
 654
 655

a






b




657
 658 **Figure 1.** Numbers of differentially expressed genes in WT and *pldζ1pldζ2* transcriptome
 659 comparisons. Venn diagrams show how gene clusters were defined. **(a)** Cluster 1 (intersection in
 660 bold) includes genes which are induced by salt stress in WT and are expressed at lower levels in the
 661 double mutant under salt stress than in WT. **(b)** Cluster 2 (intersection in bold) includes genes
 662 which are up or down regulated in *pldζ1pldζ2* compared to WT in both control and salt stress
 663 conditions.

664
 665




667 **Figure 2.** Gene expression in wild-type (WT), *pldζ1*, *pldζ2* and *pldζ1ζ2* plants treated (+) or not
668 treated (-) with 200 mM NaCl for 3 h. *DDF1* and *RD29A* (a), and *APX1*, *SOD1* and *CAT2* (b)
669 expression was estimated by RT-PCR with *APT1* as a loading control. (c) Quantitative RT-PCR of
670 *DDF1* and *RD29A* expression in WT, *pldζ1*, *pldζ2* and *pldζ1ζ2* seedlings. *DDF1* transcript
671 abundance was expressed as a ratio of the value for WT in control conditions. *APT1* gene
672 expression was used as a standard. Mean and standard errors are based on three technical repeats
673



675


676 **Figure 3.** Changes in superoxide dismutase (SOD) activity in leaves and roots of wild type and
 677 *pldζ1*, *pldζ2* and *pldζ1pldζ2* mutants of *A. thaliana* plants grown for 14 days under moderate (75
 678 mM NaCl) or severe salinity (150 mM NaCl). Data are means \pm SE of 3 replicates. Results of
 679 statistical analysis are given in each panel (ANOVA, $P \leq 0.05$). NS, not significant.

680




682 **Figure 4** Histochemical detection of hydrogen peroxide in leaves (a) and roots (b) of wild type and
683 *pldζ1*, *pldζ2* and *pldζ1pldζ2* mutants of *A. thaliana* plants grown for 14 days under moderate (75
684 mM NaCl) or severe salinity (150 mM NaCl). Brown residue from diaminobenzidine staining
685 indicates sites of H₂O₂ accumulation.


686




687
 688 **Figure 5.** Changes in hydrogen peroxide levels in leaves and roots of wild type and *pldζ1*, *pldζ2*
 689 and *pldζ1pldζ2* mutants of *A. thaliana* plants grown for 14 days with moderate (75 mM NaCl) or
 690 severe salinity (150 mM NaCl). Data are means ± SE of 3 replicates. Results of statistical analysis
 691 are given in each panel (ANOVA, $P \leq 0.05$). NS, not significant.
 692





693
 694 **Figure 6.** Changes in malondialdehyde (MDA) levels in leaves and roots of wild type and *pldζ1*,
 695 *pldζ2* and *pldζ1pldζ2* mutants of *A. thaliana* plants grown for 14 days under moderate (75 mM
 696 NaCl) or severe salinity (150 mM NaCl). Data are means \pm SE of 3 replicates. Results of statistical
 697 analysis are given in each panel (ANOVA, $P \leq 0.05$). NS, not significant.
 698




699
 700 **Figure 7.** Relative growth rate (RGR) in *A. thaliana* single and double *pldζ* mutants grown for 14
 701 days under control conditions or moderate (75 mM NaCl) or severe salinity (150 mM NaCl). Means
 702 indicated by different letters are significantly different ($P \leq 0.05$) as determined by one-way
 703 ANOVA.
 704



705
 706 **Figure 8.** Changes in the relative water content (RWC) and in F_v/F_m ratios of leaves and roots of
 707 wild type, and *pldζ1*, *pldζ2* and *pldζ1pldζ2* mutants of *A. thaliana* plants grown for 14 days under
 708 moderate (75 mM NaCl) or severe salinity (150 mM NaCl). Data are means \pm SE of 3 replicates.
 709 Results of statistical analysis (ANOVA, $P \leq 0.05$) are given in each panel. NS, not significant
 710



Supplementary Figure S1 Map and genotyping of *pldζ1*, *pldζ2* and *pldζ1pldζ2* mutants. (a) Map of T-DNA insertion site in *pldζ1* and *pldζ2* genes. Primers used for PCR are represented by arrows. (b) 1 % agarose gels of PCR products for *PLDζ1* WT allele (A+B), *pldζ1* T-DNA flanking sequence (A+C), or both (A+B+C) in hemizygous plants (top gel). PCR products for WT *PLDζ2* WT allele (D+E), *pldζ2* T-DNA flanking sequence (D+C), or both (D+E+C) in hemizygous plants (middle gel). WT allele amplification (A+B), *pldζ1* T-DNA flanking sequence (A+C) and *pldζ2* T-DNA flanking sequence (D+C) in *pldζ1pldζ2* (bottom gel).



Supplementary Figure S2 Venn diagram of WT transcriptomes under salt and mannitol stresses in comparison to control growth conditions. Numbers of differentially expressed genes for each comparison are shown.

AGI	Name	WT+NaCl / WT control		pld ζ 1pld ζ 2+NaCl / WT+NaCl	
		Log ₂ Ratio	Pvalue	Log ₂ Ratio	Pvalue
AT1G12610	DDF1 transcription factor	1.96	0.00E+0	-1.89	0.00E+0
AT1G73480	alpha/beta hydrolase family	3.40	0.00E+0	-1.04	0.00E+0
AT5G52310	COR78 (COLD REGULATED 78) (RD29A)	4.04	0.00E+0	-1.03	5.60E-12
AT2G16500	ADC1_ arginine decarboxylase 1	0.89	1.17E-4	-0.98	1.12E-10
AT1G58360	AAP1; amino acid permease 1	1.95	0.00E+0	-0.86	1.77E-7
AT2G38240	oxidoreductase. 2OG-Fe(II) oxygenase	1.23	2.24E-11	-0.81	2.50E-6
AT5G13750	ZIFL1 zinc induced facilitator-like 1	1.81	0.00E+0	-0.78	9.68E-6
AT2G46370	JAR1 (JASMONATE RESISTANT 1)	0.83	1.15E-3	-0.76	2.66E-5
AT4G26080	ABI1 (ABA INSENSITIVE 1) PP2C family	1.48	0.00E+0	-0.75	4.51E-5
AT1G51090	metal ion binding	2.99	0.00E+0	-0.74	1.05E-4
AT1G72770	HAB1; PP2C homologue to ABI1	2.53	0.00E+0	-0.73	1.22E-4
AT3G07700	Protein kinase superfamily protein	0.90	8.80E-5	-0.73	1.57E-4
AT1G01650	SPPL4(SIGNAL PEPTIDE PEPTIDASE-LIKE 4)	0.73	4.31E-2	-0.71	3.41E-4
AT3G48990	AMP-dependent synthetase and ligase	0.75	2.14E-2	-0.71	3.61E-4
AT1G29400	AML5 (ARABIDOPSIS MEI2-LIKE PROTEIN 5)	0.79	5.23E-3	-0.70	5.50E-4
AT5G17760	P-loop nucleoside triphosphate hydrolase	1.43	0.00E+0	-0.70	6.32E-4
AT1G76960	unknown protein	0.85	7.24E-4	-0.67	2.25E-3
AT2G35940	EDA29 (BEL1-like homeodomain 1)	0.80	3.45E-3	-0.65	5.59E-3
AT4G15530	PPDK pyruvate orthophosphate dikinase	0.90	1.04E-4	-0.64	8.57E-3
AT5G55970	RING/U-box superfamily protein	1.25	5.59E-12	-0.61	2.51E-2

Supplementary Figure S3 List of genes in Cluster 1. Genes are involved in abiotic or biotic stimulus and response to stress. Expression data is given as log₂ ratios.

Cluster 1 (Figure 1)**Biological process**

	Normed to Freq. In Arabidopsis set (\pm bootstrap StdDev, p-value)		
	Frequency in Cluster 1	Std dev	P value
response to abiotic or biotic stimulus (Input set freq.: 0.65; 0.13)	4.96	0.795	0.0000009492
other biological processes (Input set freq.: 0.45; 0.12)	3.65	0.874	0.0002548
response to stress (Input set freq.: 0.5; 0.14)	3.45	0.681	0.0001537
developmental processes (Input set freq.: 0.4; 0.13)	3.02	0.91	0.002149
transport (Input set freq.: 0.35; 0.12)	2.82	0.814	0.006256
signal transduction (Input set freq.: 0.15; 0.07)	2.12	1.185	0.115

Cluster 2 (Figure 4)**Biological process**

	Normed to Freq. In Arabidopsis set (\pm bootstrap StdDev, p-value)		
	Frequency in Cluster 2	Std dev	P value
electron transport or energy pathways (Input set freq.: 0.28; 0.02)	13.07	4.609	4.157e-06
response to stress (Input set freq.: 0.76; 0.13)	5.47	0.807	1.850e-10
response to abiotic or biotic stimulus (Input set freq.: 0.61; 0.13)	4.62	0.885	2.780e-07
other biological processes (Input set freq.: 0.57; 0.13)	4.28	0.815	2.565e-06
transport (Input set freq.: 0.38; 0.1)	3.57	1.227	7.785e-04
cell organization and biogenesis (Input set freq.: 0.33; 0.11)	2.81	0.94	6.493e-03
developmental processes (Input set freq.: 0.33; 0.12)	2.62	0.781	9.223e-03
signal transduction (Input set freq.: 0.14; 0.05)	2.53	1.084	0.083

http://bar.utoronto.ca/ntools/cgi-bin/ntools_classification_superviewer.cgi#annotation_list

Supplementary Figure S4 SuperViewer classification for genes listed in clusters 1 and 2.

AGI	Name	<i>pldζ1ζ2</i> / WT control		<i>pldζ1ζ2</i> +NaCl / WT+NaCl	
		Log2 Ratio	Pvalue	Log2 Ratio	Pvalue
AT5G35935	unknown protein_copia-like	1.64	0.00E+0	1.73	0.00E+0
AT1G23390	unknown protein	1.28	0.00E+0	0.78	1.40E-5
AT1G08830	CSD1; copper. zinc superoxide dismutase	1.27	0.00E+0	0.78	1.02E-5
AT1G66100	toxin receptor binding_ thionin. putative	1.23	0.00E+0	0.65	4.41E-3
AT1G12520	CCS1; superoxide dismutase copper chaperone	1.04	5.30E-9	0.81	2.04E-6
AT3G02380	COL2 transcription factor	0.81	3.05E-4	0.64	7.26E-3
AT5G48485	DIR1 lipid transfer protein (LTP)	0.74	3.99E-3	0.60	4.57E-2
AT4G09320	NDPK1 nucleoside diphosphate kinase type 1	0.72	7.93E-3	0.73	1.58E-4
AT3G48990	AMP-dependent synthetase and ligase	-0.84	7.33E-5	-1.06	0.00E+0
AT5G08530	CI5. NADH-ubiquinone oxidoreductase 51 kDa	-0.85	5.72E-5	-0.95	9.52E-10
AT3G23820	GAE6	-0.87	2.15E-5	-1.09	0.00E+0
AT5G05170	CESA3 (CELLULASE SYNTHASE 3)	-0.92	2.63E-6	-0.98	1.12E-10
AT4G32410	CESA1 (CELLULASE SYNTHASE 1)	-0.94	9.08E-7	-1.00	3.92E-11
AT3G46970	ATPHS2	-1.01	3.21E-8	-1.09	0.00E+0
AT1G59870	ABC transporter family protein	-1.08	8.23E-10	-1.04	0.00E+0
AT2G25490	EBF1 (EIN3-BINDING F BOX PROTEIN 1)	-1.15	1.12E-11	-1.31	0.00E+0
AT4G38770	PRP4 (PROLINE-RICH PROTEIN 4)	-1.29	0.00E+0	-0.94	1.15E-9
AT3G60750	transketolase_ transketolase. putative	-1.31	0.00E+0	-1.01	1.12E-11
AT3G33002	ribosomal protein S2p family	-1.33	0.00E+0	-1.11	0.00E+0
AT5G42020	ATP binding_ luminal binding protein 2 (BiP-2)	-1.38	0.00E+0	-1.53	0.00E+0
AT3G09440	HSP70-3	-1.61	0.00E+0	-1.38	0.00E+0
AT4G30650	low temperature and salt responsive protein	-1.82	0.00E+0	-1.11	0.00E+0
AT5G02500	HSC70-1	-1.83	0.00E+0	-1.39	0.00E+0

Supplementary Figure S5 List of genes in cluster 2. Genes are involved in electron transport, biotic and abiotic stress and stress responses. Expression data are given as log₂ ratios.

Supplementary Table S1 List of primer sequences used for RT-PCR analysis and genotyping.

For RT-PCR

Gene name	Accession number	5'-3' Sequence
DDF1-F	AT1G12610	GGGACTTATCCCACAGCAGA
DDF1-R	AT1G12610	ATCATTGGATTCCGGCACC
RD29AF	AT5G52310	CAAAACAGAGCACTTACACAGAGAA
RD29AR	AT5G52310	CATAATCTCTACCCGACACACTTTT
APT1F	AT1G27450	GAGACATTTTGCCTGGGATT
APT1R	AT1G27450	CGGGGATTTTAAGTGAACA
APX1F	X59600.1	CTGACATTCCTTCCACCCT
APX1R	X59600.1	CAGACCTATCCTTGTGGCAT
CSD1F	AT1G08830	GGTTCCATGTCCATGCTCT
CSD1R	AT1G08830	ATTGTGAAGGTGGCAGTTCC
CAT2F	AT4G35090.1	AACTCTGGTGCTCCTGTATGG
CAT2R	AT4G35090.1	CTCCAGTTCTCTTGGATGTG

For genotyping

A	PLDzeta1F	At3g16785	TCAGAATCACTTAAGAGGAGATGGG
B	PLDzeta1R	At3g16785	TTTTCGCATAGTCACTTGCTGT
C	LBb1	T-DNA left border	GCGTGGACCGCTTGCTGCAACT
D	PLDzeta2F	At3g05630	TCTCTGTTTTGGGCGGTACGA
E	PLDzeta2R	At3g05630	AAAATGTTTCAGCGTTCTGGAT