



**HAL**  
open science

## Application of guidelines for aminoglycosides use in French hospitals in 2013–2014

jerome robert, Y. Péan, S. Alfandari, J-P Bru, J.P. Bedos, C Rabaud, R. Gauzit

► **To cite this version:**

jerome robert, Y. Péan, S. Alfandari, J-P Bru, J.P. Bedos, et al.. Application of guidelines for aminoglycosides use in French hospitals in 2013–2014. *European Journal of Clinical Microbiology and Infectious Diseases*, 2017, 36 (7), pp.1083-1090. 10.1007/s10096-016-2892-5 . hal-01560010

**HAL Id: hal-01560010**

<https://hal.sorbonne-universite.fr/hal-01560010v1>

Submitted on 11 Jul 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1                                   **Application of guidelines for aminoglycosides use**

2                                   **in French hospitals in 2013-2014**

3  
4 Jérôme Robert<sup>1,2</sup>, Yves Péan<sup>3</sup>, Serge Alfandari<sup>4</sup>, Jean-Pierre Bru<sup>5</sup>, Jean-Pierre Bedos<sup>6</sup>, Christian  
5 Rabaud<sup>7</sup>, Rémy Gauzit<sup>8</sup> on behalf of the Société de Pathologie Infectieuse de Langue Française  
6 (SPILF), the Observatoire National de l'Epidémiologie de la Résistance Bactérienne aux  
7 Antibiotiques (ONERBA) and the Surveillance de la Prescription des Antibiotiques (SPA) group

8  
9 <sup>1</sup> Sorbonne Universités, UPMC Univ Paris 06, CR7, INSERM, U1135, Centre d'Immunologie  
10 et des Maladies Infectieuses, CIMI, team E13 (Bacteriology), F-75013, Paris, France

11 <sup>2</sup> AP-HP, Hôpitaux Universitaires Pitié Salpêtrière – Charles Foix, Bacteriology and Hygiene,  
12 F-75013 Paris, France

13 <sup>3</sup> Observatoire National de l'Epidémiologie de la Résistance Bactérienne aux Antibiotiques  
14 (ONERBA), Paris, France

15 <sup>4</sup> Service de Réanimation et Maladies Infectieuses, CH Gustave Dron, Tourcoing, France

16 <sup>5</sup> Infectious Diseases Department, CH de la Région d'Annecy, Annecy, France

17 <sup>6</sup> Intensive Care Unit, CH Henri Mignot de Versailles, Le Chesnay, France

18 <sup>7</sup> Infectious Diseases Department, CHU de Nancy, Nancy, France

19 <sup>8</sup> Intensive care Unit, CHU Cochin, APHP, Paris, France

20  
21 Running title: Aminoglycosides use in French hospitals

22 **\* Corresponding Author:** Jérôme ROBERT

23                                   Laboratoire de Bactériologie-Hygiène

24                                   Faculté de Médecine Pierre et Marie Curie (UPMC Paris 6),

25                                   91 Boulevard de l'hôpital

26                                   75634 Paris 13, France

27                                   Tel: (33) 1 40 77 97 49

28                                   Fax: (33) 1 42 16 20 72

29                                   E-mail: jerome.robert@aphp.fr

30 Word count: 2539

31  
32 Key words: aminoglycosides, amikacine, gentamicin, dosage

35 **Abstract**

36 **Purpose.** In 2011, the French Agency for Safety of Health Products issued guidelines  
37 underlining the principles of proper aminoglycosides' use. The aim of the survey was to  
38 evaluate adherence to these guidelines two years after their issue.

39 **Methods.** Characteristics of patients receiving aminoglycosides were recorded by voluntary  
40 facilities during a 3-month survey in 2013-2014. The modalities of aminoglycosides treatment  
41 were analysed by comparison with the French guidelines.

42 **Results.** 3323 patients were included by 176 facilities. Patients were mainly hospitalized in  
43 medical wards (33.0%), and treated for urinary-tract infections (24.7%). Compliance  
44 regarding the clinical indication and the daily aminoglycosides dose was observed in 65.2%  
45 and 62.9% of the cases, respectively. A 30-minute once-daily IV administration was recorded  
46 in 62.5% of the cases. Aminoglycosides treatment duration was appropriate ( $\leq 5$  days) for  
47 93.6% of the patients. When considering the four criteria together, 23.2% of the patients had a  
48 treatment regimen aligned with the guidelines. Requests for measurements of peak and trough  
49 AG serum concentrations matched the guidelines in 24.9% and 67.4% of the cases,  
50 respectively.

51 **Conclusions.** Two years after guidelines issue, aminoglycosides use remains unsatisfactory in  
52 French health-care facilities. Efforts should be made for guidelines promotion, especially  
53 regarding the issue of underdosing.

54

55 **Introduction**

56 Despite their rather old age, aminoglycosides (AG) continue to be widely used for the  
57 treatment of severe infections, including endocarditis, due to Gram-negative bacilli,  
58 staphylococci or enterococci, partly due to their broad antibacterial spectrum and the recent  
59 emergence of multi-resistant microorganisms. AG pharmacokinetic and pharmacodynamic  
60 properties include rapid concentration-dependent bactericidal activity, and a narrow  
61 therapeutic index (renal and auditory toxicity). The therapeutic effect is highest if the peak  
62 plasma concentration (C<sub>max</sub>)/minimal inhibiting concentrations (MIC) ratio is over 8 to 10  
63 [1,2]. As most broad-spectrum antibiotics, AG are used in clinical practice on an empirical  
64 basis as well as after availability of antibiotic susceptibility tests. In fact, because of their  
65 toxicity, AG are recommended only in the first days of treatment, i.e. when the bacterial  
66 inoculum is heavy, but also when the causative agent and its antibiotics susceptibility are  
67 unknown.

68 Because of AG characteristics, special attention should be given to AG daily dose  
69 determination, treatment duration, route of administration, and in some settings, to drug  
70 monitoring.

71 Although these requirements are known since the mid-1980s, AG use remained often  
72 inappropriate, in adult patients [3,4], as well as in the paediatric population [5,6].

73 In 2011, a multidisciplinary group of experts was commissioned by the French Agency for  
74 Safety of Health Products (ANSM) to develop up-to-date recommendations on the proper use  
75 of intravenous AG [7]. Two years after their issue, we decided to evaluate the appropriateness  
76 of AG prescriptions in the light of these recommendations.

77

78

79

80 **Methods**

81

82 Study design

83 Practitioners of public and private health-care facilities registered to the French society for  
84 infectious diseases (SPILF, [www.infectiologie.com](http://www.infectiologie.com)) or to the French observatory for national  
85 epidemiology of bacterial resistance to antibiotics (ONERBA, [www.onerba.org](http://www.onerba.org)) were asked  
86 to participate in an observational prospective study on AG use. From November 2013 to  
87 January 2014, each facility had to record data for at least 10 consecutive inpatients, or all  
88 inpatients if less than 10 cases were eligible, treated by AG. Topical and prophylactic uses of  
89 AG were excluded. Only the first prescription was considered in case of multiple AG  
90 regimens during the study period.

91

92 Data collection

93 Basic demographic data, renal function, prior history of hospitalization and antibiotic  
94 treatment in the previous three months, or received since admission and before the first AG  
95 administration were recorded.

96 Data regarding AG prescription included the site of infection, empirical versus documented  
97 treatment, presence of septic shock or others reasons for AG choice, and concomitant  
98 antibiotics used. Modalities of AG treatment included mode of administration, dose  
99 administered, treatment duration, and drug monitoring by determining serum concentrations.  
100 The modalities of treatment were analysed by comparison with the French recommendations  
101 for AG use issued in 2011 by the French for Safety of Health Products [7]. Briefly,  
102 appropriate administration was defined as AG administered intravenously over 30 min in a  
103 once-daily dose or multiple daily doses in case of endocarditis. Duration was considered  
104 appropriate if AG-containing treatment was  $\leq 5$  days, excepted in case of endocarditis, bone  
105 and joint infections and cystic fibrosis. Appropriate daily dose was defined as 15-30 mg/kg  
106 bodyweight for amikacine, 3-8 mg/kg bodyweight for gentamicin and tobramycin, and 4-8

107 mg/kg bodyweight for netilmicin. In case of septic shock or severe sepsis, the higher upper  
108 limits of the ranges were required. Appropriate AG indications were limited to severe  
109 infections (septic shock, complicated pyelonephritis, Gram-positive endocarditis, infections  
110 due to *P. aeruginosa*, *Acinetobacter sp.* ...), high-risk infections (late nosocomial infections  
111 and foreign-body infections) or infections in high-risk patients (cystic fibrosis, newborns, and  
112 immunosuppressed patients). Monitoring of AG peak serum concentration was not required if  
113 treatment duration was  $\leq 3$  days, except in cases of septic shock, severe burns, febrile  
114 neutropenia, intensive care units (ICU) patients with mechanical ventilation, morbid obesity,  
115 polytrauma patients, cystic fibrosis. Monitoring of AG trough concentration was required in  
116 case of planned or effective treatment duration  $> 5$  days, and in case of severe renal  
117 impairment, as declared by clinicians. In other cases, no trough monitoring was required.

118

119 Multidrug-resistant bacteria were defined as Enterobacteriaceae producing extended-spectrum  
120  $\beta$ -lactamase (ESBL), or resistant to carbapenems, and methicillin-resistant *Staphylococcus*  
121 *aureus* (MRSA). Enterobacteriaceae resistant to extended-spectrum cephalosporins but  
122 susceptible to carbapenems and ESBL-negative, and antibiotic resistance patterns of  
123 *Pseudomonas aeruginosa* and *Acinetobacter spp.* isolates were also recorded.

124

### 125 Statistical analysis

126 Continuous variables are expressed as median and range, and were compared by using the  
127 Kruskal-Wallis test. Chi<sup>2</sup> test of Fisher's exact test were used when appropriate for comparing  
128 categorical variables. For multi-level categorical variables, chi<sup>2</sup> tests for homogeneity are  
129 presented. Statistical analysis was performed by using STATA (STATA Corp, College  
130 Station, TX, USA) and  $p < 0.05$  was deemed significant.

131 A multivariate analysis model was developed in order to determine variables independently  
132 associated with a daily AG dose in the recommended ranges. Variables with  $p < 0.10$  in  
133 univariate analysis were introduced in the model, and backward analysis was performed.

134 Variables not significantly associated with the outcome were removed based on the Wald  
135 statistic. The Hosmer-Lemeshov test was used for assessing model' fitness. Only the most  
136 parsimonious model, i.e. the model with the least variables and the most significance, is  
137 presented.

138

139

## 140 **Results**

### 141 Facilities

142 A total of 215 healthcare facilities (25 teaching hospitals, 158 non-teaching or private  
143 hospitals and 32 rehabilitation or long-term care facilities) participated in the study. The  
144 participating facilities accounted for a total of 56,232 acute-care beds and 21,529  
145 rehabilitation or long-term care beds, representing 19% of all French healthcare beds. Among  
146 all facilities, 39 did not record any patient treated by AG during the study period, resulting in  
147 176 facilities that recorded at least one patient treated by AG. Among the 176 latter, 98  
148 (55.7%) declared reviewing systematically all AG-containing regimens, including 79 in all  
149 wards of the facility, and 42 by an electronic system. However, only 43 of the 98 (43.9%)  
150 facilities reviewing all prescriptions have organized an AG control feedback to the  
151 prescribers.

152

### 153 Aminoglycosides use

154 A total of 3,323 patients with a least one AG regimen were included in the study (Table 1),  
155 including 2,007 (60.4%) treated by gentamicin, 1,267 (38.1%) by amikacin, and 49 (1.5%) by  
156 another AG (Table 2).

157 Patients were mainly hospitalized in medical wards (n=1 098, 33.0%), surgical wards  
158 (n=1 002, 30.2%), or in ICU (n=600, 18.1%). The median age of the patients was 65.0  
159 (interquartile range IQR, 48-78) years, 20.9% were more than 80 years old, 1,878 (56.5%)  
160 were male, and 836 (25.2%) had renal failure (Table 1). Patients were mainly treated for  
161 urinary-tract infections (n=822, 24.7%) and digestive or respiratory tract infection (n=653,  
162 19.7% and n=601, 18.1%, respectively).

163 The use of an AG in the antibiotic regimen was justified by the presence of a septic shock in  
164 447 (13.5%) cases. In the absence of septic shock, AG-containing regimens were prescribed  
165 in case of high-risk infections (n=579, 17.4%), infection in high-risk patients (n=292, 8.8%),



166 and pyelonephritis (n=438, 13.2%). The presence or suspicion of multidrug-resistant  
167 organisms accounted for only 129 (3.9%) cases. AG were used on an empirical basis in 2568  
168 (77.3%) cases, and on a bacteriologically documented basis for 755 (22.7%) patients. Among  
169 the 755 latter, AG were used to treat infections due to Enterobacteriaceae in 352 (46.6%)  
170 patients, *Pseudomonas aeruginosa* in 133 (17.6%) cases, *Staphylococcus aureus* in 148  
171 (19.6%) cases, and streptococci or enterococci in 128 (17.0%) cases.

172 Administration by a single daily dose was the rule (n=3061, 92.1%), but its duration was over  
173 30 minutes in only 2185 (65.8%) cases. The median daily dose was in the recommended  
174 ranges for all AG, although at the lower range, and the median duration was 3 days (IQR, 2-3)  
175 days (Table 2).

176

### 177 Compliance

178 AG compliance with the French guidelines was assessed according to four main criteria.

179 The **clinical indication** for AG was respected for 2167 (65.2%) patients (Table 3).  
180 This proportion was higher for patients treated on a bacteriologically documented basis  
181 (75.8%) than for those treated on an empirical basis (62.1%; p<0.01). Pyelonephritis and  
182 community-acquired digestive tract infections represented 33.2% and 23.0% of inappropriate  
183 AG indications, respectively.

184 **Compliance regarding the total daily AG dose** was observed for 2091 (62.9%)  
185 patients (Table 3). Of interest, patients in large facilities (> 300 beds) or university hospitals  
186 were slightly more likely to receive the recommended daily AG dose (65.0%) than in the  
187 other facilities (59.6%; p<0.01). Patients in facilities claiming having a process for reviewing  
188 all AG-containing regimens, including those having an AG control feedback to the prescriber  
189 were not more likely to receive the recommended daily AG dose than those in facilities  
190 without any AG review process.

191 **Once-daily IV administration over 30 minutes** was observed for 2076 (62.5%)  
192 patients (Table 3).

193           The **overall duration of AG** treatment regimen was concordant with the guidelines,  
194 i.e. mainly 5 days or less, for 3110 (93.6%) patients. When considering all four criteria  
195 together, only 23.2% of the patients had an AG treatment regimen in full accordance with the  
196 guidelines. 2.0

197           In a logistic multivariate analysis, having a normal renal function (Odds ratio, 1.7;  
198 95% confidence interval, 1.3-2.2), and being hospitalised in a large facility (OR: 2.0) were the  
199 two variables independently associated with a daily AG dose in the recommended range  
200 (Table 4). Others factors, including age  $\geq$  75 years (OR: 0.7), overweight (OR 0.5), septic  
201 shock (OR: 0.07), and infection in high-risk patients (OR: 0.02) were inversely associated to  
202 having a dose in the recommended range. All other introduced factors, including MDR  
203 bacteria or endocarditis were not independently associated with a dose in the recommended  
204 range. When forced in the model although not significant in univariate analysis, none of the  
205 variables linked to the review process of AG in the facility were associated with the outcome  
206 variable.

207           Finally, requests for measurements of peak and trough serum concentrations matched the  
208 guidelines in 828 (24.9%) and 2241 (67.4%) cases (Table 3).

209

## 210 **Discussion**

211 The present survey aimed at evaluating adherence to AG guidelines in French healthcare  
212 facilities. The results show that AG are used in all type of wards, and that ICUs represented  
213 only 18.1% of all AG prescriptions. As expected, AG were mainly used in association with  
214 other antibiotics (97.1%) and on an empirical basis (77.3%). Indications for AG use were  
215 considered unnecessary in more than 1 out of 3 cases (34.8%). The total AG daily dose was in  
216 the recommended ranges in only 62.9% of the cases. Finally, the AG treatment duration was  
217  $\leq 5$  days for a majority of cases (93.6%).

218

219 The primary indication of AG use was concordant with the guidelines in 65.2% of the  
220 cases. This means that, for one third of the patients, the use of AG could be challenged. Such  
221 a result underlines the need for disseminating information regarding AG indications. Of  
222 interest, patients with pyelonephritis represented a large part of those with AG use that did not  
223 match guidelines criteria. The rise in Enterobacteriaceae producing extended-spectrum beta-  
224 lactamase, and in fluoroquinolone resistance in the community may explain AG overuse [8].  
225 After the issue of the French AG guidelines, the French Infectious Diseases Society updated  
226 guidelines for the management of community-acquired urinary tract infections  
227 ([www.infectiologie.com](http://www.infectiologie.com)). In the latter, AG are indicated on an empirical basis only in case of  
228 complicated pyelonephritis, i.e. with severe sepsis or with need of invasive procedure on the  
229 urinary tract. These guidelines should further decrease AG indications in pyelonephritis. On  
230 the contrary, AG are part of IDSA guidelines for the treatment of uncomplicated  
231 pyelonephritis, but usually as a single antibiotic, which is seldom the case in our study [9].

232

233 In the present survey, AG daily dose was in the recommended ranges for 62.9% of the  
234 patients. In multivariate analysis, we showed that older age, obesity, septic shock and  
235 infections in high-risk patients were factors associated to AG underdosing. Such results have  
236 been previously reported [10,11]. This discordance with the guidelines is likely to be partly

237 linked to the narrow therapeutic index of AG, that encourage prescribers to use lower doses to  
238 avoid toxicity, although pharmacokinetic/pharmacodynamic objectives have been described  
239 25 years ago [1,2]. However, AG toxicity is not directly related to peak serum concentration  
240 and toxicity remains similar for doses below or within the recommended ranges [12].  
241 Patients with weight > 100 kg are prone to receive AG doses below ranges recommended in  
242 the French guidelines. However, it should be noticed that computation of AG daily dose is  
243 complex in such patients. Indeed, guidelines are not very clear regarding computation of AG  
244 daily dose in overweight or obese patients. The use of the actual body weight, an adaptation  
245 of the ideal body weight plus a percentage of the patient's excess bodyweight, or lean weight  
246 is still debatable [13–15]. Therefore, efforts should be made to clarify AG dose computation  
247 in the overweight population, which may represent more than one third of the patients in  
248 many part of the world [16].  
249 Finally, it has been previously reported that ICU patients, and especially those with severe  
250 sepsis or septic shock, are at increased risk of AG underdosing, which consequently results in  
251 low peak serum concentrations [11,17]. This has been linked to an increase in the volume of  
252 distribution per kilogram in these patients. The recent French guidelines have been adapted to  
253 take into account the need for increasing AG daily dose in the ICU population. However, our  
254 results show that changes have not been taken into account. Despite higher recommended  
255 loading doses in the updated guidelines, it has been shown that as much as one third of  
256 patients in severe sepsis may have aminoglycosides serum peak level below the therapeutic  
257 target [11].

258

259 As recommended in French guidelines, more than 93% of the patients received AG for a  
260 duration  $\leq 5$  days, except for endocarditis and bone and joint infections. The 5-day cut-off is  
261 considered as a good compromise between efficacy and safety [18,19]. However, it is  
262 currently suggested to use a shorter duration of time, i.e.  $\leq 72$  hours of treatment. The

263 treatment duration could be prolonged to 5 days in case of unsatisfactory clinical  
264 improvement or in absence of positive bacteriological result.

265

266 Our study has some weaknesses. First it is based on a voluntary participation of facilities,  
267 and as always, representativeness could be questioned. However, the large number of patients  
268 included in a high number of facilities throughout the French territory may have limited this  
269 bias. Second, we did not record any information regarding the initial prescriber of AG-  
270 containing regimen, which could have helped to understand discrepancies with guidelines.  
271 However, we did not show any differences in overall guideline compliance between facilities  
272 with a process for reviewing AG-containing regimens and the others. This raises the question  
273 of effective AG stewardship or of facility organisation. Precise data regarding the review  
274 process, including the background training of the reviewer or consultant, were not collected.

275

276 In conclusion the use of aminoglycosides in French healthcare facilities remains inappropriate  
277 in a substantial proportion of cases although guidelines availability since more than two years.  
278 This is not surprising when considering the numerous barriers to guidelines implementation.  
279 [20] In addition, in France, guidelines diffusion is usually passive or semi-passive, while it  
280 has been shown that better antibiotic use requires multifaceted interventions [21,22]. This is  
281 especially worrisome regarding the use of an appropriate loading dose. The use of higher  
282 loading doses should be widely publicized and use of computerized system for optimized  
283 dose computation in coordination with the hospital pharmacist and infectious diseases  
284 specialist may help improving this situation.

285

286

287 **Acknowledgements**

288 In memoriam: Sandrine CLEMENT ([www.neanima.fr](http://www.neanima.fr)). Our deepest sympathies go out to her  
289 family.

290

291 The Surveillance de la Prescription des Antibiotiques (SPA) group: name (facility, city)

292

293 P. ABGUEGUEN (CHU Angers, ANGERS), A. AKPABIE (Hôpital E. Roux - APHP, LIMEIL BREVANNES),  
294 S. ALFANDARI(CH Tourcoing, TOURCOING), P. ANDRE (Clinique du Millenaire, MONTPELLIER)  
295 (Polyclinique ST-Roch, MONTPELLIER), V. ANGELUS (Clinique L'Angelus, MARSEILLE), S. ARENA-  
296 CANAULT (Maternite Catholique Provence L'Etoile, AIX EN PROVENCE), A. ARREGUY (CH St-Palais,  
297 BAYONNE), N. BACHALAT (Hôpital Joffre - APHP, DRAVEIL), MT. BANOS (CH Lannemezan,  
298 LANNEMEZAN), P. BAUNE (Hôpital Charles Richet, VILLIERS LE BEL), O. BELLON (CH Brignoles,  
299 BRIGNOLES), M. BENAÏSSA (CH Apt, APT), N. BENICHOUGRANE (Hôpital Européen, MARSEILLE), G.  
300 BENISTAND (Polyclinique Grand-Sud, NIMES), F. BERGHEAU (Ch Pays de Gier, St CHAMOND), N.  
301 BERTRAND (Clinique Chantecler, MARSEILLE), C. BERTRAND (Clinique Rhone Durance, AVIGNON), C.  
302 BIANCHI (CHR Pontchaillou, RENNES), H. BLAISE (Clinique du Parc, NANTES), C. BONNAL (Hôpital  
303 Bretonneau - APHP PARIS), S. BORDES-COUECOU (CHIC Cote Basque, BAYONNE), A. BOUMEDIENE  
304 (CHCB Kerio, PLEMET), M. BOURLEAUD (CH de Bastia, BASTIA), S. BOURZEIX DE LAROUZIERE  
305 (CH Moulin Yzeure, MOULINS), D. BREGER (Clinique Sourdille, NANTES), C. BROCARD (Clinique Les  
306 Ormeaux, LE HAVRE), JP. BRU (CH Région d'Annecy, PRINGY), P. CABARET (ES St-Vincent de Paul,  
307 LILLE), P. CABARET (Hôpital St Philibert, LOMME), P. CABARET (Clinique Ste Marie, CAMBRAI), V.  
308 CADIOU (SLD Beauséjour, HYERES), B. CASTAN (CH ND la Miséricorde, AJACCIO), B. CATTIER (CH  
309 Amboise-Chateaurenault, AMBOISE), C. CAZORLA (CHU Saint-Etienne, ST ETIENNE), D. CERVONI  
310 (Centre Les Arbelles, BOURG EN BRESSE), M. CHADAPAUD (CH MJ. Treffot, HYERES), C. CHAIX (CH  
311 Carpentras, CARPENTRAS), C. CHAUVET (CH Montélimar, MONTELMAR), P. CIPIERRE (Clinique du  
312 Parc, PERIGUEUX), M. CLIQUENOIS (Hôpital de L'Isle-Adam, L'ISLE ADAM), Y. COLIN (Centre Medical  
313 L'Arbizon, BAGNERES DE BIGORRE), S. COMPAROT (CH L. Giorgi, ORANGE), O. CORBELLI (Clinique  
314 Vignoli SALON DE PROVENCE), N. CORDAT (Clinique du Val de l'Ouest, ECULLY), S. CORMONT  
315 (Hôpital de Felleries Liesses, FELLERIES), M. CORNESSE (Hôpital Beauregard, MARSEILLE), F.  
316 COULOMB (CH Dreux, DREUX), J. CROUZET (CH Pasteur, BAGNOLS SUR CEZE), J. DARASTEANU  
317 (CH Chartres, CHARTRES), M. DAUMAS (CH Imbert Joseph, ARLES), MH. DEBOISSE (Clinique du  
318 Colombier, LIMOGES), C. DEBRUILLE (CH Douai, DOUAI), B. DECOUARD (MSPBx Bagatelle,  
319 TALENCE), S. DEFRETIN (Clinique St-Roch, RONCQ), B. DEGRENDEL (Centre Sainte Clotilde, STE  
320 CLOTILDE), J. DELHOMME (CHIC Alencon-Mamers, ALENCON), G. DELHON BUGARD (Clinique  
321 Charcot, STE FOY LES LYON), G. DEMELIN (Hôpital Thiais, THIAIS), A. DEMOUZON (EPDS de Gorze,  
322 GORZE), E. DEVAUD (CH R. Dubos, PONTOISE), S. DEWULF (CH Zuydcoote, ZUYDCOOTE), A. DINH  
323 (Hôpital R. Poincaré - APHP, GARCHES), C. DOMRAULT-TANGUY (CH Henin Beaumont, HENIN  
324 BEAUMONT), F. DOMY (Clinique Trenel, STE COLOMBE), F. DOUCET-POPULAIRE (Hôpital A. Beclere  
325 - APHP, CLAMART), L. DRIEUX-ROUZET (Hôpital Charles Foix - APHP, IVRY SUR SEINE), M.  
326 DUVIQUET (Hôpital Vaugirard - APHP, PARIS), A. EDEN (CH Joffre, PERPIGNAN), S. EDOUARD (CH  
327 Dieppe, DIEPPE), L. EL-HAJJ (Clinique Convert, BOURG EN BRESSE), C. ELOY (CH Troyes, TROYES),  
328 M. EMONET (CH Blois, BLOIS), J. EPIFANIE (Centre E. Clémentel, ENVAL), L. ESCAUT (Hôpital Bicêtre -  
329 APHP, LE KREMLIN BICETRE), F. ESPINASSE (Hôpital Ambroise Paré - APHP, BOULOGNE  
330 BILLANCOURT), C. ETIENNE (CH Gaillac, GAILLAC), M. FABRE (CH P. Oudot, BOURGOIN-JALLIEU),  
331 V. FIIHMAN (Hôpital Louis Mourier - APHP, COLOMBES), E. FORESTIER (CH Chambéry, CHAMBERY),  
332 D. FRAISSE (CH Ales, ALES), C. FRANCESCHI (Hôpital de Lunel, LUNEL), C. FUHRMANN (Centre Léon  
333 Bérard, LYON), M. GACHOT (Institut Gustave Roussy, VILLEJUIF), M. GAILLARD (CH E. Roux, LE PUY  
334 EN VELAY), B. GARO (CHRU Brest, BREST), R. GAUZIT (Hôpital Cochin - APHP, PARIS), P. GEROME  
335 (HIA Desgenettes, LYON), M. GILLMANN (CH Lemire-St-Avold, ST AVOLD), L. GIRAUDON (CH Sète,  
336 SETE), F. GLATH (Hôpital de Pompey/Lay St Christophe, POMPEY), P. GRANIER (CH Fleyriat, BOURG EN  
337 BRESSE), B. GRAVAGNA (Clinique Mutualiste de Lyon E.André et Union, LYON), A. GREDER-BELAN  
338 (CH Mignot, LE CHESNAY), F. GREIL (Centre réadaptation Revel, ST MAURICE / DARGOIRE), S.  
339 GROSSE (Hôpital St Maurice - Moyeuivre-Grande, MOYEUVRE GRANDE), C. GUERIN (SSR l'Amandier,  
340 CHATENAY MALABRY), C. GUIGNABERT (CH G. Ramon, SENS), M. GUILLAUME (CH Voiron,  
341 VOIRON), S. GUITTET (Pôle santé Léonard de Vinci, CHAMBRAY LES TOURS), S. HENARD (CHU  
342 Nancy, VANDOEUVRE LES NANCY), L. HENNEQUIN (Hôpital du Neuenberg, INGWILLER), P.

343 HONDERLICK (Hôpital Foch, SURESNES), S. HONORE (CH Auxerre, AUXERRE), A. HUAULT (EMPR  
344 Le Normandy, GRANVILLE), A. HUOT (Clinique MC Chenove, CHENOVE), L. DUCRUET (CH Ain Val de  
345 Saône, PONT DE VEYLE), L. JEANNIN (Institut de Mar Vivo, LA SEYNE SUR MER), A. JULLIAN (Centre  
346 de rééducation cardio-respiratoire, DIEULEFIT), A. JUNG (Hôpital de Sarralbe, SARRALBE), N. KASSIS-  
347 CHIKHANI (Hôpital Paul Brousse - APHP, VILLEJUIF), E. LAFEUILLE, A. ADE (CHU Pitié-Salpêtrière -  
348 APHP, PARIS), LAGNIEN-GAUME (CH L. Pasteur, DOLE), C. LANTERNIER (Clinique des Alpes,  
349 GRENOBLE), C. LARTIZIEN (Hôpital maritime, BERCK), C. LAURANS (CH Roubaix, ROUBAIX), O.  
350 LAURENT (Clinique Guillaume de Varye, ST DOULCHARD & Clinique Le Blaudy, PRECY), C. LECHICHE  
351 (CHU Caremeau, NIMES 9), D. LECOINTE (GH P. Doumer, LIANCOURT), M. LECOQ (CH W Morey,  
352 CHALON SUR SAONE), N. LEFEBVRE (CHRU Strasbourg, STRASBOURG), A. LEFORT (Hôpital Beaujon  
353 - APHP, CLICHY), AM. LELOUP (DAPT de Châtillon, CHATILLON), R. LEPEULE (CHU A.Chenevier-  
354 H.Mondor - APHP, CRETEIL), T. LEVENT (Polyclinique Vauban, VALENCIENNES & CH Maubeuge,  
355 MAUBEUGE), A. LIGNEREUX (CH Gabriel Martin, ST PAUL), J. LIVARTOWSKI (Hôpital privé d'Antony,  
356 ANTONY), B. LOCTIN (Clinique de La Part-Dieu, LYON), V. LOUBERSAC (Clinique Breteche, NANTES),  
357 JC. LUCET (CHU Bichat - APHP, PARIS), C. MAC NAB (HIA Percy, CLAMART), F. MADOUMIE  
358 (Clinique des Emailliers, LIMOGES), P. MAHE (CH F. Grall, LANDERNEAU), N. MARCHISET (CH  
359 d'Ambert, AMBERT), S. MARGUERY (Clinique Ste Marthe, DIJON), L. MARI (Centre de gérontologie de  
360 Montolivet, MARSEILLE), I. MARTIN (CH Roanne, ROANNE), C. MASSA (Clinique Montréal,  
361 CARCASSONNE), C. MATHERN (Hôpital de Creutzwald, CREUTZWALD), V. MATHIS (Association  
362 hospitalière de Joeuf, JOEUF), L. MAZZONI (Clinica Oxford, CANNES & Clinique Le Méridien, CANNES  
363 LA BOCCA), M. MELET (Polyclinique du Parc, CAEN), M.MELET (Clinique St-Jean Languedoc, Clinique  
364 Sarrus-Teinturiers & Clinique A. Pare, TOULOUSE), M. MERTZ (Hôpital de Billom, BILLOM), V.  
365 MESPLEDE (MRC Sainte Odile, BILLERE), A. MILESI-LECAT (CH Vichy, VICHY), R. MONTEIL (Hôpital  
366 Condat, CONDAT), P. MONTEIL (Médipole de Savoie, CHALLES LES EAUX), V. MOULIN (Hôpital  
367 Corentin Celton - APHP, ISSY LES MOULINEAUX), A. MOUNE (Clinique du Palais, GRASSE), E. MURET  
368 (Clinique St-George, NICE), PH. NAUDE (Maison de convalescence Les Elieux, SEICHAMPS), D. NAVAS  
369 (CHU Nantes, NANTES), G. NICOLAOS (CH Coulommiers, COULOMMIERS), JY. NIZOU (L'Institut  
370 Mutualiste Montsouris, PARIS), M. NOLL-BURGIN (GH St Vincent, STRASBOURG), C. NOWAK (CH  
371 Angoulême, ANGOULEME), M. OLEHAINI (CH Vierzon, VIERZON), O. PANTALONI (Clinique St-Pierre,  
372 PERPIGNAN), O. PATEY (CH Villeneuve-St-Georges, VILLENEUVE ST GEORGES), E. PONCET (Clinique  
373 La Source, ST LEGER LES MELEZES), F. POSPISIL (Polyclinique Synergia, CARPENTRAS), S. POULET  
374 (Polyclinique Les Fleurs, OLLIOULES) , G. RAHAL (Clinique Turin & Hôpital privé des Peupliers, PARIS), C.  
375 RAPP (HIA Begin, ST MANDE), S. RASTOUL (Clinique St-Louis, POISSY), S. RAYNAUD (Clinique  
376 médicale Monie, VILLEFRANCHE DE LAURAGAIS), J. REVEIL (CH Charleville Mezières,  
377 CHARLEVILLE MEZIERES), P. RIBELLE (Clinique Tivoli, BORDEAUX), A. RICARD (Clinique Richelieu,  
378 SAINTES), G. RONDELLOT (CHR Metz-Thionville, HAYANGE), J. ROUSSEAU (Clinique de Cognac,  
379 COGNAC), O. SABOT (CH Belley, BELLEY), L. SAFONT, Polyclinique St-Privat, BOUJAN SUR LIBRON),  
380 M. SAREM (CH Sémur-en-Auxois, SEMUR EN AUXOIS), P. SAUTIER (HP Pays de Savoie, ANNEMASSE),  
381 L. SCHANG (Clinique St-Antoine, NICE), L. SENG (Clinique de Thorigny, SERRIS), V. SIMHA (Hôpital San  
382 Salvador, HYERES), B. SIMPLOT (Hôpital de Lamarche, LAMARCHE), S. SIRE (CH Jean Rougier,  
383 CAHORS), M. SOULERIN (Clinique du Vivarais, AUBENAS), B. SOULLIE (HIA Robert Picque,  
384 VILLENAVE D'ORNON), C. SOUYRI (Centre de rééducation, CHAUDES AIGUES), JP. STAHL (CHU  
385 Grenoble, LA TRONCHE) J. TALARMIN (CH de Cornouaille, QUIMPER), V. TALPIN (Hôpital de La  
386 Clayette, LA CLAYETTE), V. TONNERRE (Clinique Mutualiste du Médoc, LESPARRE MEDOC), J.  
387 TRACOL (Centre chirurgical St-Roch, CAVAILLON), F. TURCHET (Clinique Belledonne, ST MARTIN  
388 D'HERES), F. VANDENBOS (La Maison du Mineur, VENCE), E. VAUTRIN (CH St-Dizier, ST DIZIER), N.  
389 VEISSE (Centre MGEN Pierre-Chevalier, HYERES), V. VERNET-GARNIER (CHU Reims, REIMS), M.  
390 VESANES (Hôpital du Marin, LE MARIN), R. VIAL (Hôpital de Beaujeu, BEAUJEU), F. VIELH (CH Le Secq  
391 de Crépy, BOULAY), P. VILLEMMAIN (CH de St-Flour, ST FLOUR), P. VILLEMMAIN (CH Aurillac,  
392 AURILLAC), P. VILLEMMAIN (CH Thann, THANN), J. VIOT (Centre de convalescence Wilson, ANTIBES ;  
393 Clinique du Parc Impérial, NICE 1 & MC Magnolias, ST LAURENT DU VAR), M. VOGT (Clinique Ste-  
394 Odile, STRASBOURG), C. WATELET (Clinique SSR Chateau de Gleteins, JASSANS RIOTTIER), Tamarins  
395 (Clinique les Flamboyants & Clinique Les Tamarins, LE PORT)

397

398

399 ***Funding:*** The Société de Pathologie Infectieuse de Langue Française (SPILF) provided

400 financial support for data management.

401

402 ***Competing interests:*** none declared.

403

404 ***Ethical approval:*** not required

405

406

407

408



409 **REFERENCES**

410

- 411 [1] Moore RD, Lietman PS, Smith CR. Clinical Response to Aminoglycoside Therapy: Importance of the  
412 Ratio of Peak Concentration to Minimal Inhibitory Concentration. *J Infect Dis* 1987;155:93–9.  
413 doi:10.1093/infdis/155.1.93.
- 414 [2] Zelenitsky SA, Harding GKM, Sun S, Ubhi K, Ariano RE. Treatment and outcome of *Pseudomonas*  
415 *aeruginosa* bacteraemia: an antibiotic pharmacodynamic analysis. *J Antimicrob Chemother*  
416 2003;52:668–74. doi:10.1093/jac/dkg403.
- 417 [3] Chuck SK, Raber SR, Rodvold KA, Areff D. National survey of extended-interval aminoglycoside  
418 dosing. *Clin Infect Dis* 2000;30:433–9. doi:10.1086/313692.
- 419 [4] Leong CL, Buising K, Richards M, Robertson M, Street A. Providing guidelines and education is not  
420 enough: an audit of gentamicin use at The Royal Melbourne Hospital. *Intern Med J* 2006;36:37–42.  
421 doi:10.1111/j.1445-5994.2006.01002.x.
- 422 [5] Houot M, Pilmis B, Thepot-Seegers V, Suard C, Potier C, Postaire M, et al. Aminoglycoside use in a  
423 pediatric hospital: there is room for improvement-a before/after study. *Eur J Pediatr* 2016.  
424 doi:10.1007/s00431-016-2691-0.
- 425 [6] Begg EJ, Vella-Brincat JWA, Robertshawe B, McMurtrie MJ, Kirkpatrick CMJ, Darlow B. Eight years’  
426 experience of an extended-interval dosing protocol for gentamicin in neonates. *J Antimicrob Chemother*  
427 2009;63:1043–9. doi:10.1093/jac/dkp073.
- 428 [7] Anonymous. Update on good use of injectable aminoglycosides, gentamicin, tobramycin, netilmicin,  
429 amikacin. Pharmacological properties, indications, dosage, and mode of administration, treatment  
430 monitoring. *Médecine Mal Infect* 2012;42:301–8. doi:10.1016/j.medmal.2011.07.007.
- 431 [8] Nicolas-Chanoine M-H, Gruson C, Bialek-Davenet S, Bertrand X, Thomas-Jean F, Bert F, et al. 10-Fold  
432 increase (2006-11) in the rate of healthy subjects with extended-spectrum  $\beta$ -lactamase-producing  
433 *Escherichia coli* faecal carriage in a Parisian check-up centre. *J Antimicrob Chemother* 2013;68:562–8.  
434 doi:10.1093/jac/dks429.
- 435 [9] Gupta K, Hooton TM, Naber KG, Wullt B, Colgan R, Miller LG, et al. International clinical practice  
436 guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: A 2010 update  
437 by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious  
438 Diseases. *Clin Infect Dis* 2011;52:e103–20. doi:10.1093/cid/ciq257.
- 439 [10] Fraisse T, Gras Aygon C, Paccalin M, Vitrat V, De Wazieres B, Baudoux V, et al. Aminoglycosides use  
440 in patients over 75 years old. *Age Ageing* 2014;43:676–81. doi:10.1093/ageing/afu023.

- 441 [11] Taccone FS, Laterre P-F, Spapen H, Dugernier T, Delattre I, Layeux B, et al. Revisiting the loading dose  
442 of amikacin for patients with severe sepsis and septic shock. *Crit Care* 2010;14:R53.  
443 doi:10.1186/cc8945.
- 444 [12] Barclay ML, Kirkpatrick CM, Begg EJ. Once daily aminoglycoside therapy. Is it less toxic than multiple  
445 daily doses and how should it be monitored? *Clin Pharmacokinet* 1999;36:89–98.  
446 doi:10.2165/00003088-199936020-00001.
- 447 [13] Payne KD, Hall RG. Dosing of antibacterial agents in obese adults: does one size fit all? *Expert Rev*  
448 *Anti Infect Ther* 2014;12:829–54. doi:10.1586/14787210.2014.912942.
- 449 [14] Pai MP, Nafziger AN, Bertino JS. Simplified estimation of aminoglycoside pharmacokinetics in  
450 underweight and obese adult patients. *Antimicrob Agents Chemother* 2011;55:4006–11.  
451 doi:10.1128/AAC.00174-11.
- 452 [15] Polso AK, Lassiter JL, Nagel JL. Impact of hospital guideline for weight-based antimicrobial dosing in  
453 morbidly obese adults and comprehensive literature review. *J Clin Pharm Ther* 2014;39:584–608.  
454 doi:10.1111/jcpt.12200.
- 455 [16] Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national  
456 prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for  
457 the Global Burden of Disease Study 2013. *Lancet* 2014;384:766–81. doi:10.1016/S0140-  
458 6736(14)60460-8.
- 459 [17] Gonçalves-Pereira J, Martins A, Póvoa P. Pharmacokinetics of gentamicin in critically ill patients: pilot  
460 study evaluating the first dose. *Clin Microbiol Infect* 2010;16:1258–63. doi:10.1111/j.1469-  
461 0691.2009.03074.x.
- 462 [18] Bertino JS, Booker LA, Franck PA, Jenkins PL, Franck KR, Nafziger AN. Incidence of and Significant  
463 Risk Factors for Aminoglycoside-Associated Nephrotoxicity in Patients Dosed by Using Individualized  
464 Pharmacokinetic Monitoring. *J Infect Dis* 1993;167:173–9. doi:10.1093/infdis/167.1.173.
- 465 [19] Maller R, Ahrne H, Holmen C, Lausen I, Nilsson LE, Smedjegard J. Once- versus twice-daily amikacin  
466 regimen: efficacy and safety in systemic Gram-negative infections. *J Antimicrob Chemother*  
467 1993;31:939–48. doi:10.1093/jac/31.6.939.
- 468 [20] Cabana MD, Rand CS, Powe NR, Wu AW, Wilson MH, Abboud P-AC, et al. Why Don't Physicians  
469 Follow Clinical Practice Guidelines? *JAMA* 1999;282:1458. doi:10.1001/jama.282.15.1458.
- 470 [21] Gross PA, Pujat D. Implementing practice guidelines for appropriate antimicrobial usage: a systematic  
471 review. *Med Care* 2001;39:II55–69.
- 472 [22] Barlam TF, Cosgrove SE, Abbo LM, MacDougall C, Schuetz AN, Septimus EJ, et al. Implementing an  
473 Antibiotic Stewardship Program: Guidelines by the Infectious Diseases Society of America and the

474 Society for Healthcare Epidemiology of America. Clin Infect Dis 2016;62:ciw118.

475 doi:10.1093/cid/ciw118.

476

477

478 Table 1. Characteristics of the 3 323 patients treated by aminoglycosides during the 3-month  
 479 study period

<b>Continuous variables</b>	<b>Median</b>	<b>Interquartile range</b>
Age	65	(48-78)
Weight	69	(56-80)
<b>Categorical variables</b>	<b>N</b>	<b>(%)</b>
Sex male	1 878	(56.5)
Renal insufficiency	836	(25.2)
Recent hospitalization	1 445	(43.5)
Recent antibiotic treatment	899	(27.1)
Ward of hospitalization		
- Medicine	1 098	(33.0)
- Surgery	1 002	(30.2)
- Oncology/haematology	167	(5.0)
- Paediatric	244	(7.3)
- Intensive care unit	600	(18.1)
- Rehabilitation and long-term care units	212	(6.4)
Site of infection		
- Respiratory tract	601	(18.1)
- Digestive tract	653	(19.7)
- Urinary tract	822	(24.7)
- Bone and joints	200	(6.0)
- Endocarditis	126	(3.8)
- Febrile neutropenia	92	(2.8)
- Others	829	(24.9)

480

481

482 Table 2. Characteristics of the 3 323 aminoglycosides treatment regimens

<b>Categorical variables</b>	<b>N</b>	<b>%</b>
Drug		
- Amikacin	1 267	(38.1)
- Gentamicin	2 007	(60.4)
- Tobramycin	47	(1.4)
Single daily dose	3 061	(92.1)
Intravenous administration over 30 minutes	2 185	(65.8)
AG in combination regimen	3 228	(97.1)
AG in empirical regimen	2 568	(77.3)
Primary indication for AG use		
- Septic shock	447	(13.5)
- Infection in high-risk patient	292	(8.8)
- High-risk infection (late nosocomial infection, foreign body)	579	(17.4)
- Multidrug-resistant organism (confirmed or suspected)	129	(3.9)
- <i>Pseudomonas</i> sp. or <i>Acinetobacter</i> sp. (confirmed or suspected)	189	(5.7)
- Pyelonephritis	438	(13.2)
- Community-onset digestive tract infection	284	(8.5)
- Endocarditis (confirmed or suspected)	130	(3.9)
- Positive blood culture	97	(2.9)
- Others	738	(22.2)
<b>Continuous variables</b>	<b>Median</b>	<b>Interquartile range</b>
Daily dose (mg/kg bodyweight)		
- Amikacin	15.4	(13.6-20.5)
- Gentamicin	3.3	(2.8-4.9)
- Tobramycin	5.2	(3.1-6.6)
AG treatment duration (days)	3	(2-3)

483

484

485 Table 3. Compliance with aminoglycosides guidelines

<b>Criteria for compliance</b>	<b>N</b>	<b>%</b>
Indication: treatment of severe infections or of high-risk patients	2 167	(65.2)
Daily dose in mg/kg bodyweight in the recommended range and at the upper limit in case of shock or severe sepsis	2 091	(62.9)
Once-daily intravenous administration over 30 minutes	2 076	(62.5)
Duration $\leq$ 5 days excepted for endocarditis, bone and joint infections, and cystic fibrosis	3 110	(93.6)
All four criteria above	771	(23.2)
Monitoring of aminoglycoside peak serum concentration	828	(24.9)
Monitoring of aminoglycoside trough serum concentration	2 241	(67.4)

486

487

488

489

490 Table 4. Univariate and multivariate analysis for association with daily aminoglycoside dose  
 491 in the recommended ranges

Variable	Univariate analysis		Multivariate analysis	
	OR	95% CI	OR	95% CI
Large facility	1.2	1.1-1.5	2.0	1.4-2.9
Age $\geq$ 75 years	0.6	0.56-0.74	0.7	0.56-0.87
Weight $\geq$ 100 kg	0.7	0.54-0.99	0.5	0.36-0.81
Normal renal function	2.2	1.9-2.5	1.7	1.3-2.2
Primary indication for AG use (confirmed or suspected)				
- Septic shock	0.1	0.08-0.13	0.07	0.05-0.10
- <i>Pseudomonas</i> sp. or <i>Acinetobacter</i> sp.	2.3	1.5-3.4	-	
- Multidrug-resistant organism	1.8	1.2-2.8	-	
- Infection in high-risk patient	0.05	0.03-0.07	0.02	0.01-0.04
- Endocarditis	2.3	1.5-3.5	-	

492 OR: odds ratio; CI: confidence interval

493