

Logarithmic corrections to entropy of magnetically charged AdS 4 black holes

Intak Jeon, Shailesh Lal

► **To cite this version:**

Intak Jeon, Shailesh Lal. Logarithmic corrections to entropy of magnetically charged AdS 4 black holes. Physics Letters B, Elsevier, 2017, <10.1016/j.physletb.2017.09.026>. <hal-01591923>

HAL Id: hal-01591923

<https://hal.sorbonne-universite.fr/hal-01591923>

Submitted on 22 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Contents lists available at ScienceDirect

Physics Letters B

www.elsevier.com/locate/physletb



Logarithmic corrections to entropy of magnetically charged AdS₄ black holes

Imtak Jeon^a, Shailesh Lal^b

^a Harish-Chandra Research Institute, Chhatnag Road, Jhusi, Allahabad 211019, India

^b LPTHE – UMR 7589, UPMC Paris 06, Sorbonne Universités, Paris 75005, France

ARTICLE INFO

Article history:

Received 25 August 2017

Accepted 12 September 2017

Available online xxxx

Editor: M. Cvetič

Keywords:

String theory

AdS/CFT correspondence

Black holes

Quantum entropy

AdS black hole

Logarithmic corrections

ABSTRACT

Logarithmic terms are quantum corrections to black hole entropy determined completely from classical data, thus providing a strong check for candidate theories of quantum gravity purely from physics in the infrared. We compute these terms in the entropy associated to the horizon of a magnetically charged extremal black hole in AdS₄ × S⁷ using the quantum entropy function and discuss the possibility of matching against recently derived microscopic expressions.

© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Providing a microscopic interpretation to the Bekenstein-Hawking formula in the context of certain classes of supersymmetric extremal black holes in flat space has been a main success of string theory as a theory of quantum gravity [1–6]. The expression for the microscopic entropy obtained by explicit enumeration and counting of black hole microstates in these cases contains the area law as the leading formula, but also contains higher-derivative and quantum corrections to it. We refer the reader to [7] for a review of these developments as well as more exhaustive references. Importantly, since extremal black holes expectedly possess an AdS₂ factor in the near horizon geometry, one may use the AdS₂/CFT₁ correspondence to provide an alternative, but equivalent, definition of the quantum entropy of extremal black holes in string theory. This proposal is known as the quantum entropy function, and for extremal black holes carrying charges $\vec{q} \equiv q_i$ [8,9],

$$d_{hor}(\vec{q}) \equiv \left\langle \exp \left[i \oint q_i d\theta \mathcal{A}_\theta^i \right] \right\rangle_{AdS_2}^{finite}, \quad (1)$$

where d_{hor} is the full quantum degeneracy associated with the black hole horizon, and \mathcal{A}_θ^i is the component of the i th gauge field

along the boundary of the AdS₂. In this picture the entropy associated to the horizon degrees of freedom of an extremal black hole is essentially the free energy corresponding to the partition function (1). The superscript ‘finite’ reminds us that the quantity on the right hand side of (1) is naively divergent due to the infinite volume of AdS₂ but this divergence may be regulated in accordance with general principles of the AdS/CFT correspondence and a cutoff-insensitive finite part extracted, which is then identified to d_{hor} [8–10]. The path integral is carried out over all fields that asymptote to the black hole near horizon geometry. In the context of supersymmetric extremal black holes in flat space, this has been evaluated using saddle point techniques [10–14,16,17] as well as supersymmetric localization [18–24] and the answer matched with microscopic results wherever available. Importantly, even in the cases where the full microscopic formula is unavailable, this quantity may be evaluated at least using saddle-point methods to gain some insight into the full microscopic formula [13,17].

In contrast, the situation for entropy computations for extremal black holes in AdS space is relatively in its infancy. While the Wald entropy [25] associated to the horizon may be computed for such black holes using for example the entropy function formalism, the quantum corrections to it are still unknown.¹ In this situation, it

¹ Recently the quantum entropy of the topological black hole in AdS₄ has been computed using supersymmetric localization [26], building on the analysis of [27],

E-mail address: shailesh.hri@gmail.com (S. Lal).

<https://doi.org/10.1016/j.physletb.2017.09.026>

0370-2693/© 2017 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

would clearly be of interest to evaluate (1) for such black holes to obtain the set of quantum corrections to the Wald formula. Such computations are further motivated by the recent proposal for a microscopic computation of a CFT₃ index argued to capture the quantum entropy of an AdS₄ extremal black hole [29].

In this note we will focus on the computation of (1) in the semi-classical approximation where the event horizon of the black hole has a large length scale a associated with it. In the case of extremal black holes in flat space, this large length scale arises when the charges of the black hole are taken to be large. In the present case of AdS black holes, this corresponds to taking the rank N of the gauge group in the dual CFT to be large, while the charges themselves are not scaled.

One saddle-point of the path integral (1) is the black hole's near-horizon geometry itself. By evaluating the on-shell action on this field configuration, it is possible to show that [8,9]

$$d_{hor} \simeq e^{S_{Wald}} \Rightarrow S_{BH} = \ln d_{hor} \simeq S_{Wald}. \tag{2}$$

Hence the quantum entropy function produces as the leading contribution, Wald's formula for the entropy of the horizon. In this note, we will consider subleading corrections of the form $\ln a$ to this formula, i.e.

$$S_{BH} = S_{Wald} + c \ln a + \dots, \tag{3}$$

where c is a coefficient which depends on the details of the quantum gravity that the black hole is embedded in. For example, the same four-dimensional black hole which is a quarter-BPS black hole in $\mathcal{N} = 4$ supergravity may be embedded as a one-eighth BPS black hole in $\mathcal{N} = 8$ supergravity and while the leading Bekenstein–Hawking answer is the same for the black hole, the log terms computed in both theories are different [12], and match with the microscopic computations carried out respectively in $\mathcal{N} = 4$ and $\mathcal{N} = 8$ string compactifications. This matching is an important test of the consistency of the quantum entropy function proposal.

The main reason why the log term is an important contribution to the microscopic formula is that it is a genuinely quantum correction to the Bekenstein–Hawking formula, but is determined completely from one-loop fluctuations of massless fields of two-derivate supergravity, which essentially constitute the IR data of the black hole.² To see this, let us recall some elements of a scaling argument presented in [15]. Consider the ℓ -loop free energy for a theory defined on a D dimensional background with a length scale a associated with it. A typical Feynman graph contributing to this quantity would scale as (see [15] for details)

$$\ell_p^{(D-2)(\ell-1)} a^{-(D-2)(\ell-1)} \int d^D \ell \tilde{k} \tilde{k}^{2-2\ell} F(\tilde{k}), \tag{4}$$

where $\tilde{k} = ka$ and $\{k\}$ are the loop momenta, and F is a function which approaches 1 at large values of its arguments. By focusing on the regime where all loop momenta are of the same order, and working at large \tilde{k} , we see that a $\ln a$ term arises from the $\tilde{k}^{2\ell-2-D\ell}$ term in the $\frac{1}{k}$ expansion of F at large \tilde{k} , and the full a dependence of this term is

to find a complete match with the corresponding microscopic expression. We thank Jun Nian for helpful correspondence regarding this. We also note that the quantum entropy of AdS₄ hyperbolic black holes was studied in [28].

² The term 'massless' has to be carefully defined on curved manifolds. The more precise statement is that the eigenvalues of the kinetic operator should scale as $\frac{1}{a^2}$, which is how the eigenvalues of the kinetic operator over a massless scalar, $-\Delta$ would scale.

$$\left(\frac{1}{a}\right)^{(D-2)(\ell-1)} \ln a, \tag{5}$$

which is highly suppressed in the large a limit unless $\ell = 1$. This verifies the above claim that only one-loop fluctuations give rise to the log term. Further, by considering different scaling regimes where various subsets of loop momenta scale to be much larger than the rest, one may verify the fact that only the two derivative sector of massless fields contributes to the log term. However, we do not do so here and instead refer the reader to [15] for those details.

Therefore, as argued earlier, the log term may be regarded as an IR probe of the microscopic theory, in the sense that any putative microscopic description of the black hole must correctly reproduce not only the leading Bekenstein–Hawking area law, but also the log correction to it.

In this note we shall compute the log term for a class of magnetically charged extremal black holes which asymptote to AdS₄ × S⁷ for which a complete expression for the microscopic entropy has recently been computed via the computation of a topologically twisted index in ABJM theory [29]. We omit details of the microscopic formula, referring the reader to [29] as well as the companion papers [30,31]. Further, the only features of the near horizon geometry which shall be relevant to us are that it is AdS₂ × S² × S⁷ where the S⁷ is bundled over the S² and that the AdS₂, S² and S⁷ have a common length scale a associated to them. That is, the metric over the full 11 dimensional near horizon geometry can be brought to the form $g_{\mu\nu} = a^2 g_{\mu\nu}^{(0)}$ where the metric $g^{(0)}$ and the coordinates are a independent.

Further the S² has SO(3) isometry, while the S⁷ has U(1)⁴ isometry. These inputs suffice for the macroscopic computation of the log term as a prediction for the microscopic formula of [29]. We shall finally discuss a few aspects of the proposed match.³ Details of the full black hole solution are reviewed in [29].

2. The log term from the quantum entropy function

We will now describe how the log term may be extracted from the path integral (1). To do so, it is useful to phrase the problem more generally. In particular, we consider a theory in D dimensions, with a dynamical field Φ , admitting a saddle-point which is a background with length scale a .

$$\mathcal{Z}[\Phi] = \int \mathcal{D}\Phi e^{-\frac{1}{\hbar} S[\Phi]}. \tag{6}$$

Then, as argued from Equation (4), to extract the term in the free energy proportional to $\ln a$, it is sufficient to concentrate on the one-loop partition function of the theory. The techniques for this analysis are well-known and we refer the reader to [14,38,16,17] for accounts of how these computations are carried out. Firstly, the one-loop partition function is then given by

$$\mathcal{Z}_{1-\ell} = \det' \mathcal{O}^{-\frac{1}{2}} \cdot (\mathcal{Z}_{\text{zero}})^{n_0}, \tag{7}$$

where $\det' \mathcal{O}$ is the determinant of \mathcal{O} evaluated over its non-zero modes, n_0 is the number of zero modes of \mathcal{O} , and $\mathcal{Z}_{\text{zero}}$ is the residual zero-mode integral. Therefore

$$\ln \mathcal{Z}_{1-\ell} = -\frac{1}{2} \ln \det' \mathcal{O} + n_0 \ln \mathcal{Z}_{\text{zero}}. \tag{8}$$

The final result is that the coefficient of the $\ln a$ term in the free energy computed about this saddle-point depends on $K(t; 0)$,

³ While this draft was being readied for submission, we learned of [32] where this comparison has been carried out in a numerical scheme to find a mismatch.

which is the t^0 coefficient of the heat kernel expansion of the kinetic operator \mathcal{O} about this saddle-point, and β_Φ , which determines how the zero mode contribution $\mathcal{Z}_{\text{zero}}$ to the path integral scales with a .

$$\mathcal{Z}_{\text{zero}} = a^{\beta_\Phi} \hat{\mathcal{Z}}_{\text{zero}}, \tag{9}$$

where $\hat{\mathcal{Z}}_{\text{zero}}$ does not scale with a . We eventually obtain the formula

$$\ln \mathcal{Z}_{1-\ell} = K(t; 0) \ln a + (\beta_\Phi - 1) n_0 \ln a. \tag{10}$$

It is well known that in odd-dimensional spacetimes, $K(0; t) = 0$ and hence we have the formula

$$\ln \mathcal{Z}_{1-\ell} = (\beta_\Phi - 1) n_0 \ln a. \tag{11}$$

The extension to the case of multiple fields $\{\Phi\}$ having zero modes is apparent.

$$\ln \mathcal{Z}_{1-\ell} = \sum_{\phi \in \{\Phi\}} (\beta_\phi - 1) n_0^\phi \ln a, \tag{12}$$

where n_0^ϕ is the number of zero modes of the kinetic operator over the field ϕ . The specific values for β_ϕ that will be relevant to us are for the vector, the graviton, the three-form and the gravitino. These are given by

$$\beta_v = \frac{D-2}{2}, \quad \beta_m = \frac{D}{2}, \quad \beta_f = D-1, \quad \beta_C = \frac{D}{2} - 3. \tag{13}$$

Here v denotes the vector field, m the metric or the graviton, f the gravitino, and the corresponding β values have been listed in Equation (2.37) of [14]. C denotes the three-form field and its β value may be determined exactly in the same manner as the previous fields [14,38]. We start with the expression for the normalization of the field C_{MNP} in D -dimensions.

$$\int \mathcal{D}C_{MNP} e^{\left[-\int d^d x \sqrt{g} g^{MU} g^{NV} g^{PW} C_{MNP} C_{UVW}\right]} = 1. \tag{14}$$

Here the metric scales as $g_{MN} = a^2 g_{MN}^{(0)}$ where $g^{(0)}$ does not scale with a . Therefore we have

$$\int \mathcal{D}C_{MNP} e^{\left[-a^{D-6} \int d^d x \sqrt{g^{(0)}} g^{(0)MU} g^{(0)NV} g^{(0)PW} C_{MNP} C_{UVW}\right]} = 1. \tag{15}$$

Hence the correctly normalized integration measure is

$$\prod_{x, (MNP)} d\left(a^{\frac{D}{2}-3} C_{MNP}\right). \tag{16}$$

From this we can obtain that the zero mode of C_{MNP} corresponds to

$$\beta_C = \frac{D}{2} - 3. \tag{17}$$

3. Counting the number of zero modes

From the above discussion it is clear that if we are to extract the contribution to the (logarithm of the) quantum entropy function which scales as $\ln a$, where a is the length scale associated with the radii of AdS_2 , S^2 and S^7 , it is enough to compute the zero modes of the fields appearing in the path integral (1). Zero modes can in principle appear in the massless spectrum of AdS_2 fields obtained from fields of the 11 dimensional supergravity reduced on to $S^2 \times S^7$. In this section we will enumerate these zero modes and compute their contribution to the quantum entropy of magnetically charged AdS_4 black holes. In isolating these zero modes, a special role is played by the so-called *discrete modes* of

the Laplacian for spin-1 and spin-2 fields, and the Dirac operator for spin- $\frac{3}{2}$ fields on AdS_2 explicitly enumerated in [33,34] respectively, and counting the total number of zero modes is essentially equivalent to counting the total number of discrete modes of these fields. These modes have also been listed in [12,14,17]. An important observation for us is that it turns out that naively the number of zero modes for each of these fields turns out to be infinite. However, this divergence turns out to be essentially equivalent to the volume divergence of the free energy and may be regulated in the same way. Two slightly different, but equivalent, procedures for doing this are available in [11,12,14] and [16,17] and we shall use those results for the regularized number of zero modes in the computations that follow.

3.1. Bosonic zero modes

The bosonic fields are the graviton h_{MN} and the 3-form C_{MNP} . Quantization of the graviton gives rise to a ghost vector field but this has no zero modes. Quantization of the 3-form C gives rise to a ghost 2-form B with Grassmann odd statistics, and a ghost-for-ghost 1-form A with Grassmann even statistics. We use the following conventions: M is an 11-d vector index, μ is an AdS_2 index, a is an S^2 index and i is an S^7 index. Finally α is either a or i .

Consider the metric zero modes first. The graviton h_{MN} decomposes into the AdS_2 graviton $h_{\mu\nu}$, 3 massless AdS_2 vectors $h_{\mu a}$, and 4 massless AdS_2 vectors $h_{\mu i}$, along with the AdS_2 scalars g_{ia} , g_{ij} and g_{ab} . In counting the number of massless vectors, we used the fact that these are given by

$$h_{\mu\alpha} = v_\mu k_\alpha, \tag{18}$$

where k_α is a Killing vector along an internal direction. Since the internal space has $\text{SU}(2) \otimes \text{U}(1)^{\otimes 4}$ isometry, there are $3 + 4 = 7$ Killing vectors. Each massless vector field on AdS_2 contributes

$$n_0^v = -1, \tag{19}$$

hence there are -7 zero modes from the $h_{\mu\alpha}$. Also there are -3 zero modes from the AdS_2 metric $h_{\mu\nu}$. Therefore total number of metric zero modes is

$$n_0^m = -7 - 3 = -10. \tag{20}$$

Then the contribution to the log term from the 11d metric zero modes is

$$\delta Z|_{\text{metric}} = (\beta_m - 1) n_0^m \ln a = -45 \ln a. \tag{21}$$

We next consider the 3-form field C_{MNP} . Its quantization was carried out in [35–37] and is reviewed in Section 3.1 of [38]. We will just require the following result. The quantization of a p -form requires p generalized ghost fields which are $(p-j)$ -forms, where j runs from 1 to p . Further the logarithmic contribution of all these fields to the free energy may be packaged into the expression

$$\Delta F = \sum_{j=0}^p (-1)^j (\beta_{p-j} - j - 1) n_{\mathcal{O}_{p-j}}^0 \ln a, \tag{22}$$

where \mathcal{O} is the kinetic operator. For $j = 1, \dots, p$ which are the generalized ghost fields, this is just the Hodge Laplacian. For $j = 0$, which is the physical field, this can have couplings to background fluxes as well. We now consider the reduction of C_{MNP} onto AdS_2 . This firstly leads to $C_{\mu\nu a} C_{\mu\nu i}$ which are two-forms on AdS_2 which are Hodge duals of scalars. These have no zero modes. Next, we consider a 3-form which is the wedge product of a 1-form along AdS_2 with a harmonic 2-form along $S^2 \times S^7$.

$$C^{(3)} = C_{(1)}^{(\text{AdS}_2)} \wedge C_{(2)}^{(S^2 \times S^7)}. \tag{23}$$

The number of such harmonic forms is given by the second Betti number b_2 of the manifold, which is 1 in this case, as may be readily seen from multiplying the Poincaré polynomials of S^2 and S^7 . Hence there is a single massless vector fields along AdS_2 from the 11 dimensional 3-form field. Finally we have the scalars $C_{abi}, C_{aij}, C_{ijk}$ which don't have zero modes. Hence the 3-form contributes

$$\delta Z|_C = (\beta_C - 1)n_0^C \ln a = -\frac{3}{2} \ln a \quad (24)$$

to the log term. We next consider the ghost B_{MN} which arises from the quantization of C . This decomposes into the following massless fields on AdS_2 . First we have $B_{\mu\nu}$ which contributes no zero modes. Next we may obtain a massless 1-form on AdS_2 from this field by decomposing B_{MN} into a wedge product of an AdS_2 1-form with a harmonic 1-form on $S^2 \times S^7$. The number of such harmonic 1-forms is the first Betti number of $S^2 \times S^7$ which is zero. Finally we have the scalars B_{ab}, B_{ai} and B_{ij} , which contribute no zero modes. Therefore the contribution to the log term is

$$\delta Z|_B = -(\beta_B - 2)n_0^B \ln a = 0. \quad (25)$$

The overall minus sign is on account of Grassmann odd statistics of this field. Finally we have the ghost-for-ghost field A_M from the quantization of B . This leads to one massless vector field on AdS_2 and therefore

$$\delta Z|_A = (\beta_A - 3)n_0^A \ln a = -\frac{3}{2} \ln a. \quad (26)$$

We therefore add (21), (24), (25) and (26) to obtain

$$\delta Z = \left(-45 - \frac{3}{2} - \frac{3}{2}\right) \ln a = -48 \ln a. \quad (27)$$

The residual scalar generalized ghost has no zero modes.

3.2. Fermionic zero modes

To count fermion zero modes, we need to compute the regularized number of discrete modes $\xi_{\mu}^{(k)+}$ and $\hat{\xi}_{\mu}^{(k)+}$, $k = 1, 2, \dots$, on AdS_2 . The relevant computations are available in [12,14,17] and we only mention final results. Firstly, it may be shown that the regularized number of modes of both ξ and $\hat{\xi}$ is given by -1 . Further, these modes should be tensored with the spinors associated with directions transverse to AdS_2 . This will give rise to additional multiplicity factors. To determine the multiplicity, we note first that the near horizon geometry of the black hole has a superconformal symmetry $su(1, 1|1)$ with fermionic generators G_n^α where $\alpha = \pm$ and $n \in \mathbb{Z} + \frac{1}{2}$. The gravitino zero modes we consider are associated with the generators G_n^α where $|n| \geq \frac{3}{2}$. In particular, we identify G_n^α with $n \geq \frac{3}{2}$ with ξ^k where $n = k + \frac{1}{2}$ and G_n^α with $n \leq -\frac{3}{2}$ with $\hat{\xi}^k$ where $n = -k - \frac{1}{2}$. Hence, there is an overall multiplicity factor of 2 coming from α taking values $+$ and $-$. Therefore the number of fermion zero modes is

$$n_0^f = (-1 - 1) \times 2 = -4. \quad (28)$$

Then the contribution to the log term from the fermionic zero modes is

$$\delta Z|_{\text{fermions}} = -(\beta_f - 1)n_0^f \ln a = +36 \ln a. \quad (29)$$

The overall minus is on account of Grassmann odd statistics. Adding (27) and (29) we see that the log term is given by

$$\Delta F = (-48 + 36) \ln a = -12 \ln a. \quad (30)$$

4. On the comparison with microscopics

We have so far computed the log term for magnetically charged AdS_4 extremal black holes using the quantum entropy function. In this section we shall very briefly discuss how in principle a match with the proposed microscopic answer of [29] may be carried out. At first glance one may expect that the large N expansion of the logarithm of the topologically twisted index in the CFT computed in [29] may be matched term by term with the large a expansion of the logarithm of the quantum entropy function. We note however, that several steps should in principle be necessary to carry out before a match can be meaningfully proposed.

Firstly, the index computed by [29] measures the black hole entropy in the grand canonical ensemble as it employs the $\text{AdS}_4/\text{CFT}_3$ correspondence. In contrast, the natural boundary conditions for the quantum entropy function (1) pick out the *microcanonical* ensemble [9]. While the choice of ensemble is irrelevant in the strict large N limit, it is typically important when finite N effects are taken into account. Indeed examples exist where the choice of ensemble is explicitly shown to affect the value of the log term [15]. Therefore, as a first step, we expect that one would need to go to the microcanonical ensemble when computing the CFT answer to match with the quantum entropy function.

Second, it is not obvious how the naive large N scalings of the log of the index are to be reproduced by the quantum entropy function. In particular, it appears from the analysis of [29] that $\ln Z$ scales as

$$\ln Z \sim N^{3/2} + \mathcal{O}(N \ln N). \quad (31)$$

The $N^{3/2}$ term precisely matches with the Bekenstein-Hawking entropy of the black hole, but the possible subleading term of order $N \ln N$ is particularly surprising from the point of view of the general scalings of contributions to the quantum entropy as expected from the quantum entropy function. In particular, on going through the scaling analysis of Equation (4), it is apparent that we do not expect a term of the form $N \ln N$, and that this term should drop from the CFT_3 answer if a match with the quantum entropy function is to be possible.

Remarkably, it seems that a numerical estimate of the large N behavior of the index produces the pattern one would naturally expect from the quantum entropy function scalings, including an absence of the $N \ln N$ term, and the presence of the $\ln N$ term, albeit with a mismatching coefficient [32]. This remarkable feature should certainly be better understood by carrying out a systematic large N expansion of the CFT index while accounting for the choice of ensemble as we have indicated above. It would be interesting to return to these questions in the future.

Acknowledgements

We would like to thank Rajesh Gupta, Nick Halmagyi, Euihun Joung, Tomoki Nosaka, Chiung Hwang, Leopoldo Pando Zayas, Ashoke Sen, Piljin Yi and Alberto Zaffaroni for helpful discussions. SL thanks the Korea Institute of Advanced Study and Kyung Hee University for hospitality while this work was carried out. SL's work is supported by a Marie Skłodowska Curie 2014 Individual Fellowship.

References

- [1] A. Strominger, C. Vafa, Microscopic origin of the Bekenstein-Hawking entropy, Phys. Lett. B 379 (1996) 99, arXiv:hep-th/9601029.
- [2] R. Dijkgraaf, E.P. Verlinde, H.L. Verlinde, Counting dyons in $N=4$ string theory, Nucl. Phys. B 484 (1997) 543, [http://dx.doi.org/10.1016/S0550-3213\(96\)00640-2](http://dx.doi.org/10.1016/S0550-3213(96)00640-2), arXiv:hep-th/9607026.

- [3] B. Pioline, BPS black hole degeneracies and minimal automorphic representations, *J. High Energy Phys.* 0508 (2005) 071, <http://dx.doi.org/10.1088/1126-6708/2005/08/071>, arXiv:hep-th/0506228.
- [4] D. Shih, A. Strominger, X. Yin, Counting dyons in N=8 string theory, *J. High Energy Phys.* 0606 (2006) 037, <http://dx.doi.org/10.1088/1126-6708/2006/06/037>, arXiv:hep-th/0506151.
- [5] J.R. David, A. Sen, CHL dyons and statistical entropy function from D1–D5 system, *J. High Energy Phys.* 0611 (2006) 072, <http://dx.doi.org/10.1088/1126-6708/2006/11/072>, arXiv:hep-th/0605210.
- [6] A. Sen, N=8 dyon partition function and walls of marginal stability, *J. High Energy Phys.* 0807 (2008) 118, <http://dx.doi.org/10.1088/1126-6708/2008/07/118>, arXiv:0803.1014 [hep-th].
- [7] A. Sen, Black hole entropy function, attractors and precision counting of microstates, *Gen. Relativ. Gravit.* 40 (2008) 2249, <http://dx.doi.org/10.1007/s10714-008-0626-4>, arXiv:0708.1270 [hep-th].
- [8] A. Sen, Entropy function and AdS(2) / CFT(1) correspondence, *J. High Energy Phys.* 0811 (2008) 075, <http://dx.doi.org/10.1088/1126-6708/2008/11/075>, arXiv:0805.0095 [hep-th].
- [9] A. Sen, Quantum entropy function from AdS(2)/CFT(1) correspondence, *Int. J. Mod. Phys. A* 24 (2009) 4225, <http://dx.doi.org/10.1142/S0217751X09045893>, arXiv:0809.3304 [hep-th].
- [10] A. Sen, Arithmetic of quantum entropy function, *J. High Energy Phys.* 0908 (2009) 068, <http://dx.doi.org/10.1088/1126-6708/2009/08/068>, arXiv:0903.1477 [hep-th].
- [11] S. Banerjee, R.K. Gupta, A. Sen, Logarithmic corrections to extremal black hole entropy from quantum entropy function, *J. High Energy Phys.* 1103 (2011) 147, arXiv:1005.3044 [hep-th].
- [12] S. Banerjee, R.K. Gupta, I. Mandal, A. Sen, Logarithmic corrections to N=4 and N=8 black hole entropy: a one loop test of quantum gravity, *J. High Energy Phys.* 1111 (2011) 143, arXiv:1106.0080 [hep-th].
- [13] A. Sen, Logarithmic corrections to N=2 black hole entropy: an infrared window into the microstates, *Gen. Relativ. Gravit.* 44 (5) (2012) 1207, <http://dx.doi.org/10.1007/s10714-012-1336-5>, arXiv:1108.3842 [hep-th].
- [14] A. Sen, Logarithmic corrections to rotating extremal black hole entropy in four and five dimensions, *Gen. Relativ. Gravit.* 44 (2012) 1947, <http://dx.doi.org/10.1007/s10714-012-1373-0>, arXiv:1109.3706 [hep-th].
- [15] A. Sen, Logarithmic corrections to Schwarzschild and other non-extremal black hole entropy in different dimensions, *J. High Energy Phys.* 1304 (2013) 156, [http://dx.doi.org/10.1007/JHEP04\(2013\)156](http://dx.doi.org/10.1007/JHEP04(2013)156), arXiv:1205.0971 [hep-th].
- [16] R.K. Gupta, S. Lal, S. Thakur, Heat kernels on the AdS(2) cone and logarithmic corrections to extremal black hole entropy, *J. High Energy Phys.* 1403 (2014) 043, [http://dx.doi.org/10.1007/JHEP03\(2014\)043](http://dx.doi.org/10.1007/JHEP03(2014)043), arXiv:1311.6286 [hep-th].
- [17] R.K. Gupta, S. Lal, S. Thakur, Logarithmic corrections to extremal black hole entropy in $\mathcal{N}=2, 4$ and 8 supergravity, *J. High Energy Phys.* 1411 (2014) 072, [http://dx.doi.org/10.1007/JHEP11\(2014\)072](http://dx.doi.org/10.1007/JHEP11(2014)072), arXiv:1402.2441 [hep-th].
- [18] A. Dabholkar, J. Gomes, S. Murthy, Quantum black holes, localization and the topological string, *J. High Energy Phys.* 1106 (2011) 019, [http://dx.doi.org/10.1007/JHEP06\(2011\)019](http://dx.doi.org/10.1007/JHEP06(2011)019), arXiv:1012.0265 [hep-th].
- [19] A. Dabholkar, J. Gomes, S. Murthy, Localization & exact holography, *J. High Energy Phys.* 1304 (2013) 062, [http://dx.doi.org/10.1007/JHEP04\(2013\)062](http://dx.doi.org/10.1007/JHEP04(2013)062), arXiv:1111.1161 [hep-th].
- [20] R.K. Gupta, S. Murthy, All solutions of the localization equations for N=2 quantum black hole entropy, *J. High Energy Phys.* 1302 (2013) 141, [http://dx.doi.org/10.1007/JHEP02\(2013\)141](http://dx.doi.org/10.1007/JHEP02(2013)141), arXiv:1208.6221 [hep-th].
- [21] A. Dabholkar, J. Gomes, S. Murthy, Nonperturbative black hole entropy and Kloosterman sums, *J. High Energy Phys.* 1503 (2015) 074, [http://dx.doi.org/10.1007/JHEP03\(2015\)074](http://dx.doi.org/10.1007/JHEP03(2015)074), arXiv:1404.0033 [hep-th].
- [22] S. Murthy, V. Reys, Functional determinants, index theorems, and exact quantum black hole entropy, *J. High Energy Phys.* 1512 (2015) 028, [http://dx.doi.org/10.1007/JHEP12\(2015\)028](http://dx.doi.org/10.1007/JHEP12(2015)028), arXiv:1504.01400 [hep-th].
- [23] R.K. Gupta, Y. Ito, I. Jeon, Supersymmetric localization for BPS black hole entropy: 1-loop partition function from vector multiplets, *J. High Energy Phys.* 1511 (2015) 197, [http://dx.doi.org/10.1007/JHEP11\(2015\)197](http://dx.doi.org/10.1007/JHEP11(2015)197), arXiv:1504.01700 [hep-th].
- [24] S. Murthy, V. Reys, Single-centered black hole microstate degeneracies from instantons in supergravity, *J. High Energy Phys.* 1604 (2016) 052, [http://dx.doi.org/10.1007/JHEP04\(2016\)052](http://dx.doi.org/10.1007/JHEP04(2016)052), arXiv:1512.01553 [hep-th].
- [25] R.M. Wald, Black hole entropy is the Noether charge, *Phys. Rev. D* 48 (1993) 3427, arXiv:gr-qc/9307038.
- [26] J. Nian, X. Zhang, Entanglement entropy of ABJM theory and entropy of topological black hole, *J. High Energy Phys.* 1707 (2017) 096, [http://dx.doi.org/10.1007/JHEP07\(2017\)096](http://dx.doi.org/10.1007/JHEP07(2017)096), arXiv:1705.01896 [hep-th].
- [27] A. Dabholkar, N. Drukker, J. Gomes, Localization in supergravity and quantum AdS₄/CF₃ holography, *J. High Energy Phys.* 1410 (2014) 90, [http://dx.doi.org/10.1007/JHEP10\(2014\)090](http://dx.doi.org/10.1007/JHEP10(2014)090), arXiv:1406.0505 [hep-th].
- [28] A. Cabo-Bizet, V.I. Giraldo-Rivera, L.A. Pando Zayas, Microstate counting of AdS₄ hyperbolic black hole entropy via the topologically twisted index, *J. High Energy Phys.* 1708 (2017) 023, [http://dx.doi.org/10.1007/JHEP08\(2017\)023](http://dx.doi.org/10.1007/JHEP08(2017)023), arXiv:1701.07893 [hep-th].
- [29] F. Benini, K. Hristov, A. Zaffaroni, Black hole microstates in AdS₄ from supersymmetric localization, *J. High Energy Phys.* 1605 (2016) 054, [http://dx.doi.org/10.1007/JHEP05\(2016\)054](http://dx.doi.org/10.1007/JHEP05(2016)054), arXiv:1511.04085 [hep-th].
- [30] F. Benini, A. Zaffaroni, A topologically twisted index for three-dimensional supersymmetric theories, *J. High Energy Phys.* 1507 (2015) 127, [http://dx.doi.org/10.1007/JHEP07\(2015\)127](http://dx.doi.org/10.1007/JHEP07(2015)127), arXiv:1504.03698 [hep-th].
- [31] F. Benini, K. Hristov, A. Zaffaroni, Exact microstate counting for dyonic black holes in AdS₄, *Phys. Lett. B* 771 (2017) 462, <http://dx.doi.org/10.1016/j.physletb.2017.05.076>, arXiv:1608.07294 [hep-th].
- [32] J.T. Liu, L.A. Pando Zayas, V. Rathee, W. Zhao, Toward microstate counting beyond large N in localization and the dual one-loop quantum supergravity, arXiv:1707.04197 [hep-th].
- [33] R. Camporesi, A. Higuchi, Spectral functions and zeta functions in hyperbolic spaces, *J. Math. Phys.* 35 (1994) 4217, <http://dx.doi.org/10.1063/1.530850>.
- [34] R. Camporesi, A. Higuchi, On the eigenfunctions of the Dirac operator on spheres and real hyperbolic spaces, *J. Geom. Phys.* 20 (1996) 1, [http://dx.doi.org/10.1016/0393-0440\(95\)00042-9](http://dx.doi.org/10.1016/0393-0440(95)00042-9), arXiv:gr-qc/9505009.
- [35] W. Siegel, Hidden ghosts, *Phys. Lett. B* 93 (1980) 170, [http://dx.doi.org/10.1016/0370-2693\(80\)90119-7](http://dx.doi.org/10.1016/0370-2693(80)90119-7).
- [36] J. Thierry-Mieg, BRS structure of the antisymmetric tensor gauge theories, *Nucl. Phys. B* 335 (1990) 334, [http://dx.doi.org/10.1016/0550-3213\(90\)90497-2](http://dx.doi.org/10.1016/0550-3213(90)90497-2).
- [37] E.J. Copeland, D.J. Toms, Quantized antisymmetric tensor fields and self-consistent dimensional reduction in higher dimensional space-times, *Nucl. Phys. B* 255 (1985) 201, [http://dx.doi.org/10.1016/0550-3213\(85\)90134-8](http://dx.doi.org/10.1016/0550-3213(85)90134-8).
- [38] S. Bhattacharyya, A. Grassi, M. Marino, A. Sen, A one-loop test of quantum supergravity, *Class. Quantum Gravity* 31 (2014) 015012, <http://dx.doi.org/10.1088/0264-9381/31/1/015012>, arXiv:1210.6057 [hep-th].