

Does habitat specialization shape the evolutionary potential of wild bird populations?

I. Martinossi-Allibert, Joanne Clavel, S. Ducatez, I. Le Viol, C. Teplitsky

▶ **To cite this version:**

I. Martinossi-Allibert, Joanne Clavel, S. Ducatez, I. Le Viol, C. Teplitsky. Does habitat specialization shape the evolutionary potential of wild bird populations?. Journal of Avian Biology, 2017, 48 (8), pp.1158 - 1165. 10.1111/jav.01011 . hal-01611788

HAL Id: hal-01611788

https://hal.sorbonne-universite.fr/hal-01611788

Submitted on 6 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Does habitat specialization shape the evolutionary potential of wild bird populations?

Martinossi-Allibert I. [1,2] imartinossi@gmail.com

Clavel J. [1,3] joanne.clavel@gmail.com

Ducatez S.[4] simon.ducatez@gmail.com

Le Viol I. [1] ileviol@mnhn.fr

Teplitsky C. [1] celine.TEPLITSKY@cefe.cnrs.fr

[1] **Centre pour les Sciences de la conservation, UMR7204 CNRS / MNHN / UPMC**

[2] **Department of Ecology and Genetics, Evolutionary Biology Center, University of Uppsala, Sweden**

[3] **Department of Environmental Science, Policy and Management, University of California, Berkeley, USA**

[4] **School of Biological Sciences, University of Sydney, NSW 2006 Australia**

Running title: Specialization and evolutionary potential.

Type of article: Research paper **Abstract**: 156 words, **Main text**: 4127 words. **References**: 57, **Table count:** 3**, Figures count:** 2**.**

To whom correspondence should be sent:

Ivain Martinossi-Allibert, imartinossi@gmail.com, 01 40 79 57 38

Author contributions: C.T, I.L, J.C designed the project. C.T. and I.M. collected genetic data and performed statistical analyzes. S.D provided specialization data. I.M wrote the first draft of the manuscript and all authors contributed substantially to revisions.

Abstract

Because specialist species evolved in more temporally and spatially homogeneous environments than generalist species, they are supposed to experience less fluctuating selection. For this reason, we expect specialists to show lower overall genetic variation as compared to generalists. We also expect populations from specialist species to be smaller and more fragmented, with lower neutral genetic diversity. We tested these hypotheses by investigating patterns of genetic diversity along a habitat specialization gradient in wild birds, based on estimates of heritability, coefficients of variation of additive genetic variance, and heterozygosity available in the literature. We found no significant effect of habitat specialization on any of the quantitative genetic estimators but generalists had higher heterozygosity. This effect was mainly a consequence of the larger population size of generalists. Our results suggest that evolutionary potential does not differ at the population level between generalist and specialist species, but the trend observed in heterozygosity levels and population sizes may explain their difference in susceptibility to extinction.

Keywords: heritability; additive genetic variance; neutral diversity; individual specialization; niche width

Introduction

Native specialist species are becoming less abundant in disturbed landscape to the benefit of generalist species (Shultz et al. 2005, Le Viol et al. 2012). There is some evidence that climate (Davey et al. 2013) and land use (Devictor et al. 2008a, b) changes are driving this decline of specialization in wild communities, leading to a functional biotic homogenization (Clavel et al. 2011). The higher sensitivity of specialists to global changes is expected to result from their narrower ecological niche, which is associated with smaller and more fragmented populations, more vulnerable to the destruction of their habitat. The persistence of species will ultimately depend upon two mechanisms: range shift or adaptation to environmental changes, through phenotypic plasticity or microevolution (Lynch and Lande 1993). Anthropogenic changes are now considered as the greatest evolutionary force (Palumbi 2001, Steffen et al. 2011) and it therefore becomes crucial to determine how specialist and generalist species differ in their ability to respond to these changes. Assessing whether generalist and specialist species differ in terms of evolutionary potential should give valuable information on their respective ability to defer extinction through evolution.

The two alternative strategies (specialist vs generalist) are generally thought to arise depending on fitness trade-offs between alternate phenotypes across environments (Richmond et al. 2005, Poisot et al. 2011): spatially and temporally variable environments should favour generalist strategies, because generalists can achieve higher average performance than specialists across multiple environments. In turn, stable environments should favour specialists which are able to exploit more efficiently a single resource (see Moran 1992 for theoretical evidence, and Kassen 2002 for experimental evidence). Expressed in terms of selection pressures, heterogeneous environments generate fluctuating selection, while in a homogeneous or stable environment, selection is more likely to be stabilising, or persistently directional (Keightley and Hill 1988). Additionally, generalist strategies may become less

profitable under high levels of local stabilizing selection, because trade-offs for the performance in different niches become accordingly strong (Svardal et al. 2014). Specialists can therefore be expected to evolve under stronger and more consistent selection pressures that could deplete overall genetic variability. In contrast, the exposure of generalists to a variety of environments, and to spatially and temporally fluctuating selection, could help maintain a higher genetic diversity. For these reasons, we expect to find higher evolutionary potential in generalist species than in specialist ones, which could partly explain the highest sensitivity of specialists to environmental changes.

Only a scarce number of empirical studies support this prediction so far. Hägele and Rowell-Rahier (2000) showed in a herbivorous insect genera that host specialist species have a lower heritability (and thus a lower evolutionary potential) of growth rate than host generalists. A review of selection experiments in bacteria by Kassen (2002) also suggests that environmental heterogeneity may be driving both the quantity of genetic variation in a population and the niche width of individuals, resulting in the evolution of generalists and the maintenance of genetic variation in heterogeneous environments. However, the link between specialization and evolutionary potential is largely unknown in wild populations.

Here, we review available estimates of evolutionary potential in relation to habitat specialisation and population size in wild bird populations. As emphasised in the univariate breeder's equation, $R = h^2 S$ (Lande 1979), the predicted evolutionary response (R) depends on the selection gradient (S) and on the evolutionary potential of a trait, measured as its heritability (h^2, the ratio between the amount of genetic additive variance and total phenotypic variance). We thus mainly focused on estimates of heritability and additive genetic variance as they also represent the most widely reported estimates of evolutionary potential for quantitative traits (Postma 2014). We predict higher genetic variance for generalist species (heritability,

hereafter h^2, and the additive genetic coefficient of variation, hereafter CV_A, are the two standardized estimates of genetic variance we used).

We also reviewed estimates of heterozygosity (hereafter Hz), a widely available measure of genetic diversity known to depend on demographic processes such as inbreeding level, drift and population size (Soulé 1976, Frankham et al. 2002). Hz is not a direct estimator of evolutionary potential, as it does not relate to additive genetic variance, but rather a proxy for overall genetic diversity, which is often considered in conservation study as an indicator of population health. More specifically, Hz has been correlated with extinction risks (Evans and Sheldon 2008) and population fitness (Reed and Frankham 2003). On this topic again, little empirical work can be found, but a study carried out on the bark beetle (Kelley et al. 2000) indicated that specialist sister-species indeed had reduced levels of Hz when compared to generalist ones. Simply because of a narrower ecological niche, specialist species have access to a more limited range of habitats than generalist ones. For this reason, populations of specialists may be smaller and more fragmented which is likely to negatively impact genetic diversity by limiting effective population size and gene flow (Whitlock and Barton 1997). We therefore predict that specialists should have lower Hz than generalists, mainly as a result of smaller population size.

Material & Methods

Data collection

Habitat specialization indexes

Assessing specialization by categorizing habitat types has been criticized as too subjective and unrealistic (Bazzaz 1991, Thompson et al. 1998, Fridley et al. 2007). Therefore, we used two different indices based on independent methods: The Species Specialization Index (SSI), as defined by Julliard et al. (2006) and the relative niche width measurement (θ) as introduced in plants by Fridley et al (2007) and in birds by Ducatez et al. (2014). Both indices use different approaches and are therefore not strongly correlated, but rather represent the diversity of methods available to measure habitat specialisation.

The SSI is defined as the coefficient of variation of a species abundance across a set of predefined habitats (Julliard et al. 2004). A high SSI thus implies that a species abundance varies strongly across habitats. Specialist species have higher values of SSI, generalists have lower values. SSI was calculated for 234 species at the European scale, the territory being partitioned into 98 habitat categories defined in the EUNIS database (see Le Viol et al., 2012 for details). Affinity of each species for nesting and foraging in a given habitat is coded by experts on three levels (3: primary habitats; 2: secondary; 1: others). Note that because the coefficient of variation is the standard deviation scaled to the mean, SSI does not depend on population size.

The niche width (Θ) is based on the expectation that a generalist species occurs in a range of habitat categories that vary considerably in species composition, whereas a specialist species is found in habitats that contain a consistent suite of other species. The idea of using the compositional diversity of species that co-occur with a target species to measure the degree of specialization was introduced by Colwell and Futuyma (1971). Specialist species have lower values of θ, generalists have higher values. To avoid a bias due to differences in species range, this index is based on habitat co-occurrence (rather than realized species co-

occurrence), meaning that two species that occupy the same habitat type in different places are considered as co-occurring, even if their range does not overlap, e.g. this avoids island species to be automatically considered as specialists. The niche width was estimated using species co-occurrence data within each of the 82 IUCN habitat subcategories, and considering all extant 9870 species listed by the IUCN (see Ducatez et al., 2014 for details). For a given species, habitat width was measured as $\theta = \gamma / \mu(\alpha)$, where γ is the cumulative number of species in the habitats used by a given species and $\mu(\alpha)$ is the mean habitat species richness calculated over the different habitats inhabited by that species. This specialization index is positively correlated with population size (Pearson's product-moment correlation $\rho = 0.25$, 95 % Confidence Interval [0.044, 0.43], p-value = 0.02, see supplementary material S1 for the correlation of both specialization indexes with population size).

Habitat specialization was chosen over other available specialization indices as it gathers the greatest number of niche axes in one parameter and is consequently the most integrative. Indeed, temperature and food availability depend on habitat, which makes habitat specialization linked with thermic and food specialization (Barnagaud et al. 2012). Our dataset of quantitative estimates of evolutionary potential (i.e. quantitative genetic parameters) includes species that cover a range of 3.1 in SSI (from 2.3 to 5.4) and of 6.4 in niche width (from 1.58 to 8), and the neutral diversity data (i.e. Hz) covers a range of 4.6 in SSI (from 2.3 to 6.9) and of 10.6 in niche width (from 1 to 10.6), when the total range available for SSI is 8, ranging from 1.8 to 9.9 and the total range for niche width is 13.54, ranging from 1 to 14.54. The distributions of both specialization index for each of our datasets can be found in supplementary figure S2.

166 ***Population size***

167 Information on local population size was not available so the population size data we used
168 was extracted from the BirdLife International website in May 2014 (*http://www.birdlife.org/*).
169 Population sizes correspond to the total number of individuals in the world according to the
170 censuses published by BirdLife (2005, 2008).

171

172 ***Quantitative data: heritability and coefficient of additive genetic variance***

173 We chose the bird taxon because it gathers the greatest number of quantitative estimators
174 across all taxa (Postma, 2014). In order to gather estimates of evolutionary potential (h^2 and
175 CV_A) in bird populations, we performed a search in the *Web Of Knowledge* research engine.
176 We used the key-words: Topic= (« bird* ») AND (« heritabil* » OR « genetic* estimate* »)
177 and the search was performed in February 2013. When studies aimed at quantifying the
178 response of quantitative parameters to different environmental conditions (e.g. food quality),
179 only parameters estimated under natural conditions were used. These references were further
180 checked against the list gathered by Postma (2014), in order to maximize the number of
181 references providing estimates of heritability and coefficients of additive genetic variance. h^2
182 and CV_As can be estimated with two different methods: the parent offspring regression and
183 the animal model. The former is the earliest method and has been used since the end of the
184 19th century. It estimates heritability by the phenotypic correlation between parents and
185 offspring (Lynch and Walsh 1998). The latter, the *animal model*, has been applied in ecology
186 for merely more than a decade. The animal model is a mixed model using information about
187 relatedness among individuals to infer the part of phenotypic variance due to additive genetic
188 variance (Henderson 1973, Kruuk 2004). Because parent offspring regression may
189 overestimate heritability (shared environments increase the resemblance between parents and
190 offspring), the method used to estimate quantitative parameters was specified in our dataset
191 and subsequent analyses.

Only life-history and morphological traits (See supplementary material, table S3 for trait classification) were kept, as for behavioural traits the type of traits were extremely heterogeneous and for physiological traits, data were available for only four traits from two studies. We considered here that pooling heritabilities or CV_As from different traits measured in a population is relevant, as we were not looking for an accurate estimate of evolutionary potential, but for an estimate of the general level of additive genetic variance in the population. Both h^2 and CV_A represent standardised measures of additive genetic variance, respectively to the total phenotypic variance and to the mean.

Genetic diversity

We used the heterozygosity (Hz) data published by Evans and Sheldon (2008). They collected Hz values measured at microsatellite loci for wild bird species in the literature and used expected Hz because it is less sensitive to sample size (Nei 1978) and the occurrence of null alleles (Callen et al. 1993). When multiple studies were available for one species, Evans & Sheldon (2008) kept the study with the largest sample size and the highest number of microsatellites loci used to estimate Hz. Thus, while the quantitative genetics estimates are available for several populations in some species, the estimates for Hz are only available for one population per species. We used the Hz values for the 54 species that matched our SSI specialization index and for the 183 species in their dataset that matched our estimation of the niche width (θ).

Phylogeny

To account for the non-independence of species, we used species relatedness extracted from phylogenetic trees. Phylogenetic trees were obtained from the website *www.birdtree.org* that provides subsets of the first complete bird phylogeny (Jetz et al. 2012). Each model was run with 100 alternative trees.

Statistical analyses

To assess the effect of habitat specialization on evolutionary potential (h^2, CV_A) or neutral genetic diversity (Hz) we used linear mixed models. Ideally, models should include as fixed effects population size (log-transformed) and an estimator of habitat specialization (SSI or niche width Θ). However, for h^2 and CV_A, population size was only available for few species, resulting in a too small sample size to run our models. We thus included population size in models considering Hz but not in models considering h^2 and CV_A. In terms of random variables, a phylogeny effect was included in all models to account for species non-independence.

Because of a more complex data structure, analyses of h^2 and CV_A included several additional fixed and random effects compared to models run for heterozygosity. First, we expected systematic differences between certain types of traits (morphological or life history traits, Postma, 2014). Consequently, we added "Type of trait" as a fixed effect in models with two levels, "morphological" and "life-history". Similarly, as the method used to estimate heritabilities (animal model or parent-offspring regression) entails a systematic difference, we added an "Estimation method" fixed effect in models investigating heritability.

Second, h^2 and CV_A estimates are trait and population specific because they are based on observations of some characteristics of individuals sampled from a given population. Therefore it was necessary to include in the models two additional random effects: "population identity" and "trait", with a nested structure. "Trait" was nested within "population identity" and "population identity" within "phylogeny". The MCMCglmm R package (Hadfield 2013) does not model nested effects, but the variables were designed in order to account for this nested structure: each species/population/trait combination was given a unique ID allowing to take into account each level in a meaningful way (phylogeny

(species), population within species, trait within population within species). Some species (e.g. *Fulica atra*) were represented by one estimator for only one population when others (e.g. *Passer domesticus*) were represented by multiple populations, and multiple traits for each population. The random effects "population identity" and "trait" allow taking this lack of balance into account.

Finally, the standard error associated with heritability estimates was included as a random effect to account for measurement error. Unfortunately, this was not possible with CV_A, due to the paucity of errors reported, nor with heterozygosity for which standard errors were not available. The random effects were assumed to follow normal distributions with a mean of 0 and a variance equal to $\sigma_a^2 A$ for the phylogeny **(**where σ_a^2 is the phylogenetic variance and **A** is the phylogenetic relatedness matrix), a variance equal to $\sigma_m^2 M$ for the error effect (where σ_m^2 is the measurement error variance and **M** is a diagonal matrix containing the square standard error of the published estimates; we fixed the variance σ_m^2 to 1 as the sampling variances are known, see Hadfield & Nakagawa 2009 for details) and variances equal to $\sigma^2 I$ for the remaining effects (where σ^2 represents the population variance within species, the trait variance within population within species or the residual variance; **I** is the identity matrix).

These models were run using Bayesian methods as implemented in the MCMCglmm R package. 1 000 000 iterations of each model were run. The first 50 000 iterations were used to initiate the chain, not to estimate parameters, and the thinning interval was set to 10. The amount of iterations was considered sufficient as autocorrelations were low (<0.06) for fixed and random effects.

An inverse-gamma prior for random effects was used (V = 1 and nu = 0.002), in order to minimise the effect of prior information on posterior distribution. To assess prior sensitivity, the same models were run using a parameter expanded prior (V = 1, nu = 1,

alpha.mu = 0, alpha.V = 10000) and a weakly informative prior (V = σ/4, n = 1 where σ is the total variance of the dependent variable). Using different priors did not affect the results (Supplementary Table S5 and S6).

To account for phylogeny uncertainty, each model was run with each of the 100 alternative trees provided by Jetz et al. (2012), and the estimates presented in the result section represent the average over the 100 models.

Results

Available data for quantitative genetics estimates

For the analysis using the SSI index, 520 estimates of h^2 and 150 estimates of CV_A from respectively 64 and 13 studies were gathered, with an average of 3.52 traits per study for h^2 and 6.62 per studies for CV_A. Variance among studies in number of estimates is large, ranging from 1 to 48 for h^2 and from 1 to 14 for CV_A. The number of estimates calculated from parent offspring regression is 363 for h^2 against 157 from "animal model" and 39 for CV_A against 111 from "animal model". Only life-history and morphological traits were kept, with respectively 132 and 388 estimates for h^2, and 43 and 107 for CV_A.

For the analysis using the niche width measurement (θ), a total of 923 h^2 and 190 CV_A estimates from respectively 64 and 20 studies were gathered, with an average of 2.6 traits per study for h^2 and 5.7 per study for CV_A. Variance among studies in the number of estimates was large, ranging from 1 to 48 for h^2 and from 1 to 14 for CV_A. 707 h^2 estimates came from parent-offspring regressions and 216 from "animal models". For CV_A, 52 were estimated from parent offspring regression and 139 from "animal models". Only life-history and morphological traits were kept in the analyses (due to the low number of physiological and behavioural traits), with respectively 176 and 747 estimates for h^2, and 61 and 129 for CV_A.

Effect of specialization on additive genetic variance (h^2, CVA)

Heritability was not predicted by specialization, whether considering the SSI (Table 1, posterior mean = 0.054, 95% CI [-0.022, 0.14], pMCMC = 0.13, Fig 2a) or the niche width (θ, Table 2, posterior mean = -0.018 95% CI [-0.057, 0.023], pMCMC = 0.36, Fig 3a) as indices of habitat specialization. Morphological traits had higher heritability than life-history traits in both data sets (Table1 & Table 2), and estimates from the animal model were lower than estimates from parent-offspring regressions for the SSI data (Table1).

Similarly, CV_A was not predicted by either the SSI (posterior mean = 0.66, 95% CI [-1.3, 2.7], pMCMC = 0.48, Fig 2b) or the niche width (θ) (Table2, posterior mean = -0.26, 95% CI [-1.9, 1.5], pMCMC = 0.71, Fig 3b).

Effect of specialization and population size on genetic diversity (Hz)

Heterozygosity was not predicted by either the SSI (Table 3, posterior mean=0.014, 95% CI [-0.025, 0.053], pMCMC = 0.48) or the niche width (Table 3, posterior mean=0.051, 95% CI [-0.021, 0.12], pMCMC = 0.16), but population size did have a significant effect on heterozygosity (Table 3, posterior mean=0.044, 95% CI [0.023, 0.064], P <0.001, and posterior mean=0.036, 95% CI [0.020, 0.054], P <0.001). Species with larger global population sizes had higher levels of heterozygosity. Because niche width and population size are positively correlated (Pearson's product-moment correlation=0.25, df=88, P=0.020), an effect of niche width on heterozygosity appears when population size is not part of the model (posterior mean=0.071, 95% CI [0.022, 0.12], pMCMC =0.0045). The SSI however was not correlated to population size (Pearson's product-moment correlation=-0.27, df=35, P=0.10)

Discussion

The ability of wild populations to persist in a changing environment depends largely on their evolutionary potential (Hoffmann and Sgrò 2011, Vander Wal et al. 2013). We expected a

316 higher evolutionary potential in generalists as compared to specialists, based on the different
317 selection pressures their ecology suggests they are exposed to. In contrast with this, across 36
318 wild bird populations and using two different specialization indexes, we found no significant
319 effect of specialization on evolutionary potential -as estimated by quantitative genetics
320 estimates, h^2 and CV_A. Specialization did not correlate with genetic diversity measured by Hz
321 either, but we found a significant positive relationship between population size and
322 heterozygosity. In turn, population size correlates positively with niche width (but not with
323 the SSI index), indicating that species with larger niches (generalists) also tend to exhibit
324 more genetic diversity, as a consequence of their larger population sizes. The fact that
325 populations of generalist species are larger than populations of specialist species can be
326 explained by several factors, such as larger available habitat patches, better connectivity
327 between patches, or higher dispersal rates (Kattan 1992, Stireman 2005). In turn, higher
328 genetic diversity in larger populations is in line with theoretical and empirical expectations
329 (Soulé 1976, Frankham 1996). Our results thus suggest that the effect of specialization on
330 genetic diversity results from demographic processes rather than evolutionary processes such
331 as long-term differences in selection pressures. Indeed, we found that specialization had an
332 effect on heterozygosity through population size, but not on quantitative genetic estimates:
333 neutral markers are tightly linked to demography, migration, drift and mutation while
334 selection is expected to shape variance of quantitative traits (Sgrò et al. 2011).

335 The prediction that specialists should have in general lower evolutionary potential than
336 generalists due to stronger and more consistent selection pressures was not confirmed by this
337 study. So far, the empirical support for this hypothesis was limited, to our knowledge, to two
338 local scale study in insects (Hagele & Rowell-Rahier 2000 and Kelley et al. 2000) and to
339 experimental evolution in bacteria, as reviewed by Kassen (2002). Several reasons (outlined

below) may explain the absence of pattern, but in any case, the generality of this result will need to be confirmed (or infirmed) by more empirical studies from wild populations.

First, we know little about the origin of generalism. For example, if a species niche is fully covered by specialized populations, each population exploiting a part of the niche only, the species will be described as generalist. In this case, high genetic diversity is expected among populations, but not necessarily within populations, making the pattern undetectable when genetic parameters are sampled at the population level (as it is the case for h^2 and CV_A). This particular genetic structure may have evolved under large spatial scale environmental variation. The ability of populations to respond to new selection pressures may then depend on the amount of gene flow among populations. To provide further insights in this direction, an informative index would be the F_{ST}, which compares within population genetic variation to among-population genetic variation, a core question in our investigation. Unfortunately, estimating F_{ST} requires heterozygosity measures in multiple populations for each studied species, which would be available only after implementing new field studies.

A second major issue is the definition of "evolutionary potential". Because it was the most readily provided estimate of additive genetic variance in natural populations and because of its central role in the breeder's equation, we used heritability. However, because it is a ratio and because of its univariate definition, heritability can be a misleading estimate of evolutionary potential. Heritability is a ratio of variances and as such its direction of variation can be interpreted in multiple ways, i.e. higher heritability can be the result of either higher additive genetic variance or lower environmental variance. This is of fundamental importance here since environmental variance could be larger for generalist species dealing with more heterogeneous environments than specialist species, hence masking any potential increase in additive genetic variance. A solution around this issue is the use of coefficients of variation of additive genetic variance rather than heritability (Houle 1992, Hansen et al. 2003, 2011,

Hansen and Houle 2008). Unfortunately, CV_A was reported for only 15 species, and often published without associated standard errors. Moreover, 45% of those estimators are likely to be biased (Garcia-Gonzalez et al. 2012). It was thus not possible to reliably assess the effect of specialisation on CV_A but this is an avenue worthwhile pursuing in the future. Another aspect through which heritability may not be a suitable estimate of evolutionary potential is its univariate and trait specific nature as it may not reflect overall evolvability and constraints (e.g. genetic correlations, Sih et al. 2004). Multivariate approaches are therefore be needed to bring further light on these issues (Walsh and Blows 2009, Hansen et al. 2011, Teplitsky et al. 2014).

Third, on a more functional note, the absence of patterns for quantitative genetic estimates in our study may also be explained by a lack of knowledge about the functionality of the traits. Indeed, in the study of Hagele & Rowell-Rahier (2000), growth rate heritability was related to a diet specialization index. As a selection pressure resulting from food availability is expected to directly impact growth rate, the niche parameter used to assess specialization index was thus tightly linked to the trait for which evolutionary potential was measured. In this way, the effect of selection pressure is more likely to be detected when measuring heritability. However, this would not mean overall lower evolutionary potential in specialists, but only for a subset of traits closely associated to the specialization parameter. In our case, with the dataset obtained from the literature, such a clear functional link between traits and habitat specialization is not available: we chose a integrative specialization parameters and investigated the additive genetic variance for all available traits together. However, a population can be submitted to strong selection pressures, for instance on wing size (Hall et al. 2004) or beak shape (Grant, B., Grant 2003), that would reduce genetic variance for these traits, a pattern that would not be detected in our analyses because we pooled genetic variance estimates irrespectively of the traits' function. In the case of reduced

overall evolutionary potential, we would expect a decreased genetic variance in traits closely associated to fitness. We thus tried to restrict the data to heritability estimates related to fitness traits (e.g. clutch size, breeding success) but the sample size was drastically reduced in this case (90 heritabilities for 24 species), which didn't allow us to estimate effects properly.

In conclusion, our study indicates that there is no detectable effect of habitat specialization on evolutionary potential in birds, at least when considering estimates of additive genetic variance currently available in the literature. However, this does not mean that specialist and generalist species stand the same chances when facing environmental changes: our study suggests that species with narrower habitat niches have smaller populations with reduced levels of neutral genetic diversity, which indicate increased extinction risks (Evans and Sheldon 2008). In the close future, new estimates of additive genetic variance for wild populations may be available, such as multivariate ones, allowing for further exploration of the relationship between specialization and evolutionary potential.

Acknowledgements

This project was funded by the French ANR (ANR-12-ADAP-0006). We also thank two anonymous reviewers and Darren Irwin for comments that greatly improved the clarity of our manuscript.

Supporting information

S1 Correlation between both specialization indexes and population size.

S2 Specialization indexes distributions

S3 Trait Classification

S4 Phylogenetic trees

S5 Results with weakly informative priors

S6 Results with parameter expanded priors

S7 References for heritability

S8 References for CV_As

S9 List of species taken from Evans &Sheldon (2007)

S10 Visualization for heritability data

References

Barnagaud, J. Y., Devictor, V., Jiguet, F., Barbet-Massin, M., Viol, I. and Archaux, F. 2012. Relating habitat and climatic niches in birds. - PLoS One 7: e32819.

Bazzaz, F. A. 1991. Habitat Selection in Plants. - Am. Nat. 137: S116–S130.

BirdLife, F. 2005. Birds in Europe: population estimates, trends and conservation status. - Br. Birds 98: 269–271.

Callen, D. F., Thompson, a D., Shen, Y., Phillips, H. a, Richards, R. I., Mulley, J. C. and

Sutherland, G. R. 1993. Incidence and origin of “null” alleles in the (AC)n microsatellite markers. - Am. J. Hum. Genet. 52: 922–927.

Clavel, J., Julliard, R. and Devictor, V. 2011. Worldwide decline of specialist species: Toward a global functional homogenization? - Front. Ecol. Environ. 9: 222–228.

Colwell, R. and Futuyma, D. 1971. On the measurement of niche breadth and overlap.pdf. - Ecology 52: 567–576.

Davey, C., Devictor, V. and Jonzen, N. 2013. Impact of climate change on communities: revealing species’ contribution. - J. Anim. Ecol. 82: 551–561.

Devictor, V., Julliard, R., Couvet, D. and Jiguet, F. 2008a. Birds are tracking climate warming, but not fast enough. - Proc. R. Soc. London B Biol. Sci. 275: 2743–2748.

Devictor, V., Julliard, R., Clavel, J., Jiguet, F., Lee, a and Couvet, D. 2008b. Functional biotic homogenization of bird communities in disturbed landscapes. - Glob. Ecol. Biogeogr. 17: 252–261.

Ducatez, S. D., Ingley, R. T. and SHine, R. 2014. Using species co-occurrence patterns to quantify relative habitat breadth in terrestrial vertebrates. - Ecosphere 5: 1–12.

Evans, S. R. and Sheldon, B. C. 2008. Interspecific patterns of genetic diversity in birds: correlations with extinction risk. - Conserv. Biol. 22: 1016–25.

Frankham, R. 1996. Relationship of genetic variation to population size in wildlife. - Conserv. Biol. 10: 1500–1508.

Frankham, R., Ballou, J. D. and Briscoe, D. A. 2002. Introduction to Conservation Genetics.

Fridley, J. D., Vandermast, D. B., Kuppinger, D. M., Manthey, M. and Peet, R. K. 2007. Co-occurrence based assessment of habitat generalists and specialists: A new approach for the measurement of niche width. - J. Ecol. 95: 707–722.

Garcia-Gonzalez, F., Simmons, L. W., Tomkins, J. L., Kotiaho, J. S. and Evans, J. P. 2012. Comparing evolvabilities: Common errors surrounding the calculation and use of

coefficients of additive genetic variation. - Evolution (N. Y). 66: 2341–2349.

Grant, B., Grant, P. 2003. What Darwin's Finches can Teach Us about the Evolutionary Origin and Regualtion of Biodiversity. - Bioscience 53: 965–975.

Hadfield, A. J. 2013. Package " MCMCglmm ." in press.

Hägele, B. F. and Rowell-Rahier, M. 2000. Choice, performance and heritability of performance of specialist and generalist insect herbivores towards cacalol and seneciphylline, two allelochemicals of Adenostyles alpina (Asteraceae). - J. Evol. Biol. 13: 131–142.

Hall, K. S. S., Ryttman, H., Fransson, T. and Stolt, B. O. 2004. Stabilising selection on wing length in reed warblers Acrocephalus scirpaceus. - J. Avian Biol. 35: 7–12.

Hansen, T. F. and Houle, D. 2008. Measuring and comparing evolvability and constraint in multivariate characters. - J. Evol. Biol. 21: 1201–19.

Hansen, T. F., Armbruster, W., Carlson, M. and Pelabon, C. 2003. Evolvability and genetic constraints in Dalechampia Blossoms: genetic correlations and conditional evolvability. - J. Evol. Biol. 16: 754–766.

Hansen, T. F., Pélabon, C. and Houle, D. 2011. Heritability is not Evolvability. - Evol. Biol. 38: 258–277.

Henderson, C. R. 1973. Sire evaluation and genetic trends. - Proc. Anim. Breed. Genet. Symp. Honour J.L. Lush.: 10–41.

Hoffmann, A. and Sgrò, C. 2011. Climate change and evolutionary adaptation. - Nature 470: 479–485.

Houle, D. 1992. Comparing evolvability and variability of quantitative traits. - Genetics 130: 195–204.

International, B. 2008. State of the World's Birds - indicators for our changing world. - Database

478 Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K. and Mooers, a O. 2012. The global
479 diversity of birds in space and time. - Nature 491: 444–8.
480 Julliard, R., Jiguet, F. and Couvet, D. 2004. Common birds facing global changes: What
481 makes a species at risk? - Glob. Chang. Biol. 10: 148–154.
482 Julliard, R., Clavel, J., Devictor, V., Jiguet, F. and Couvet, D. 2006. Spatial segregation of
483 specialists and generalists in bird communities. - Ecol. Lett. 9: 1237–44.
484 Kassen, R. 2002. The experimental evolution of specialists, generalists, and the maintenance
485 of diversity. - J. Evol. Biol. 15: 173–190.
486 Kattan, G. H. 1992. Rarity and vulnerability: the birds of the cordilheira central of Colombia.
487 - Conserv. Biol. 6: 64–70.
488 Keightley, P. D. and Hill, W. G. 1988. Quantitative genetic variability maintained by
489 mutation-stabilizing selection balance in finite populations. - Genet. Res. 54: 33–43.
490 Kelley, S. T., Farrell, B. D. and Mitton, J. B. 2000. Effects of specialization on genetic
491 differentiation in sister species of bark beetles. Heredity 84: 218 Ð227. - Heredity
492 (Edinb). in press.
493 Kruuk, L. 2004. Estimating genetic parameters in natural populations using the “animal
494 model.” - Philos. Trans. R. Soc. London B Biol. Sci. 359: 873–890.
495 Lande, R. 1979. Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain:
496 Body Size Allometry. - Genet. Anal. 33: 402–416.
497 Lynch, M. and Lande, R. 1993. Evolution and extinction in response to envrionmental
498 change. - In: Biotic interactions and global change. pp. 234–250.
499 Lynch, M. and Walsh, B. 1998. Genetics and analysis of quantitative traits.
500 Moran, N. A. 1992. The evolutionary maintenance of alternative phenotypes. - Am. Nat. 139:
501 971–989.
502 Nei, M. 1978. Estimation of Average Heterozygosity and genetice distance from a small

number of individuals. - Genetics 89: 583–590.

Palumbi, S. R. 2001. Evolution - Humans as the world's greatest evolutionary force. - Science (80-.). 293: 1786–1790.

Poisot, T., Bever, J. D., Nemri, A., Thrall, P. H. and Hochberg, M. E. 2011. A conceptual framework for the evolution of ecological specialization. - Ecol. Lett. 14: 841–851.

Postma, E. 2014. Four decades of estimating heritabilities in wild vertebrate populations: Improved methods, more data, better estimates? - In: Quantitative Genetics in the Wild. pp. 16–33.

Reed, D. H. and Frankham, R. 2003. Correlations between fitness and genetic diversity. - Conserv. Biol. 17: 230–237.

Richmond, C., Breitburg, D. L. and Rose, K. A. 2005. The role of environmental generalist s ecies in ecosystem function. - Ecol. Modell. 188: 279–295.

Sgrò, C. M., Lowe, A. J. and Hoffmann, A. A. 2011. Building evolutionary resilience for conserving biodiversity under climate change: {Conserving} biodiversity under climate change. - Evol. Appl. 4: 326–337.

Shultz, S., Bradbury, R. and Evans, K. 2005. Brain size and resource specialization predict long-term population trends in British birds. - Proc. R. Soc. London B Biol. Sci. 272: 2305–2311.

Sih, A., Bell, A. and Johnson, J. 2004. Behaviroal syndroms: an ecological and evolutionary overview. - Trends Ecol. Evol. 19: 372–378.

Soulé, M. 1976. Allozyme variation, its determinants in space and time. - In: Ayala, F. (ed), Molecular evolution. Sinauer Associates, Sunderland, Massachusetts., pp. 60–77.

Steffen, W., Persson, Å., Deutsch, L. and Zalasiewicz, J. 2011. The Anthropocene: From Global Change to Planetary Stewardship. - Ambio 19: 142–151.

Stireman, J. O. 2005. The evolution of generalization? Parasitoid flies and the perils of

528 inferring host range evolution from phylogenies. - J. Evol. Biol. 18: 325–336.
529 Svardal, H., Rueffler, C. and Doebeli, M. 2014. Organismal complexity and the potential for
530 evolutionary diversification. - Evolution (N. Y). 68: 3248–3259.
531 Teplitsky, C., Robinson, M. R. and Merilä, J. 2014. Evolutionary potential and constraints in
532 wild populations. - In: Quantitative genetics in the wild. pp. 190–208.
533 Thompson, K., Hodgson, J. G. and Gaston, K. J. 1998. Abundance-range size relationships in
534 the herbaceous flora of central England. - J. Ecol. 86: 439–448.
535 Vander Wal, E., Garant, D., Festa-Bianchet, M. and Pelletier, F. 2013. Evolutionary rescue in
536 vertebrates: evidence, applications and uncertainty. - Philos. Trans. R. Soc. Lond. B.
537 Biol. Sci. 368: 20120090.
538 Viol, I. Le, Jiguet, F. and Brotons, L. 2012. More and more generalists: two decades of
539 changes in the European avifauna. - Biol. Lett. 8: 780–782.
540 Walsh, B. and Blows, M. W. 2009. Abundant Genetic Variation + Strong Selection =
541 Multivariate Genetic Constraints: A Geometric View of Adaptation. - Annu. Rev. Ecol.
542 Evol. Syst. 40: 41–59.
543 Whitlock, M. C. and Barton, N. H. 1997. The effective size of subdivided population. -
544 Genetics 146: 427–441.
545
546
547
548
549
550
551

Table1. Effect of Species Specialization Index (SSI) on two quantitative estimates of evolutionary potential: heritability and coefficient of variation of additive genetic variance (CV_A).. Estimates of effects are given by posterior means (Post. Mean) and their associated 95% confidence intervals (CI 95%). pMCMC values are also given for fixed effects. For the effect of type of trait, the reference level is 'life history' as compared to 'morphological'. For the effect of the estimation method, the reference level is 'animal model' as compared to 'parent offspring regression'.

Explanatory var.		**Heritability**			**CV_A**		
	Effect	**Post. mean**	**CI 95%**	**Pc**	**Post.mean**	**CI 95%**	**Pc**
Fixed	**Intercept**	0.0299	-0.261/0.281	0.866	9.81	2.79/16.5	0.013**
	SSI	0.0535	-0.0218/0.139	0.128	0.664	-1.31/2.67	0.479
	Type of trait: morphological	0.192	0.130/0.252	<0.001***	-8.25	-11.3/-5.03	<0.001***
	Estimation method: regression	0.0452	0.00283/0.0871	0.036*	-0.848	3.71/1.96	0.549
Random	**Phylogeny**	0.004	0.000/0.017		4.07	0.000/18.7	
	Population	0.002	0.000/0.013		0.677	0.000/3.39	
	Trait	0.021	0.016/0.029		17.3	8.25/30.7	
	Residual	0.004	0.003/0.006		25.8	19.0/36.0	

Table2 Effect of species niche width Index (θ) on two estimates of genetic variance: heritability and coefficient of variation of additive genetic variance (CV_A). Estimates of effects are given by posterior means (Post. Mean) and their associated 95% confidence intervals (CI 95%). pMCMC values are also given for fixed effects. For the effect of type of trait, the reference level is 'life history' as compared to 'morphological'. For the effect of the estimation method, the reference level is 'animal model' as compared to 'parent offspring regression'.

	Explanatory var.	Heritability			CVa		
	Effect	**Post.mean**	**CI 95%**	**Pc**	**Post.mean**	**CI 95%**	**Pc**
Fixed	**Intercept**	0.34	0.086 / 0.59	0.011	12	4 / 20	0.012
	Θ	-0.018	-0.057 / 0.023	0.36	-0.26	-1.9 / 1.5	0.71
	Type of trait : Morphological	0.21	0.17 / 0.25	<0.001	-7.0	-10 / -4.0	<0.001
	Estimation method : regression	-0.00097	-0.045 / 0.044	0.97	-1.5	-5.4 / 2.2	0.44
Random	**Phylogeny**	0.040	0.012 / 0.074		3.8	0.00013 / 19	
	Population	0.0060	0.0026 / 0.0096		1.7	0.00012 / 8.1	
	Trait	0.0061	0.0020 / 0.011		4.8	0.00014 / 21	
	Residual	0.017	0.014 / 0.021		74	54 / 93	

Table3. Effect of species niche width (θ) and Species Specialization Index (SSI) on heterozygosity (Hz). Estimates of effects are given by posterior means (Post. Mean) and their associated 95% confidence intervals (CI 95%). pMCMC values are also given for fixed effects. The second model includes population size as a fixed effect.

	Explanatory var.	Heterozygosity			Response var.	Heterozygosity		
	Effect	**Post.mean**	**CI 95%**	**Pc**	**Effect**	**Post.mean**	**CI 95%**	**Pc**
Fixed	**Intercept**	0.047	-0.21 / 0.29	0.71	**Intercept**	0.058	-0.28 / 0.39	0.72
	θ	0.051	-0.021 / 0.12	0.16	**SSI**	0.014	-0.025 / 0.053	0.48
	Log Population size	0.044	0.023 / 0.064	<0.001	**Log Population size**	0.036	0.020 / 0.054	<0.001
Random	**Phylogeny**	0.0047	0.00023 / 0.012		**Phylogeny**	0.010	0.00020 / 0.030	
	Residual	0.020	0.014 / 0.028		**Residual**	0.014	0.0044 / 0.024	

569

570

571 **Fig1** Distribution of coefficients of variation of genetic additive variance (CV_A, a), heritabilities (b) and heterozygosity (c) along gradient of

572 habitat specialization (SSI).

573

574 **Fig2** Distribution of coefficients of variation of genetic additive variance (CV_A, a), heritabilities (b) and heterozygosity (c) along gradient of

575 habitat niche width (θ).

576

Fig 1 Distribution of coefficient of additive genetic variance (a), heritability (b), and heterozygosity (c) along a habitat specialization gradient. Each SSi value represents a species which name appears below






Fig 2 Distribution of coefficient of additive genetic variance (a), heritability (b), and heterozygosity (c) along a habitat niche width gradient. Each Θ value represents a species.



583

S1. Pearson's product-moment correlation of both the Species Specialization Index and Niche width with population size



584

585

586

S2. Distribution of specialization indexes, SSI and niche width, for available data.





Table S3. List of traits for heritability and coefficient of additive genetic variance data sorted in "life-history" and "morphology" trait categories.

Life-history (h^2)		
actual egg weight	female first-year recruits	mean chroma (egg colour trait)
actual egg weight adjusted for annual variation	female life span	mean egg size
adult survival probability	female lifetime fledging success	mean egg weight
age at first reproduction	female lifetime reproductive success	mean lightness (egg colour trait)
age at last reproduction	first egg volume	median egg size
anual productivity	hatch date	nr of eggs fledged
arrival date	hatch weight	nr of eggs hatched
breeding time	laying date	number of fledglings
calendar hatching date	laying date elevation	number of nesting attempts per year
clutch size	laying date plasticity	relative hatching date
clutch size adjusted for annual variation	life span	SD chroma (egg colour trait)
delay before breeding	lifetime fledgling production	SD lightness (egg colour trait)
egg breadth	lifetime reproductive success	settling date
egg length	lifetime reproductive success	sex ratio at 6 days old
egg mass	log10-transformed recruiting age	sex ratio at independence
egg shape	longevity	standardized laying date
egg size	male annual fitness	start of wing moult
egg volume	male extra-pair reproductive success	time of laying
elevation of the laying date-temperature reaction norm	male first-year fledging success	total number of eggs laid per year
estimated egg weight	male first-year recruits	total number of young raised to 30 days of age
estimated egg weight adjusted for annual variation	male life span	total number of young raised to 6 days of age
femal annual fitness	male lifetime fledging success	.
female annual fitness	male lifetime reproductive success	.
female extra-pair paternity rate	max chroma (egg colour trait)	.
female first-year fledging success	max lightness (egg colour trait)	.


Morphology (h^2)			
Backcolour	culmen depth	head-length poor condition	PC2-bill (excl. Extra pair youngs)
badge size	culmen length	height of white patch	PC2-body
bill depth	culmen width	male bill depth	PC2-body (excl. Extra pair youngs)
bill depth (excl. Extra pair youngs)	Drostian colour	male bill length	skull length
bill length	fat breeding season	male body condition index	tail length
bill length (excl. Extra pair youngs)	fat winter	male body mass	tail white
bill length . fall-spring	feather growth bar width	male nestling mass	tarsus length
bill length . summer	female bill depth	male tarsus length	tarsus length (excl. Extra pair youngs)
bill length at 2 days	female bill length	male wing length	tarsus length . fall-spring
bill width	female body condition index	mate's forehead patch size	tarsus length . summer
bill width (at a depth of 4mm)	female body mass	muscle breeding season	tarsus length at day 10
bill width (excl. Extra pair youngs)	female nestling mass	muscle insertion length left	tarsus length at day 5
bill width . fall-spring	female tarsus length	muscle insertion length right	total badge size (black throat badge; sexual ornament)
bill width . summer	female wing length	muscle winter	UV/bluevsGreen/Red
body condition index	first primary length	nestling mass	UVvsBlue
body mass	flattened wing chord	nestling tarsus length	visible badge size
body mass . fall-spring	forehead patch size	offspring condition	weight
body mass . summer	forehead patch size old	outer tail length	Weight (excl. Extra pair youngs)
body mass at day 10	forehead patch size young	patch heigth	width of white patch
body mass at day 5	forhead patch size of a female's mate	patch width	Wing
Brownness scores	GreenvsRed	PC1-bill	Wing (excl. Extra pair youngs)
Capcolour	head length	PC1-bill (excl. Extra pair youngs)	wing length
central tail length	head size	PC1-body	wing patch size old
Chestcolour	head-length good condition	PC1-body (excl. Extra pair youngs)	wing patch size young
chick condition	head-length normal condition	PC2-bill	wing projection

Life-history (C_{Va})		
age at first reproduction	egg laying date standardized spring temperature 1	lifetime reproductive success
age at last reproduction	egg laying date standardized spring temperature -1	lifetime reproductive success
annual fitness	egg mass	log10-transformed recruiting age
annual reproductive success	female fitness	longevity
breeding time	female mate choice	male extra-pair reproductive success
clutch size	first egg volume	male fitness
clutch size standardized spring temperature 0	first-year fledging success	mean egg weight
clutch size standardized spring temperature 1	first-year recruits	nr of eggs fledged
clutch size standardized spring temperature -1	laying date	nr of eggs hatched
clutch size standardized spring temperature NA	life span	nr of seasons bred
egg laying date	lifetime fledging success	.
egg laying date standardized spring temperature 0	lifetime fledgling production	.

Morphology (C_{Va})		
adult body mass	forehead patch size	SD lightness (egg colour trait)
adult tarsus length	head-length good condition	skull length
bill depth	head-length normal condition	tail length
bill length	head-length poor condition	tarsus length
bill width	max chroma (egg colour trait)	total badge size (black throat badge; sexual ornament)
body mass	max lightness (egg colour trait)	visible badge size
culmen depth	mean chroma (egg colour trait)	weight
culmen length	mean lightness (egg colour trait)	wing length
culmen width	nestling mass	wing patch size
feather growth bar width	patch heigth	wing projection
first primary length	patch width	
fledgling mass	SD chroma (egg colour trait)	



S4.a **Phylogenetic tree for heritability data**



S4.b **Phylogenetic tree for coefficient of additive genetic variance data**

592

Table S3 Estimators (Post.mean), confidence intervals (CI) and critical probabilities (Pc) for the fixed and random effects of the three mixed models corresponding to each estimator of evolutionary potential: heritability, CVa and Heterozygosity. **For weakly informative priors.**

	Response var.	**Heritability**			**CVa**			**Heterozygosity**		
	Effect	**Post.mean**	**CI 95%**	**pMCMC**	**Post.mean**	**CI 95%**	**Pc**	**Post.mean**	**CI 95%**	**pMCMC**
Fixed	**Intercept**	0.050	-0.210/0.309	0.680	9.21	-1.63/20.1	0.088^{**}	0.657	0.419/0.887	$<10^{-5***}$
	SSI	0.046	-0.023/0.117	0.139	0.740	-2.24/3.82	0.607	-0.003	-0.047/0.041	0.158
	Type of Trait : morphologique	0.198	0.140/0.256	$<10^{-5***}$	-7.77	-11.28/-4.24	2.10^{-5***}			
	Calcul method : regression	0.046	0.005/0.088	0.030^{*}	-1.40	-4.36/1.61	0.355			
Random	**Phylogeny**	0.008	0.001/0.021		13.7	1.01/39.7		0.025	0.000/0.072	
	Population	0.007	0.001/0.014		5.75	1.02/13.53				
	Trait	0.021	0.015/0.027		18.74	8.18/30.12				
	Residual	0.004	0.003/0.006		27.19	19.1/36.2		0.016	0.015/0.031	

Table S4 Estimators (Post.mean), confidence intervals (CI) and critical probabilities (Pc) for the fixed and random effects of the three mixed models corresponding to each estimator of evolutionary potential: heritability, CVa and Heterozygosity. For parameter expanded prior.

	Response var.	Heritability			CVa			Heterozygosity		
	Effect	**Post.mean**	**CI 95%**	**pMCMC**	**Post.mean**	**CI 95%**	**Pc**	**Post.mean**	**CI 95%**	**pMCMC**
Fixed	**Intercept**	0.053	-0.162/0.270	0.579	9.59	-0.194/19.9	0.057.	0.695	0.411/0.980	$<10^{-5***}$
	SSI	0.044	-0.018/0.103	0.136	0.656	--2.05/3.37	0.583	-0.012	-0.078/0.052	0.704
	Type of trait : morphological	0.198	0.142/0.255	$<10^{-5***}$	-7.93	-11.2/-4.49	$<10^{-5***}$			
	Estimation method : regression	0.048	0.007/0.090	0.022^{*}	--1.17	-4.05/1.74	0.426			
Random	**Phylogeny**	0.004	0.000/0.017		14.8	0.000/55.8		0.007	0.000/0.026	
	Population	0.005	0.000/0.013		2.03	0.000/7.81				
	Trait	0.022	0.015/0.028		19.6	9.03/31.6				
	Résidual	0.004	0.002/0.006		26.6	18.8/35.3		0.052	0.030/0.077	

Fig S7. References for heritability data

Akesson, M., Bensch, S., & Hasselquist, D. (2007). Genetic and phenotypic associations in morphological traits : a long term study of great reed warblers Acrocephalus arundinaceus. Journal of Avian Biology, 38, 58–72. doi:10.1111/j.2007.0908-8857.03669.x

Alatalo, R,Lundberg, A, (1989) Heritability and selection on tarsus length in the Pied Flycatcher (Ficedula hypoleuca). Evolution, 40(3), 574-583.

Alatalo, R. V., Gustafsson, L., & Lundberg, A. (1990). Phenotypic Selection on Heritable Size Traits : Environmental Variance and Genetic Response. The American Naturalist, 135(3), 464–471.

Barbraud . C. (2000). Natural selection on body size traits in a long-lived bird, the snow petrel Pagodroma nivea. Journal of Evolutionary Biology (13). 81–88. Retrieved from http://www.jstor.org/stable/2409649

Björklund, M. (1997). Variation in growth in the blue tit (Parus caeruleus). Journal of Evolutionary Biology, 10, 139 – 155.

Boag . P. T. (1983). The Heritability of External Morphology in Darwin's Ground Finches (Geospiza) on Isla Daphne Major . Galapagos. Evolution . 37(5) . 877.

Bonneaud, C., Sinsheimer, J. S., Richard, M., Chastel, O., & Sorci, G. (2009). Mhc polymorphisms fail to explain the heritability of phytohaemagglutinin-induced skin swelling in a wild passerine. Biology letters, 5(6), 784–7. doi:10.1098/rsbl.2009.0435

Boulinier . T. . Sorci . G. . Monnat . J.-Y. . & Danchin . E. (1997). Parent–offspring regression suggests heritable susceptibility to ectoparasites in a natural population of kittiwake. Journal of Evolutionary Biology . 10(1) . 77. Retrieved from http://apps.webo

Brommer, J. E., Merilä, J., Sheldon, B. C., & Gustafsson, L. (2005). Natural selection and genetic variation for reproductive reaction norms in a wild bird population. Evolution; international journal of organic evolution, 59(6), 1362–71.

Brommer . J. E. . & Rattiste . K. (2008). "Hidden" reproductive conflict between mates in a wild bird population. Evolution; international journal of organic evolution . 62(9) . 2326–33. doi:10.1111/j.1558-5646.2008.00451.x

Caro, S. P., Charmantier, A., Lambrechts, M. M., Blondel, J., Balthazart, J., & Williams, T. D. (2009). Local adaptation of timing of reproduction: females are in the driver's seat. Functional Ecology, 23(1), 172–179. doi:10.1111/j.1365-2435.2008.01486.x

Charmantier, A., & Réale, D. (2005). How do misassigned paternities affect the estimation of heritability in the wild ? Molecular Ecology, 14, 2839–2850. doi:10.1111/j.1365-294X.2005.02619.x

Charmantier . A. . Keyser . A. J. . & Promislow . D. E. L. (2007). First evidence for heritable variation in cooperative breeding behaviour. Proceedings. Biological sciences / The Royal Society . 274(1619) . 1757–61. doi:10.1098/rspb.2007.0012

Charmantier, A., Kruuk, L. E. B., & Lambrechts, M. M. (2004). PARASITISM REDUCES THE POTENTIAL FOR EVOLUTION IN A WILD BIRD POPULATION. Evolution, 58(1), 203–206.

Charmantier, A., Perrins, C., McCleery, R. H., & Sheldon, B. C. (2006). Evolutionary response to selection on clutch size in a long-term study of the mute swan. The American naturalist, 167(3), 453–65. doi:10.1086/499378

Cuervo, J.J., Moller, A.P., De Lope, F., (2007), Haematocrit is weakly related to condition in nestling Barn Swallow Hirundo rustica,Ibis,149, 128-134.

Dhondt, A. A. (1982). Heritability of Blue Tit Tarsus Length from Normal and Cross-Fostered Broods. Evolution, 36(2), 418. doi:10.2307/2408064

Duckworth . R. a . & Kruuk . L. E. B. (2009). Evolution of genetic integration between dispersal and colonization ability in a bird. Evolution; international journal of organic evolution . 63(4) . 968–77. doi:10.1111/j.1558-5646.2009.00625.x

Findlay . C. S. . & Cooke . F. (1982). Breeding Synchrony in the Lesser Snow Goose (Anser caerulescens caerulescens). I. Genetic and Environmental Components of Hatch Date Variability and their Effects on Hatch Synchrony. Evolution . 36(2) . 342. doi:10.2307/2

Findlay . C. S. . & Cooke . F. (1983). Genetic and Environmental Components of Clutch Size Variance in a Wild Population of Lesser Snow Geese (Anser caerulescens caerulescens). Evolution . 37(4) . 724–734.

Flux . J. E. C. . & Flux . M. M. (1982). Artificial selection and gene flow in wild starlings . Sturnus vulgaris. Naturwissenschaften . 69(2) . 96–97. doi:10.1007/BF00441236
Freeman-Gallant . C. . & Rothstein . M. (1999). Apparent heritability of parental care in Savannah Sparrows. Auk . 116(4) . 1132–1136. Retrieved from http://apps.webofknowledge.com/full record.do?product=UA&search mode=CitingArticles&qid=41&SID=N2kchoHKHEHl5ja
Frentiu . F. D. . Clegg . S. M. . Blows . M. W. . & Owens . I. P. F. (2007). Large body size in an island-dwelling bird: a microevolutionary analysis. Journal of evolutionary biology . 20(2) . 639–49. doi:10.1111/j.1420-9101.2006.01242.x
Garant, D., Sheldon, B. C., & Gustafsson, L. (2004). CLIMATIC AND TEMPORAL EFFECTS ON THE EXPRESSION OF SECONDARY SEXUAL CHARACTERS : GENETIC AND ENVIRONMENTAL COMPONENTS. Evolution, 58(3), 634–644.
Garant, D., Hadfield, J. D., Kruuk, L. E. B., & Sheldon, B. C. (2008). Stability of genetic variance and covariance for reproductive characters in the face of climate change in a wild bird population. Molecular ecology, 17(1), 179–88. doi:10.1111/j.1365-2
Garant, D., Kruuk, L. E. B., McCleery, R. H., & Sheldon, B. C. (2004). Evolution in a changing environment: a case study with great tit fledging mass. The American naturalist, 164(5), E115–29. doi:10.1086/424764
Garant, D., Kruuk, L. E. B., Wilkin, T. a, McCleery, R. H., & Sheldon, B. C. (2005). Evolution driven by differential dispersal within a wild bird population. Nature, 433(7021), 60–5. doi:10.1038/nature03051
Gibbs . H. L. (1988). Heritability and Selection on Clutch Size in Darwin's Medium Ground Finches (Geospiza fortis). Evolution . 42(4) . 750–762.
Gienapp, P., Postma, E., & Visser, M. E. (2006). WHY BREEDING TIME HAS NOT RESPONDED TO SELECTION FOR EARLIER BREEDING IN A SONGBIRD POPULATION. Evolution, 60(11), 2381–2388.
Gosler, A. G., & Harper, D. G. C. (2000). Assessing the heritability of body condition in birds: a challenge exemplified by the great tit Parus major L. (Aves). Biological Journal of the Linnean Society, 71(1), 103–117.
Grant . P. R. . & Grant . B. R. (1995). Predicting Microevolutionary Responses to Directional Selection on Heritable Variation. Evolution . 49(2) . 241. doi:10.2307/2410334
Griffith, SC Owens, IPF Burke, T. (1999). Environmental determination of a sexually selected trait. Nature, 400(6742), 358–560. doi:10.1038/22536
Gustafsson, L. (1986). Lifetime Reproductive Success and Heritability: Empirical Support for Fisher's Fundamental Theorem. The American Naturalist, 128(5), 761. doi:10.1086/284601
Hadfield, J.D, Nutall,A., Osorio,D., Owens, I.P.F., (2006), Testing the phenotypic gambit: phenotypic, genetic and environmental correlations of colour, European society for evolutionary biology, 20, 549-557.
Hadfield, J.D., Owens,I.P.F., (2006), strong environmental determination of carotenoid-based plumage trait is not mediated by carotenoid availability, european society for evolutionary biology, 19, 1104-1114.
Hasselquist, D., Bensch, S., & Schantz, T. V. (1995). Estimating cuckoldry in birds : the heritability method and DNA fingerprinting give different results. Oikos, 72(2), 173–178.
Hegyi, G., Herényi, M., Wilson, A. J., Garamszegi, L. Z., Rosivall, B., Eens, M., & Török, J. (2010). Breeding experience and the heritability of female mate choice in collared flycatchers. PloS one, 5(11), e13855. doi:10.1371/journal.pone.0013855
Husby, A., Hille, S. M., & Visser, M. E. (2011). Testing mechanisms of Bergmann's rule: phenotypic decline but no genetic change in body size in three passerine bird populations. The American naturalist, 178(2), 202–13. doi:10.1086/660834
Jakober, H., Stauber,W., (2000) Werden die neutöter (Lanius collurio) kleiner?, J.Ornithol, 141, 408-417.
Jensen, H., Steinsland, I., Ringsby, T. H., & Saether, B.-E. (2008). Evolutionary dynamics of a sexual ornament in the house sparrow (Passer domesticus): the role of indirect selection within and between sexes. Evolution;

Jensen, H., Saether, B.-E., Ringsby, T. H., Tufto, J., Griffith, S. C., & Ellegren, H. (2003). Sexual variation in heritability and genetic correlations of morphological traits in house sparrow (Passer domesticus). Journal of Evolutionary Biology, 16(6),
Johnson . S. G. . & Johnston . R. F. (1990). Environmental variation and quantitative genetic parameters in the feral pigeon . Columba livia. Biological Journal of the Linnean Society . 40(4) . 321–332. doi:10.1111/j.1095-8312.1990.tb00542.x
Keller . L. F. . Grant . P. R. . Grant . B. R. . & Petren . K. (2001). Heritability of morphological traits in Darwin ' s Finches : misidenti ® ed paternity and maternal effects. Heredity . 87 . 325–336.
Kim . S.-Y. . Drummond . H. . Torres . R. . & Velando . a. (2011). Evolvability of an avian life history trait declines with father's age. Journal of evolutionary biology . 24(2) . 295–302. doi:10.1111/j.1420-9101.2010.02165.x
Kontiainen, P., Brommer, J.E., Karell, P., Pieieäinen, H., (2007), Heritability, plasticity and canalization of Ural owlegg size in a cyclic environment, Journal of Evolutionary Biology, 21, 88-97.
Kvalnes, T., Ringsby, T.H., Jensen, H., Saether, B.-E., (2013) Correlates of egg size variation in a population of house sparrow Passer domesticus, Oecologia, 171(2), 391-402.
Larsson, K., & Forslund, P. (1992). Genetic and Social Inheritance of Body and Egg Size in the Barnacle Goose (Branta leucopsis). Evolution, 46(1), 235–244.
Larsson . K. . Rattiste . K. . & Lilleleht . V. (1997). Heritability of head size in the common gull Larus canus in relation to environmental conditions during offspring growth. Heredity . 79 . 201–207.
Larsson, K. (1993). Inheritance of body size in the Barnacle Goose under different environmental conditions. Journal of Evolutionary Biology, 6(2), 195–208
Lessells . C. M. . Cooke . F. . & Rockwell . R. F. (1989). Is there a trade-off between egg weight and clutch size in wild Lesser Snow Geese (Anser c . caerulescens)? Journal of Evolutionary Biology . 2 . 457–472.
Lessells, C. M., & Ovenden, G. N. (1989). Heritability of Wing Length and Weight in European Bee-Eaters (Merops apiaster). The Condor, 91(1), 210. doi:10.2307/1368167
Liedvogel, M., Cornwallis, C. K., Sheldon, B.C., (2012) Integrating candidate gene and quantitative genetic approaches to understand variation in timing of breeding in wild tit populations, Journal of evolutionary Biology, 25(5), 813-823
Lifjeld, T., Slagsvold, T., (1990) Manipulations of male parental investment in polygynous pied flycatchers, Ficedula hypoleuca, Behavioral Ecology, 1(1), 48-54.
McCleery, R. H., Pettifor, R. a, Armbruster, P., Meyer, K., Sheldon, B. C.,

Fig S8. List of References for C_{Va} data.

Akesson, M., Bensch, S., Hasselquist, D., Tarka, M., & Hansson, B. (2008). Estimating heritabilities and genetic correlations: comparing the "animal model" with parent-offspring regression using data from a natural population. PloS one, 3(3), e1739. doi:1
Akesson, M., Bensch, S., & Hasselquist, D. (2007). Genetic and phenotypic associations in morphological traits : a long term study of great reed warblers Acrocephalus arundinaceus. Journal of Avian Biology, 38, 58–72. doi:10.1111/j.2007.0908-8857.03669.x
MacColl, A. D. C., & Hatchwell, B. J. (2003). HERITABILITY OF PARENTAL EFFORT IN A PASSERINE BIRD. Evolution, 57(9), 2191. Retrieved from http://apps.webofknowledge.com/full_record.do?product=UA&search_mode=GeneralSearch&qid=3&SID=U2A@flBJ8oI6n889kJf&page
Charmantier, A., Kruuk, L. E. B., & Lambrechts, M. M. (2004). PARASITISM REDUCES THE POTENTIAL FOR EVOLUTION IN A WILD BIRD POPULATION. Evolution, 58(1), 203–206.
Nilsson, J., Akesson, M., & Nilsson, J. F. (2009). Heritability of resting metabolic rate in a wild population of blue tits. Journal of evolutionary biology, 22(9), 1867–74. doi:10.1111/j.1420-9101.2009.01798.x

760 Charmantier, A., Perrins, C., McCleery, R. H., & Sheldon, B. C. (2006). Quantitative genetics of age at
761 reproduction in wild swans: support for antagonistic pleiotropy models of senescence. Proceedings
762 of the National Academy of Sciences of the United States.
763 Charmantier, A., Keyser, A. J., & Promislow, D. E. L. (2007). First evidence for heritable variation in
764 cooperative breeding behaviour. Proceedings. Biological sciences / The Royal Society, 274(1619),
765 1757–61. doi:10.1098/rspb.2007.0012
766 Charmantier, A., Perrins, C., McCleery, R. H., & Sheldon, B. C. (2006). Evolutionary response to
767 selection on clutch size in a long-term study of the mute swan. The American naturalist, 167(3), 453–
768 65. doi:10.1086/499378 Merilä, J., & Sheldon, B. (2000). Lifetime Reproductive Success and
769 Heritability in Nature. The American naturalist, 155(3), 301–310. doi:10.1086/303330
770 Qvarnström, A., Brommer, J. E., & Gustafsson, L. (2006). Testing the genetics underlying the co-
771 evolution of mate choice and ornament in the wild. Nature, 441(7089), 84–6.
772 doi:10.1038/nature04564
773 Garant, D., Sheldon, B. C., & Gustafsson, L. (2004). CLIMATIC AND TEMPORAL EFFECTS ON THE
774 EXPRESSION OF SECONDARY SEXUAL CHARACTERS : GENETIC AND ENVIRONMENTAL COMPONENTS.
775 Evolution, 58(3), 634–644.
776 Tschirren, B., Sendecka, J., Groothuis, T. G. G., Gustafsson, L., & Doligez, B. (2009). Heritable variation
777 in maternal yolk hormone transfer in a wild bird population. The American naturalist, 174(4), 557–64.
778 doi:10.1086/605379
779 Morales, J., Kim, S.-Y., Lobato, E., Merino, S., Tomás, G., Martínez-de la Puente, J., & Moreno, J.
780 (2010). On the heritability of blue-green eggshell coloration. Journal of evolutionary biology, 23(8),
781 1783–91. doi:10.1111/j.1420-9101.2010.02044.x
782 Brommer, J. E., Rattiste, K., & Wilson, a. (2010). The rate of ageing in a longlived bird is not heritable.
783 Heredity, 104(4), 363–70. doi:10.1038/hdy.2009.125
784 Larsson, K., Rattiste, K., & Lilleleht, V. (1997). Heritability of head size in the common gull Larus canus
785 in relation to environmental conditions during offspring growth. Heredity, 79, 201–207.
786 Teplitsky, C., Mills, J. a, Yarrall, J. W., & Merilä, J. (2009). Heritability of fitness components in a wild
787 bird population. Evolution; international journal of organic evolution, 63(3), 716–26.
788 doi:10.1111/j.1558-5646.2008.00581.x
789 Reid, J. M., Arcese, P., Sardell, R. J., & Keller, L. F. (2011). Additive genetic variance, heritability, and
790 inbreeding depression in male extra-pair reproductive success. The American naturalist, 177(2), 177–
791 87.
792 doi:10.1086/657977
793 Husby, A., Hille, S. M., & Visser, M. E. (2011). Testing mechanisms of Bergmann's rule: phenotypic
794 decline but no genetic change in body size in three passerine bird populations. The American
795 naturalist, 178(2), 202–13.
796 doi:10.1086/660834
797 Husby, A., Nussey, D. H., Visser, M. E., Wilson, A. J., Sheldon, B. C., & Kruuk, L. E. B. (2010).
798 Contrasting patterns of phenotypic plasticity in reproductive traits in two great tit (Parus major)
799 populations. Evolution; international journal of organic
800 Garant, D., Hadfield, J. D., Kruuk, L. E. B., & Sheldon, B. C. (2008). Stability of genetic variance and
801 covariance for reproductive characters in the face of climate change in a wild bird population.
802 Molecular ecology, 17(1), 179–88. doi:10.1111/j.1365-2
803 McCleery, R. H., Pettifor, R. a, Armbruster, P., Meyer, K., Sheldon, B. C., & Perrins, C. M. (2004).
804 Components of variance underlying fitness in a natural population of the great tit Parus major. The
805 American naturalist, 164(3), E62–
806 72. doi:10.1086/42266
807 Garant, D., Kruuk, L. E. B., Wilkin, T. a, McCleery, R. H., & Sheldon, B. C. (2005).
808 Evolution driven by differential dispersal within a wild bird population. Nature, 433(7021), 60–5.
809 doi:10.1038/nature03051

Garant, D., Kruuk, L. E. B., McCleery, R. H., & Sheldon, B. C. (2004). Evolution in a changing environment: a case study with great tit fledging mass. The American naturalist, 164(5), E115–29. doi:10.1086/424764

Jensen, H., Steinsland, I., Ringsby, T. H., & Saether, B.-E. (2008). Evolutionary dynamics of a sexual ornament in the house sparrow (Passer domesticus): the role of indirect selection within and between sexes. Evolution; international journal of organic

Jensen, H., Saether, B.-E., Ringsby, T. H., Tufto, J., Griffith, S. C., & Ellegren, H. (2003). Sexual variation in heritability and genetic correlations of morphological traits in house sparrow (Passer domesticus). Journal of Evolutionary Biology, 16(6),

Gienapp, P., & Merilä, J. (2010). Genetic and environmental effects on a condition-dependent trait: feather growth in Siberian jays. Journal of evolutionary biology, 23(4), 715–23. doi:10.1111/j.1420-9101.2010.01949.x

Duckworth, R. a, & Kruuk, L. E. B. (2009). Evolution of genetic integration between dispersal and colonization ability in a bird. Evolution; international journal of organic evolution, 63(4), 968–77. doi:10.1111/j.15585646.2009.00625.x

Kim, S.-Y., Drummond, H., Torres, R., & Velando, a. (2011). Evolvability of an avian life history trait declines with father's age. Journal of evolutionary biology, 24(2), 295–302. doi:10.1111/j.1420-9101.2010.02165.x

Frentiu, F. D., Clegg, S. M., Blows, M. W., & Owens, I. P. F. (2007). Large body size in an island-dwelling bird: a microevolutionary analysis. Journal of evolutionary biology, 20(2), 639–49. doi:10.1111/j.1420-9101.2006.01242.x

Fig S9. List of references from Evans & Sheldon (2008)

Abbott, C. L., and M. C. Double. 2003. Genetic structure, conservation genetics and evidence of speciation by range expansion in shy and white-capped albatrosses. Molecular Ecology 12: 2953-2962.

Adcock, G. J., and R. A. Mulder. 2002. Polymorphic microsatellite loci for paternity analysis in the Madagascar paradise flycatcher (*Terpsiphone mutata* : Aves). Molecular Ecology Notes 2: 287-289.

Adcock, G. J., N. E. Langmore, R. A. Mulder, and R. M. Kilner. 2005. Microsatellite loci for population and behavioural studies of Horsfield's bronzecuckoo (*Chalcites basalis* : Aves). Molecular Ecology Notes 5: 619-621.

Akst, E. P., P. D. Boersma, and R. C. Fleischer. 2002. A comparison of genetic diversity between the Galapagos Penguin and the Magellanic Penguin. Conservation Genetics 3: 375-383.

Asai, S., C. Shimoda, I. Nishiumi, K. Eguchi, and S. Yamagishi. 1999. Isolation of microsatellite loci for paternity testing in the rufous vanga *Schetba rufa*. Molecular Ecology 8: 513-514.

Bardeleben, C. 2004. Isolation of polymorphic tetranucleotide microsatellite markers for the little greenbul (*Andropadus virens*). Molecular Ecology Notes 4: 698-700.

Bardeleben, C., and M. M. Gray. 2005. Isolation of polymorphic tetranucleotide microsatellite markers for the pygmy kingfisher *Ceyx picta*. Molecular Ecology Notes 5: 478-480.

Bardeleben, C., R. L. Moore, J. A. Nicholls, J. J. Austin, and T. E. Robson. 2005. Isolation of polymorphic tetranucleotide microsatellite markers in the satin bowerbird, *Ptilonorhynchus violaceus*. Molecular Ecology Notes 5: 305-307.

Bardeleben, C., M. M. Gray, J. Austin, and I. A. do Rosario. 2005. Isolation of polymorphic tetranucleotide microsatellite for the large-billed scrubwren (*Sericomis magnirostris*). Molecular Ecology Notes 5: 143-145.

Bardeleben, C., K. S. Delaney, J. Austin, and R. L. Moore. 2005. Isolation of polymorphic tetranucleotide microsatellite markers for the grey-headed robin (*Poecilodryas albispecularis*). Molecular Ecology Notes 5: 146-148.

Beck, N., R. Peakall, and R. Heinsohn. 2003. Isolation and characterization of polymorphic microsatellite markers in the white-winged chough (*Corcorax melanorhamphos*). Molecular Ecology Notes 3: 586-588.

Benson, D. P. 2002. Low extra-pair paternity in White-tailed Ptarmigan. Condor 104:192-197.

Blanchard, L., and J. S. Quinn. 2001. The characterization of microsatellite loci in the communally breeding smooth-billed ani (*Crotophaga ani*). Molecular Ecology Notes 1: 152-154.

Bouzat, J. L., H. H. Cheng, H. A. Lewin, R. L. Westemeier, J. D. Brawn, and K. N. Paige. 1998. Genetic evaluation of a demographic bottleneck in the Greater Prairie Chicken. Conservation Biology 12: 836-843.

Bouzat, J. L., and K. Johnson. 2004. Genetic structure among closely spaced leks in a peripheral population of lesser prairie-chickens. Molecular Ecology 13: 499505.

Bulgin, N. L., H. L. Gibbs, P. Vickery, and A. J. Baker. 2003. Ancestral polymorphisms in genetic markers obscure detection of evolutionarily distinct populations in the endangered Florida grasshopper sparrow (*Ammodramus savannarum floridanus*). Molecular Ecology 12: 831-844.

Burg, T. M. 1999. Isolation and characterization of microsatellites in albatrosses. Molecular Ecology 8: 338-341.

Burg, T. M., A. J. Gaston, K. Winker, and V. L. Friesen. 2005. Rapid divergence and postglacial colonization in western North American Steller's jays (*Cyanocitta stelleri*). Molecular Ecology 14: 3745-3755.

Burgess, S. L., and R. C. Fleischer. 2006. Isolation and characterization of polymorphic microsatellite loci in the Hawaiian flycatcher, the elepaio (*Chasiempis sandwichensis*). Molecular Ecology Notes 6: 14-16.

912 Busch, J. D., T. E. Katzner, E. Bragin, and P. Keim. 2005. Tetranucleotide microsatellites for aquila and
913 haliaeetus eagles. Molecular Ecology Notes 5: 3941.
914 Byers, B. E., H. L. Mays, I. R. K. Stewart, and D. F. Westneat. 2004. Extrapair paternity increases
915 variability in male reproductive success in the chestnutsided warbler (*Dendroica pensylvanica*), a
916 socially monogamous songbird. Auk 121: 788-795.
917 Cabe, P. R., and K. E. Marshall. 2001. Microsatellite loci from the house wren (*Troglodytes aedon*).
918 Molecular Ecology Notes 1: 155-156.
919 Caizergues, A., O. Ratti, P. Helle, L. Rotelli, L. Ellison, and J. Y. Rasplus. 2003. Population genetic
920 structure of male black grouse (*Tetrao tetrix* L.) in fragmented vs. continuous landscapes. Molecular
921 Ecology 12: 2297-2305.
922 Caizergues, A., A. Bernard-Laurent, J. F. Brenot, L. Ellison, and J. Y. Rasplus. 2003. Population genetic
923 structure of rock ptarmigan *Lagopus mutus* in Northern and Western Europe. Molecular Ecology 12:
924 2267-2274.
925 Caparroz, R., C. Y. Miyaki, and A. J. Baker. 2003. Characterization of microsatellite loci in the Blue-
926 and-gold Macaw, *Ara ararauna* (Psittaciformes : Aves). Molecular Ecology Notes 3: 441-443.
927 Cathey, J. C., J. A. DeWoody, and L. M. Smith. 1998. Microsatellite markers in Canada geese (*Branta*
928 *canadensis*). Journal of Heredity 89: 173-175.
929 Chbel, F., D. Broderick, Y. Idaghdour, A. Korrida, and P. McCormick. 2002. Characterization of 22
930 microsatellites loci from the endangered Houbara bustard (*Chlamydotis undulata undulata*).
931 Molecular Ecology Notes 2: 484-487.
932 Clegg, S. M., J. F. Kelly, M. Kimura, and T. B. Smith. 2003. Combining genetic markers and stable
933 isotopes to reveal population connectivity and migration patterns in a Neotropical migrant, Wilson's
934 warbler (*Wilsonia pusilla*). Molecular Ecology 12: 819-830.
935 Dasmahapatra, K. K., C. M. Lessells, A. C. Mateman, and W. Amos. 2004. Microsatellite loci in the
936 European bee-eater, *Merops apiaster*. Molecular Ecology Notes 4: 500-502.
937 Dawson, D. A., O. Hanotte, C. Greig, I. R. K. Stewart, and T. Burke. 2000. Polymorphic microsatellites
938 in the blue tit *Parus caeruleus* and their crossspecies utility in 20 songbird families. Molecular Ecology
939 9: 1941-1944.
940 Dawson, D. A., F. M. Hunter, J. Pandhal, R. Buckland, A. Parham, I. L. Jones, M.
941 Bradshaw, R. Jehle, and T. Burke. 2005. Assessment of 17 new whiskered auklet (*Aethia pygmaea*)
942 microsatellite loci in 42 seabirds identifies 5-15 polymorphic markers for each of nine Alcinae species.
943 Molecular Ecology Notes 5: 289-297. Delaney, K. S., and R. K. Wayne. 2005. Adaptive units for
944 conservation:
945 Population distinction and historic extinctions in the Island Scrub-Jay.
946 Conservation Biology 19: 523-533.
947 Denk, A. G., B. Gautschi, K. Carter, and B. Kempenaers. 2004. Seven polymorphic microsatellite loci
948 for paternity assessment in the mallard (*Anas platyrhynchos*). Molecular Ecology Notes 4: 506-508.
949 Doerr, E. D. 2005. Characterization of microsatellite loci in the brown treecreeper (*Climacteris*
950 *picumnus*) and cross-species amplification in the white-throated treecreeper (*Cormobates*
951 *leucophaeus*). Molecular Ecology Notes 5: 654-656.
952 Dowling, D. K., J. Adcock, and R. A. Mulder. 2003. Novel polymorphic microsatellite markers for
953 paternity analysis in the red-capped robin (*Petroica goodenovii* : Aves). Molecular Ecology Notes 3:
954 517-519.
955 Duval, E. H., and K. J. Nutt. 2005. Isolation and characterization of polymorphic microsatellite loci in
956 the lance-tailed manakin (*Chiroxiphia lanceolata*). Molecular Ecology Notes 5: 112-114.
957 Eggert, L. S., and R. C. Fleischer. 2004. Isolation of polymorphic microsatellite loci in the Hawaii
958 amakihi (*Hemignathus virens*) and their use in other honeycreeper species. Molecular Ecology Notes
959 4: 725-727.
960 Estoup, A., and S. M. Clegg. 2003. Bayesian inferences on the recent island colonization history by the
961 bird *Zosterops lateralis lateralis*. Molecular Ecology 12: 657-674.
962 Fields, R. L., and K. T. Scribner. 1997. Isolation and characterization of novel waterfowl microsatellite
963 loci: Cross-species comparisons and research applications. Molecular Ecology 6: 199-202.

964 Francisco, M. R., P. M. Galetti, and L. Gibbs. 2004. Isolation and characterization of microsatellite loci
965 in the blue manakin, *Chiroxiphia caudata* (Aves, Pipridae). Molecular Ecology Notes 4: 758-760.
966 Galbusera, P., M. Githiru, L. Lens, and E. Matthysen. 2004. Genetic equilibrium despite habitat
967 fragmentation in an Afrotropical bird. Molecular Ecology 13: 1409-1421.
968 Gautschi, B., I. Tenzer, J. P. Muller, and B. Schmid. 2000. Isolation and characterization of
969 microsatellite loci in the bearded vulture (*Gypaetus barbatus*) and cross-amplification in three Old
970 World vulture species. Molecular Ecology 9: 2193-2195.
971 Gautschi, B., M. K. Arter, R. E. Husi, W. Wettstein, and B. Schmid. 2002. Isolation and characterization
972 of microsatellite loci in the globally endangered Corncrake, *Crex crex* Linne. Conservation Genetics 3:
973 451-453.
974 Gay, L., P. D. Du Rau, J. Y. Mondain-Monval, and P. A. Crochet. 2004.
975 Phylogeography of a game species: the red-crested pochard (*Netta rufina*) and consequences for its
976 management. Molecular Ecology 13: 1035-1045.
977 Gibbs, H. L., L. M. Tabak, and K. Hobson. 1999. Characterization of microsatellite DNA loci for a
978 neotropical migrant songbird, the Swainson's thrush (*Catharus ustulatus*). Molecular Ecology 8: 1551-
979 1552.
980 Gibbs, H. L., R. J. G. Dawson, and K. A. Hobson. 2000. Limited differentiation in microsatellite DNA
981 variation among northern populations of the yellow warbler: evidence for male-biased gene flow?
982 Molecular Ecology 9: 2137-2147.
983 Goetz, J. E., K. P. McFarland, and C. C. Rimmer. 2003. Multiple paternity and multiple male feeders in
984 Bicknell's Thrush (*Catharus bicknelli*). Auk 120: 10441053.
985 Gonzalez, E. G., A. M. Castilla, and R. Zardoya. 2005. Novel polymorphic microsatellites for the red-
986 legged partridge (*Alectoris rufa*) and cross-species amplification in Alectoris graeca. Molecular
987 Ecology Notes 5: 449-451.
988 Grant, P. R., B. R. Grant, J. A. Markert, L. F. Keller, and K. Petren. 2004. Convergent evolution of
989 Darwin's finches caused by introgressive hybridization and selection. Evolution 58: 1588-1599.
990 Griffith, S. C., I. R. K. Stewart, D. A. Dawson, I. P. F. Owens, and T. Burke. 1999. Contrasting levels of
991 extra-pair paternity in mainland and island populations of the house sparrow (*Passer domesticus*): is
992 there an 'island effect'? Biological Journal of the Linnean Society 68: 303-316.
993 Guay, P. J., and R. A. Mulder. 2005. Isolation and characterization of microsatellite markers in musk
994 duck (*Biziura lobata* : Aves), and their application to other waterfowl species. Molecular Ecology
995 Notes 5: 249-252. Haavie, J., G. P. Saetre, and T. Moum. 2000. Discrepancies in population
996 differentiation at microsatellites, mitochondrial DNA and plumage colour in the pied flycatcher -
997 inferring evolutionary processes. Molecular Ecology 9: 11371148.
998 Hale, A. M., and C. R. Hughes. 2003. Characterization of polymorphic microsatellite loci in a
999 neotropical wood-quail, *Odontophorus leucolaemus*. Molecular Ecology Notes 3: 508-510.
1000 Hamao, S., and D. S. Saito. 2005. Extrapair fertilization in the black-browed reed warbler
1001 (*Acrocephalus bistrigiceps*): Effects on mating status and nesting cycle of cuckolded and cuckolder
1002 males. Auk 122: 1086-1096.
1003 Hammond, E. L., A. J. Lymbery, G. B. Martin, D. Groth, and J. D. Wetherall. 2002. Microsatellite
1004 analysis of genetic diversity in wild and farmed emus (Dromaius *novaehollandiae*). Journal of
1005 Heredity 93: 376-380.
1006 Hanotte, O., C. Zanon, A. Pugh, C. Greig, A. Dixon, and T. Burke. 1994. Isolation and Characterization
1007 of Microsatellite Loci in a Passerine Bird - the Reed Bunting *Emberiza schoeniclus*. Molecular Ecology
1008 3: 529-530.
1009 Hansson, B., H. Westerdahl, D. Hasselquist, M. Akesson, and S. Bensch. 2004. Does linkage
1010 disequilibrium generate heterozygosity-fitness correlations in great reed warblers? Evolution 58:
1011 870-879.
1012 Hasegawa, O., Y. Ishibashi, and S. Abe. 2000. Isolation and characterization of microsatellite loci in
1013 the red-crowned crane *Grus japonensis*. Molecular Ecology 9: 1677-1678.
1014 Hasegawa, O., Y. Ishibashi, and S. Abe. 2005. Polymorphic microsatellite DNA markers for the
1015 rhinoceros auklet (*Cerorhinca monocerata*). Molecular Ecology Notes 5: 637-638.

Hawley, D. M. 2005. Isolation and characterization of eight microsatellite loci from the house finch (*Carpodacus mexicanus*). Molecular Ecology Notes 5: 443445.
Hille, S. M., M. Nesje, and G. Segelbacher. 2003. Genetic structure of kestrel populations and colonization of the Cape Verde archipelago. Molecular Ecology 12: 2145-2151.
Hsu, Y.-C., Severinghaus, L. L., Lin, Y.-S. & Li, S.-H. 2003. Isolation and characterization of microsatellite DNA markers from the Lanyu scops owl (*Otus elegans botelensis*). Molecular Ecology Notes 3: 595-597.
Huang, Y. J., C. S. Chen, and S. H. Li. 2004. Polymorphic tetranucleotide microsatellite loci in the Hwamei (*Garrulax canorus canorus*) (Timaliidae).
Molecular Ecology Notes 4: 170-172.
Hughes, C. R., and D. M. Deloach. 1997. Developing microsatellites when they are rare: Trinucleotide repeat loci in the northern mockingbird *Mimus polyglottos*. Molecular Ecology 6: 1099-1102.
Hughes, C. R., and T. R. Robinson. 2001. Characterization of microsatellite loci developed for song *wrens Cyphorhinus phaeocephalus*. Molecular Ecology Notes 1: 165-167.
Hughes, J. M., P. B. Mather, A. Toon, J. Ma, I. Rowley, and E. Russell. 2003. High levels of extra-group paternity in a population of Australian magpies *Gymnorhina tibicen*: evidence from microsatellite analysis. Molecular Ecology 12: 3441-3450.
Ishibashi, Y., O. Mikami, and S. Abe. 2000. Isolation and characterization of microsatellite loci in the Japanese marsh warbler *Locustella pryeri*. Molecular Ecology 9: 373-375.
Jeffery, K. J., L. F. Keller, P. Arcese, and M. W. Bruford. 2001. The development of microsatellite loci in the song sparrow, *Melospiza melodia* (Aves) and genotyping errors associated with good quality DNA. Molecular Ecology Notes 1: 11-13.
Ji, Y. J., Y. D. Liu, C. Q. Ding, and D. X. Zhang. 2004. Eight polymorphic microsatellite loci for the critically endangered crested ibis, *Nipponia nippon* (Ciconiiformes : Threskiornithidae). Molecular Ecology Notes 4: 615-617.
Johnson, P. C. D., M. K. Fowlie, and W. Amos. 2005. Isolation of microsatellite loci from the common buzzard, *Buteo buteo* (Aves : Accipitridae). Molecular Ecology Notes 5: 208-211.
King, T. L., M. S. Eackles, A. P. Henderson, C. I. Bocetti, D. Currie, and J. M. Wunderle. 2005. Microsatellite DNA markers for delineating population
structure and kinship among the endangered Kirtland's warbler (*Dendroica kirtlandii*). Molecular Ecology Notes 5: 569-571.
Koopman, M. E., N. A. Schable, and T. C. Glenn. 2004. Development and optimization of microsatellite DNA primers for boreal owls (*Aegolius funereus*). Molecular Ecology Notes 4: 376-378.
Kraaijeveld, K., P. J. Carew, T. Billing, G. J. Adcock, and R. A. Mulder. 2004. Extra-pair paternity does not result in differential sexual selection in the mutually ornamented black swan (*Cygnus atratus*). Molecular Ecology 13:
1625-1633.
Kretzmann, M. B., N. Capote, B. Gautschi, J. A. Godoy, J. A. Donazar, and J. J. Negro. 2003. Genetically distinct island populations of the Egyptian vulture (*Neophron percnopterus*). Conservation Genetics 4: 697-706.
Lambert, D. M., T. King, L. D. Shepherd, A. Livingston, S. Anderson, and J. L. Craig. 2005. Serial population bottlenecks and genetic variation: Translocated populations of the New Zealand Saddleback (*Philesturnus carunculatus rufusater*). Conservation Genetics 6: 1-14.
Latch, E. K., E. J. Smith, and O. E. Rhodes. 2002. Isolation and characterization of microsatellite loci in wild and domestic turkeys (*Meleagris gallopavo*). Molecular Ecology Notes 2: 176-178.
Li, S. H., Y. J. Huang, and J. L. Brown. 1997. Isolation of tetranucleotide microsatellites from the Mexican jay *Aphelocoma ultramarina*. Molecular Ecology 6: 499-501.
Lieckfeldt, D., A. Schmidt, and C. Pitra. 2001. Isolation and characterization of microsatellite loci in the great bustard, *Otis tarda*. Molecular Ecology Notes 1: 133-134.
Loyau, A., B. Moureau, M. Richard, P. Christe, P. Heeb, and G. Sorci. 2005. Cross-amplification of polymorphic microsatellites reveals extra-pair paternity and brood parasitism in *Sturnus vulgaris*. Molecular Ecology Notes 5: 135-139.

Maak, S., K. Wimmers, S. Weigend, and K. Neumann. 2003. Isolation and characterization of 18 microsatellites in the Peking duck (*Anas platyrhynchos*) and their application in other waterfowl species. Molecular Ecology Notes 3: 224-227.
MacDougall-Shackleton, E. A., and S. A. MacDougall-Shackleton. 2001. Cultural and genetic evolution in mountain white-crowned sparrows: Song dialects are associated with population structure. Evolution 55: 2568-2575.
Mank, J. E., J. E. Carlson, and M. C. Brittingham. 2004. A century of hybridization: Decreasing genetic distance between American black ducks and mallards. Conservation Genetics 5: 395-403.
Martinez, J. G., T. Burke, D. Dawson, J. J. Soler, M. Soler, and A. P. Moller. 1998. Microsatellite typing reveals mating patterns in the brood parasitic great spotted cuckoo (*Clamator glandarius*). Molecular Ecology 7: 289-297.
Martinez, J. G., J. J. Soler, M. Soler, A. P. Moller, and T. Burke. 1999. Comparative population structure and gene flow of a brood parasite, the great spotted cuckoo (*Clamator glandarius*), and its primary host, the magpie (*Pica pica*). Evolution 53: 269-278.
Martinez-Cruz, B., V. A. David, J. A. Godoy, J. J. Negro, S. J. O'Brien, and W. E. Johnson. 2002. Eighteen polymorphic microsatellite markers for the highly endangered Spanish imperial eagle (*Aquila adalberti*) and related species. Molecular Ecology Notes 2: 323-326.
Maurer, G., M. L. Hale, M. H. Verduijn, and K. Wolff. 2005. Polymorphic microsatellite loci in pheasant coucal (*Centropus phasianinus*). Molecular Ecology Notes 5: 337-339.
McDonald, D. B., and W. K. Potts. 1994. Cooperative Display and Relatedness among Males in a Lek-Mating Bird. Science 266: 1030-1032.
McGuire, H. L., and M. A. F. Noor. 2002. Microsatellite loci for great white herons and great blue herons (Aves, Ardeidae, *Ardea herodias*). Molecular Ecology Notes 2: 170-172.
McInnes, L. M., I. R. Dadour, M. E. Stewart, W. G. F. Ditcham, P. Mawson, and P. B. S. Spencer. 2005. Characterization of polymorphic microsatellite markers for the Carnaby's cockatoo (*Calyptorhynchus latirostris*) and related black cockatoo species. Molecular Ecology Notes 5: 504-506.
McRae, S. B., and W. Amos. 1999. Characterization of hypervariable microsatellites in the cooperatively breeding white-browed sparrow weaver *Plocepasser mahali*. Molecular Ecology 8: 903-904.
McRae, S. B., S. T. Emlen, D. R. Rubenstein, and S. M. Bogdanowicz. 2005. Polymorphic mircosatellite loci in a plural breeder, the grey-capped social weaver *(Pseudonigrita arnaudi*), isolated with an improved enrichment protocol using fragment size-selection. Molecular Ecology Notes 5: 16-20.
Mira, S., C. Billot, T. Guillemaud, L. Palma, and M. L. Cancela. 2002. Isolation and characterization of polymorphic microsatellite markers in Eurasian vulture *Gyps fulvus*. Molecular Ecology Notes 2: 557-558.
Mira, S., K. Wolff, and M. L. Cancela. 2005. Isolation and characterization of microsatellite markers in Bonelli's eagle (*Hieraaetus fasciatus*). Molecular Ecology Notes 5: 493-495.
Muniz, L. S. B., R. H. F. Macedo, and J. Graves. 2003. Isolation and characterization of dinucleotide microsatellite loci in communally breeding Guira cuckoos (Aves : Cuculidae). Molecular Ecology Notes 3: 209-211.
Munoz-Fuentes, V., N. Gyllenstrand, J. J. Negro, A. J. Green, and C. Vila. 2005. Microsatellite markers for two stifftail ducks: the white-headed duck, Oxyura *leucocephala*, and the ruddy duck, *O. jamaicensis*. Molecular Ecology Notes 5: 263-265.
Nesje, M., and K. H. Roed. 2000. Microsatellite DNA markers from the gyrfalcon (*Falco rusticolus*) and their use in other raptor species. Molecular Ecology 9: 1438-1440.
Nesje, M., K. H. Roed, J. T. Lifjeld, P. Lindberg, and O. F. Steen. 2000. Genetic relationships in the peregrine falcon (*Falco peregrinus*) analysed by microsatellite DNA markers. Molecular Ecology 9: 53-60.
Ost, M., E. Vitikainen, P. Waldeck, L. Sundstrom, K. Lindstrom, T. Hollmen, J. C. Franson, and M. Kilpi. 2005. Eider females form non-kin brood-rearing coalitions. Molecular Ecology 14: 3903-3908.
Otsuka, R., I. Nishiumi, and M. Wada. 2003. Characterization of 12 polymorphic microsatellite loci in the Japanese bush warbler *Cettia diphone*. Molecular Ecology Notes 3: 44-46.

Oyler-McCance, S. J., J. St John, S. E. Taylor, A. D. Apa, and T. W. Quinn. 2005. Population genetics of Gunnison sage-grouse: Implications for management. Journal of Wildlife Management 69: 630-637.
Pearce, J. M., S. L. Talbot, B. J. Pierson, M. R. Petersen, K. T. Scribner, D. L. Dickson, and A. Mosbech. 2004. Lack of spatial genetic structure among nesting and wintering King Eiders. Condor 106: 229-240.
Petren, K., P. R. Grant, B. R. Grant, and L. F. Keller. 2005. Comparative landscape genetics and the adaptive radiation of Darwin's finches: the role of peripheral isolation. Molecular Ecology 14: 2943-2957.
Piertney, S. B., A. D. C. MacColl, P. J. Bacon, and J. F. Dallas. 1998. Local genetic structure in red grouse (*Lagopus lagopus scoticus*): evidence from microsatellite DNA markers. Molecular Ecology 7: 1645-1654.
Piertney, S. B., A. Goostrey, J. F. Dallas, and D. N. Carss. 1998. Highly polymorphic microsatellite markers in the great cormorant *Phalacrocorax carbo*. Molecular Ecology 7: 138-140.
Piertney, S. B., L. Shorey, and J. Hoglund. 2002. Characterization of microsatellite DNA markers in the white-bearded manakin (*Manacus manacus*). Molecular Ecology Notes 2:504-505.
Primmer, C. R., A. P. Moller, and H. Ellegren. 1995. Resolving GeneticRelationships with Microsatellite Markers - a Parentage Testing System for the Swallow *Hirundo rustica*. Molecular Ecology 4: 493-498.
Proudfoot, G., R. Honeycutt, and R. D. Slack. 2005. Development and characterization of microsatellite DNA primers for ferruginous pygmy-owls (*Glaucidium brasilianum*). Molecular Ecology Notes 5: 90-92.
Randi, E., Tabarroni, C., Rimondi, V. L. & Sfougaris, A. 2003. Phylogeography of the rock partridge (*Alectoris graeca*). Molecular Ecology 12: 2201-2214.
Rasner, C. A., P. Yeh, L. S. Eggert, K. E. Hunt, D. S. Woodruff, and T. D. Price. 2004. Genetic and morphological evolution following a founder event in the dark-eyed junco, *Junco hyemalis thurberi*. Molecular Ecology 13: 671-681.
Richardson, D. S., F. L. Jury, D. A. Dawson, P. Salgueiro, J. Komdeur, and T. Burke. 2000. Fifty Seychelles warbler (*Acrocephalus sechellensis*) microsatellite loci polymorphic in Sylviidae species and their cross-species amplification in other passerine birds. Molecular Ecology 9: 2226-2231.
Roeder, A. D., R. K. Marshall, A. J. Mitchelson, T. Visagathilagar, P. A. Ritchie, D. R. Love, T. J. Pakai, H. C. McPartlan, N. D. Murray, N. A. Robinson, K. R. Kerry, and D. M. Lambert. 2001. Gene flow on the ice: genetic differentiation among Adelie penguin colonies around Antarctica. Molecular Ecology 10: 1645-1656.
Russello, M., D. Calcagnotto, R. DeSalle, and G. Amato. 2001. Characterization of microsatellite loci in the endangered St. Vincent Parrot, *Amazona guildingii*. Molecular Ecology Notes 1: 162-164.
Sainsbury, J. P., E. S. Macavoy, and G. K. Chambers. 2004. Characterization of microsatellite loci in the Kaka, *Nestor meridionalis*. Molecular Ecology Notes 4: 623-625.
Saito, D., I. Nishiumi, and M. Nakamura. 2001. Characterization of nine polymorphic microsatellite loci from the alpine accentor *Prunella collaris*. Molecular Ecology Notes 1:258-259.
Saito, D. S., T. Saitoh, and I. Nishiumi. 2005. Isolation and characterization of microsatellite markers in Ijima's leaf warbler, *Phylloscopus ijimae* (Aves : Sylviidae). Molecular Ecology Notes 5: 666-668.
Saladin, V., D. Bonfils, T. Binz, and H. Richner. 2003. Isolation and characterization of 16 microsatellite loci in the Great Tit *Parus major*. Molecular Ecology Notes 3: 520-522.
Schable, N. A., B. C. Faircloth, W. E. Palmer, J. P. Carroll, L. W. Burger, L. A. Brennan, C. Hagen, and T. C. Glenn. 2004. Tetranucleotide and dinucleotide microsatellite loci from the northern bobwhite (*Colinus virginianus*). Molecular Ecology Notes 4:415-419.
Schlosser, J. A., T. W. J. Garner, J. M. Dubach, and A. G. McElligott. 2003. Characterization of microsatellite loci in Humboldt penguin (*Spheniscus humboldti*) and cross-amplification in other penguin species. Molecular Ecology Notes 3: 62-64.
Scribner, K. T., M. R. Petersen, R. L. Fields, S. L. Talbot, J. M. Pearce, and R. K. Chesser. 2001. Sex-biased gene flow in spectacled eiders (anatidae): Inferences from molecular markers with contrasting modes of inheritance. Evolution 55: 2105-2115.


Sefc, K. M., R. B. Payne, and M. D. Sorenson. 2001. Characterization of microsatellite loci in village indigobirds *Vidua chalybeata* and cross-species amplification in estrildid and ploceid finches. Molecular Ecology Notes 1: 252254.
Segelbacher, G., R. J. Paxton, G. Steinbruck, P. Trontelj, and I. Storch. 2000. Characterization of microsatellites in capercaillie *Tetrao urogallus* (AVES). Molecular Ecology 9:1934-1935.
Strausberger, B. M., and M. V. Ashley. 2003. Breeding biology of brood parasitic brown-headed cowbirds (*Molothrus ater*) characterized by parent-offspring and sibling-group reconstruction. Auk 120: 433-445.
Szczys, P., C. R. Hughes, and R. V. Kesseli. 2005. Novel microsatellite markers used to determine the population genetic structure of the endangered Roseate Tern*, Sterna dougallii*, in Northwest Atlantic and Western Australia. Conservation Genetics 6: 461-466.
Tarof, S. A., L. M. Ratcliffe, and P. T. Boag. 2001. Polymorphic microsatellite loci for assigning parentage in least flycatchers (*Empidonax minimus*). Molecular Ecology Notes 1:146-148.
Tarr, C. L., S. Conant, and R. C. Fleischer. 1998. Founder events and variation at microsatellite loci in an insular passerine bird, the Laysan finch (*Telespiza cantans*). Molecular Ecology 7: 719-731.
Tarr, C. L., and R. C. Fleischer. 1999. Population boundaries and genetic diversity in the endangered Mariana crow (*Corvus kubaryi*). Molecular Ecology 8: 941-949.
Thuman, K. A., F. Widemo, and S. B. Piertney. 2002. Characterization of polymorphic microsatellite DNA markers in the ruff (*Philomachus pugnax*). Molecular Ecology Notes 2: 276-277.
Tirard, C., F. Helfenstein, and E. Danchin. 2002. Polymorphic microsatellites in the black-legged kittiwake *Rissa tridactyla*. Molecular Ecology Notes 2: 431433.
Tomasulo-Seccomandi, A. M., N. A. Schable, A. L. Bryan, I. L. Brisbin, S. N. Del Lama, and T. C. Glenn. 2003. Development of microsatellite DNA loci from the wood stork (Aves, Ciconiidae, *Mycteria americana*). Molecular Ecology Notes 3:563-566.
Topinka, J. R., and B. May. 2004. Development of polymorphic microsatellite loci in the northern goshawk (*Accipiter gentilis*) and cross-amplification in other raptor species. Conservation Genetics 5: 861-864.
Van Dongen, W. F. D., and R. A. Mulder. 2005. Isolation and characterization of microsatellite markers for paternity assessment in the golden whistler (*Pachycephala pectoralis*: Aves). Molecular Ecology Notes 5: 4-6.
Van Treuren, R., R. Bijlsma, J. M. Tinbergen, D. Heg, and L. Van de Zande. 1999. Genetic analysis of the population structure of socially organized oystercatchers (*Haematopus ostralegus*) using microsatellites. Molecular Ecology 8: 181-187.
Veit, M. L., R. J. Robertson, P. B. Hamel, and V. L. Friesen. 2005. Population genetic structure and dispersal across a fragmented landscape in cerulean warblers (*Dendroica cerulea*). Conservation Genetics 6: 159-174.
Wang, M. T., Y. C. Hsu, C. T. Yao, and S. H. Li. 2005. Isolation and characterization of 12 tetranucleotide repeat microsatellite loci from the greenbacked tit (*Parus monticolus*). Molecular Ecology Notes 5: 439-442.
Watson, C. J. W., A. A. Beheler, and O. E. Rhodes. 2002. Development of hypervariable microsatellite loci for use in eastern phoebes (*Sayornis phoebe*) and related Tyrannids. Molecular Ecology Notes 2: 117-118.
Westneat, D. F., and H. L. Mays. 2005. Tests of spatial and temporal factors influencing extra-pair paternity in red-winged blackbirds. Molecular Ecology 14: 2155-2167.
Williams, D. A., E. C. Berg, A. M. Hale, and C. R. Hughes. 2004. Characterization of microsatellites for parentage studies of white-throated magpie-jays (*Calocitta formosa*) and brown jays (*Cyanocorax morio*). Molecular Ecology Notes 4: 509-511.
Williams, C. L., A. M. Fedynich, D. B. Pence, and O. E. Rhodes. 2005. Evaluation of allozyme and microsatellite variation in Texas and Florida Mottled Ducks. Condor 107: 155-161.


Winker, K., T. C. Glenn, and G. R. Graves. 1999. Dinucleotide microsatellite loci in a migratory wood warbler (Parulidae : *Limnothlypis swainsonii*) and amplification among other songbirds. Molecular Ecology 8: 1553-1556.
Wright, T. F., A. M. Rodriguez, and R. C. Fleischer. 2005. Vocal dialects, sexbiased dispersal, and microsatellite population structure in the parrot *Amazona auropalliata*. Molecular Ecology 14: 1197-1205.
Yeung, C., Y. J. Huang, and S. H. Li. 2004. Development of polymorphic microsatellite markers for the Steere's Liocichla (*Liocichla steerii*). Molecular Ecology Notes 4: 420-422.
Yodogawa, Y., I. Nishiumi, D. Saito, and K. Okanoya. 2003. Characterization of eight polymorphic microsatellite loci from the Bengalese finch (*Lonchura striata* var. *domestica*). Molecular Ecology Notes 3: 183-185.

S10. Data visualisation: heritability of body mass and tarsus length as a function of SSI and Niche width


In this figure we present heritability of tarsus length and body mass, the two most represented traits in our dataset, for families that had more than one species represented. Species belonging to the same family are linked by hand-drawn ellipses.


Heritability of Tarsus length





Heritability of Body mass





1274