



HAL
open science

Latent HIV-1 TAR Regulates 7SK-responsive P-TEFb Target Genes and Targets Cellular Immune Responses in the Absence of Tat

Sebastian Eilebrecht, Bernd-Joachim Benecke, Arndt Benecke

► **To cite this version:**

Sebastian Eilebrecht, Bernd-Joachim Benecke, Arndt Benecke. Latent HIV-1 TAR Regulates 7SK-responsive P-TEFb Target Genes and Targets Cellular Immune Responses in the Absence of Tat. *Genomics, Proteomics and Bioinformatics*, 2017, 15 (5), pp.313-323. 10.1016/j.gpb.2017.05.003 . hal-01617224

HAL Id: hal-01617224

<https://hal.sorbonne-universite.fr/hal-01617224v1>

Submitted on 16 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Accepted Manuscript

ORIGINAL RESEARCH

Latent HIV-1 TAR Regulates 7SK-responsive P-TEFb Target Genes and Targets Cellular Immune Responses in the Absence of Tat

Sebastian Eilebrecht, Bernd-Joachim Benecke, Arndt G. Benecke


PII: S1672-0229(17)30140-7
DOI: <https://doi.org/10.1016/j.gpb.2017.05.003>
Reference: GPB 272

To appear in: *Genomics, Proteomics & Bioinformatics*

Received Date: 4 March 2017
Revised Date: 10 May 2017
Accepted Date: 24 May 2017

Please cite this article as: S. Eilebrecht, B-J. Benecke, A.G. Benecke, Latent HIV-1 TAR Regulates 7SK-responsive P-TEFb Target Genes and Targets Cellular Immune Responses in the Absence of Tat, *Genomics, Proteomics & Bioinformatics* (2017), doi: <https://doi.org/10.1016/j.gpb.2017.05.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Latent HIV-1 TAR Regulates 7SK-responsive P-TEFb Target Genes and Targets Cellular Immune Responses in the Absence of Tat

Sebastian Eilebrecht^{1,2,*a}, Bernd-Joachim Benecke^{2,b}, Arndt G. Benecke^{1,3,*c}

¹CNRS UMR8246, Université Pierre et Marie Curie, Paris 75005, France

²ACSIOMA GmbH, Technologiezentrum Ruhr, Bochum 44799, Germany

³Center for Innate Immunity and Immune Disease, University of Washington School of Medicine, Seattle, Washington 98195, USA

* Corresponding authors.

E-mail: seilebrecht@acsioma.com (Eilebrecht S); arndt.benecke@upmc.fr (Benecke AG).

^a ORCID: 0000-0002-8111-950X.

^b ORCID: 0000-0001-5436-2267.

^c ORCID: 0000-0002-7121-823X.

Running title: *Eilebrecht S et al / HIV-1 TAR Expression Targets Host Cell Immune Response*

Total word count: 3974

Number of figures: 7

Number of supplemental tables: 3

Abstract

The transactivating response element (TAR) structure of the nascent HIV-1 transcript is critically involved in the recruitment of inactive positive transcription elongation factor b (P-TEFb) to the promoter proximal paused RNA polymerase II. The viral transactivator Tat is responsible for subsequent P-TEFb activation in order to start efficient viral transcription elongation. In the absence of the viral transactivator of transcription (Tat), *e.g.*, during latency or in early stages of HIV transcription, TAR mediates an interaction of P-TEFb with its inhibitor hexamethylene bis-acetamide-inducible proteins 1 (HEXIM1), keeping P-TEFb in its inactive form. In this study, we address the function of HIV-1 TAR in the absence of Tat by analyzing consequences of HIV-1 TAR overexpression on host cellular gene expression. An RNA chimera consisting of Epstein-Barr virus-expressed RNA 2 (EBER2) and HIV-1 TAR was developed to assure robust overexpression of TAR in HEK293 cells. The overexpression results in differential expression of more than 800 human genes. A significant proportion of these genes is involved in the suppression of cellular immune responses, including a significant set of 7SK-responsive P-TEFb target genes. Our findings identify a novel role for HIV-1 TAR in the absence of Tat, involving the interference with host cellular immune responses by targeting 7SK RNA-mediated gene expression and P-TEFb inactivation.

Keywords: HIV-1; Latency; TAR; 7SK RNA; Cellular immunity

Introduction

HIV-1 latency is established by an interplay of different mechanisms, which render the HIV provirus transcriptionally silent. A major step for the activation of HIV-1 gene expression is the switch from transcriptional stalling to productive transcription elongation. The HIV-1 core promoter contains binding sites for a number of different transcription factors, such as nuclear factor of activated T cells (NFAT), nuclear factor kappaB (NF- κ B), and specificity protein 1 (SP1) [1–3], which are involved in the regulation of viral transcription. Upon initial transcription initiation, RNA polymerase II (RNAPII) produces a nascent viral transcript, before it is stalled proximal to the transcription start site (TSS) due to its non-processive phosphorylation status as well as the transcription inhibitory activities of 5,6-dichloro-1- β -D-ribofuranosylbenzimidazole (DRB) sensitivity-inducing factor (DSIF) and negative elongation factor (NELF) [4–6]. The nascent transcript forms a stable hairpin structure, the

transactivating response element (TAR), which serves as a binding platform for a number of cellular and viral proteins [7–10].

An essential host cellular factor for the switch towards productive transcription elongation is the positive transcription elongation factor b (P-TEFb), whose active form consists of the cyclin-dependent kinase 9 (CDK9) and one of the cyclins, T1 or T2 [11,12]. Active P-TEFb phosphorylates the carboxy-terminal domain (CTD) of RNAPII, as well as DSIF and NELF, turning DSIF into an elongation-promoting factor and releasing NELF from the stalled RNAPII [13,14]. A large proportion of cellular P-TEFb is sequestered in an inactive complex with the hexamethylene bis-acetamide-inducible proteins 1 or 2 (HEXIM1/2), the COUP-TF interacting protein 2 (CTIP2) and the small nuclear RNA (snRNA) 7SK [15–17]. We have recently shown that the inactive P-TEFb complex is recruited to the HIV-1 core promoter by the high mobility group protein A1 (HMGA1) that interacts with 7SK [18–22]. As soon as sufficient amounts of the viral transactivator of transcription (Tat) are expressed, the protein is able to release 7SK and HEXIM1/2 from the inactive complex, resulting in P-TEFb activation [23,24]. Subsequently, the resulting active Tat/P-TEFb complex is recruited to the promoter proximal paused RNAPII by specific interactions with TAR [13,25], thus initiating the efficient viral transcription elongation reaction. Sedore and co-workers have previously shown that P-TEFb and HEXIM1 bind to TAR in the absence of Tat [26]. In this case, P-TEFb is inactive due to the inhibitory function of the HEXMI1 protein present.

Apart from that, TAR has a number of additional cellular interacting partners, such as the TAR RNA binding protein (TRBP), HMGA1, or the La auto-antigen [7–9]. These interactions point at additional functions of TAR besides HIV-1 transactivation. In this study, we try to examine such functions of HIV-1 TAR, utilizing heterologous fusions with the Epstein-Barr virus-expressed RNA 2 (EBER2) [27]. Similar strategies have previously been proven effective for investigating the HMGA1 regulation by 7SK L2 substructures [18]. Here we show that heterologous TAR expression in the absence of Tat results in significant changes in the expression of host genes involved in immune regulation. These modulating activities can be ascribed to interference with 7SK and P-TEFb mediated gene expression, which would contribute to a better understanding of HIV-1 latency and early reactivation.

Results and discussion

HIV-1 TAR can be efficiently expressed as EBER2 RNA chimera

EBER2 is a non-coding nuclear RNA transcribed by RNAPIII. About 10^6 – 10^7 copies of

EBER2 can be detected in one EBV-infected host cell [27,28], making the *EBER2* promoter highly attractive for overexpression of small RNA structures [18,29,30]. Previous studies have quantified the levels of multiply spliced viral mRNAs during HIV-1 proviral latency in peripheral blood mononuclear cells (PBMCs) from HIV-1-seropositive patients. These mRNAs could amount to about 5×10^3 copies per PBMC [31] and about 1×10^4 copies per CD4⁺ T cell, which represent the major reservoir of latent HIV-1 in the blood and account for about 40% of the PBMCs [32]. During reactivation, Tat induces efficient HIV-1 transcription by a factor of approximately 10–60 [19,33], resulting in up to 6×10^5 RNA molecules per infected cell. Thus, the use of the *EBER2* promoter-driven HIV-1 TAR chimera would confer a robust 100–1000-fold overexpression as compared to the latency state and a 2–20-fold overexpression as compared to the state of reactivation. Such overexpression should be sufficient to efficiently titrate potential binding factors and obtain a maximal downstream effect on gene expression.

We have previously utilized the *EBER2* promoter to overexpress the HMGA1-binding 7SK L2 substructure in HEK293 cell line, resulting in the regulation of the vast majority of cellular HMGA1 target genes [18]. Besides gene-external promoter elements such as an Sp1 site, an activating transcription factor (ATF)-binding site, and the *EBER* TATA box (ETAB), the *EBER2* promoter also consists of gene internal box A and box B elements. Given these elements are essential for promoter activity, an *EBER2* backbone has to be kept for RNA fusion approaches [34] (**Figure 1A**). Moreover, expression of this *EBER2* backbone has to be used as an appropriate negative control in transcriptome analyses in order to identify specific target genes for the RNA structure of interest.

The 172-nt-long *EBER2* RNA folds into a Y-like structure with two stem-loops, named loop a (La) and loop b (Lb), attached to a basal hairpin. The gene-internal promoter elements are located in La, which ends at nt 74, suggesting the exchange of Lb by the RNA of interest. Therefore, based on our positive experiences with the 7SK L2 chimeras [29,34], we generated an *EBER2*-HIV-1 TAR chimeric RNA by replacing *EBER2* Lb with nt 10 to 49 of HIV-1 TAR RNA. RNA secondary structure prediction suggested that both the *EBER2* and the HIV-1 TAR part of the resulting chimeric construct fold similar to each RNA in its wild type context (**Figure 1A**). Importantly, the *EBER2*-Lb construct lacking the *EBER2* Lb structure completely, which was used as the *EBER2* backbone negative control in these studies, folds the same way as in the context of wild type *EBER2* (**Figure 1A**).

We next verified the transcriptional activity of the generated *EBER2* TAR construct by semi-quantitative RT-PCR (**Figure 1B**). We have shown previously that the *EBER2*-Lb

control construct is transcribed at comparable levels as the wild type EBER2 RNA [29]. Also EBER2 RNA chimeras containing the HMGA1-binding 7SK L2 RNA substructure have shown a robust expression in HEK293 cells [18,19]. Using *ACTB* (encoding beta actin) as a control, semi-quantitative measurements of the EBER2 TAR construct revealed no significant differences in transcription rates when compared to the EBER2-Lb construct (Figure 1B). Notably, expression of HIV-1 TAR as an EBER2 chimeric RNA enables TAR expression without the context of the entire HIV-1 genome, ensuring that no HIV-1 Tat can be produced due to the lack of Tat coding sequence.

HIV-1 TAR induced host cellular gene expression changes

In order to analyze the effect of HIV-1 TAR on the host cellular transcriptome in the absence of Tat, we identified HIV-1 TAR target genes in HEK293 cells overexpressing HIV-1 TAR. As mentioned above, major host cellular players involved in HIV gene expression and latency, such as CDK9, 7SK, and HMGA1, have been extensively characterized previously in HEK293 cell line with respect to their regulatory functions [18,29,30,35]. The expression of the EBER2 HIV-1 TAR chimera in HEK293 cells results in the significantly differential regulation of 859 cellular genes ($P < 0.01$), of which expression of 352 (41%) genes is up-regulated and expression of 507 (59%) genes is down-regulated, when compared to HEK293 cells expressing the EBER2-Lb control construct (Figure 2A and Table S1).

Pathway analysis revealed several cellular pathways significantly enriched for these HIV-1 TAR target genes. These include T helper cell differentiation, IL-4 signaling, lipid antigen representation by CD1, inducible T-cell costimulatory (iCOS)-iCOS ligand (iCOSL) signaling in T helper cells, JAK1 and JAK3 in cytokine signaling, low affinity immunoglobulin gamma Fc region receptor II-b (FcγRIIb) signaling in B lymphocytes, IL-15 signaling, and granulocyte-macrophage colony-stimulating factor (GM-CSF) signaling, all of which are involved in cellular immune responses (Figure 2B). A subset of these genes collaboratively suppress the activation of leukocytes and lymphocytes (Figure 2C). The shRNA-mediated knockdown of *FOXP3*, which is significantly down-regulated by HIV-1 TAR expression (\log_2 fold change = -2.94), has been shown to suppress the activity of naive human T-regulatory cells [36]. iCOS, encoded by another HIV-1 TAR suppressed gene, mediates the activation of T lymphocytes [37]. The interleukins IL-4 and IL-5, both of which are down-regulated by HIV-1 TAR, increase the activation of human lymphocytes and eosinophils, respectively [38,39]. The TYRO protein tyrosine kinase-binding protein encoded by *TYROBP*, a HIV-1 TAR-suppressed gene, mediates the activation of natural killer cells together with natural

cytotoxicity triggering receptor 2 (NCR2) [40]. All mentioned genes are down-regulated upon EBER2 HIV-1 TAR overexpression and are part of a regulated gene network, which results in a global repression of immune cell activation and proliferation, as well as an inhibition of active HIV-1 replication (**Figure 3**).

A cellular pathway, which is particularly repressed by EBER2 HIV-1 TAR overexpression at different nodes, is the iCOS-iCOSL signaling pathway in T helper cells (**Figure 4**). iCOS (CD278) is expressed in activated T cells and mediates a signaling cascade via phosphatidylinositol-3-kinase (PI3K), which further downstream leads to the NF- κ B-mediated expression of immune response genes. HIV-1 TAR significantly down-regulates several key players involved in this pathway, including iCOS itself, PI3K, major histocompatibility complex (MHC) class II, but also the NF- κ B inhibitor I κ B (Figure 2C). Taken together, our results suggest a TAR regulatory function involved in the repression of the human immune response in the absence of Tat.

A subset of HIV-1 target genes consists of HMGA1 and P-TEFb targets

HIV-1 TAR has a variety of cellular interacting partners, such as TRBP, P-TEFb, and HMGA1 [7–10], which all are involved in the regulation of HIV-1 TAR target genes. In a previous study, we have identified HMGA1 target genes in HEK293 cells overexpressing a FLAG-tagged version of HMGA1 [40]. Over-expression of HMGA1 results in the significant regulation of 3096 genes ($P < 0.05$), of which expression of 2159 genes is up-regulated and expression of 937 genes is down-regulated. Of these HMGA1 targets, 92 genes are also significantly regulated by overexpression of HIV-1 TAR chimera, as shown here (**Figure 5A**). However, the direction of such regulation is ambiguous, suggesting that interference of HMGA1 function by HIV-1 TAR, as previously demonstrated for the 7SK L2 substructure [18], to the overall TAR-mediated gene expression changes is minor. Binding of P-TEFb to HIV-1 TAR in the absence of Tat can retain P-TEFb in its inactive state [26]. Taking this into account, at least a subset of HIV-1 TAR target genes may be regulated by this HIV-1 TAR-mediated P-TEFb inactivation.

We have previously identified P-TEFb target genes by transcriptome profiling of HEK293 cells expressing a dominant negative, kinase-inactivated point-mutant of Cdk9 [40]. Such overexpression results in the significant regulation of a total of 1776 genes ($P < 0.001$), of which expression of 440 genes is up-regulated and expression of 1336 genes is down-regulated. Out of these P-TEFb targets, 131 genes are also regulated upon EBER2 HIV-1 TAR overexpression (Figure 5A). Strikingly, a majority of 114 genes of this common subset

are concomitantly regulated upon HIV-1 TAR expression and dnCdk9 expression (Figure 5B) (hypergeometric distribution; $P = 1.197e^{-28}$). Moreover, expression of 110 of these concomitantly-regulated genes is down-regulated in both conditions, which is in line with a HIV-1 TAR-mediated inhibition of P-TEFb.

There exists a gene-specific cooperativity of HMGA1 and P-TEFb in gene expression regulation [40]. HIV-1 TAR overexpression also targets a subset of these genes, represented as the common subset of all three conditions, EBER2 HIV-1 TAR overexpression, dnCdk9 expression, and HMGA1 overexpression (Figure 5A and 5C). 13 out of 16 genes of this common subset are concomitantly regulated when comparing the three conditions, with expression of these genes down-regulated upon HIV-1 TAR overexpression and dnCdk9 expression and up-regulated upon HMGA1-FLAG overexpression (Figure 5C). These 13 genes may thus be regulated by a HMGA1-mediated recruitment of P-TEFb and represent genes that are likely regulated by HIV-1 TAR in two ways: the TAR-mediated titration of 7SK RNA-free HMGA1, and the TAR-mediated inactivation of P-TEFb [18,30,40].

The TAR-regulated P-TEFb target genes are responsive to 7SK

Cellular P-TEFb activity is tightly controlled by 7SK and HEXIM1, which repress Cdk9 kinase activity in a small nuclear ribonucleo protein particle (snRNP) [15,16]. Thereby, 7SK RNA acts as a scaffold, bringing Cdk9 and Cyclin T1/T2 into close proximity of the HEXIM1 protein, which then mediates the repression of Cdk9 kinase activity. As both P-TEFb and HEXIM1 have been shown to bind HIV-1 TAR also in the absence of Tat, TAR may in this case act similar to 7SK, mediating an interaction of P-TEFb with its inhibitor HEXIM1 [26]. Thus, we compared the HIV-1 TAR target genes and the dnCdk9-regulated genes with 7SK-responsive genes. The 7SK-responsive genes were defined as genes whose expression in HEK293 cells is concordantly significantly ($P < 0.01$) regulated upon 7SK knockdown or overexpression (Figure 6A and 6B). As a result, we found that 109 of the EBER2 HIV-1 TAR-regulated genes are also responsive to 7SK RNA (Figure 6A) (hypergeometric distribution; $P = 1.040e^{-67}$), while 251 of the dnCdk9 target genes are also regulated by 7SK RNA. The common subset of all three conditions contains 88 genes (67%) of the dnCdk9-regulated HIV-1 TAR target genes. Furthermore, the common subset of HIV-1 TAR and 7SK target genes shows significant correlations in differential expression when compared to either 7SK knockdown ($R = -0.53$) or overexpression ($R = 0.65$) (Figure 6B and C).

To test whether the total set of HIV-1 TAR-regulated P-TEFb target genes are also regulated by the 7SK-mediated inactivation of P-TEFb, we monitored behavior of these genes

in transcriptome profiles upon 7SK RNA knockdown or overexpression [40] (Figure 6D, E, and F). Out of 110 TAR-responsive P-TEFb target genes, whose expression is down-regulated upon dnCdk9 expression, expression of 108 genes (98%) are up-regulated upon 7SK knockdown (Figure 6E) and concomitantly down-regulated upon 7SK overexpression (Figure 6F). These data reveal that HIV-1 TAR indeed regulates a set of classic P-TEFb target genes, most likely by repressing Cdk9 kinase activity via HEXIM1 and by interfering with cellular functions of 7SK RNA.

Conclusions

Our study globally identifies cellular genes, which probably are highly relevant during HIV transcriptional latency in the absence of Tat. Expression of these genes is affected by HIV-1 TAR. Previous overexpression studies of HIV-1 TAR have demonstrated a significant reduction of HIV-1 replication [41,42]. However, these studies have not investigated the effects of HIV-1 TAR expression on the host cellular gene expression. Here, by expressing a HIV-1 TAR EBER2 chimera in HEK293 cells, we observe the TAR-induced regulation of host genes involved in HIV-1 replication, which leads to a predicted significant down-regulation of HIV-1 replication (see Figure 4). Thus, the HIV-1 TAR-mediated gene regulation even in the absence of Tat may be responsible for the previously-observed reduction in HIV-1 replication. This phenomenon may be of importance for the maintenance of viral latency.

HIV-1 TAR has been shown to have various cellular binding partners and the regulation of the activity or availability of these binding partners by the overexpressed HIV-1 TAR may be the critical mechanism by which this gene expression regulation takes place. Indeed, our results suggest a TAR-mediated inactivation of P-TEFb as one potential route for the regulation of this subset of the HIV-1 TAR-responsive genes, a function that is endogenously fulfilled by 7SK RNA (**Figure 7**). A large proportion of these genes is implicated in central pathways involved in cellular immune responses, suggesting that HIV-1 TAR plays a role during host immune response modulation in the absence of Tat, and thus opening a path to reactivation after HIV-1 latency.

A significant set of HIV-1 TAR target genes identified here could not be assigned directly to known cellular interaction partners. For instance, TRBP has recently been shown to interact with Dicer and to contribute in that way to microRNA-mediated gene silencing [43]. Moreover, HIV-1 TAR has been suggested to sequester TRBP, resulting in an inhibition of

Dicer-RISC activity [44]. Thus, a part of the HIV-1 TAR-regulated gene set may show altered expression due to an impaired Dicer activity, thus leading to significant consequences on the downstream regulatory activity. Also HIV-1 TAR itself has been reported to be processed to virus-derived microRNAs in a Tat-dependent or -independent manner [45,46], which can alter cellular gene expression to protect against apoptosis [47] and inhibit viral replication [48]. However, as our EBER2 HIV-1 TAR chimera does not contain the full sequence encoding these microRNAs, and our approach does not include the expression of Tat, the observed gene regulatory effects cannot be a consequence of viral microRNA production. Thus, our study rather points at further, yet unidentified, specific interacting partners. Importantly, the observed significant effects of HIV-1 TAR on cellular 7SK and P-TEFb-mediated gene expression of cellular innate immune responses provide critical information on the mechanisms underlying HIV-1 latency and reactivation.

Materials and methods

Constructs

The EBER2 gene containing the endogenous sequence from -171 nt to +226 nt, including the complete viral promoter sequence, was inserted into the EcoRI and HindIII restriction sites of pUC18 vector (Fermentas) [29]. The resulting plasmid was then used to construct the EBER2 HIV-1 TAR RNA chimera. To obtain the EBER2 HIV-1 TAR construct, +81 nt to +140 nt sequence of wild type EBER2 RNA were replaced with +10 nt to +49 nt sequence of HIV-1 TAR RNA. To obtain the EBER2-Lb construct, the sequence from +80 nt to +145 nt of wild type EBER2 RNA was deleted as described previously [29]. Constructs used for 7SK knockdown and overexpression as well as for dnCdk9 expression have been described previously [30].

RNA secondary structure prediction

The secondary structure of RNA molecules was calculated using RNAstructure software (Version 4.4) [49].

Cell culture

HEK293 cells were cultured as described previously [18,29]. Transfections were performed using FugeneHD (Roche) according to the manufacturer's instructions. Total RNA from

transfected cells was extracted using the RNeasy Midi Kit (Qiagen), as recommended by the manufacturer.

RT-PCR

RT-PCR analyses were conducted using SuperScript™ 2 Reverse Transcriptase (Invitrogen), Lightcycler (Roche), and Fast Start DNA MasterPLUS SYBR Green I reaction mix (Roche) as recommended by the manufacturer. Primers used to quantify the expression of reference gene β -actin are 5'-TCT TCC AGC CTT CCT TCC TG-3' (sense) and 5'-CAC GGA GTA CTT GCG CTC AG-3' (antisense). Primers used for the quantification of the EBER2 chimeras are 5'-CCG TTG CCC TAG TGG TTT C-3' (sense) and 5'-ACA GCG GAC AAG CCG AAT A-3'.

Transcriptome analyses

For microarray analyses, RNA amplification, labeling, hybridization, and detection were performed following the protocols supplied by Applied Biosystems using the corresponding kits (Applied Biosystems, Prod Nos: 4339628 and 4336875). The resulting data were further analyzed as described previously [18,34]. Briefly, the raw data were quality controlled [50], NeoNORM normalized using $k = 0.2$ [51], and then analyzed as outlined previously [52–54]. For each condition, three biological replicates were merged and compared to the merged signals of three biological replicates of the corresponding control condition. Canonical pathway enrichment studies were performed using Ingenuity Pathway Analysis® software (Ingenuity® Systems, Qiagen) as recommended by the manufacturer. Transcriptome data, annotated to the minimum information about a microarray experiment (MIAME) I+II standards, were deposited in the public database MACE (<http://mace.ihes.fr>) using accession numbers 2529043676 (EBER2 TAR expression), 2791359734 (dnCdk9 expression), 2655275478 (7SK overexpression), 2784969974 (7SK knockdown) and 3189818614 (HMGA1-FLAG overexpression). These data will also be made available soon through a dedicated database at 7SK.acsioma.com.

Authors' contributions

SE has performed the experiments, analyzed and interpreted the data, and written the manuscript. BJB has designed the study and interpreted the data. AGB has designed the study,

analyzed and interpreted the data, and contributed to manuscript writing. All authors read and approved the final manuscript.

Competing interests

All authors have declared to have no competing interests.

Acknowledgments

SE received a post-doctoral fellowship of the Agence Nationale de Recherches sur le SIDA et les hépatites virales (ANRS), France. This work was supported by funds from the Genopole Evry, the Institut des Hautes Études Scientifiques, and the Centre National de la Recherche Scientifique (CNRS), France, awarded to AGB.

References

- [1] Lu YC, Touzjian N, Stenzel M, Dorfman T, Sodroski JG, Haseltine WA. Identification of *cis*-acting repressive sequences within the negative regulatory element of human immunodeficiency virus type 1. *J Virol* 1990;64:5226–9.
- [2] Kretzschmar M, Meisterernst M, Scheidereit C, Li G, Roeder RG. Transcriptional regulation of the HIV-1 promoter by NF-kappaB *in vitro*. *Genes Dev* 1992;6:761–74.
- [3] Majello B, De Luca P, Hagen G, Suske G, Lania L. Different members of the Sp1 multigene family exert opposite transcriptional regulation of the long terminal repeat of HIV-1. *Nucleic Acids Res* 1994;22:4914–21.
- [4] Heidemann M, Hintermair C, Voß K, Eick D. Dynamic phosphorylation patterns of RNA polymerase II CTD during transcription. *Biochim Biophys Acta* 2013;1829:55–62.
- [5] Yamaguchi Y, Inukai N, Narita T, Wada T, Handa H. Evidence that negative elongation factor represses transcription elongation through binding to a DRB sensitivity-inducing factor/RNA polymerase II complex and RNA. *Mol Cell Biol* 2002;22:2918–27.
- [6] Yamaguchi Y, Shibata H, Handa H. Transcription elongation factors DSIF and NELF: promoter-proximal pausing and beyond. *Biochim Biophys Acta* 2013;1829:98–104.
- [7] Gatignol A, Buckler-White A, Berkhout B, Jeang KT. Characterization of a human TAR RNA-binding protein that activates the HIV-1 LTR. *Science* 1991;251:1597–600.
- [8] Chang YN, Kenan DJ, Keene JD, Gatignol A, Jeang KT. Direct interactions between autoantigen La and human immunodeficiency virus leader RNA. *J Virol* 1994;68:7008–20.
- [9] Eilebrecht S, Wilhelm E, Benecke BJ, Bell B, Benecke AG. HMGA1 directly interacts with TAR to modulate basal and Tat-dependent HIV transcription. *RNA Biol* 2013;10:436–44.
- [10] Weeks KM, Ampe C, Schultz SC, Steitz TA, Crothers DM. Fragments of the HIV-1 Tat protein specifically bind TAR RNA. *Science* 1990;249:1281–5.

- [11] Yang X, Gold MO, Tang DN, Lewis DE, Aguilar-Cordova E, Rice AP, et al. TAK, an HIV Tat-associated kinase, is a member of the cyclin-dependent family of protein kinases and is induced by activation of peripheral blood lymphocytes and differentiation of promonocytic cell lines. *Proc Natl Acad Sci U S A* 1997;94:12331–6.
- [12] Papanicolas NF, Durvale MC, Canduri F. The emerging picture of CDK9/P-TEFb: more than 20 years of advances since PITALRE. *Mol Biosyst* 2017;13:246–76.
- [13] Wie P, Garber ME, Fang SM, Fischer WH, Jones KA. A novel CDK9-associated C-type cyclin interacts directly with HIV-1 Tat and mediates its high-affinity, loop-specific binding to TAR RNA. *Cell* 1998;92:451–62.
- [14] Fujinaga K, Irwin D, Huang Y, Taube R, Kurosu T, Peterlin BM. Dynamics of human immunodeficiency virus transcription: P-TEFb phosphorylates RD and dissociates negative effectors from the transactivation response element. *Mol Cell Biol* 2004;24:787–95.
- [15] Yang Z, Zhu Q, Luo K, Zhou Q. The 7SK small nuclear RNA inhibits the CDK9/cyclin T1 kinase to control transcription. *Nature* 2001;414:317–22.
- [16] Nguyen VT, Kiss T, Michels AA, Bensaude O. 7SK small nuclear RNA binds to and inhibits the activity of CDK9/cyclin T complexes. *Nature* 2001;414:322–5.
- [17] Cherrier T, Le Douce V, Eilebrecht S, Riclet R, Marban C, Dequiedt F, et al. CTIP2 is a negative regulator of P-TEFb. *Proc Natl Acad Sci U S A* 2013;110:12655–60.
- [18] Eilebrecht S, Brysbaert G, Wegert T, Urlaub H, Benecke BJ, Benecke A. 7SK small nuclear RNA directly affects HMGA1 function in transcription regulation. *Nucleic Acids Res* 2011;39:2057–72.
- [19] Eilebrecht S, Le Douce V, Riclet R, Targat B, Hallay H, Van Driessche B, et al. HMGA1 recruits CTIP2-repressed P-TEFb to the HIV-1 and cellular target promoters. *Nucleic Acids Res* 2014;42:4962–71.
- [20] Eilebrecht S, Schwartz C, Rohr O. Non-coding RNAs: novel players in chromatin-regulation during viral latency. *Curr Opin Virol* 2013;3:387–93.
- [21] Benecke AG, Eilebrecht S. RNA-mediated regulation of HMGA1 function. *Biomolecules* 2015;5:943–57.
- [22] Le Douce V, Forouzanfar F, Eilebrecht S, Van Driessche B, Ait-Ammar A, Verdikt R, et al. HIC1 controls cellular- and HIV-1- gene transcription via interactions with CTIP2 and HMGA1. *Sci Rep* 2016;11:34920.
- [23] Muniz L, Egloff S, Ughy B, Jady BE, Kiss T. Controlling cellular P-TEFb activity by the HIV-1 transcriptional transactivator Tat. *PLoS Pathog* 2010;6:e1001152.
- [24] Barboric M, Yik JH, Czudnochowski N, Yang Z, Chen R, Contreras X, et al. Tat competes with HEXIM1 to increase the active pool of P-TEFb for HIV-1 transcription. *Nucleic Acids Res* 2007;35:2003–12.
- [25] Garber ME, Wie P, Jones KA. HIV-1 Tat interacts with cyclin T1 to direct the P-TEFb CTD kinase complex to TAR RNA. *Cold Spring Harb Symp Quant Biol* 1998;63:371–80.
- [26] Sedore SC, Byers SA, Biglione S, Price JP, Maury WJ, Price DH. Manipulation of P-TEFb control machinery by HIV: recruitment of P-TEFb from the large form by Tat and binding of HEXIM1 to TAR. *Nucleic Acids Res* 2007;35:4347–58.

- [27] Dümpelmann E, Mittendorf H, Benecke BJ. Efficient transcription of the EBER2 gene depends on the structural integrity of the RNA. *RNA* 2003;9:432–42.
- [28] Lerner MR, Andrews NC, Miller G, Steitz JA. Two small RNAs encoded by Epstein-Barr virus and complexed with protein are precipitated by antibodies from patients with systemic lupus erythematosus. *Proc Natl Acad Sci U S A* 1981;78:805–9.
- [29] Eilebrecht S, Pellay FX, Odenwälder P, Brysbaert G, Benecke BJ, Benecke A. EBER2 RNA-induced transcriptome changes identify cellular processes likely targeted during Epstein Barr Virus infection. *BMC Res Notes* 2008;1:100.
- [30] Eilebrecht S, Bécavin C, Léger H, Benecke BJ, Benecke A. HMGA1-dependent and independent 7SK RNA gene regulatory activity. *RNA Biol* 2011;8:143–57.
- [31] Seshamma T, Bagasra O, Trono D, Baltimore D, Pomerantz RJ. Blocked early-stage latency in the peripheral blood cells of certain individuals infected with human immunodeficiency virus type 1. *Proc Natl Acad Sci U S A* 1992;89:10663–7.
- [32] Yukl SA, Shergill AK, Girling V, Li Q, Killian M, Epling L, et al. Site-Specific Differences in T Cell Frequencies and Phenotypes in the Blood and Gut of HIV-Uninfected and ART-Treated HIV+ Adults. *PLoS One* 2015;10:e0121290.
- [33] Jordan A, Defechereux P, Verdin E. The site of HIV-1 integration in the human genome determines basal transcriptional activity and response to Tat transactivation. *EMBO J* 2001;20:1726–38.
- [34] Howe JG, Shu MD. Epstein-Barr virus small RNA (EBER) genes: unique transcription units that combine RNA polymerase II and III promoter elements. *Cell* 1989;57:825–34.
- [35] Eilebrecht S, Benecke BJ, Benecke A. 7SK snRNA-mediated, gene-specific cooperativity of HMGA1 and P-TEFb. *RNA Biol* 2011;8:1084–93.
- [36] Zheng Y, Josefowicz SZ, Kas A, Chu TT, Gavin MA, Rudensky AY. Genome-wide analysis of Foxp3 target genes in developing and mature regulatory T cells. *Nature* 2007;445:936–40.
- [37] Dong C, Juedes AE, Temann UA, Shresta S, Allison JP, Ruddle NH, et al. ICOS costimulatory receptor is essential for T-cell activation and function. *Nature* 2001;409:97–101.
- [38] Spits H, Yssel H, Paliard X, Kastelein R, Figdor C, de Vries JE. IL-4 inhibits IL-2-mediated induction of human lymphokine-activated killer cells, but not the generation of antigen-specific cytotoxic T lymphocytes in mixed leukocyte cultures. *J Immunol* 1988;141:29–36.
- [39] Walsh GM, Hartnell A, Wardlaw AJ, Kurihara K, Sanderson CJ, Kay AB. IL-5 enhances the *in vitro* adhesion of human eosinophils, but not neutrophils, in a leucocyte integrin (CD11/18)-dependent manner. *Immunology* 1990;71:258–65.
- [40] Wu J, Cherwinski H, Spies T, Phillips JH, Lanier LL. DAP10 and DAP12 form distinct, but functionally cooperative, receptor complexes in natural killer cells. *J Exp Med* 2000;192:1059–68.
- [41] Sullenger BA, Gallardo HF, Ungers GE, Gilboa E. Overexpression of TAR sequences renders cells resistant to human immunodeficiency virus replication. *Cell* 1990;63:601–8.
- [42] Michienzi A, Li S, Zaia JA, Rossi JJ. A nucleolar TAR decoy inhibitor of HIV-1

- replication. *Proc Natl Acad Sci U S A* 2002;99:14047–52.
- [43] Haase AD, Jaskiewicz L, Zhang H, Lainé S, Sack R, Gatignol A, et al. TRBP, a regulator of cellular PKR and HIV-1 virus expression, interacts with Dicer and functions in RNA silencing. *EMBO Rep* 2005;6:961–7.
- [44] Bennasser Y, Yeung ML, Jeang KT. HIV-1 TAR RNA subverts RNA interference in transfected cells through sequestration of TAR RNA-binding protein, TRBP. *J Biol Chem* 2006;281:27674–8.
- [45] Ouellet DL, Plante I, Landry P, Barat C, Janelle ME, Flamand L, et al. Identification of functional microRNAs released through asymmetrical processing of HIV-1 TAR element. *Nucleic Acids Res* 2008;36:2353–65.
- [46] Harwig A, Jongejan A, van Kampen AH, Berkhout B, Das AT. Tat-dependent production of an HIV-1 TAR-encoded miRNA-like small RNA. *Nucleic Acids Res* 2016;44: 4340–53.
- [47] Klase Z, Winograd R, Davis J, Carpio L, Hildreth R, Heydarian M, et al. HIV-1 TAR miRNA protects against apoptosis by altering cellular gene expression. *Retrovirology* 2009;6:18.
- [48] Li L, Feng H, Da Q, Jiang H, Chen L, Xie L, et al. Expression of HIV-encoded microRNA-TAR and its inhibitory effect on viral replication in human primary macrophages. *Arch Virol* 2016;161:1115–23.
- [49] Zuker M. Computer prediction of RNA structure. *Methods Enzymol* 1989;180:262–88.
- [50] Brysbaert G, Pelly FX, Noth S, Benecke A. Quality assessment of transcriptome data using intrinsic statistical properties. *Genomics Proteomics Bioinformatics* 2010;8:57–71.
- [51] Noth S, Brysbaert G, Benecke A. Normalization using weighted negative second order exponential error functions (NeONORM) provides robustness against asymmetries in comparative transcriptome profiles and avoids false calls. *Genomics Proteomics Bioinformatics* 2006;4:90–109.
- [52] Noth S, Benecke A. Avoiding inconsistencies over time and tracking difficulties in Applied Biosystems AB1700/Panther probe-to-gene annotations. *BMC Bioinformatics* 2005;6:307.
- [53] Noth S, Brysbaert G, Pelly FX, Benecke A. High-sensitivity transcriptome data structure and implications for analysis and biologic interpretation. *Genomics Proteomics Bioinformatics* 2006;4:212–29.
- [54] Benecke A. Genomic plasticity and information processing by transcription coregulators. *ComPlexUs* 2003;1:65–76.

Figure legends

Figure 1 Schematic representation and expression of EBER2 TAR fusion RNAs

A. Schematic representation of the EBER2-TAR fusion transcript for the expression of HIV-1 TAR RNA structure (EBER2 TAR). The second hairpin (loop b, Lb) of wild type EBER2 RNA of Epstein-Barr Virus was replaced with the terminal hairpin structure of HIV-1 TAR (highlighted in grey and in red in the schematic structure). EBER2-Lb, the EBER2 backbone

lacking the Lb structure completely, was used as a control. Gene internal promoter elements of the EBER2 backbone are highlighted in green. Nucleotide numbers are indicated. **B.** Expression of EBER2-Lb and EBER2 TAR in HEK293 cells. HEK293 cells were transfected with the constructs EBER2-Lb or EBER2 TAR. Expression levels of both constructs were monitored 48 hours post transfection by RT-PCR. *ACTB* was used as a reference gene.

Figure 2 Identification of HIV-1 TAR target genes in HEK293 cells

A. Heatmap of significant DEGs upon expression of the EBER2 TAR construct as compared to the expression of EBER2-Lb ($P < 0.01$). The \log_2 fold change (L) as well as the merged signals (S) of three biological replicates for each condition are indicated as a color scale. **B.** Pathway enrichment analyses for genes indicated in panel A. Significantly-enriched pathways are indicated together with the corresponding P values (bars) and ratios of the number of significantly-regulated genes per total number of genes involved in each pathway (orange line). The P value threshold (0.05) is indicated as dotted line. z -scores were calculated using Ingenuity Pathway Analysis. Negative z -scores (blue) indicate a repression and positive z -scores (orange) indicate an activation of the corresponding pathway. **C.** Fold changes in expression of immune-related genes upon EBER2 TAR expression. DEG, differentially-expressed gene.

Figure 3 HIV-1 TAR targets the cellular immune response in HEK293 cells

Genes regulated by HIV-1 TAR, which are involved in the repression of activation and proliferation of blood cells and in the repression of HIV-1 replication. Genes repressed and upregulated by TAR are highlighted in green and red, respectively, with the corresponding \log_2 fold change values given in color gradients. Genes are grouped according to their function by symbols. Blue arrows indicate repressive roles, inconsistent regulatory roles are indicated by yellow arrows, whereas grey arrows represent effects that are not predicted. Repressed pathways are indicated in blue, with the corresponding activation z -scores given in a color gradient.

Figure 4 iCOS pathway associated with HIV-1 TAR target genes

iCOS signaling in T helper cells. Proteins encoded by genes with up-regulated and down-regulated expression are highlighted in red and green, respectively.

Figure 5 HIV-1 TAR regulates subsets of Cdk9 and HMGA1 target genes

A. Venn diagram of significant genes targeted by EBER2 TAR ($P < 0.01$; green), dnCdk9 ($P < 0.001$; red), and HMGA1-Flag ($P < 0.05$; blue), respectively. The common subset of HIV-1 TAR targets and dnCdk9 targets is shown in purple, the common subset of HIV-1 TAR targets and HMGA1-Flag targets is shown in brown, the common subset of HMGA1-FLAG targets and dnCdk9 targets is shown in pink and the common subset of all three conditions is shown in black. The hypergeometric distribution P values for each common subset are indicated. **B.** Scatter plot of the EBER2 TAR target genes and dnCdk9 target genes as well as the common subset shown in panel A. The \log_2 fold change in gene expression upon dnCdk9 expression is indicated on the Y-axis, and the \log_2 fold change upon EBER2 TAR expression is indicated on the X-axis. The Pearson correlation coefficient (R) is displayed for the common subset. **C.** Genes of the common subset of all three conditions shown in panel A.

Figure 6 HIV-1 TAR-regulated Cdk9 target genes are 7SK-responsive

A. Venn diagram of significant genes targeted by EBER2 TAR ($P < 0.01$; green), dnCdk9 ($P < 0.001$; red) and 7SK RNA ($P < 0.01$; yellow), respectively. The common subset of HIV-1 TAR targets and dnCdk9 targets is shown in purple, the common subset of HIV-1 TAR targets and 7SK RNA targets is shown in dark red, the common subset of 7SK RNA targets and dnCdk9 targets is shown in orange, and the common subset of all three conditions is shown in black. The hypergeometric distribution P -value for the common subset of HIV-1 TAR and 7SK RNA target genes is indicated. **B.** Scatter plot of the genes targeted by EBER2 TAR and 7SK RNA, as well as the common subset shown in panel A. The \log_2 fold change in gene expression upon 7SK knockdown is indicated on the Y-axis, and the \log_2 fold change upon EBER2 TAR expression is indicated on the X-axis. The Pearson correlation coefficient (R) is displayed for the common subset. **C.** Scatter plot of the genes targeted by EBER2 TAR and 7SK RNA, as well as the common subset shown in panel A. The \log_2 fold change in gene expression upon 7SK overexpression is indicated on the Y-axis and the \log_2 fold change upon EBER2 TAR expression is indicated on the X-axis. The Pearson correlation coefficient (R) is displayed for the common subset. The expression change of the 131 genes commonly targeted by HIV-1 TAR and dnCdk9 upon dnCdk9 overexpression (**D**), upon 7SK knockdown (**E**), and upon 7SK overexpression (**F**), respectively, are sorted from the most negative to the most positive value.

Figure 7 Model for the HIV-1 TAR-mediated regulation of 7SK-responsive P-TEFb target genes




The HIV-1 TAR substructure binds to inactive, HEXIM1-associated P-TEFb in the absence of Tat – either as an RNA chimera in the context of the EBER2 RNA (EBER2 TAR) or as the nascent transcript of the HIV-1 LTR. P-TEFb is thus sequestered in its inactive form, which is endogenously stabilized by 7SK RNA. In both cases, the equilibrium between active (red) and inactive P-TEFb is shifted towards its inactive form, resulting in the repression of P-TEFb target genes. TAR, transactivating response element; HEXIM1, hexamethylene bis-acetamide-inducible protein 1; P-TEFb, positive transcription elongation factor b; Tat, transactivator of transcription; LTR, long terminal repeat; RNAPII, RNA polymerase II; DSIF, 5,6-dichloro-1- β -D-ribofuranosylbenzimidazole (DRB) sensitivity-inducing factor; NELF, negative elongation factor.

Supplemental material



Table S1 List of genes significantly ($P < 0.01$) differentially expressed upon EBER2 HIV-1 TAR expression in HEK293 cells


Table S2 List of the common target genes of EBER2 HIV-1 TAR ($P < 0.01$), dnCdk9 ($P < 0.001$), and 7SK RNA knockdown ($P < 0.01$) in HEK293 cells


Table S3 List of the common target genes of EBER2 HIV-1 TAR ($P < 0.01$), dnCdk9 ($P < 0.001$), and 7SK RNA overexpression ($P < 0.01$) in HEK293 cells




A**B**






CRIP



A**B****C**







A**B****C**

A**B****C****D****E****F**



Host cellular gene



HIV-1 LTR