



**HAL**  
open science

# The binding efficiency of RPA to telomeric G-strands folded into contiguous G-quadruplexes is independent of the number of G4 units

Astrid Lancrey, Loyal Safa, Jean Chatain, Emmanuelle Delagoutte,  
Jean-François Riou, Patrizia Alberti, Carole Saintomé

## ► To cite this version:

Astrid Lancrey, Loyal Safa, Jean Chatain, Emmanuelle Delagoutte, Jean-François Riou, et al.. The binding efficiency of RPA to telomeric G-strands folded into contiguous G-quadruplexes is independent of the number of G4 units. *Biochimie*, 2018, 146, pp.68-72. 10.1016/j.biochi.2017.11.017. hal-01668767

**HAL Id: hal-01668767**

**<https://hal.sorbonne-universite.fr/hal-01668767>**

Submitted on 20 Dec 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Short communication

# The binding efficiency of RPA to telomeric G-strands folded into contiguous G-quadruplexes is independent of the number of G4 units

Astrid Lancrey<sup>a</sup>, Loyal Safa<sup>a</sup>, Jean Chatain<sup>a</sup>, Emmanuelle Delagoutte<sup>a</sup>,  
Jean-François Riou<sup>a</sup>, Patrizia Alberti<sup>a, \*\*</sup>, Carole Saintomé<sup>a, b, \*</sup>

<sup>a</sup> Structure et Instabilité des Génomes, Sorbonne Universités, Muséum National d'Histoire Naturelle, CNRS UMR 7196, INSERM U1154, 43 rue Cuvier, F-75005, Paris, France

<sup>b</sup> Sorbonne Universités, UPMC Univ Paris 06, UFR927, F-75005, Paris, France

---

## A B S T R A C T

Replication protein A (RPA) is a single-stranded DNA binding protein involved in replication and in telomere maintenance. During telomere replication, G-quadruplexes (G4) can accumulate on the lagging strand template and need to be resolved. It has been shown that human RPA is able to unfold a single G4. Nevertheless, the G-strand of human telomeres is prone to fold into higher-order structures formed by contiguous G-quadruplexes. To understand how RPA deals with these structures, we studied its interaction with telomeric G-strands folding into an increasing number of contiguous G4s. The aim of this study was to determine whether the efficiency of binding/unfolding of hRPA to telomeric G-strands depends on the number of G4 units. Our data show that the number  $n$  of contiguous G4 units ( $n \geq 2$ ) does not affect the efficiency of hRPA to coat transiently exposed single-stranded telomeric G-strands. This feature may be essential in preventing instability due to G4 structures during telomere replication.

---

### Keywords:

G-quadruplex

Telomere

Replication protein A

Fluorescence resonance energy transfer (FRET)

Gel electrophoresis

---

## 1. Introduction

The G-rich telomeric sequence can adopt non-canonical DNA conformations known as G-quadruplexes (G4s). These four-stranded structures are based on guanine quartets stabilized by Hoogsteen hydrogen bonds and coordinated by central cations. Evidence is accumulating that G4s can form at telomeres during lagging strand DNA replication and at the 3' G-overhang [1–4]. Several helicases (such as the Werner helicase, WRN) able to unwind G4s are essential for telomere stability and replication [5–7]. In addition, several non-helicase single-stranded DNA binding proteins, involved in telomere maintenance (replication and capping), have been shown to unfold G4s. These proteins are the POT1-TPP1 complex [8,9] (two Shelterin components [10]), RPA [11–14] (a key player in DNA replication, recombination and repair,

that is highly conserved in eukaryotes [15]), and the CST complex [16] (a RPA-like complex [17]). The recruitment of these proteins is tightly coordinated through the cell cycle, ensuring telomere replication and telomere protection [18,19]. Here, we focus on human Replication Protein A (hRPA). In human cells, RPA associates with telomeres [20]; it has been proposed that it binds telomeric single-stranded DNA at replication forks (or telomere ends) during early to middle S-phase [18]. A recent study shows that yeast RPA ensures correct replication of the telomeric lagging strand, and suggests a role of RPA in preventing DNA secondary structures on the telomeric lagging strand template [21]. So far, studies concerning the interaction of hRPA with G4s have focused on a single quadruplex formed by four repetitions of the human telomeric motif (GGGTTA) [11,22–24]. Results obtained with a single G4 may not necessarily reflect the binding properties of hRPA to long telomeric G-strands, which are prone to fold into contiguous (or multimeric) G4s, like beads-on-a-string [25–29]. In the present work, we address, for the first time, the question of how hRPA deals with contiguous (or multimeric) telomeric G4s, with the aim of determining whether its efficiency of binding/unfolding long telomeric G-strands folded into contiguous G4 depends on the number of G4 units.

---

\* Corresponding author. Structure et Instabilité des Génomes, Sorbonne Universités, Muséum National d'Histoire Naturelle, CNRS UMR 7196, INSERM U1154, 43 rue Cuvier, F-75005, Paris, France.

\*\* Corresponding author.

E-mail addresses: [alberti@mnhn.fr](mailto:alberti@mnhn.fr) (P. Alberti), [carole.saintome@mnhn.fr](mailto:carole.saintome@mnhn.fr) (C. Saintomé).

## 2. Materials and methods

### 2.1. Materials

Recombinant hRPA was expressed in the *Escherichia coli* BL21 (DE3) strain transformed with the plasmid pET<sub>11a</sub>hRPA that permits the co-expression of RPA1, RPA2 and RPA3. hRPA was purified as previously described [21]. Oligonucleotides were purchased from Eurogentec.

### 2.2. Electrophoretic mobility shift assays

Oligonucleotides were labeled with  $\gamma$ [<sup>32</sup>P]ATP using T4 polynucleotide kinase. For all EMSA experiments, hRPA was diluted and pre-incubated (20 min at 4 °C) in a buffer containing 50 mM Tris-HCl pH 7.5, 100 mM KCl, 1 mM DTT, 10% glycerol, 0.2 mg/mL BSA and 0.1 mM EDTA. In a standard reaction, radiolabeled oligonucleotide (2 nM) (slowly annealed from high to low temperature) was incubated with various amounts of protein in 10  $\mu$ L reaction buffer [50 mM HEPES pH 7.9, 0.1 mg/ml BSA, 100 mM KCl or NaCl and 2% glycerol] for 20 min at 20 °C. Samples were then loaded on a native 1% agarose in 0.5 $\times$  TBE buffer. After electrophoresis, the gel was dried and exposed on a phosphorimager screen, scanned with a Typhoon instrument (Molecular Dynamics). Quantification was made with ImageQuant version 5.1. For each hRPA concentration, the fraction of radiolabelled oligonucleotide bound to hRPA was calculated as follow:  $I_{DNA\ bound\ to\ hRPA} / (I_{free\ DNA} + I_{DNA\ bound\ to\ hRPA})$ .

### 2.3. Fluorescence titrations

Fluorescence titrations were carried out on a SPEX Fluorolog spectrofluorimeter (Jobin Yvon) equipped with a circulating water bath to regulate the temperature of the cell holder. Fluorescence spectra of 100 nM H69-Q1,2,3 oligonucleotides (doubly-labeled with a FAM and a TAMRA) were recorded at 20 °C in a buffer containing 50 mM KCl, 2 mM MgCl<sub>2</sub>, and 5 mM lithium cacodylate (pH 7.2). H69-Q1,2,3 were pre-incubated 10 min at 20 °C. The protein (from 100 to 1000 nM) was directly added to the solutions containing the oligonucleotides. The spectra were collected after 2 min of incubation while exciting at 470 nm. The fluorescence intensity of the donor FAM ( $I_D$ ) was measured at 518 nm, the fluorescence intensity of the acceptor TAMRA ( $I_A$ ) was measured at 586 nm; the percentage of opened labeled G4 units at 20 °C was calculated as:

$$(P - P_f) \times 100 / (P_u - P_f),$$

where:

$$\begin{aligned} P &= I_D / (I_D + I_A) \text{ at a given hRPA concentration at } 20^\circ\text{C}, \\ P_f &= I_D / (I_D + I_A) \text{ in the absence of hRPA at } 20^\circ\text{C}, \\ P_u &= I_D / (I_D + I_A) \text{ in the absence of hRPA at } 80^\circ\text{C}. \end{aligned}$$


We assumed that, in the absence of hRPA, at 20 °C all the labeled G4 units are folded, while at 80 °C all the labeled G4 units are unfolded. Under these assumptions,  $P_f$  is the  $P$  value corresponding to 0% of unfolded labeled G4 units, while  $P_u$  is the  $P$  value corresponding to 100% of unfolded labeled G4 units. The expression  $(P - P_f) \times 100 / (P_u - P_f)$  provides, hence, the percentage of unfolded labeled G4 units for a given hRPA concentration.

## 3. Results and discussion

We carried out our study in the presence of the biological relevant potassium cation. We first investigated if hRPA is able to

unfold contiguous G4s. To this purpose, we carried out fluorescent resonance energy transfer (FRET) titration experiments using the telomeric sequence H69 ((G<sub>3</sub>T<sub>2</sub>A)<sub>3</sub>G<sub>3</sub>), which folds into three contiguous G4 units. Three modified H69 oligonucleotides, doubly labeled with a 6-carboxyfluorescein (FAM) and a tetramethylrhodamine (TAMRA) at different positions, were used. In H69-Q1, the fluorophores are placed at both sides of the first G4 unit; in H69-Q2, they are placed at both sides of the second G4 unit; in H69-Q3, they are placed at both sides of the third G4 unit (Supplementary Information, Table S1). The doubly labeled G4 units of these three constructs have temperatures of half-denaturation in the range of 50–60 °C, as previously assessed by melting experiments followed by fluorescence [26]. For all H69-Q1,2,3, fluorimetric titration with hRPA revealed an increase of FAM fluorescence and a decrease in TAMRA fluorescence upon hRPA addition, indicating that hRPA is able to open each of the three G4 units, independently of its position (Fig. 1). These results demonstrate that hRPA is able to unfold all the G4s present in a multimeric structure.

Having shown that hRPA unfolds multimeric G4s, we next investigated whether its binding efficiency depended on the number of G4 units. To this purpose, we carried out electrophoretic mobility shift assays (EMSA) using four oligonucleotides: H27 (T<sub>2</sub>A(G<sub>3</sub>T<sub>2</sub>A)<sub>3</sub>G<sub>3</sub>T<sub>2</sub>A), H51 (T<sub>2</sub>A(G<sub>3</sub>T<sub>2</sub>A)<sub>7</sub>G<sub>3</sub>T<sub>2</sub>A), H75 (T<sub>2</sub>A(G<sub>3</sub>T<sub>2</sub>A)<sub>11</sub>G<sub>3</sub>T<sub>2</sub>A) and H99 (T<sub>2</sub>A(G<sub>3</sub>T<sub>2</sub>A)<sub>15</sub>G<sub>3</sub>T<sub>2</sub>A). We previously characterized the structures formed by these oligonucleotides: we showed that H51, H75 et H99 fold into two, three and four contiguous G4 units, respectively, which are non-interacting and have identical stabilities; while the single G4 formed by the shortest sequence H27 has a higher stability [26] (Supplementary Information, Table S1). We point out that the binding constant of hRPA to dT<sub>50</sub>, dT<sub>60</sub> and dT<sub>70</sub> oligonucleotides (longer than the size of the 30-nt RPA extended-binding mode and able in principle to bind more than one RPA, up to three RPA for dT<sub>70</sub>) does not depend on their length [30]. Hence, any significant difference in the binding of RPA to H51, H75 and H99 should not be due to oligonucleotide length, but solely on the number of G4 units. H27, H51, H75, H99 oligonucleotides in the presence of increasing hRPA concentrations showed retarded bands on agarose gel (Fig. 2A). The number of bands increased with the length of the oligonucleotide, revealing hRPA:DNA complexes of increasing stoichiometry, where one, two, three, four (or more) hRPA are bound. These results show that hRPA is able to bind telomeric strands structured in multimeric G4s and to form complexes where several proteins are present along the DNA strand. For each oligonucleotide, we quantified the percentage of DNA bound to hRPA as a function of hRPA concentration and determined the hRPA concentration to obtain 50% of bound oligonucleotide (Fig. 2B and C). The binding curves showed two main features: (i) hRPA bound more efficiently the multimeric G4 sequences (H51, H75 and H99) than the monomeric G4 sequence (H27); (ii) the efficiency of hRPA binding to the multimeric G4 sequences (H51, H75 and H99) was independent of the number of formed G4 units (two, three or four). In addition, the non-sigmoidal shape of the binding curves, the coexistence of complexes of different stoichiometries (CI, CII, CIII and CIII+) up to 25–50 nM hRPA concentrations (quantification of the percentage of each complex from Fig. 1 is shown in Supplementary Information, Fig. S1) and data fitting according to a model of binding to a one-dimensional homogeneous lattice [31] (Supplementary Information, Fig. S2) further suggest a poorly cooperative binding mode of hRPA to H51, H75 and H99, as previously observed with long non-structured single-stranded oligonucleotides [30,32]. The higher hRPA binding efficiency ([hRPA]<sub>50</sub> around 5 nM) toward multimeric G4s compared to the single G4 ([hRPA]<sub>50</sub> around 20 nM) can be explained by the lower stability of these higher-order structures




**Fig. 1. Fluorescence titrations.** (A) Fluorescence emission spectra of H69-Q1, H69-Q2 and H69-Q3 with increasing amounts of hRPA in  $K^+$  solution (excitation wavelength 470 nm). Spectra of the doubly labeled oligonucleotides in the absence of hRPA are in bold line. Dotted arrows indicate the evolution of emission spectra with increasing hRPA concentrations. (B) Percentages of unfolded labeled G4 units in H69-Q1 (diamonds), H69-Q2 (circles) and H69-Q3 (squares) as a function of hRPA/oligonucleotide concentration ratio  $r$ . The percentage of unfolded labeled G4 units was calculated as detailed in Methods section. The oligonucleotide concentration was 100 nM and the protein concentration varied from 100 to 1100 or 1300 nM.

(temperature of thermal transition  $T_t$  50 °C [26], Supplementary Information, Table S1) compared to the single G4 formed by H27 ( $T_t$  55 °C [26], Supplementary Information, Table S1). Indeed, we previously reported an inverse relationship between the efficiency of binding of hRPA to a G4 sequence and the stability of the formed G4 [23]. In addition, the lower binding efficiency of hRPA to H27 may also be due to its length (27 nt), which is 3 nt shorter than the optimal length to obtain the more stable hRPA/DNA complex (30 nt-binding mode) [33]. The independence of hRPA binding efficiency of the number of contiguous G4 units may result from the poorly cooperative binding mode of hRPA. We asked if this feature is also due to the peculiar character of multimeric G4 structures formed by telomeric G-strands in potassium, where the contiguous G4 units are non-interacting and have similar stabilities, disregarding of their number [26]. To answer to this question, we studied the binding of hRPA to H21 ((G<sub>3</sub>T<sub>2</sub>A)<sub>3</sub>G<sub>3</sub>), H45 ((G<sub>3</sub>T<sub>2</sub>A)<sub>7</sub>G<sub>3</sub>), H69 ((G<sub>3</sub>T<sub>2</sub>A)<sub>11</sub>G<sub>3</sub>) and H93 ((G<sub>3</sub>T<sub>2</sub>A)<sub>15</sub>G<sub>3</sub>) sequences in sodium, folding into one, two, three and four G4 units, respectively [34], but lacking the T<sub>2</sub>A tails at their extremities. We previously showed that, among contiguous G4 structures formed by telomeric DNA, the structure formed by H45 in sodium is unique, since its two G4 units interact with each other [34]. In sodium, H45 behaves as a single structural entity more stable than the beads-on-a-string structures formed by the longer H69 and H93 sequences [34] (Supplementary Information, Table S1). hRPA displayed a low

binding efficiency for the sequence H21, compared to H69 and H93 (Fig. 2D–E); this is due to the higher stability of the single G4 formed by H21 ( $T_t$  58 °C for H21,  $T_t$  43/45 °C for H69/H93 [34], Supplementary Information, Table S1) and to its length (9 nt shorter than the optimal length to obtain the more stable hRPA/DNA complex), similarly to what observed in potassium with H27. Among contiguous G4s, the binding curves obtained by EMSA showed that, in sodium, hRPA had similar binding efficiencies for the two sequences H69 and H93, but a lower binding efficiency for the sequence H45 (Fig. 2E–D). The lower binding efficiency of hRPA to H45 in sodium might result not solely from its higher stability ( $T_t$  53 °C [34], Supplementary Information, Table S1), but also from conformational factors related to the interaction between its two G4 units. Overall, EMSA on H51, H75 and H99 oligonucleotides in potassium demonstrate that the number of contiguous G4s does not affect the efficiency of binding of hRPA to telomeric G-strands. The result obtained with the H45 counterexample in sodium supports that this feature is related to the peculiar character of telomeric G4 units in potassium: non-interacting G4 units with similar stabilities, disregarding of their number.

We point out that, based on results obtained with the shortest oligonucleotide H27 (and H21), it would be wrong to argue that hRPA unfolds less efficiently a single G4 formed within a telomeric G-strand than contiguous G4s. Indeed, the single G4s formed by the 21mer sequence (G<sub>3</sub>T<sub>2</sub>A)<sub>3</sub>G<sub>3</sub> (H21) or the extended 27mer sequence



**Fig. 2. Electrophoretic mobility shift assays.**  $P^{32}$ -oligonucleotides (2 nM) were incubated with increasing amounts of hRPA in the presence of 100 mM KCl (A, B, C) or 100 mM NaCl (D, E). Electrophoresis was run on a native 1% agarose gel. (A) EMSA gels in KCl. (B) Percentages of oligonucleotide bound to hRPA in KCl: H27 (diamonds), H51 (circles), H75 (squares), H99 (triangles) as a function of hRPA concentration. Error bars correspond to standard deviations calculated from four independent experiments for points corresponding to 5, 10, 25 and 50 nM of hRPA, and from two independent experiments for points 2 and 100 nM hRPA. (C) Mean concentration of hRPA to obtain 50% of bound oligonucleotide ( $[hRPA]_{50}$ ) from four independent experiments in KCl. (D) Percentages of oligonucleotide bound to hRPA in NaCl: H21 (diamonds), H45 (circles), H69 (squares), H93 (triangles) as a function of hRPA concentration. Error bars correspond to standard deviations calculated from two independent experiments. (E) Mean concentration of hRPA to obtain 50% of bound oligonucleotide ( $[hRPA]_{50}$ ) from two independent experiments in NaCl. EMSA gels in NaCl are shown in Supplementary Information, Fig. S3.



TTA-(G<sub>3</sub>T<sub>2</sub>A)<sub>3</sub>G<sub>3</sub>-TTA (H27) are not suitable models for the binding of RPA to a single G4 unit within a telomeric G-strand, both for their length and for their higher stability [26].

#### 4. Conclusion

In this study, we addressed for the first time, the question of how a single-stranded DNA binding protein involved in telomere replication, hRPA, deals with multimeric G4s that may form on the lagging strand template during telomere replication. We showed that hRPA binds telomeric G-strands structured into contiguous G4s, unfolding them. Importantly, we showed that the efficiency of binding of hRPA to telomeric G-strands is independent of the number of contiguous G4 units. In other words, if contiguous G4s form during replication, their number does not affect the efficiency of hRPA to coat transiently exposed single-stranded telomeric G-strands. This feature of hRPA may be crucial during telomere replication in suppressing or preventing multiple G4 accumulation on the lagging strand template and in ensuring telomere stability.

#### Author agreement

All the co-authors have contributed collectively to this study in order named as full participants. All the co-authors approved the final version of the manuscript and its submission to *BIOCHIMIE*. This work has not previously been published and is not being considered for publication elsewhere.

#### Acknowledgments

We thank the members of the laboratory team “Nucleic Acid Structure, Telomeres and Evolution”, in particular Anthony Bugaut, for helpful discussions. This work was supported by the “Muséum National d’Histoire Naturelle” (MNHN), the “Centre National de la Recherche Scientifique” (CNRS) and the “Institut National de la Santé et de la Recherche Médicale” (Inserm). LS was supported by a doctoral fellowship from the “Ligue Nationale contre le Cancer”.

#### Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.biochi.2017.11.017>.

#### References

- [1] L. Crabbe, R.E. Verdun, C.I. Haggblom, J. Karlseder, Defective telomere lagging strand synthesis in cells lacking WRN helicase activity, *Science* 306 (2004) 1951–1953.
- [2] D. Gomez, R. Paterski, T. Lemarteleur, K. Shin-Ya, J.L. Mergny, J.F. Riou, Interaction of telomestatin with the telomeric single-strand overhang, *J. Biol. Chem.* 279 (2004) 41487–41494.
- [3] A. Sfeir, S.T. Kosiyatrakul, D. Hockemeyer, S.L. MacRae, J. Karlseder, C.L. Schildkraut, T. de Lange, Mammalian telomeres resemble fragile sites and require TRF1 for efficient replication, *Cell* 138 (2009) 90–103.
- [4] J.B. Vannier, V. Pavicic-Kaltenbrunner, M.I. Petalcorin, H. Ding, S.J. Boulton, RTEL1 dismantles T loops and counteracts telomeric G4-DNA to maintain telomere integrity, *Cell* 149 (2012) 795–806.
- [5] P. Mohaghegh, J.K. Karow, R.M. Brosh Jr., V.A. Bohr, I.D. Hickson, The Bloom’s and Werner’s syndrome proteins are DNA structure-specific helicases, *Nucleic Acids Res.* 29 (2001) 2843–2849.
- [6] C. Ribeyre, J. Lopes, J.B. Boule, A. Piazza, A. Guedin, V.A. Zakian, J.L. Mergny, A. Nicolas, The yeast Pif1 helicase prevents genomic instability caused by G-quadruplex-forming CEB1 sequences in vivo, *PLoS Genet.* 5 (2009) e1000475.
- [7] J.B. Vannier, G. Sarek, S.J. Boulton, RTEL1: functions of a disease-associated helicase, *Trends Cell Biol.* 24 (2014) 416–425.
- [8] A.J. Zaugg, E.R. Podell, T.R. Cech, Human POT1 disrupts telomeric G-quadruplexes allowing telomerase extension in vitro, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 10864–10869.
- [9] H. Hwang, N. Buncher, P.L. Opresko, S. Myong, POT1-TPP1 regulates telomeric overhang structural dynamics, *Structure* 20 (2012) 1872–1880.
- [10] I. Schmutz, T. de Lange, Shelterin, *Curr. Biol.* CB 26 (2016) R397–R399.
- [11] T.R. Salas, I. Petruseva, O. Lavrik, A. Bourdoncle, J.L. Mergny, A. Favre, C. Saintome, Human replication protein A unfolds telomeric G-quadruplexes, *Nucleic acids Res.* 34 (2006) 4857–4865.
- [12] T. Kibe, Y. Ono, K. Sato, M. Ueno, Fission yeast Taz1 and RPA are synergistically required to prevent rapid telomere loss, *Mol. Biol. Cell* 18 (2007) 2378–2387.
- [13] V. Schramke, P. Luciano, V. Brevet, S. Guillot, Y. Corda, M.P. Longhese, E. Gilson, V. Geli, RPA regulates telomerase action by providing Est1p access to chromosome ends, *Nat. Genet.* 36 (2004) 46–54.
- [14] J. Smith, H. Zou, R. Rothstein, Characterization of genetic interactions with RFA1: the role of RPA in DNA replication and telomere maintenance, *Biochimie* 82 (2000) 71–78.
- [15] M.S. Wold, Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism, *Annu. Rev. Biochem.* 66 (1997) 61–92.
- [16] N.F. Lue, R. Zhou, L. Chico, N. Mao, O. Steinberg-Neifach, T. Ha, The telomere capping complex CST has an unusual stoichiometry, makes multipartite interaction with G-Tails, and unfolds higher-order G-tail structures, *PLoS Genet.* 9 (2013), e1003145.
- [17] C. Rice, E. Skordalakes, Structure and function of the telomeric CST complex, *Comput. Struct. Biotechnol. J.* 14 (2016) 161–167.
- [18] R.L. Flynn, R.C. Centore, R.J. O’Sullivan, R. Rai, A. Tse, Z. Songyang, S. Chang, J. Karlseder, L. Zou, TERRA and hnRNP A1 orchestrate an RPA-to-POT1 switch on telomeric single-stranded DNA, *Nature* 471 (2011) 532–536.
- [19] L.Y. Chen, S. Redon, J. Lingner, The human CST complex is a terminator of telomerase activity, *Nature* 488 (2012) 540–544.
- [20] J. Dejjardin, R.E. Kingston, Purification of proteins associated with specific genomic loci, *Cell* 136 (2009) 175–186.
- [21] J. Audry, L. Maestroni, E. Delagoutte, T. Gauthier, T.M. Nakamura, Y. Gachet, C. Saintome, V. Geli, S. Coulon, RPA prevents G-rich structure formation at lagging-strand telomeres to allow maintenance of chromosome ends, *EMBO J.* 34 (2015) 1942–1958.
- [22] M.H. Qureshi, S. Ray, A.L. Sewell, S. Basu, H. Balci, Replication protein A unfolds G-quadruplex structures with varying degrees of efficiency, *J. Phys. Chem. B* 116 (2012) 5588–5594.
- [23] L. Safa, E. Delagoutte, I. Petruseva, P. Alberti, O. Lavrik, J.F. Riou, C. Saintome, Binding polarity of RPA to telomeric sequences and influence of G-quadruplex stability, *Biochimie* 103 (2014) 80–88.
- [24] L. Safa, N.M. Gueddouda, F. Thiebaut, E. Delagoutte, I. Petruseva, O. Lavrik, O. Mendoza, A. Bourdoncle, P. Alberti, J.F. Riou, C. Saintome, 5’ to 3’ unfolding directionality of DNA secondary structures by replication protein A: G-quadruplexes and duplexes, *J. Biol. Chem.* 291 (2016) 21246–21256.
- [25] J. Abraham Punnoose, Y. Cui, D. Koirala, P.M. Yangyuoru, C. Ghimire, P. Shrestha, H. Mao, Interaction of G-quadruplexes in the full-length 3’ human telomeric overhang, *J. Am. Chem. Soc.* 136 (2014) 18062–18069.
- [26] A. Bugaut, P. Alberti, Understanding the stability of DNA G-quadruplex units in long human telomeric strands, *Biochimie* 113 (2015) 125–133.
- [27] L. Petraccone, C. Spink, J.O. Trent, N.C. Garbett, C.S. Mekmaysy, C. Giancola, J.B. Chaires, Structure and stability of higher-order human telomeric quadruplexes, *J. Am. Chem. Soc.* 133 (2011) 20951–20961.
- [28] Y. Xu, T. Ishizuka, K. Kurabayashi, M. Komiyama, Consecutive formation of G-quadruplexes in human telomeric-overhang DNA: a protective capping structure for telomere ends, *Angew. Chem.* 48 (2009) 7833–7836.
- [29] H.Q. Yu, D. Miyoshi, N. Sugimoto, Characterization of structure and stability of long telomeric DNA G-quadruplexes, *J. Am. Chem. Soc.* 128 (2006) 15461–15468.
- [30] C. Kim, B.F. Paulus, M.S. Wold, Interactions of human replication protein A with oligonucleotides, *Biochemistry* 33 (1994) 14197–14206.
- [31] J.D. McGhee, P.H. von Hippel, Theoretical aspects of DNA-protein interactions: co-operative and non-co-operative binding of large ligands to a one-dimensional homogeneous lattice, *J. Mol. Biol.* 86 (1974) 469–489.
- [32] C. Kim, M.S. Wold, Recombinant human replication protein A binds to polynucleotides with low cooperativity, *Biochemistry* 34 (1995) 2058–2064.
- [33] I.M. Wyka, K. Dhar, S.K. Binz, M.S. Wold, Replication protein A interactions with DNA: differential binding of the core domains and analysis of the DNA interaction surface, *Biochemistry* 42 (2003) 12909–12918.
- [34] C. Saintome, S. Amrane, J.L. Mergny, P. Alberti, The exception that confirms the rule: a higher-order telomeric G-quadruplex structure more stable in sodium than in potassium, *Nucleic acids Res.* 44 (2016) 2926–2935.