Dinuclear ($N \wedge C \wedge N$) Pincer Pt(II) Complexes with Bridged Organometallic Linkers: Synthesis, Structures, Self-Aggregation, and Photophysical Properties

Hugo Sesolis, Carmen Ka-Man Chan, Geoffrey Gontard, Heidi Li-Ki-Fu,
Vivian Wing-Wah Yam, Hani Amouri

► To cite this version:

Hugo Sesolis, Carmen Ka-Man Chan, Geoffrey Gontard, Heidi Li-Ki-Fu, Vivian Wing-Wah Yam, et al.. Dinuclear ($N \wedge C \wedge N$) Pincer Pt(II) Complexes with Bridged Organometallic Linkers: Synthesis, Structures, Self-Aggregation, and Photophysical Properties. *Organometallics*, 2017, 36 (24), pp.4794 - 4801. 10.1021/acs.organomet.7b00680 . hal-01675266

HAL Id: hal-01675266

<https://hal.sorbonne-universite.fr/hal-01675266>

Submitted on 4 Jan 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Dinuclear (N[^]C[^]N) Pincer Pt(II) Complexes with Bridged Organometallic Linkers: Synthesis, Structures, Self-Aggregation, and Photophysical Properties

Hugo Sesolis,[‡] Carmen Ka-Man Chan,[†] Geoffrey Gontard,[‡] Heidi Li-Ki Fu,[†] Vivian Wing-Wah Yam,^{*,†,‡,Ⓛ} and Hani Amouri^{*,‡,Ⓛ}

[‡]Sorbonne Universités, UPMC Université Paris 06, and CNRS, IPCM UMR 8232, 4 place Jussieu, Paris CEDEX 05 75252, France

[†]Institute of Molecular Functional Materials (Areas of Excellence Scheme, University Grants Committee, Hong Kong) and Department of Chemistry, The University of Hong Kong, Pokfulam Road, Hong Kong, PR China

Supporting Information

ABSTRACT: A new family of cationic dinuclear cyclometalated Pt(II) complexes containing an organometallic assembling ligand has been reported. The general formulas for these compounds is as follow: $[R-(N^{\wedge}C^{\wedge}N)Pt-L-L-Pt(N^{\wedge}C^{\wedge}N)-R][X]_2$, where $L-L = [Cp^*Ir-p-(\eta^4-C_6H_4S_2)]$, $R = H$, $X = OTf$ (**5**); $R = CF_3$, $X = OTf$ (**6a**), SbF_6 (**6b**); $L-L = [Cp^*Ir-p-(\eta^4-C_6H_4Se_2)]$, $R = CF_3$, $X = OTf$ (**7a**), SbF_6 (**7b**). In these coordination assemblies two cyclometalated $R-(N^{\wedge}C^{\wedge}N)Pt$ moieties are held by either η^4 -dithio-*p*-benzoquinone complex $[Cp^*Ir-p-(\eta^4-C_6H_4S_2)]$ (**3**) or η^4 -diseleno-*p*-benzoquinone complex $[Cp^*Ir-p-(\eta^4-C_6H_4Se_2)]$ (**4**). The molecular structures of the known complex $[4-CF_3-(N^{\wedge}C^{\wedge}N)-PtCl]$ as well as two compounds of the above family $[(N^{\wedge}C^{\wedge}N)Pt-Cp^*Ir-p-(\eta^4-C_6H_4S_2)-Pt(N^{\wedge}C^{\wedge}N)][CF_3SO_3]_2$ (**5**) and $[CF_3-(N^{\wedge}C^{\wedge}N)Pt-Cp^*Ir-p-(\eta^4-C_6H_4S_2)-Pt(N^{\wedge}C^{\wedge}N)-CF_3][CF_3SO_3]_2$ (**6a**) were ascertained by single crystal X-ray diffraction study and confirmed the formation of the target molecules. The solid-state packing of **5** and **6a** confirms the presence $\pi-\pi$ and short $Pt\cdots Pt$ interactions among individual units providing 1D supramolecular chains. To our knowledge, these are the only known examples where two cyclometalated $Pt(N^{\wedge}C^{\wedge}N)$ units are assembled by a bridging ligand (vide infra). All compounds show phosphorescence in the bluish-green region (486–521 nm) in the solution at room temperature and exhibit higher luminescence quantum yields relative to the analogous compounds containing a $Pt(terpy)$ chromophore in thin film.

Molecular structure of **6a** showing self-aggregation due to $Pt\cdots Pt$ interactions.

■ INTRODUCTION

Cyclometalated platinum(II) complexes containing the pincer-type $N^{\wedge}C^{\wedge}N$ ligands have received considerable attention.^{1–4} They provide rich photophysical properties^{5,6} and unusually high phosphorescence quantum yields, for example, a photoluminescence quantum yield (PLQY) of up to 0.68 in solution was reported by Williams and co-workers.⁷ Such compounds form stable square-planar complexes with d^8 electronic configuration, which promotes metal–metal and $\pi-\pi$ interactions that profoundly impact the properties of the electronic excited states.^{8,9} Moreover, the introduction of substituents into the aryl or pyridyl units impacts the excited state energy and eventually the color of such compounds. Due to their charge neutrality and intense phosphorescence, cyclometalated platinum(II) complexes can be viewed as attractive candidates for applications for optical devices and OLEDs.^{10–13}

Despite these advances, cationic cyclometalated $Pt(N^{\wedge}C^{\wedge}N)$ compounds are less known; the synthesis and luminescent properties of some mononuclear cationic species have been reported by Che and co-workers and others.^{14–17} Moreover,

luminescent coordination assemblies containing two cationic $Pt(N^{\wedge}C^{\wedge}N)$ units are even more scarce.¹⁶ Thus, we sought to design novel classes of luminescent coordination assemblies where appropriate bridging linkers are capable to attach two $(N^{\wedge}C^{\wedge}N)Pt$ units, in the hopes that such novel compounds might promote strong intermolecular $Pt\cdots Pt$ and $\pi-\pi$ interactions which in turn would affect the luminescent properties. In a previous work, the synthesis of luminescent heterotrinnuclear coordination assemblies, $[Pt(terpy)\{Cp^*Ir-p-(\eta^4-C_6H_4E_2)\}Pt(terpy)][OTf]_4$ was reported, in which an organometallic linker $[Cp^*Ir(\eta^4-C_6H_4E_2)]$ ($E = S, Se$) holds two $Pt(terpy)$ units through coordination via the sulfur or selenium centers (Figure 1).^{18,19} The molecular structures showed that individual units exhibit $\pi-\pi$ and $Pt\cdots Pt$ interactions to give a 1D chain in the solid state. The heterotrinnuclear coordination assembly showed unusual UV–vis absorption and luminescence properties at low temperature,

Figure 1. $[\text{Pt}(\text{terpy})\{\text{Cp}^*\text{Ir}-p-(\eta^4\text{-C}_6\text{H}_4\text{S}_2)\}\text{Pt}(\text{terpy})][\text{OTf}]_4$ and $[\text{Pt}(\text{terpy})\{\text{Cp}^*\text{Ir}-p-(\eta^4\text{-C}_6\text{H}_4\text{Se}_2)\}\text{Pt}(\text{terpy})][\text{OTf}]_4$ reported previously.^{18,19}

Figure 2. Schematic drawings of the thio- and seleno-quinone intermediates (1 and 2), the isolated metalated thio- and seleno-quinones (3 and 4), and the luminescent cyclometalated platinum salts (5, 6a, 6b, 7a, and 7b).

imparted from self-aggregation of the individual subunits mediated by π - π and Pt...Pt interactions.

Pursuing our research in this area, we wish to report the synthesis and luminescent properties of rare examples where two cyclometalated Pt chromophores are now attached to a thio- or seleno- organometallic linker p -[Cp*Ir(η^4 -C₆H₄E₂), E = S (3) and Se (4) (Figure 2). To the best of our knowledge, these are the first dicationic Pt(N^{^C}N) compounds to be prepared displaying quinonoid π -complexes as assembling ligands. The reason for the scarcity of such compounds is that unlike quinones which are a prominent class of compounds that play an important role in chemistry and biology the related thio- and seleno-quinones “C₆H₄Se₂” (1) (Figure 2), are in contrast, unstable^{20–22} and consequently do not exist in nature; hence, their chemical properties remain little known.

Thus, we anticipated that metalated thio- and seleno-quinone might be appropriate assembling ligands for a new class of luminescent Pt(II) cyclometalated complexes (Figure 2). The latter represents a novel strategy to obtain cationic luminescent coordination assemblies containing cyclometalated Pt(N^{^C}N) chromophores.

Pursuing our research program in the area of metal-stabilized reactive intermediates, we demonstrated that the Cp*Ir fragment is a powerful stabilizing entity.^{23–26} For instance, in 1998 we reported the synthesis of the first stable *ortho*-quinone methide metal complex, in which the Cp*Ir unit is attached to the internal diene moiety through η^4 -coordination.^{27–29} Later, in 2006 we described the synthesis of the first stable iridium *p*-dithiobenzoquinone complex, p -[Cp*Ir(η^4 -C₆H₄S₂)], and the related *ortho* congener, o -[Cp*Ir(η^4 -C₆H₄S₂)].^{30,31} Both compounds were used as successful organometallic linkers to construct functional coordination assemblies.^{32,33} Moreover, we

reported the synthesis and the X-ray molecular structure of the *p*-diselenobenzoquinone (2) as a metal complex, p -[Cp*Ir(η^4 -C₆H₄Se₂)] (4) (Figure 2).³⁴

In this work, the preparation of several luminescent cationic dinuclear cyclometalated Pt(II) complexes containing an organometallic linker is described. For instance, [(N^{^C}N)-Pt-Cp*Ir- p -(η^4 -C₆H₄S₂)-Pt(N^{^C}N)][OTf]₂ (5) and those containing CF₃-cyclometalated platinum counterparts, [CF₃-(N^{^C}N)Pt-L-L-Pt(N^{^C}N)-CF₃][X]₂, with L-L = [Cp*Ir- p -(η^4 -C₆H₄S₂)], X = OTf (6a), SbF₆ (6b); L-L = [Cp*Ir- p -(η^4 -C₆H₄Se₂)], X = OTf (7a), SbF₆ (7b). The molecular structures of 5 and 6a were determined by single crystal X-ray diffraction, which confirmed the formation of the target complexes. Remarkably, the crystal packing shows the presence of π - π and short Pt...Pt interactions among individual units, which induce the formation of 1D supramolecular chains. These complexes exhibit interesting luminescent properties in the solid state and in thin film matrix at room temperature and showed higher luminescence quantum yields compared to those obtained with Pt(terpy) chromophores. This behavior might arise from the nature of the cyclometalated Pt(N^{^C}N) luminophore.¹⁹ To the best of our knowledge, our compounds are the first examples of dicationic luminescent Pt(N^{^C}N) complexes bearing an ancillary ligand coordinated through sulfur and selenium centers.³

RESULTS AND DISCUSSION

Synthesis of the Cationic (N^{^C}N)Pt(II) Complexes with the Organometallic Linkers (5, 6a and 7a). The pincer-type cyclometalated starting material (N^{^C}N)Pt-Cl was first prepared by Cardenas and co-workers³⁵ in 1999 and the high emission quantum yield was revealed by Williams and

Scheme 1. General Synthesis Procedure to Luminescent Heterotrinnuclear Species 5, 6a, 6b, 7a and 7b

Figure 3. Molecular structures of the free ligand 4- $\text{CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N}$, metal complex $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$, and the related head-to-tail dimer.

co-workers⁷ in 2003, while the synthesis and luminescent properties of the functionalized cyclometalated platinum complex $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ was reported by Che et al. in 2009.¹⁴

Treatment of $[(\text{N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ or $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ in acetone with silver salt AgX and subsequent reaction with the organometallic linkers, $[\text{Cp}^*\text{Ir-}p\text{-(}\eta^4\text{-C}_6\text{H}_4\text{E}_2\text{)}]$ ($\text{E} = \text{S}$ (**3**) or Se (**4**)), provided an orange-red solution which after workup gave target compounds **5**, **6a**, **6b**, **7a** and **7b** (Scheme 1). For instance, the ^1H NMR spectrum of **5** recorded in CD_3CN shows the presence of a singlet at δ 7.23 ppm attributed to the metalated *p*-quinonoid unit and a singlet at δ 1.92 ppm assigned to the Cp^*Ir moiety. Furthermore, we note the presence of several multiplets in the range of δ 7.72–9.10 ppm assigned to the protons of the fragment “ $(\text{N}^{\wedge}\text{C}^{\wedge}\text{N})\text{Pt}$ ”. We noticed however that upon standing at room temperature in CD_3CN solution the complex undergoes changes and decoordination of the chromophore occurs slowly on the NMR time-scale. Perhaps this is due to the σ -*trans* effect of the carbon center, which weakens the coordination of the sulfur ligands in **5**. Such behavior had also been reported in cyclometalated Pt(II) complexes containing a sulfur coligand.³ We anticipated that the related CF_3 -cyclometalated platinum compounds would be more stable since the electron-withdrawing inductive effect of the CF_3 - group might alleviate the *trans* effect of the carbon center *trans* to the sulfur or selenium nucleophile in **6a**, **6b**, **7a** and **7b**. Complex **5** was fully characterized and in addition its X-ray molecular structure was determined (vide infra). Compounds **6a**, **6b**, **7a** and **7b** gave pattern in the ^1H NMR spectra similar to that of **5** and turned out to be more stable in solution. Moreover, the presence of the CF_3 - functional group is observed in the ^{19}F NMR spectra (Experimental Section). The infrared data of the above compounds confirm the presence of the respective counter-anion (Experimental Section).

X-ray Molecular Structures of 4- $\text{CF}_3\text{-N}^{\wedge}\text{CH}^{\wedge}\text{N}$ Ligand, $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$, and $[\text{R-(N}^{\wedge}\text{C}^{\wedge}\text{N})\text{Pt-Cp}^*\text{Ir-}p\text{-(}\eta^4\text{-C}_6\text{H}_4\text{S}_2\text{)}\text{-Pt(N}^{\wedge}\text{C}^{\wedge}\text{N)-R}][\text{CF}_3\text{SO}_3\text{]}_2$ ($\text{R} = \text{H}$ (5**) and CF_3 (**6a**)).** Convenient crystals of 4- $\text{CF}_3\text{-N}^{\wedge}\text{CH}^{\wedge}\text{N}$ ligand for X-ray analysis were obtained by slow diffusion of diethyl ether into CH_3CN solution of the ligand. The crystal structure of the 2,2'-(5-trifluoromethyl-1,3-phenylene)dipyridine (dpyb- CF_3) shows a head–tail arrangement of the two pyridyl rings in a similar fashion to that reported for the analogous 2,2'-(5-bromo-1,3-phenylene)dipyridine (dpyb-Br) (Figure 3). Moreover, one of the nitrogen atoms displays weak $\text{C-H}\cdots\text{N}$ intermolecular interaction [$\text{C16-H16}\cdots\text{N1} = 3.551(4) \text{ \AA}$] with a carbon of the pyridyl moiety of another ligand which might favor the *transoid* configuration.³⁶ We also note that the three aromatic rings are not coplanar with angles of $3.30(8)$ and $22.53(9)^\circ$ between the two pyridyl units and the central benzene ring, mainly, the pyridyl unit pointing toward the CF_3 - group displaying the larger angle.

Crystals of the starting material $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ were also obtained from DMSO/MeOH mixture. The structure shows that the complex displays a distorted square planar geometry around the platinum center as expected for this type of $\text{N}^{\wedge}\text{C}^{\wedge}\text{N}$ compounds (Figure 3).³⁶ The Pt1-C1 bond distance is $1.909(3) \text{ \AA}$ which is short and typical of such compounds suggesting higher rigidity of the molecular skeleton compared to the related $\text{Pt(N}^{\wedge}\text{N}^{\wedge}\text{C)}$ and $\text{Pt(N}^{\wedge}\text{N}^{\wedge}\text{N)}$ related compounds.³⁷ Examining the crystal packing showed the formation of a dimer species through π - π interaction; however, the $\text{Pt}\cdots\text{Pt}$ distance, $5.086(1) \text{ \AA}$, is long to suggest any interaction.^{8,9,36}

On the other hand, crystals of our target molecules **5** and **6a** were successfully obtained. The crystal structure of **5** shows indeed that the *p*-dithiobenzoquinone iridium complex bridges two $\text{Pt(N}^{\wedge}\text{C}^{\wedge}\text{N)}$ units through the two sulfur centers with Pt-S bond distances of $2.421(2) \text{ \AA}$ (Pt1-S1) and $2.410(2) \text{ \AA}$ (Pt2-S).

Figure 4. (a) View of the cationic part of complex **5**. Selected bond distances (Å) and angles (deg) for **5**: Ir1...C1 = 2.348(4), Ir1...C2 = 2.220(4), Ir1...C3 = 2.218(4), Ir1...C4 = 2.234(4), Ir1...C5 = 2.234(4), Ir1...C6 = 2.219(4), C1...S = 1.720(4), C4...S2 = 1.719(5), S1...Pt1 = 2.421(2), S2...Pt2 = 2.410(2), C7...Pt1 = 1.930(5), C23...Pt2 = 1.919(5), N1-Pt1-N2 = 160.4(2), N3-Pt2-N4 = 160.7(2). (b) 1D supramolecular chain formed through π - π (3.53 Å) and Pt...Pt (3.300 Å) contacts between individual molecules.

Figure 5. (a) View of the cationic part of complex **6a**. Selected bond distances (Å) and angles (deg) for **6a**: Ir1...C1 = 2.328(11), Ir1...C2 = 2.229(10), Ir1...C3 = 2.221(10), Ir1...C4 = 2.316(12), Ir1...C5 = 2.197(11), Ir1...C6 = 2.221(12), C1...S1 = 1.728(11), C4...S2 = 1.744(12), S1...Pt1 = 2.410(5), S2...Pt2 = 2.413(4), C7...Pt1 = 1.917(16), C24...Pt2 = 1.889(17), N1-Pt1-N2 = 161.7(6), N3-Pt2-N4 = 160.8(5). (b) 1D supramolecular chain formed through π - π (3.40 Å), Pt1...Pt1 (3.306(1) Å) and Pt2...Pt2 (3.355(1) Å) contacts between individual molecules.

S2) (Figure 4a). The coordination geometry around each Pt center is a distorted square planar, formed by two nitrogen atoms of the two lateral pyridyl units, one carbon of the central arene ring and one sulfur center from bridging organometallic linker **3**. The Pt1-C7 bond distance is 1.930(5) Å, while Pt2-C23 = 1.919(5) Å, Pt1-N1 = 2.049(4) Å, Pt1-N2 = 2.048(4) Å, Pt2-N3 = 2.037(4) Å, and Pt2-N4 = 2.039(4) Å, respectively. The aromatic ring of the organometallic linker is almost planar, with the thioquinone carbons bent out of the diene plane (hinge angles $\theta_1 = 7.98^\circ$, $\theta_2 = 7.66^\circ$) while acting as a bidentate ligand and connecting two Pt(N⁺C⁻N) units.

These angles are slightly bigger than those reported for the free organometallic linker [Cp*Ir-*p*-(η^4 -C₆H₄S₂)] (**3**). The C-S bond distances in **5** are 1.720(4) and 1.719(4) Å, slightly longer than those reported for **3**.³⁰

The two Pt(N⁺C⁻N) units in **5** are not linearly disposed but are puckered and lean backward with Pt...Pt distance in [(N⁺C⁻N)Pt-Cp*Ir-*p*-(η^4 -C₆H₄S₂)-Pt(N⁺C⁻N)][CF₃SO₃]₂ (**5**) of 7.838(1) Å. Analysis of the crystal packing of **5** reveals important information such as the molecules being stacked in a head-to-tail fashion with a set of π - π /Pt1...Pt2 interactions occurring at each side of one Pt(N⁺C⁻N) arm with the adjacent

Table 1. Photophysical Data for 6a,b and 7a,b

complex	λ_{abs} [nm] (ϵ [dm ³ mol ⁻¹ cm ⁻¹]) ^a	medium (T [K])	λ_{em} [nm] (τ_0 [μ s])	Φ_{lum} ^b
6a	358 (19230), 375 (19180), 400 sh (13650), 474 (750)	acetone (298)	485, 517 (3.9)	0.014
		solid (298)	680 (0.2)	0.004
		solid (77)	625 (0.5)	^c
		thin film (298) ^d	486	0.35
6b	358 (19570), 374 (19720), 400 sh (14130), 474 (750)	acetone (298)	485, 517 (3.5)	0.016
		solid (298)	648 (0.2)	0.002
		solid (77)	623 (0.5)	^c
		thin film (298) ^d	486	0.40
7a	362 sh (18880), 374 (19240), 402 sh (12360), 430 sh (7560)	acetone (298)	485, 517 (4.0)	0.032
		solid (298)	685 (0.2)	0.005
		solid (77)	652 (<0.1)	^c
		thin film (298) ^d	486	0.37
7b	362 sh (19210), 374 (19860), 402 sh (13020), 430 sh (7960)	acetone (298)	485, 517 (3.6)	0.018
		solid (298)	685 (0.3)	0.002
		solid (77)	640 (<0.1)	^c
		thin film (298) ^d	486	0.37

^aMeasured in acetone at 298 K. ^bThe luminescence quantum yield, measured at room temperature using quinine sulfate in 0.5 M H₂SO₄ as the reference (excitation wavelength = 365 nm, Φ_{lum} = 0.55). ^cNot determined. ^dA 5% complex was doped into 1,3-bis(carbazole-9-yl)benzene (MCP), excitation wavelength at 340 nm.

molecule. The Pt1...Pt2 distance is 3.300(1) Å suggesting a strong interaction.⁹ These Pt...Pt/ π - π interactions among individual subunits of **5** construct a 1D supramolecular chain (Figure 4b). Similar examples of Pt(N^{^C}^{^N}) complexes showing a combination of π - π /Pt...Pt interactions among the individual units were previously reported.^{9,16}

The crystal structure of complex **6a** was similar to that **5** suggesting little influence of CF₃ group on the packing. The structure of **6a** shows similar features to that observed for **5**, i.e., the organometallic linker [Cp*Ir-*p*-(η^4 -C₆H₄S₂)] (**3**) holds two 4-CF₃-(N^{^C}^{^N})Pt units through the two sulfur centers (Figure 5a). The 4-CF₃-(N^{^C}^{^N})Pt units also lean backward in an analogous fashion to that of **5** with Pt...Pt distance of 7.922(1) Å. The crystal packing of **6a** also shows the formation of 1D chain as seen for complex **5**; however, head-to-head and tail-to-tail arrangements are visible. Thus, two combinations of π - π /Pt1...Pt1 and π - π /Pt2...Pt2 interactions occur among individual 4-CF₃-(N^{^C}^{^N})Pt units. The crystal structure shows π - π interactions with distances of 3.40 Å and strong metal-metal interactions with Pt1...Pt1 3.306(1) Å and Pt2...Pt2 at 3.355(1) Å, respectively (Figure 5b).⁹ Although we were able to obtain only two molecular structures of the above series with and without the CF₃ functional group, it is obvious that the same OTf counterion provides a similar packing of these molecules in the solid state. We then examined their optical and luminescent properties.

Photophysical Properties of 6a, 6b, 7a and 7b. The photochemistry of all compounds was investigated except for **5** due to its limited stability in solution. The electronic absorption spectra of **6a**, **6b**, **7a** and **7b** in acetone at room temperature showed an intense absorption band at 358–375 nm ($\epsilon > 10^4$ dm³ mol⁻¹ cm⁻¹) with vibrational progression spacings of ca. 1300 cm⁻¹, corresponding to the skeletal vibrational frequency of the N^{^C}^{^N} ligand in the excited state. In addition, a shoulder at 400–430 nm ($\epsilon > 10^3$ dm³ mol⁻¹ cm⁻¹) and weak bands at 446–476 nm ($\epsilon = 140$ –750 dm³ mol⁻¹ cm⁻¹) are observed. The photophysical data are summarized in Table 1,

and the electronic absorption of these complexes are depicted in Figure 6. With reference to previous spectroscopic studies on

Figure 6. Electronic absorption spectra of **6a** (—), **6b** (.....), **7a** (---), and **7b** (-.-) in acetone at room temperature.

the organometallogands Cp*Ir(η^4 -dithiobenzoquinone)¹⁸ and Cp*Ir(η^4 -diselenobenzoquinone)¹⁹ as well as [Pt(N^{^C}^{^N})-Cl],⁷ the high-energy bands are attributed to admixtures of $\pi \rightarrow \pi^*$ intraligand (IL) transitions of Cp*Ir(η^4 -dithiobenzoquinone)/Cp*Ir(η^4 -diselenobenzoquinone) and metal-perturbed IL transitions of the N^{^C}^{^N} ligand, while the shoulders are tentatively assigned as admixtures of $d\pi(\text{Pt}) \rightarrow \pi^*(\text{N}^{\wedge}\text{C}^{\wedge}\text{N})$ metal-to-ligand charge transfer (MLCT) transitions and metal-perturbed $\pi \rightarrow \pi^*$ IL transitions of Cp*Ir(η^4 -dithiobenzoquinone)/Cp*Ir(η^4 -diselenobenzoquinone) systems.

An additional low-energy absorption shoulder at 430 nm is observed for **7a** and **7b**. On the basis of the strong electron-donating behavior of the diselenobenzoquinone group, such a low-energy absorption shoulder is assigned to the $p\pi(\text{Se}) \rightarrow \pi^*(\text{N}^{\wedge}\text{C}^{\wedge}\text{N})$ ligand-to-ligand charge transfer (LLCT) transition with mixing of MLCT character.¹⁹ Compared to their *p*-dithiobenzoquinone analogues, **6a** and **6b**, no such additional low-energy absorption shoulder is observed, probably due to the lower electron-donating ability of the dithiobenzoquinone

counterparts.¹⁹ In contrast, the low-energy weak absorption band at 474 nm for **6a** and **6b** is unlikely to be due to d–d transitions, since the N[^]C[^]N ligand is a strong σ donor and it is unlikely that they would occur at such low energies. In view of the large spin–orbital coupling resulting from the presence of heavy Pt atom, a $^3\pi \rightarrow \pi^*(N^{\wedge}C^{\wedge}N)$ IL transition is suggested.⁷ Similar assignments have also been reported for the related platinum(II) complexes.^{38,39}

Upon excitation at $\lambda > 350$ nm, these complexes show vibronically structured emission with emission maximum of 485 nm in acetone at room temperature (Figure 7). The vibronic

Figure 7. Normalized emission spectra of **6b** (—) and **7b** (---) in acetone at room temperature.

progression of 1300 cm^{-1} is in agreement with the skeletal vibrational frequency of the N[^]C[^]N ligand in the ground state. The pronounced vibronic structure is typical of emission from a $\pi \rightarrow \pi^*$ excited state, indicating that the emission originates from a state of primarily $^3\text{IL } \pi \rightarrow \pi^*$ character. The relatively long lifetimes of several microseconds are typical of the phosphorescence of complexes of this class. These emissions, which are also observed in the related Pt(N[^]C[^]N) complexes are tentatively assigned as derived from states of metal-perturbed $^3\text{IL } \pi \rightarrow \pi^*(N^{\wedge}C^{\wedge}N)$ origin.^{38,39} Solid-state emission of cyclometalated Pt(II) complexes could be significantly affected by the packing of molecules in their crystal structures.^{40,41} Figure 8 shows the normalized emission spectra

Figure 8. Normalized emission spectra of **6b** (—) and **7b** (---) in the solid state at 77 K.

of **6b** and **7b** in the solid state at 77 K. Compounds **6a**, **6b**, **7a** and **7b** were found to give emission at 623–685 nm in the solid state. The structureless emission is attributed to $^3\text{MMLCT}$ excited states associated with Pt...Pt interactions,⁴² which are further confirmed by the solid-state packing of **6a** in the X-ray crystallography. Higher luminescence quantum yields of up to 40% are observed in the thin film samples of the complexes doped in 1,3-bis(carbazole-9-yl)benzene (MCP), which could

be attributed to the reduced nonradiative decay as a result of the increased rigidity in the film.

■ CONCLUDING REMARKS

In conclusion, we reported the synthesis of the first series of binuclear Pt(N[^]C[^]N) compounds with π -bonded thio- and seleno-quinone as bridging ligands. The organic ligand 4-CF₃-N[^]CH[^]N and two of the platinum(II) complexes have also been structurally characterized by X-ray crystallography. Vibronic-structured emission bands in the bluish-green region have been observed for these platinum(II) complexes that are indicative of the involvement of the cyclometalated ligand in the excited state in solution at room temperature. Interestingly, these complexes are red emitters in their solid state, and their photophysical properties are attributed to $^3\text{MMLCT}$ excited states that are associated with Pt...Pt interactions, which have been further confirmed by the solid-state packing of **6a** in the X-ray crystallography. These interesting behaviors might open up a new strategy to develop new photoluminescent materials.

■ EXPERIMENTAL SECTION

All experimental manipulations were carried out under argon using Schlenk tube techniques except the chlorides/anions metathesis workup which was performed under aerobic conditions. All solvents were purified by standard techniques. The ¹H NMR and ¹³C NMR spectra were recorded in CD₃NO₂ using a Bruker Avance 300 NMR or 400 NMR spectrometer. Infrared spectra were recorded from neat samples on a Bruker FT-IR spectrometer Tensor 27 equipped with an Attenuated Total Reflection (ATR) Harrick accessory. The organometallic linkers $\eta^4\text{-[Cp}^*\text{Ir}(\eta^5\text{-C}_6\text{H}_4\text{S}_2\text{)]}$,³⁰ $\eta^4\text{-[Cp}^*\text{Ir}(\eta^5\text{-C}_6\text{H}_4\text{Se}_2\text{)]}$ ³⁴ (**2**), [Pt(N[^]C[^]N)Cl], and [(4-CF₃-N[^]C[^]N)Pt–Cl] were prepared according to published procedures.^{14,35} UV–visible spectra were obtained on a Hewlett-Packard 8452A diode array spectrophotometer. Steady-state excitation and emission spectra in the solid state (298 and 77 K) and in solution at room temperature were recorded on a Spex Fluorolog-3 Model FL3-211 fluorescence spectrofluorometer equipped with a R2658P PMT detector. Solid-state photophysical studies were carried out with solid samples contained in a quartz tube inside a quartz-walled Dewar flask. Measurements of the solid-state sample at 77 K were similarly conducted with liquid nitrogen filled in the optical Dewar flask. All solutions for photophysical studies were degassed on a high-vacuum line in a two-compartment cell consisting of a 10 mL Pyrex bulb and a 1 cm path length quartz cuvette and sealed from the atmosphere by a Bibby Rotaflo HP6 Teflon stopper. The solutions were rigorously degassed with at least four successive freeze–pump–thaw cycles. Emission lifetime measurements were performed using a conventional laser system. The excitation source was the 355 nm output (third harmonic) of a Spectra-Physics Quanta-Ray Q-switched GCR-150–10 pulsed Nd:YAG laser. Luminescence decay signals were recorded on a Tektronix model TDS620A digital oscilloscope and analyzed using a program for exponential fits. Relative luminescence quantum yields were measured by the optical dilute method reported by Demas and Crosby.⁴³ The solid and thin films at room temperature were measured on a Hamamatsu C9920–03 Absolute Photoluminescence Yield Measurement System.

Synthesis of [(N[^]C[^]N)₂Pt₂{Cp^{*}Ir- η^4 -(p-C₆H₄S₂)}][OTf]₂ (5**).** To a solution of [(N[^]C[^]N)PtCl] (90 mg, 0.19 mmol) in acetone (30 mL) was added a solution of AgOTf (50 mg, 0.19 mmol), and the mixture was stirred for 18 h. The resulting solution was filtered into a dry Schlenk tube kept under argon, and the AgCl salt was removed. To this solution was added p-[Cp^{*}Ir(η^4 -C₆H₄S₂)] (**3**) (50 mg, 0.11 mmol) in CH₂Cl₂ (10 mL) and the mixture was stirred for 2 h at room temperature. The solvent was then reduced to roughly 5 mL and distilled Et₂O (30 mL) was added. The orange-red precipitate was recovered on a glass frit and washed three times with Et₂O (15 mL), then dried under vacuum to give [(N[^]C[^]N)₂Pt₂{Cp^{*}Ir- η^4 -(p-C₆H₄S₂)}][OTf]₂ as an orange microcrystalline solid (48 mg, 0.03

mmol). Yield: 31%. The compound was not very stable; hence, no elemental analyses were obtained. X-ray molecular structure was determined. ^1H NMR (300.13 MHz, CD_3NO_2): δ 9.01 (d, $J = 5.8$ Hz, $^3J_{\text{Pt-H}} = 34$ Hz, 4H; H_a), 8.17 (td, $J = 7.7$ and 1.4 Hz, 4H; H_c), 7.94 (dd, $J = 6.9$ and 0.6 Hz, 4H; H_d), 7.65 (d, $J = 7.7$ Hz, 4H; H_e), 7.39 (td, $J = 6.1$ and 2.0 Hz, 4H; H_b), 7.36 (t, $J = 7.4$ Hz, 2H; H_f), 6.98 (s, 4H; arene), 1.98 (s, 15H; Cp^*). ^{13}C $\{^1\text{H}\}$ NMR (75.4 MHz, CD_3NO_2): δ 166.6, 151.0, 150.3, 146.4, 141.4, 140.1, 139.2, 130.8, 123.3, 119.3, 99.9, 90.8, 6.6. IR (neat ATR Harricks, cm^{-1}): $\nu(\text{CF}_3\text{SO}_3)$ 1254; 1025.

Synthesis of $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})_2\text{Pt}_2\{\text{Cp}^*\text{Ir-}\eta^4\text{-(p-C}_6\text{H}_4\text{S}_2)\}][\text{OTf}]_2$ (6a). This compound was prepared following the procedure described for **5** but using $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ instead of $[(\text{N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$.

6a was obtained as orange microcrystalline solid (149 mg, 0.084 mmol). Yield: 90% starting with $[(\text{CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ (100 mg, 0.186 mmol), AgOTf (48 mg, 0.186 mmol), and $p\text{-}[\text{Cp}^*\text{Ir}(\eta^4\text{-C}_6\text{H}_4\text{S}_2)]$ (**3**) (43 mg, 0.093 mmol). Anal. Calcd for $\text{C}_{52}\text{H}_{39}\text{F}_{12}\text{IrN}_4\text{O}_6\text{Pt}_2\text{S}_4\cdot\text{CH}_2\text{Cl}_2$: C, 34.60; H, 2.23; N, 3.05. Found: C, 34.25; H, 2.24; N, 3.13. ^1H NMR (400 MHz, CD_3NO_2): δ 8.87 (d, $J = 6.0$ Hz, $^3J_{\text{Pt-H}} = 34.1$ Hz, 4H; H_c), 8.17 (td, $J = 7.8$ and 1.2 Hz, 4H; H_c), 7.85 (d, $J = 8.0$ Hz, 4H; H_d), 7.71 (s, 4H; H_e), 7.42 (td, $J = 7.2$ and 1.2 Hz, 4H; H_b), 6.85 (s, 4H; arene), 1.85 (s, 15H; Cp^*). ^{13}C NMR $\{^1\text{H}\}$ (100.61 MHz, CD_3NO_2): δ 165.9, 152.3, 141.4, 140.8, 126.1, 125.6, 123.4, 121.7 (q, $^1J_{\text{C-F}} = 192.8$ Hz), 121.1, 119.5, 104.3, 101.2, 92.0, 7.8. ^{19}F NMR (282 MHz, CD_3NO_2): δ -63.9 (6F; $-\text{CF}_3$), -80.7 (6F; OTf). IR (neat ATR Harricks, cm^{-1}): $\nu(\text{CF}_3\text{SO}_3)$ 1256; 1027.

Synthesis of $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})_2\text{Pt}_2\{\text{Cp}^*\text{Ir-}\eta^4\text{-(p-C}_6\text{H}_4\text{S}_2)\}][\text{SbF}_6]_2$ (6b). This compound was prepared following the general procedure described for **6a** but using AgSbF_6 instead of AgOTf .

6b was obtained as orange microcrystalline solid (158 mg, 0.082 mmol). Yield: 88% starting with $[(\text{CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ (100 mg, 0.186 mmol), AgSbF_6 (63 mg, 0.186 mmol), and $p\text{-}[\text{Cp}^*\text{Ir}(\eta^4\text{-C}_6\text{H}_4\text{S}_2)]$ (**3**) (43 mg, 0.093 mmol). Anal. Calcd for $\text{C}_{50}\text{H}_{39}\text{SbF}_{18}\text{IrN}_4\text{Pt}_2\text{S}_2\cdot 1/2(\text{CH}_3)_2\text{CO}$: C, 31.61; H, 2.14; N, 2.86. Found: C, 31.75; H, 2.04; N, 2.80. ^1H NMR (400.13 MHz, CD_3NO_2): δ 9.00 (d, $J = 5.6$ Hz, $^3J_{\text{Pt-H}} = 32.6$ Hz, 4H; H_a), 8.23 (td, $J = 7.2$ and 1.2 Hz, 4H; H_c), 8.00 (d, $J = 8.0$ Hz, 4H; H_d), 7.87 (s, 4H; H_e), 7.48 (td, $J = 6.8$ and 2.0 Hz, 4H; H_b), 6.94 (s, 4H; arene), 1.93 (s, 15H; Cp^*). ^{13}C $\{^1\text{H}\}$ NMR (75.4 MHz, CD_3NO_2): δ 165.9, 152.2, 141.3, 140.8, 139.2, 126.3, 126.1, 126.0, 125.5, 123.4, 121.1, 120.8, 92.0, 7.9. ^{19}F NMR (375 MHz, CD_3NO_2): δ -63.9 (6F; $-\text{CF}_3$). IR (neat ATR Harricks, cm^{-1}): $\nu(\text{SbF}_6)$ 651.

Synthesis of $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})_2\text{Pt}_2\{\text{Cp}^*\text{Ir-}\eta^4\text{-(p-C}_6\text{H}_4\text{S}_2)\}][\text{OTf}]_2$ (7a). This compound was prepared following the general procedure described for **6a** but using the organometallic ligand $p\text{-}[\text{Cp}^*\text{Ir}(\eta^4\text{-C}_6\text{H}_4\text{S}_2)]$ (**4**).

7a was obtained as a burgundy microcrystalline solid (159 mg, 0.086 mmol). Yield: 91%, starting with $[(\text{CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ (100 mg, 0.186 mmol), AgOTf (48 mg, 0.186 mmol), and $p\text{-}[\text{Cp}^*\text{Ir}(\eta^4\text{-C}_6\text{H}_4\text{S}_2)]$ (**4**) (52 mg, 0.093 mmol). Anal. Calcd for $\text{C}_{52}\text{H}_{39}\text{F}_{12}\text{IrN}_4\text{O}_6\text{Pt}_2\text{S}_2\cdot 2\text{CH}_2\text{Cl}_2$: C, 32.14; H, 2.15; N, 2.78. Found: C, 31.90; H, 2.10; N, 2.74. ^1H NMR (300.13 MHz, CD_3NO_2): δ 9.27 (d, $J = 5.2$ Hz, $^3J_{\text{Pt-H}} = 33.8$ Hz, 4H; H_a), 8.23 (t, $J = 7.8$ Hz, 4H; H_c), 8.03 (d, $J = 7.2$ Hz, 4H; H_d), 7.91 (s, 4H; H_e), 7.48 (t, $J = 7.0$ Hz, 4H; H_b), 7.08 (s, 4H; arene), 1.93 (s, 15H; Cp^*). ^{13}C $\{^1\text{H}\}$ NMR (75.4 MHz, CD_3NO_2): δ 152.5, 140.3, 140.0, 139.5, 124.7, 121.1, 120.0, 119.8, 118.9, 116.9, 6.4. ^{19}F NMR (375 MHz, CD_3NO_2): δ -64.0 (6F; $-\text{CF}_3$), -80.7 (6F; OTf). IR (neat ATR Harricks, cm^{-1}): $\nu(\text{CF}_3\text{SO}_3)$ 1251; 1025.

Synthesis of $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})_2\text{Pt}_2\{\text{Cp}^*\text{Ir-}\eta^4\text{-(p-C}_6\text{H}_4\text{S}_2)\}][\text{SbF}_6]_2$ (7b). This compound was prepared following the general procedure described for **7a** but using AgSbF_6 instead of AgOTf .

7b was obtained as burgundy microcrystalline solid (148 mg, 0.073 mmol). Yield: 78%, starting with $[(\text{CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$ (100 mg, 0.186 mmol), AgSbF_6 (63 mg, 0.186 mmol), and $p\text{-}[\text{Cp}^*\text{Ir}(\eta^4\text{-C}_6\text{H}_4\text{S}_2)]$ (**4**) (52 mg, 0.093 mmol). Anal. Calcd for $\text{C}_{50}\text{H}_{39}\text{SbF}_{18}\text{IrN}_4\text{Pt}_2\text{S}_2\cdot\text{CH}_2\text{Cl}_2$: C, 29.07; H, 1.94; N, 2.65. Found: C, 29.27; H, 1.96; N, 2.70. ^1H NMR (300.13 MHz, CD_3NO_2): δ 9.12 (d, $J = 5.6$ Hz, $^3J_{\text{Pt-H}} = 31.1$ Hz, 4H; H_a), 8.14 (t, $J = 8.0$ Hz, 4H; H_c), 7.85 (d, $J = 8.4$ Hz, 4H; H_d), 7.72 (s, 4H; H_e), 7.35 (t, $J = 7.2$ Hz, 4H; H_b), 6.96 (s, 4H; arene), 1.85

(s, 15H; Cp^*). ^{13}C $\{^1\text{H}\}$ NMR (75.4 MHz, CD_3NO_2): δ 164.9, 152.5, 140.4, 139.6, 124.6, 119.9, 119.5, 100.6, 94.9, 6.3. ^{19}F NMR (375 MHz, CD_3NO_2): δ -64.0 (6F; $-\text{CF}_3$). IR (neat ATR Harricks, cm^{-1}): $\nu(\text{SbF}_6)$ 635.

X-ray Crystal Structure Determination. Single crystals were selected, mounted, and transferred into a cold nitrogen gas stream. Intensity data was collected with Bruker Kappa-APEX2 systems using microsource Cu $K\alpha$ for $4\text{-CF}_3\text{-N}^{\wedge}\text{CH}^{\wedge}\text{N}$ or fine-focus sealed tube Mo $K\alpha$ radiation for $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{Pt-Cl}]$, **5**, **6a**. Unit-cell parameters determination, data collection strategy, integration, and absorption correction were carried out with the Bruker APEX2 suite of programs. The structures were solved with SHELXT-2014⁴⁴ [$(4\text{-CF}_3\text{-N}^{\wedge}\text{CH}^{\wedge}\text{N})$, $(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{Pt-Cl}$] or SIR92⁴⁵ (**5**, **6a**) and refined anisotropically by full-matrix least-squares methods with SHELXL-2014⁴⁴ using the WinGX suite.⁴⁶ The structures were deposited at the Cambridge Crystallographic Data Centre with numbers CCDC 1538568 for ligand $(4\text{-CF}_3\text{-N}^{\wedge}\text{CH}^{\wedge}\text{N})$, CCDC 1538569 for complex $(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{Pt-Cl}$, CCDC 1538570 for **5**, and CCDC 1538571 for **6a** and can be obtained free of charge via www.ccdc.cam.ac.uk.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.organo- met.7b00680.

Crystallographic data of $4\text{-CF}_3\text{-N}^{\wedge}\text{CH}^{\wedge}\text{N}$, $[(4\text{-CF}_3\text{-N}^{\wedge}\text{C}^{\wedge}\text{N})\text{PtCl}]$, **5**, and **6a**. ^1H and ^{13}C NMR spectra of compounds **5**, **6a**, **6b**, **7a** and **7b** (PDF)

Accession Codes

CCDC 1538568–1538571 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: hani.amouri@upmc.fr.

*E-mail: wvyam@hku.hk. Tel.: (+852) 2859-2153. Fax: (+852) 2857-1586.

ORCID

Vivian Wing-Wah Yam: 0000-0001-8349-4429

Hani Amouri: 0000-0001-6955-8406

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the CNRS, by the Université Pierre et Marie Curie-Paris 6 and by French ANR grant (project OPTOELECTR-OM ANR-11-BS07-001-01), which we gratefully acknowledged. V.W.W.Y. acknowledges support from the University Grants Committee Areas of Excellence Scheme (AoE/P-03/08).

■ REFERENCES

- (1) Williams, J. A. G. Photochemistry and photophysics of coordination compounds: Platinum. *Photochemistry and Photophysics of Coordination Compounds Ii* 2007, 281, 205–268.
- (2) Williams, J. A. G. *Chem. Soc. Rev.* 2009, 38, 1783–1801.
- (3) Tarran, W. A.; Freeman, G. R.; Murphy, L.; Benham, A. M.; Katay, R.; Williams, J. A. G. *Inorg. Chem.* 2014, 53, 5738–5749.
- (4) Kitano, M.; Tanaka, T.; Osuka, A. *Organometallics* 2017, 36, 2559–2564.

- (5) Sivchik, V. V.; Grachova, E. V.; Melnikov, A. S.; Smirnov, S. N.; Ivanov, A. Y.; Hirva, P.; Tunik, S. P.; Koshevoy, I. O. *Inorg. Chem.* **2016**, *55*, 3351–3363.
- (6) Gong, Z.-L.; Zhong, Y.-W. *Inorg. Chem.* **2016**, *55*, 10143–10151.
- (7) Williams, J. A. G.; Beeby, A.; Davies, E. S.; Weinstein, J. A.; Wilson, C. *Inorg. Chem.* **2003**, *42*, 8609–8611.
- (8) Rossi, E.; Colombo, A.; Dragonetti, C.; Roberto, D.; Demartin, F.; Cocchi, M.; Brulatti, P.; Fattori, V.; Williams, J. A. G. *Chem. Commun.* **2012**, *48*, 3182–3184.
- (9) Wang, Z. X.; Turner, E.; Mahoney, V.; Madakuni, S.; Groy, T.; Li, J. A. *Inorg. Chem.* **2010**, *49*, 11276–11286.
- (10) Rossi, E.; Murphy, L.; Brothwood, P. L.; Colombo, A.; Dragonetti, C.; Roberto, D.; Ugo, R.; Cocchi, M.; Williams, J. A. G. *J. Mater. Chem.* **2011**, *21*, 15501–15510.
- (11) Tam, A. Y. Y.; Tsang, D. P. K.; Chan, M. Y.; Zhu, N. Y.; Yam, V. W. *Chem. Commun.* **2011**, *47*, 3383–3385.
- (12) Lam, E. S. H.; Tsang, D. P. K.; Lam, W. H.; Tam, A. Y. Y.; Chan, M. Y.; Wong, W. T.; Yam, V. W. W. *Chem. - Eur. J.* **2013**, *19*, 6385–6397.
- (13) Huckaba, A. J.; Cao, B.; Hollis, T. K.; Valle, H. U.; Kelly, J. T.; Hammer, N. I.; Oliver, A. G.; Webster, C. E. *Dalton Trans.* **2013**, *42*, 8820–8826.
- (14) Lu, W.; Chen, Y.; Roy, V. A. L.; Chui, S. S. Y.; Che, C. M. *Angew. Chem., Int. Ed.* **2009**, *48*, 7621–7625.
- (15) Li, K.; Chen, Y.; Lu, W.; Zhu, N. Y.; Che, C. M. *Chem. - Eur. J.* **2011**, *17*, 4109–4112.
- (16) Chen, Y.; Lu, W.; Che, C. M. *Organometallics* **2013**, *32*, 350–353.
- (17) Wang, Z.; Sun, Z.; Hao, X.-Q.; Niu, J.-L.; Wei, D.; Tu, T.; Gong, J.-F.; Song, M.-P. *Organometallics* **2014**, *33*, 1563–1573.
- (18) Moussa, J.; Wong, K. M.-C.; Chamoreau, L.-M.; Amouri, H.; Yam, V. W.-W. *Dalton Transactions* **2007**, 3526–3530.
- (19) Moussa, J.; Wong, K. M.-C.; Le Goff, X. F.; Rager, M. N.; Chan, C. K.-M.; Yam, V. W.-W.; Amouri, H. *Organometallics* **2013**, *32*, 4985–4992.
- (20) Suzuki, R.; Matsumoto, K.; Kurata, H.; Oda, M. *Chem. Commun.* **2000**, 1357–1358.
- (21) Wirth, T. *Organoselenium Chemistry: Modern Developments*. In *Organic Synthesis*; Springer: Berlin, 2000; Vol. 208.
- (22) Beer, L.; Reed, R. W.; Robertson, C. M.; Oakley, R. T.; Tham, F. S.; Haddon, R. C. *Org. Lett.* **2008**, *10*, 3121–3123.
- (23) Le Bras, J.; Amouri, H.; Vaissermann, J. *Organometallics* **1998**, *17*, 1116–1121.
- (24) Amouri, H.; Le Bras, J. *Acc. Chem. Res.* **2002**, *35*, 501–510.
- (25) Moussa, J.; Guyard-Duhayon, C.; Herson, P.; Amouri, H.; Rager, M. N.; Jutand, A. *Organometallics* **2004**, *23*, 6231–6238.
- (26) Moussa, J.; Chamoreau, L. M.; Boubekour, K.; Amouri, H.; Rager, M. N.; Grotjahn, D. B. *Organometallics* **2008**, *27*, 67–71.
- (27) Amouri, H.; Besace, Y.; Le Bras, J.; Vaissermann, J. *J. Am. Chem. Soc.* **1998**, *120*, 6171–6172.
- (28) Amouri, H.; Vaissermann, J.; Rager, M. N.; Grotjahn, D. B. *Organometallics* **2000**, *19*, 1740–1748.
- (29) Amouri, H.; Vaissermann, J.; Rager, M. N.; Grotjahn, D. B. *Organometallics* **2000**, *19*, 5143–5148.
- (30) Moussa, J.; Lev, D. A.; Boubekour, K.; Rager, M. N.; Amouri, H. *Angew. Chem., Int. Ed.* **2006**, *45*, 3854–3858.
- (31) Moussa, J.; Rager, M. N.; Boubekour, K.; Amouri, H. *Eur. J. Inorg. Chem.* **2007**, *2007*, 2648–2653.
- (32) Moussa, J.; Amouri, H. *Angew. Chem., Int. Ed.* **2008**, *47*, 1372–1380.
- (33) Moussa, J.; Rager, M. N.; Chamoreau, L. M.; Ricard, L.; Amouri, H. *Organometallics* **2009**, *28*, 397–404.
- (34) Amouri, H.; Moussa, J.; Renfrew, A. K.; Dyson, P. J.; Rager, M. N.; Chamoreau, L.-M. *Angew. Chem., Int. Ed.* **2010**, *49*, 7530–7533.
- (35) Cardenas, D. J.; Echavarren, A. M.; de Arellano, M. C. R. *Organometallics* **1999**, *18*, 3337–3341.
- (36) Farley, S. J.; Rochester, D. L.; Thompson, A. L.; Howard, J. A. K.; Williams, J. A. G. *Inorg. Chem.* **2005**, *44*, 9690–9703.
- (37) Hofmann, A.; Dahlenburg, L.; van Eldik, R. *Inorg. Chem.* **2003**, *42*, 6528–6538.
- (38) Connick, W. B.; Miskowski, V. M.; Houlding, V. H.; Gray, H. B. *Inorg. Chem.* **2000**, *39*, 2585–2592.
- (39) Mroz, W.; Botta, C.; Giovannella, U.; Rossi, E.; Colombo, A.; Dragonetti, C.; Roberto, D.; Ugo, R.; Valore, A.; Williams, J. A. G. *J. Mater. Chem.* **2011**, *21*, 8653–8661.
- (40) Lai, S. W.; Che, C. M. *Top. Curr. Chem.* **2004**, *241*, 27–63.
- (41) Wong, K. M. C.; Yam, V. W. W. *Acc. Chem. Res.* **2011**, *44*, 424–434.
- (42) Cheng, G.; Chen, Y.; Yang, C.; Lu, W.; Che, C.-M. *Chem. - Asian J.* **2013**, *8*, 1754–1759.
- (43) Demas, J. N.; Crosby, G. A. *J. Phys. Chem.* **1971**, *75*, 991–1024.
- (44) Sheldrick, G. M. *Acta Crystallogr., Sect. A: Found. Crystallogr.* **2008**, *64*, 112–122.
- (45) Altomare, A.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A. *J. Appl. Crystallogr.* **1993**, *26*, 343–350.
- (46) Farrugia, L. J. *J. Appl. Crystallogr.* **1999**, *32*, 837–838.