



$W^{2,p}$ -estimates for surfaces in terms of their two fundamental forms

Philippe G. Ciarlet, Cristinel Mardare

► To cite this version:

Philippe G. Ciarlet, Cristinel Mardare. $W^{2,p}$ -estimates for surfaces in terms of their two fundamental forms. Comptes Rendus. Mathématique, 2018, 356 (1), pp.85-91. 10.1016/j.crma.2017.12.003 . hal-01727386

HAL Id: hal-01727386

<https://hal.sorbonne-universite.fr/hal-01727386>

Submitted on 9 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Differential geometry/Mathematical problems in mechanics

$W^{2,p}$ -estimates for surfaces in terms of their two fundamental forms



Estimations dans $W^{2,p}$ pour des surfaces à partir de leurs deux formes fondamentales

Philippe G. Ciarlet^a, Cristinel Mardare^b^a Department of Mathematics, City University of Hong Kong, 83 Tat Chee Avenue, Kowloon, Hong Kong^b Sorbonne Universités, Université Pierre-et-Marie-Curie, Laboratoire Jacques-Louis-Lions, 75005 Paris, France

ARTICLE INFO

Article history:

Received 7 December 2017

Accepted 7 December 2017

Available online 19 December 2017

Presented by Philippe G. Ciarlet

ABSTRACT

Let $p > 2$. We show how the fundamental theorem of surface theory for surfaces of class $W_{\text{loc}}^{2,p}(\omega)$ over a simply-connected open subset of \mathbb{R}^2 established in 2005 by S. Mardare can be extended to surfaces of class $W^{2,p}(\omega)$ when ω is in addition bounded and has a Lipschitz-continuous boundary. Then we establish a nonlinear Korn inequality for surfaces of class $W^{2,p}(\omega)$. Finally, we show that the mapping that defines in this fashion a surface of class $W^{2,p}(\omega)$, unique up to proper isometries of \mathbb{E}^3 , in terms of its two fundamental forms is locally Lipschitz-continuous.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

R É S U M É

Soit $p > 2$. Nous montrons comment le théorème fondamental de la théorie des surfaces de classe $W_{\text{loc}}^{2,p}(\omega)$ sur un ouvert simplement connexe ω de \mathbb{R}^2 établi par S. Mardare en 2005 peut être étendu à des surfaces de classe $W^{2,p}(\omega)$ lorsque ω est de plus borné et de frontière lipschitzienne. Ensuite, nous établissons une inégalité de Korn non linéaire pour des surfaces de classe $W^{2,p}(\omega)$. Nous établissons enfin que l'application qui définit une surface de classe $W^{2,p}(\omega)$ à une isométrie propre de \mathbb{E}^3 près en fonction de ses deux formes fondamentales est localement lipschitzienne.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Preliminaries

In what follows, Greek indices and exponents, except ε and δ , vary in the set $\{1, 2\}$, Latin indices vary in the set $\{1, 2, 3\}$, and the summation convention for repeated indices and exponents is used. Boldface letters denote vector and matrix fields.

E-mail addresses: mapgc@cityu.edu.hk (P.G. Ciarlet), mardare@ann.jussieu.fr (C. Mardare).

<https://doi.org/10.1016/j.crma.2017.12.003>

1631-073X/© 2017 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

The three-dimensional Euclidean space is denoted \mathbb{E}^3 . The inner product, exterior product, and norm, in \mathbb{E}^3 are respectively denoted \cdot , \wedge , and $|\cdot|$. The set of all proper isometries of \mathbb{E}^3 is denoted and defined by

$$\mathbf{Isom}_+(\mathbb{E}^3) := \{\mathbf{r} : \mathbb{E}^3 \rightarrow \mathbb{E}^3, \mathbf{r}(x) = \mathbf{R}x + \mathbf{a}, x \in \mathbb{E}^3; \mathbf{R} \in \mathbb{O}_+^3, \mathbf{a} \in \mathbb{E}^3\},$$

where \mathbb{O}_+^3 denotes the set of all real 3×3 proper orthogonal matrices.

Remark 1. The set $\mathbf{Isom}_+(\mathbb{E}^3)$ is in effect a smooth submanifold of dimension six of the space of all 3×3 real matrices and its tangent space at the identity mapping $\mathbf{id} \in \mathbf{Isom}_+(\mathbb{E}^3)$ is the space of all “infinitesimal rigid displacements of \mathbb{E}^3 ”, which is denoted and defined by

$$\mathbf{Rig}(\mathbb{E}^3) = \mathcal{T}_{\mathbf{id}}\mathbf{Isom}_+(\mathbb{E}^3) := \{\boldsymbol{\zeta} : \mathbb{E}^3 \rightarrow \mathbb{E}^3, \boldsymbol{\zeta}(x) = \mathbf{A}x + \mathbf{b}, x \in \mathbb{E}^3; \mathbf{A} \in \mathbb{A}^3, \mathbf{b} \in \mathbb{E}^3\},$$

where \mathbb{A}^3 denotes the set of all real 3×3 antisymmetric matrices. \square

Given an open subset ω of \mathbb{R}^2 , we let $y = (y_\alpha)$ denote a generic point in ω , and we let $\partial_\alpha := \partial/\partial y_\alpha$ and $\partial_{\alpha\beta} := \partial^2/\partial y_\alpha \partial y_\beta$.

The space of distributions over an open subset ω of \mathbb{R}^2 is denoted $\mathcal{D}'(\omega)$. For each integer $m \geq 1$ and each real number $p \geq 1$, $C^m(\omega)$ denotes the subspace of $C^0(\omega)$ of functions that possess continuous partial derivatives up to order m , and $W^{m,p}(\omega)$ denotes the usual Sobolev space.

The notation $L_{\text{loc}}^p(\omega)$, resp. $W_{\text{loc}}^{m,p}(\omega)$, denotes the space of functions $f : \omega \rightarrow \mathbb{R}$ such that $f|_U \in L^p(U)$, resp. $f|_U \in W^{m,p}(U)$, for all open sets $U \Subset \omega$, where $f|_U$ denotes the restriction of f to U and the notation $U \Subset \omega$ means that the closure of the set U is a compact subset of ω . Given any finite dimensional real space \mathbb{Y} , the notation $L_{\text{loc}}^p(\omega; \mathbb{Y})$, resp. $W_{\text{loc}}^{1,p}(\omega; \mathbb{Y})$, denotes the space of \mathbb{Y} -valued fields with components in $L_{\text{loc}}^p(\omega)$, resp. $W_{\text{loc}}^{1,p}(\omega)$. Other similar notations with self-explanatory definitions will be used.

An immersion from ω into \mathbb{E}^3 is a smooth enough mapping $\boldsymbol{\theta} : \omega \rightarrow \mathbb{E}^3$ such that the two vector fields $\partial_\alpha \boldsymbol{\theta} : \omega \rightarrow \mathbb{E}^3$ are linearly independent at each point of ω . Given an immersion $\boldsymbol{\theta} : \omega \rightarrow \mathbb{E}^3$, define the functions

$$\hat{a}_{\alpha\beta}(\boldsymbol{\theta}) := \hat{\mathbf{a}}_\alpha(\boldsymbol{\theta}) \cdot \hat{\mathbf{a}}_\beta(\boldsymbol{\theta}) \text{ and } \hat{b}_{\alpha\beta}(\boldsymbol{\theta}) := \partial_\alpha \hat{\mathbf{a}}_\beta(\boldsymbol{\theta}) \cdot \hat{\mathbf{a}}_3(\boldsymbol{\theta}),$$

where

$$\hat{\mathbf{a}}_\alpha(\boldsymbol{\theta}) := \partial_\alpha \boldsymbol{\theta} \text{ and } \hat{\mathbf{a}}_3(\boldsymbol{\theta}) := \frac{\partial_1 \boldsymbol{\theta} \wedge \partial_2 \boldsymbol{\theta}}{|\partial_1 \boldsymbol{\theta} \wedge \partial_2 \boldsymbol{\theta}|}.$$

The image $S = \boldsymbol{\theta}(\omega)$ is thus a surface in \mathbb{E}^3 and the functions $\hat{a}_{\alpha\beta}(\boldsymbol{\theta})$ and $\hat{b}_{\alpha\beta}(\boldsymbol{\theta})$ are the covariant components of the first and second fundamental forms of S .

The space of real 2×2 symmetric matrices is denoted \mathbb{S}^2 ; its subset formed by all positive-definite matrices is denoted $\mathbb{S}_{>}^2$.

An open subset ω of \mathbb{R}^2 satisfies the *uniform interior cone property* if there exists a bounded open cone $V \subset \mathbb{R}^2$ such that any point $y \in \omega$ is the vertex of a cone V_y congruent with V and contained in ω . An open subset ω of \mathbb{R}^2 is a *domain* if it is bounded and has a Lipschitz-continuous boundary.

Detailed proofs of the results announced here will be found in [4].

2. The fundamental theorem of surface theory in the spaces $W_{\text{loc}}^{2,p}(\omega)$ and $W^{2,p}(\omega)$

The *fundamental theorem of surface theory*, which is classically established in the spaces of continuously differentiable functions (cf., e.g., [5, Theorem 3.8.8], [1, Appendix to Chapter 4], [2, Theorems 8.16-1 and 8.17-1]), has been shown to hold in function spaces with little regularity, according to the following remarkable result, due to S. Mardare [6, Theorem 9]:

Theorem 1. Let ω be a simply-connected open subset of \mathbb{R}^2 , let $p > 2$, and let a matrix field $(a_{\alpha\beta}) \in W_{\text{loc}}^{1,p}(\omega; \mathbb{S}_{>}^2)$ and a matrix field $(b_{\alpha\beta}) \in L_{\text{loc}}^p(\omega; \mathbb{S}^2)$ be given that satisfy the Gauss and Codazzi–Mainardi equations, viz.

$$R_{\alpha\beta\tau}^\sigma := \partial_\tau \Gamma_{\alpha\beta}^\sigma - \partial_\beta \Gamma_{\alpha\tau}^\sigma + \Gamma_{\alpha\beta}^\gamma \Gamma_{\tau\gamma}^\sigma - \Gamma_{\alpha\tau}^\gamma \Gamma_{\beta\gamma}^\sigma - b_{\alpha\beta} b_\tau^\sigma + b_{\alpha\tau} b_\beta^\sigma = 0 \text{ in } \mathcal{D}'(\omega)$$

and

$$R_{\alpha\beta\tau}^3 := \partial_\tau b_{\alpha\beta} - \partial_\beta b_{\alpha\tau} + \Gamma_{\alpha\beta}^\gamma b_{\tau\gamma} - \Gamma_{\alpha\tau}^\gamma b_{\beta\gamma} = 0 \text{ in } \mathcal{D}'(\omega),$$

where the functions $\Gamma_{\alpha\beta}^\sigma \in L_{\text{loc}}^p(\omega)$ and $b_\alpha^\sigma \in L_{\text{loc}}^p(\omega)$ are defined by

$$\Gamma_{\alpha\beta}^\sigma := \frac{1}{2} a^{\sigma\tau} (\partial_\alpha a_{\beta\tau} + \partial_\beta a_{\alpha\tau} - \partial_\tau a_{\alpha\beta}) \text{ and } b_\beta^\sigma := a^{\sigma\tau} b_{\tau\beta}, \text{ where } (a^{\sigma\tau}) := (a_{\alpha\beta})^{-1}.$$

Then there exists an immersion $\theta \in W_{\text{loc}}^{2,p}(\omega; \mathbb{E}^3)$ such that

$$\hat{a}_{\alpha\beta}(\theta) = a_{\alpha\beta} \text{ and } \hat{b}_{\alpha\beta}(\theta) = b_{\alpha\beta} \text{ a.e. in } \omega.$$

Besides, an immersion $\psi \in W_{\text{loc}}^{2,p}(\omega; \mathbb{E}^3)$ satisfies

$$\hat{a}_{\alpha\beta}(\psi) = \hat{a}_{\alpha\beta}(\theta) \text{ and } \hat{b}_{\alpha\beta}(\psi) = \hat{b}_{\alpha\beta}(\theta) \text{ a.e. in } \omega$$

if and only if there exists an isometry $\mathbf{r} \in \mathbf{Isom}_+(\mathbb{E}^3)$ such that

$$\psi = \mathbf{r} \circ \theta \text{ in } \omega. \quad \square$$

Our first objective (Theorem 2) consists in showing that an existence and uniqueness theorem similar to Theorem 1 holds in the spaces $W^{m,p}(\omega)$ instead of the spaces $W_{\text{loc}}^{m,p}(\omega)$ if the open set ω is in addition a domain.

Theorem 2. Let ω be a simply-connected domain in \mathbb{R}^2 , let $p > 2$, and let a matrix field $(a_{\alpha\beta}) \in W^{1,p}(\omega; \mathbb{S}_{>}^2)$ and a matrix field $(b_{\alpha\beta}) \in L^p(\omega; \mathbb{S}^2)$ be given that satisfy the equations

$$R_{\alpha\beta\tau}^\sigma = 0 \text{ and } R_{\alpha\beta\tau}^3 = 0 \text{ in } \mathcal{D}'(\omega).$$

Then there exists an immersion $\theta \in W^{2,p}(\omega; \mathbb{E}^3)$ such that

$$\hat{a}_{\alpha\beta}(\theta) = a_{\alpha\beta} \text{ and } \hat{b}_{\alpha\beta}(\theta) = b_{\alpha\beta} \text{ a.e. in } \omega.$$

Besides, an immersion $\psi \in W^{2,p}(\omega; \mathbb{E}^3)$ satisfies

$$\hat{a}_{\alpha\beta}(\psi) = \hat{a}_{\alpha\beta}(\theta) \text{ and } \hat{b}_{\alpha\beta}(\psi) = \hat{b}_{\alpha\beta}(\theta) \text{ a.e. in } \omega$$

if and only if there exists an isometry $\mathbf{r} \in \mathbf{Isom}_+(\mathbb{E}^3)$ such that

$$\psi = \mathbf{r} \circ \theta \text{ in } \omega. \quad \square$$

Sketch of proof. Since $p > 2$ and ω is a domain, $W^{1,p}(\omega)$ is a Banach algebra and the canonical injection from $W^{1,p}(\omega)$ into $C^0(\overline{\omega})$ is continuous. Combining these two observations with the Gauss equations

$$\partial_\alpha \hat{a}_\beta(\theta) = \Gamma_{\alpha\beta}^\sigma \hat{a}_\sigma(\theta) + b_{\alpha\beta} \hat{a}_3(\theta) \text{ a.e. in } \omega$$

and the relations

$$|\hat{a}_\alpha(\theta)| = \sqrt{a_{\alpha\alpha}} \text{ (no summation on } \alpha \text{ here) and } |\hat{a}_3(\theta)| = 1 \text{ a.e. in } \omega,$$

where $\theta \in W_{\text{loc}}^{2,p}(\omega; \mathbb{E}^3)$ denotes the immersion found in Theorem 1 and the functions $\Gamma_{\alpha\beta}^\sigma$ are defined as in Theorem 1 (in effect the Christoffel symbols associated with θ), shows that the three vector fields $\hat{a}_i(\theta)$ belong to $L^\infty(\omega; \mathbb{E}^3)$, which in turn implies that $\partial_\alpha \theta \in L^\infty(\omega; \mathbb{E}^3)$ and $\partial_{\alpha\beta} \theta \in L^p(\omega; \mathbb{E}^3)$. It is then an easy matter to conclude that $\theta \in L^p(\omega; \mathbb{E}^3)$, hence that $\theta \in W^{2,p}(\omega; \mathbb{E}^3)$. The uniqueness up to isometries follows immediately from Theorem 1. \square

3. A nonlinear Korn inequality for surfaces of class $W^{2,p}$

The second objective of this Note is to complement the existence and uniqueness result of Theorem 2 by a stability result (Theorem 3 below), showing that the distance modulo a proper isometry between two surfaces in $W^{2,p}$ -norm is bounded by the distance between their first fundamental forms in the $W^{1,p}$ -norm and the distance between their second fundamental forms in the L^p -norm. A notation such as $c = c(\omega, p, \varepsilon)$ means that c is a real constant that depends on ω , p and ε .

Theorem 3. Let ω be a bounded and connected open subset of \mathbb{R}^2 that satisfies the uniform interior cone property. Given any $p > 2$ and $\varepsilon > 0$, let

$$V_\varepsilon(\omega; \mathbb{E}^3) := \left\{ \theta \in W^{2,p}(\omega; \mathbb{E}^3); \|\theta\|_{W^{2,p}(\omega; \mathbb{E}^3)} \leq 1/\varepsilon \text{ and } |\partial_1 \theta \wedge \partial_2 \theta| \geq \varepsilon \text{ in } \omega \right\}.$$

Then there exists a constant $c = c(\omega, p, \varepsilon)$ such that

$$\inf_{\mathbf{r} \in \mathbf{Isom}_+(\mathbb{E}^3)} \|\varphi - \mathbf{r} \circ \psi\|_{W^{2,p}(\omega; \mathbb{E}^3)} \leq c \left\{ \|(\hat{a}_{\alpha\beta}(\varphi) - \hat{a}_{\alpha\beta}(\psi))\|_{W^{1,p}(\omega; \mathbb{S}^2)} + \|(\hat{b}_{\alpha\beta}(\varphi) - \hat{b}_{\alpha\beta}(\psi))\|_{L^p(\omega; \mathbb{S}^2)} \right\}$$

for all $\varphi \in V_\varepsilon(\omega; \mathbb{E}^3)$ and $\psi \in V_\varepsilon(\omega; \mathbb{E}^3)$. \square

Remark 2. The above inequality can indeed be seen as a *nonlinear* Korn inequality for surfaces of class $W^{2,p}$, since a *formal linearization* (such a linearization consists first in letting in the above nonlinear inequality $\boldsymbol{\varphi} := \boldsymbol{\theta} + \boldsymbol{\eta}$ and $\boldsymbol{\psi} := \boldsymbol{\theta}$, where $\boldsymbol{\theta} \in W^{2,p}(\omega; \mathbb{E}^3)$ is a given immersion considered as “fixed”, and $\boldsymbol{\eta} \in W^{2,p}(\omega; \mathbb{E}^3)$ is an arbitrary vector field, then in canceling all the terms that depend nonlinearly on $\boldsymbol{\eta}$) yields the following *linear* Korn inequality on the surface $S = \boldsymbol{\theta}(\omega)$: There exists a constant $c_0 = c_0(\boldsymbol{\theta}, \omega)$ such that (the space $\mathbf{Rig}(\mathbb{E}^3)$ is defined in Remark 1)

$$\inf_{\boldsymbol{\zeta} \in \mathbf{Rig}(\mathbb{E}^3)} \|\boldsymbol{\eta} - \boldsymbol{\zeta}\|_{W^{2,p}(\omega; \mathbb{E}^3)} \leq c_0 \left\{ \|(\gamma_{\alpha\beta}(\boldsymbol{\eta}))\|_{W^{1,p}(\omega; \mathbb{S}^2)} + \|(\rho_{\alpha\beta}(\boldsymbol{\eta}))\|_{L^p(\omega; \mathbb{S}^2)} \right\} \text{ for all } \boldsymbol{\eta} \in W^{2,p}(\omega; \mathbb{E}^3),$$

where

$$\gamma_{\alpha\beta}(\boldsymbol{\eta}) := \frac{1}{2} [\hat{a}_{\alpha\beta}(\boldsymbol{\theta} + \boldsymbol{\eta}) - \hat{a}_{\alpha\beta}(\boldsymbol{\theta})]^{\text{lin}} \text{ and } \rho_{\alpha\beta}(\boldsymbol{\eta}) := [\hat{b}_{\alpha\beta}(\boldsymbol{\theta} + \boldsymbol{\eta}) - \hat{b}_{\alpha\beta}(\boldsymbol{\theta})]^{\text{lin}}$$

designate the linear parts with respect to $\boldsymbol{\eta}$ of the tensors appearing in the right-hand side of the inequality of Theorem 3. \square

The proof of Theorem 3 relies on a *comparison theorem between solutions to general Pfaff systems* due to the first author and S. Mardare (see Theorem 3.1 and Remark 3.1 in [3] and Theorem 4.1 in [7]), which we state below only in the particular case needed here. The notations \mathbb{M}^3 and $|\cdot|$ used in the next theorem respectively denote the space of 3×3 real matrices and the Frobenius norm in this space. The notation $(\mathbf{a} \mid \mathbf{b} \mid \mathbf{c})$ denotes the matrix in \mathbb{M}^3 with column vectors $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{E}^3$.

Theorem 4. Let ω be a bounded and connected open subset of \mathbb{R}^2 that satisfies the uniform interior cone property. Given any $p > 2$, $\varepsilon > 0$, and $y_0 \in \omega$, there exists a constant $c_1 = c_1(\omega, p, \varepsilon, y_0)$ such that

$$\|\mathbf{F} - \tilde{\mathbf{F}}\|_{W^{1,p}(\omega; \mathbb{M}^3)} \leq c_1 \left(|\mathbf{F}(y_0) - \tilde{\mathbf{F}}(y_0)| + \sum_{\alpha} \|\Gamma_{\alpha} - \tilde{\Gamma}_{\alpha}\|_{L^p(\omega; \mathbb{M}^3)} \right)$$

for all matrix fields $\mathbf{F}, \tilde{\mathbf{F}} \in W^{1,p}(\omega; \mathbb{M}^3)$ and $\Gamma_{\alpha}, \tilde{\Gamma}_{\alpha} \in L^p(\omega; \mathbb{M}^3)$ that satisfy

$$|\mathbf{F}(y_0)| + \sum_{\alpha} \|\Gamma_{\alpha}\|_{L^p(\omega; \mathbb{M}^3)} \leq \frac{1}{\varepsilon} \text{ and } |\tilde{\mathbf{F}}(y_0)| + \sum_{\alpha} \|\tilde{\Gamma}_{\alpha}\|_{L^p(\omega; \mathbb{M}^3)} \leq \frac{1}{\varepsilon},$$

and

$$\partial_{\alpha} \mathbf{F} = \mathbf{F} \Gamma_{\alpha} \text{ and } \partial_{\alpha} \tilde{\mathbf{F}} = \tilde{\mathbf{F}} \tilde{\Gamma}_{\alpha} \text{ a.e. in } \omega. \quad \square$$

Sketch of the proof of Theorem 3. With any immersion $\boldsymbol{\varphi} \in W^{2,p}(\omega; \mathbb{E}^3)$, we associate: the proper isometry $\mathbf{r}(\boldsymbol{\varphi}, y_0)$ of \mathbb{E}^3 defined by

$$\mathbf{r}(\boldsymbol{\varphi}, y_0)(x) := (\mathbf{B}^T \mathbf{B})^{1/2} \mathbf{B}^{-1} (x - \boldsymbol{\varphi}(y_0)) \text{ for all } x \in \mathbb{E}^3,$$

where

$$\mathbf{B} := (\hat{\mathbf{a}}_1(\boldsymbol{\varphi})(y_0) \mid \hat{\mathbf{a}}_2(\boldsymbol{\varphi})(y_0) \mid \hat{\mathbf{a}}_3(\boldsymbol{\varphi})(y_0));$$

the immersion

$$\boldsymbol{\theta}(\boldsymbol{\varphi}, y_0) := \mathbf{r}(\boldsymbol{\varphi}, y_0) \circ \boldsymbol{\varphi} \in W^{2,p}(\omega; \mathbb{E}^3);$$

and the matrix fields

$$\mathbf{F}(\boldsymbol{\varphi}, y_0) := (\hat{\mathbf{a}}_1(\boldsymbol{\theta}(\boldsymbol{\varphi}, y_0)) \mid \hat{\mathbf{a}}_2(\boldsymbol{\theta}(\boldsymbol{\varphi}, y_0)) \mid \hat{\mathbf{a}}_3(\boldsymbol{\theta}(\boldsymbol{\varphi}, y_0)))$$

and

$$\mathbf{A}(\boldsymbol{\varphi}) := \begin{pmatrix} a_{11} & a_{12} & 0 \\ a_{21} & a_{22} & 0 \\ 0 & 0 & 1 \end{pmatrix} \text{ and } \Gamma_{\alpha}(\boldsymbol{\varphi}) := \begin{pmatrix} \Gamma_{\alpha 1}^1 & \Gamma_{\alpha 2}^1 & -b_{\alpha}^1 \\ \Gamma_{\alpha 1}^2 & \Gamma_{\alpha 2}^2 & -b_{\alpha}^2 \\ b_{\alpha 1} & b_{\alpha 2} & 0 \end{pmatrix},$$

where

$$a_{\alpha\beta} := \hat{a}_{\alpha\beta}(\boldsymbol{\varphi}), \quad b_{\alpha\beta} := \hat{b}_{\alpha\beta}(\boldsymbol{\varphi}), \quad b_{\beta}^{\alpha} := a^{\alpha\sigma} b_{\sigma\beta}, \quad (a^{\sigma\tau}) := (a_{\alpha\beta})^{-1},$$

and

$$\Gamma_{\alpha\beta}^{\sigma} := \frac{1}{2} a^{\sigma\tau} (\partial_{\alpha} a_{\beta\tau} + \partial_{\beta} a_{\alpha\tau} - \partial_{\tau} a_{\alpha\beta}).$$

These matrix fields satisfy the *Pfaff system*

$$\partial_\alpha \mathbf{F}(\boldsymbol{\varphi}, y_0) = \mathbf{F}(\boldsymbol{\varphi}, y_0) \boldsymbol{\Gamma}_\alpha(\boldsymbol{\varphi}) \text{ a.e. in } \omega,$$

and the “initial condition”

$$(\mathbf{F}(\boldsymbol{\varphi}, y_0))(y_0) = (\mathbf{A}(\boldsymbol{\varphi})(y_0))^{1/2} \in \mathbb{S}_>^3.$$

Note in passing that the above Pfaff system is equivalent to the *equations of Gauss and Weingarten* associated with the immersion $\boldsymbol{\theta}(\boldsymbol{\varphi}, y_0)$.

In addition, if $\boldsymbol{\varphi} \in V_\varepsilon(\omega; \mathbb{E}^3)$ for some $\varepsilon > 0$ (the set $V_\varepsilon(\omega; \mathbb{E}^3)$ is defined in the statement of [Theorem 3](#)), then

$$\mathbf{F}(\boldsymbol{\varphi}, y_0) \in W^{1,p}(\omega; \mathbb{S}^3) \text{ and } \boldsymbol{\Gamma}_\alpha(\boldsymbol{\varphi}) \in L^p(\omega; \mathbb{M}^3),$$

and there exists a constant $c_1 = c_1(\omega, p, \varepsilon)$ such that

$$|(\mathbf{F}(\boldsymbol{\varphi}, y_0))(y_0)| + \|\boldsymbol{\Gamma}_\alpha(\boldsymbol{\varphi})\|_{L^p(\omega; \mathbb{M}^3)} \leq c_1.$$

This allows us to apply [Theorem 4](#) and to deduce that there exists a constant $c_2 = c_2(\omega, y_0, p, \varepsilon)$ such that

$$\|\mathbf{F}(\boldsymbol{\varphi}, y_0) - \mathbf{F}(\boldsymbol{\psi}, y_0)\|_{W^{1,p}(\omega; \mathbb{M}^3)} \leq c_2 \left(|(\mathbf{A}(\boldsymbol{\varphi}))(y_0) - (\mathbf{A}(\boldsymbol{\psi}))(y_0)| + \sum_{\alpha} \|\boldsymbol{\Gamma}_\alpha(\boldsymbol{\varphi}) - \boldsymbol{\Gamma}_\alpha(\boldsymbol{\psi})\|_{L^p(\omega; \mathbb{M}^3)} \right)$$

for all immersions $\boldsymbol{\varphi}$ and $\boldsymbol{\psi}$ that belong to the set $V_\varepsilon(\omega; \mathbb{E}^3)$.

Next, using the expressions of the matrix fields appearing in the right-hand side of the above inequality in terms of the fundamental forms associated with the immersions $\boldsymbol{\varphi}$ and $\boldsymbol{\psi}$, we deduce after a series of straightforward, but somewhat technical, computations that there exist two constants $c_3 = c_3(\omega, p, \varepsilon)$ and $c_4 = c_4(\omega, p, \varepsilon)$ such that

$$|(\mathbf{A}(\boldsymbol{\varphi}))(y_0) - (\mathbf{A}(\boldsymbol{\psi}))(y_0)| \leq c_3 \|(\hat{a}_{\alpha\beta}(\boldsymbol{\varphi}) - \hat{a}_{\alpha\beta}(\boldsymbol{\psi}))\|_{W^{1,p}(\omega; \mathbb{S}^2)},$$

and

$$\|\boldsymbol{\Gamma}_\alpha(\boldsymbol{\varphi}) - \boldsymbol{\Gamma}_\alpha(\boldsymbol{\psi})\|_{L^p(\omega; \mathbb{M}^3)} \leq c_4 \left(\|(\hat{a}_{\alpha\beta}(\boldsymbol{\varphi}) - \hat{a}_{\alpha\beta}(\boldsymbol{\psi}))\|_{W^{1,p}(\omega; \mathbb{S}^2)} + \|(\hat{b}_{\alpha\beta}(\boldsymbol{\varphi}) - \hat{b}_{\alpha\beta}(\boldsymbol{\psi}))\|_{L^p(\omega; \mathbb{S}^2)} \right).$$

Finally, the definition of the immersions $\boldsymbol{\theta}(\boldsymbol{\varphi}, y_0)$ and $\boldsymbol{\theta}(\boldsymbol{\psi}, y_0)$ implies that the vector field

$$\boldsymbol{\eta} := \boldsymbol{\theta}(\boldsymbol{\varphi}, y_0) - \boldsymbol{\theta}(\boldsymbol{\psi}, y_0) \in W^{2,p}(\omega; \mathbb{E}^3)$$

satisfies the *Poincaré system* (the notation $[\cdot]_\alpha$ denotes the α -th column vector of the matrix appearing between the brackets)

$$\partial_\alpha \boldsymbol{\eta} = [\mathbf{F}(\boldsymbol{\varphi}, y_0) - \mathbf{F}(\boldsymbol{\psi}, y_0)]_\alpha \text{ in } \omega$$

and the “initial condition”

$$\boldsymbol{\eta}(y_0) = \mathbf{0}.$$

Using an *inequality of Poincaré’s type*, we infer from the above system and initial condition that there exists a constant $c_5 = c_5(\omega, p)$ such that

$$\|\boldsymbol{\eta}\|_{W^{2,p}(\omega; \mathbb{E}^3)} \leq c_5 \|\mathbf{F}(\boldsymbol{\varphi}, y_0) - \mathbf{F}(\boldsymbol{\psi}, y_0)\|_{W^{1,p}(\omega; \mathbb{M}^3)}.$$

The conclusion follows by combining the above inequalities and by noting that, thanks to the invariance under rotations of the Euclidean and Frobenius norms,

$$\|\boldsymbol{\eta}\|_{W^{2,p}(\omega; \mathbb{E}^3)} = \|\boldsymbol{\theta}(\boldsymbol{\varphi}, y_0) - \boldsymbol{\theta}(\boldsymbol{\psi}, y_0)\|_{W^{2,p}(\omega; \mathbb{E}^3)} \geq \inf_{\mathbf{r} \in \text{Isom}_+(\mathbb{E}^3)} \|\boldsymbol{\varphi} - \mathbf{r} \circ \boldsymbol{\psi}\|_{W^{2,p}(\omega; \mathbb{E}^3)}. \quad \square$$

4. Local Lipschitz-continuity of the mapping defining a surface of class $W^{2,p}$, $p > 2$, in terms of its fundamental forms

Let ω be an open subset of \mathbb{R}^2 . Given two symmetric matrix fields

$$\mathbf{A} = (a_{\alpha\beta}) \in W_{\text{loc}}^{1,p}(\omega; \mathbb{S}^2) \text{ and } \mathbf{B} = (b_{\alpha\beta}) \in L_{\text{loc}}^p(\omega; \mathbb{S}^2), \quad p > 2,$$

such that $\mathbf{A}(y) \in \mathbb{S}_{>}^2$ for all $y \in \overline{\omega}$, define the distributions

$$R_{\alpha\beta\tau}^\sigma(\mathbf{A}, \mathbf{B}) := \partial_\tau \Gamma_{\alpha\beta}^\sigma - \partial_\beta \Gamma_{\alpha\tau}^\sigma + \Gamma_{\alpha\beta}^\gamma \Gamma_{\tau\gamma}^\sigma - \Gamma_{\alpha\tau}^\gamma \Gamma_{\beta\gamma}^\sigma - b_{\alpha\beta} b_\tau^\sigma + b_{\alpha\tau} b_\beta^\sigma \in \mathcal{D}'(\omega),$$

$$R_{\alpha\beta\tau}^3(\mathbf{A}, \mathbf{B}) := \partial_\tau b_{\alpha\beta} - \partial_\beta b_{\alpha\tau} + \Gamma_{\alpha\beta}^\gamma b_{\tau\gamma} - \Gamma_{\alpha\tau}^\gamma b_{\beta\gamma} \in \mathcal{D}'(\omega),$$

where

$$\Gamma_{\alpha\beta}^\sigma = \Gamma_{\alpha\beta}^\sigma(\mathbf{A}) := \frac{1}{2} a^{\sigma\tau} (\partial_\alpha a_{\beta\tau} + \partial_\beta a_{\alpha\tau} - \partial_\tau a_{\alpha\beta}) \in L_{\text{loc}}^p(\omega),$$

$$b_\beta^\sigma := a^{\sigma\tau} b_{\tau\beta} \in L_{\text{loc}}^p(\omega), \text{ and } (a^{\sigma\tau}) := (a_{\alpha\beta})^{-1} \in W_{\text{loc}}^{1,p}(\omega).$$

Remark 3. The above regularity assumptions on the fields \mathbf{A} and \mathbf{B} are the minimal possible in order that the definitions of the distributions $R_{\alpha\beta\tau}^j(\mathbf{A}, \mathbf{B})$ make sense: combined with the Sobolev embedding $W_{\text{loc}}^{1,p}(\omega) \subset C^0(\omega)$, they ensure that $\det \mathbf{A}$ is a continuous positive function over ω , which in turn implies that $a^{\sigma\tau} \in C^0(\omega)$ and so the products appearing in the definitions of $\Gamma_{\alpha\beta}^\sigma$ and b_α^σ belong to $L_{\text{loc}}^p(\omega)$; this allows to define the partial derivatives of $\Gamma_{\alpha\beta}^\sigma$ and b_α^σ appearing in the above definition of $R_{\alpha\beta\tau}^j(\mathbf{A}, \mathbf{B})$ as distributions in $\mathcal{D}'(\omega)$. \square

The *third objective* of this Note is to establish, as a consequence of the nonlinear Korn inequality of [Theorem 3](#), the following “existence, uniqueness, and stability theorem” for the reconstruction of a surface from its fundamental forms in the spaces $W^{1,p}(\omega; \mathbb{S}^2)$ and $L^p(\omega; \mathbb{S}^2)$.

In [Theorem 5](#) below, the set $\dot{W}^{2,p}(\omega; \mathbb{E}^3)$ is the quotient set of the space $W^{2,p}(\omega; \mathbb{E}^3)$ by the *equivalence relation* between isometrically equivalent immersions, and the set $\mathbb{T}(\omega)$ is the subset of the space $W^{1,p}(\omega; \mathbb{S}^2) \times L^p(\omega; \mathbb{S}^2)$ formed by all pairs of a positive-definite symmetric matrix field and a symmetric matrix field that satisfy together the *equations of Gauss and Codazzi–Mainardi* in the distributional sense. As such, the sets $\dot{W}^{2,p}(\omega; \mathbb{E}^3)$ and $\mathbb{T}(\omega)$ are *metric spaces* equipped respectively with the distances defined by

$$\text{dist}_{\dot{W}^{2,p}(\omega; \mathbb{E}^3)}(\dot{\boldsymbol{\theta}}, \dot{\boldsymbol{\psi}}) := \inf_{\tilde{\boldsymbol{\theta}} \in \dot{\boldsymbol{\theta}}, \tilde{\boldsymbol{\psi}} \in \dot{\boldsymbol{\psi}}} \|\tilde{\boldsymbol{\theta}} - \tilde{\boldsymbol{\psi}}\|_{W^{2,p}(\omega; \mathbb{E}^3)} = \inf_{\mathbf{r} \in \mathbf{Isom}_+(\mathbb{E}^3)} \|\boldsymbol{\theta} - \mathbf{r} \circ \boldsymbol{\psi}\|_{W^{2,p}(\omega; \mathbb{E}^3)}$$

for all $\dot{\boldsymbol{\theta}}$ and $\dot{\boldsymbol{\psi}}$ in $\dot{W}^{2,p}(\omega; \mathbb{E}^3)$, and by

$$\text{dist}_{\mathbb{T}(\omega)}((\mathbf{A}, \mathbf{B}), (\tilde{\mathbf{A}}, \tilde{\mathbf{B}})) := \|\mathbf{A} - \tilde{\mathbf{A}}\|_{W^{1,p}(\omega; \mathbb{S}^2)} + \|\mathbf{B} - \tilde{\mathbf{B}}\|_{L^p(\omega; \mathbb{S}^2)}$$

for all (\mathbf{A}, \mathbf{B}) and $(\tilde{\mathbf{A}}, \tilde{\mathbf{B}})$ in $\mathbb{T}(\omega)$.

Theorem 5. Let ω be a domain in \mathbb{R}^2 . Given any $p > 2$, define the sets

$$\dot{W}^{2,p}(\omega; \mathbb{E}^3) := \{\dot{\boldsymbol{\theta}} = \{\mathbf{r} \circ \boldsymbol{\theta}; \mathbf{r} \in \mathbf{Isom}_+(\mathbb{E}^3)\}; \boldsymbol{\theta} \in W^{2,p}(\omega; \mathbb{E}^3)\}$$

and

$$\mathbb{T}(\omega) := \{(\mathbf{A}, \mathbf{B}) \in W^{1,p}(\omega; \mathbb{S}^2) \times L^p(\omega; \mathbb{S}^2); \mathbf{A}(y) \in \mathbb{S}_{>}^2 \text{ at each } y \in \overline{\omega}, R_{\alpha\beta\tau}^j(\mathbf{A}, \mathbf{B}) = 0 \text{ in } \mathcal{D}'(\omega)\}.$$

Then the following assertions are true:

(a) Two matrix fields $\mathbf{A} = (a_{\alpha\beta})$ and $\mathbf{B} = (b_{\alpha\beta})$ satisfy

$$(\mathbf{A}, \mathbf{B}) \in \mathbb{T}(\omega)$$

if and only if there exists an immersion $\boldsymbol{\theta} \in W^{2,p}(\omega; \mathbb{E}^3)$ such that

$$\hat{a}_{\alpha\beta}(\boldsymbol{\theta}) = a_{\alpha\beta} \text{ in } \omega \text{ and } \hat{b}_{\alpha\beta}(\boldsymbol{\theta}) = b_{\alpha\beta} \text{ a.e. in } \omega.$$

(b) Two immersions $\boldsymbol{\theta} \in W^{2,p}(\omega; \mathbb{E}^3)$ and $\boldsymbol{\psi} \in W^{2,p}(\omega; \mathbb{E}^3)$ satisfy the relations

$$\hat{a}_{\alpha\beta}(\boldsymbol{\theta}) = \hat{a}_{\alpha\beta}(\boldsymbol{\psi}) \text{ in } \omega \text{ and } \hat{b}_{\alpha\beta}(\boldsymbol{\theta}) = \hat{b}_{\alpha\beta}(\boldsymbol{\psi}) \text{ a.e. in } \omega$$

if and only if there exists a proper isometry \mathbf{r} of \mathbb{E}^3 such that

$$\boldsymbol{\psi} = \mathbf{r} \circ \boldsymbol{\theta} \text{ in } \omega.$$

(c) The mapping defined by (a) and (b), namely

$$\mathcal{G} : (\mathbf{A}, \mathbf{B}) \in \mathbb{T}(\omega) \rightarrow \mathcal{G}((\mathbf{A}, \mathbf{B})) := \dot{\boldsymbol{\theta}} \in \dot{W}^{2,p}(\omega; \mathbb{E}^3),$$

where $\boldsymbol{\theta} \in W^{2,p}(\omega; \mathbb{E}^3)$ is any immersion that satisfies

$$(\hat{a}_{\alpha\beta}(\boldsymbol{\theta})) = \mathbf{A} \text{ and } (\hat{b}_{\alpha\beta}(\boldsymbol{\theta})) = \mathbf{B} \text{ a.e. in } \omega,$$

is locally Lipschitz-continuous. \square

Sketch of proof. Parts (a) and (b) are just a re-statement of [Theorem 2](#). Otherwise, the rest of the proof follows a strategy introduced by the first author and S. Mardare in [\[3\]](#). More precisely, part (c) of [Theorem 5](#) is deduced from [Theorem 3](#) as follows.

On the one hand, the Sobolev embedding $W^{1,p}(\omega) \subset C^0(\overline{\omega})$ implies that, given any $(\mathbf{A}, \mathbf{B}) \in \mathbb{T}(\omega)$, there exists $\delta = \delta(\mathbf{A}, \mathbf{B}) > 0$ such that the set

$$\mathbb{T}_\delta(\omega) := \left\{ (\tilde{\mathbf{A}}, \tilde{\mathbf{B}}) \in \mathbb{T}(\omega); \det \tilde{\mathbf{A}} \geq \delta \text{ in } \omega, \|\tilde{\mathbf{A}}\|_{W^{1,p}(\omega; \mathbb{S}^2)} \leq 1/\delta, \text{ and } \|\tilde{\mathbf{B}}\|_{L^p(\omega; \mathbb{S}^2)} \leq 1/\delta \right\}$$

is a neighborhood of (\mathbf{A}, \mathbf{B}) in the metric space $\mathbb{T}(\omega)$. It also implies that

$$\mathbb{T}(\omega) = \bigcup_{\delta > 0} \mathbb{T}_\delta(\omega).$$

Besides, for each $\delta > 0$, there exists $\varepsilon(\delta) > 0$ such that

$$\mathcal{G}(\mathbb{T}_\delta(\omega)) \subset \{\dot{\boldsymbol{\theta}} \in \dot{W}^{2,p}(\omega; \mathbb{E}^3); \boldsymbol{\theta} \in V_{\varepsilon(\delta)}(\omega; \mathbb{E}^3)\},$$

where \mathcal{G} denotes the mapping defined in part (c) of the statement of the theorem and $V_{\varepsilon(\delta)}(\omega; \mathbb{E}^3)$ is defined as in [Theorem 3](#).

On the other hand, [Theorem 3](#) implies that there exists a constant $c = c(\omega, p, \varepsilon(\delta))$ such that

$$\inf_{\mathbf{r} \in \text{Isom}_+(\mathbb{E}^3)} \|\boldsymbol{\varphi} - \mathbf{r} \circ \boldsymbol{\psi}\|_{W^{2,p}(\omega; \mathbb{E}^3)} \leq c \left\{ \|(\hat{a}_{\alpha\beta}(\boldsymbol{\varphi}) - \hat{a}_{\alpha\beta}(\boldsymbol{\psi}))\|_{W^{1,p}(\omega; \mathbb{S}^2)} + \|(\hat{b}_{\alpha\beta}(\boldsymbol{\varphi}) - \hat{b}_{\alpha\beta}(\boldsymbol{\psi}))\|_{L^p(\omega; \mathbb{S}^2)} \right\}$$

for all mappings $\boldsymbol{\varphi} \in V_{\varepsilon(\delta)}(\omega; \mathbb{E}^3)$ and $\boldsymbol{\psi} \in V_{\varepsilon(\delta)}(\omega; \mathbb{E}^3)$ (note that [Theorem 3](#) can be applied under the assumptions of [Theorem 5](#) since a domain satisfies the uniform interior cone property).

We then infer from the observations above that, given any mappings $\boldsymbol{\varphi} \in V_{\varepsilon(\delta)}(\omega; \mathbb{E}^3)$ and $\tilde{\boldsymbol{\varphi}} \in V_{\varepsilon(\delta)}(\omega; \mathbb{E}^3)$ such that $\dot{\boldsymbol{\varphi}} = \mathcal{G}(\mathbf{A}, \mathbf{B})$ and $\dot{\tilde{\boldsymbol{\varphi}}} = \mathcal{G}(\tilde{\mathbf{A}}, \tilde{\mathbf{B}})$ for some $(\mathbf{A}, \mathbf{B}) \in \mathbb{T}_\delta(\omega)$ and $(\tilde{\mathbf{A}}, \tilde{\mathbf{B}}) \in \mathbb{T}_\delta(\omega)$,

$$\text{dist}_{\dot{W}^{2,p}(\omega; \mathbb{E}^3)}(\dot{\boldsymbol{\varphi}}, \dot{\tilde{\boldsymbol{\varphi}}}) \leq c \text{dist}_{\mathbb{T}(\omega)}((\mathbf{A}, \mathbf{B}), (\tilde{\mathbf{A}}, \tilde{\mathbf{B}})).$$

This shows that the restriction of the mapping \mathcal{G} to the set $\mathbb{T}_\delta(\omega)$ is Lipschitz-continuous. \square

Acknowledgements

The work described in this paper was substantially supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region, China [Project No. 9042388, CityU 11305716].

References

- [1] M.P. do Carmo, *Differential Geometry of Curves and Surfaces*, Prentice-Hall, Englewood Cliffs, 1976.
- [2] P.G. Ciarlet, *Linear and Nonlinear Functional Analysis with Applications*, SIAM, Philadelphia, 2013.
- [3] P.G. Ciarlet, S. Mardare, Nonlinear Korn inequalities in \mathbb{R}^n and immersions in $W^{2,p}$, $p > n$, considered as functions of their metric tensors in $W^{1,p}$, *J. Math. Pures Appl.* 105 (2016) 873–906.
- [4] P.G. Ciarlet, C. Mardare, A surface in $W^{2,p}$ is a locally Lipschitz-continuous function of its fundamental forms in $W^{1,p}$ and L^p , $p > 2$, in preparation.
- [5] W. Klingenberg, *Eine Vorlesung über Differentialgeometrie*, Springer, Berlin, 1973. English translation: *A Course in Differential Geometry*, Springer, Berlin, 1978.
- [6] S. Mardare, On Pfaff systems with L^p coefficients and their applications in differential geometry, *J. Math. Pures Appl.* 84 (2005) 1659–1692.
- [7] S. Mardare, On systems of first order linear partial differential equations with L^p coefficients, *Adv. Differ. Equ.* 12 (2007) 301–360.