Organic matter deposition in the Ghadames Basin (Libya) during the Late Devonian-A multidisciplinary approach

Armelle Riboulleau, Amalia Spina, Marco Vecoli, Laurent Riquier, Melesio Quijada, Nicolas Tribovillard, Olivier Averbuch

► To cite this version:

Armelle Riboulleau, Amalia Spina, Marco Vecoli, Laurent Riquier, Melesio Quijada, et al.. Organic matter deposition in the Ghadames Basin (Libya) during the Late Devonian-A multidisciplinary approach. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 2018, 497, pp.37-51. 10.1016/j.palaeo.2018.02.004 . hal-01780055

HAL Id: hal-01780055

<https://hal.sorbonne-universite.fr/hal-01780055>

Submitted on 27 Apr 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Organic matter deposition in the Ghadames Basin (Libya) during the Late Devonian—a multidisciplinary approach

¹ Armelle Riboulleau *

² Amalia Spina

³ Marco Vecoli¹

⁴ Laurent Riquier

¹ Melesio Quijada

¹ Nicolas Tribouvillard

¹ Olivier Averbuch

¹ Laboratoire d'Océanologie et de Géosciences, UMR 8187, Université de Lille, CNRS, Université du Littoral Côte d'Opale, 59655 Villeneuve d'Ascq, France.

² Department of Physics and Geology, University of Perugia, Perugia, Italy

³ Evolution, Ecologie et Paléontologie - UMR CNRS 8198, Université de Lille, 59655 Villeneuve d'Ascq, France

⁴ Sorbonne Universités, UPMC Univ. Paris 06, CNRS, UMR IStEP, 75252 Paris, France

* corresponding author:

e-mail address: armelle.riboulleau@univ-lille1.fr (Armelle Riboulleau)

Abstract

The organic rich deposits of Late Devonian age are the second most important petroleum source rocks in the Ghadames Basin (Libya). From available stratigraphy, the peak of organic matter deposition, corresponding to a level of radioactive shales easily recognized in gamma ray logs, occurred at the Frasnian-Famennian transition. The present study was focused on samples from borehole D1-26, located in the central part of the Ghadames Basin, belonging to the Aouinet Ouenine III and IV formations (Frasnian to Famennian). The mineral and organic content of the rocks were analyzed in order to determine the origin of organic matter accumulation and estimate if the organic matter enrichment could be related to

¹ Present address: Biostratigraphy Group, Exploration Technical Services Department, Saudi Aramco, Dhahran, Saudi Arabia

the well-known Upper Kellwasser event. The results indicate that during the Frasnian, the planktonic primary productivity was moderate, but anoxic conditions reaching the euphotic zone were frequent and allowed efficient preservation of the organic matter. At the Frasnian-Famennian transition, the deposition of the radioactive shales level can be related to a eustatic rise, which allowed incursion of nutrient-rich water in the basin, leading to increase in primary productivity and photic zone anoxia. During the Famennian, the conditions changed, associated with a relative sea level fall and/or a progradation of detrital sediments originating from the south. Photic zone anoxia was only episodic and plankton-derived organic matter was poorly preserved. Nevertheless, the important influx of organic matter of terrestrial origin allowed significant enrichment of the sediment in organic matter. Several indications suggest the radioactive shales interval could be coeval with the Upper Kellwasser level, nevertheless further stratigraphic work is needed to confirm this hypothesis.

Keywords

Biomarkers, palynofacies, major and trace elements, Frasnian-Famennian

1. Introduction

The Late Devonian was a period of extensive organic matter (OM) deposition along continental margins (Ulmishek and Klemme, 1990). The reasons for such widespread deposition of OM within oceanic realm is still a matter of discussion but a combination of several favorable factors has been proposed, including: (1) an overall rise of the sea level and a warmer climate favoring water stratification (Ulmishek and Klemme, 1990; Bond and Wignall, 2008; Riquier et al., 2010); (2) a tectonic context marked by the onset of the Variscan orogeny inducing both an increase of the nutrient supply from continents and the reduction of sea-water communications along the former Paleotethysian seaways (Tribovillard et al., 2004; Averbuch et al., 2005); (3) the development of root systems in terrestrial plants during this interval that may have also contributed to an increase in nutrient delivery to oceanic basins, promoting planktonic productivity (Algeo and Scheckler, 1998). The maximum convergence of these favoring conditions resulted in several anoxic events associated with biotic crises, the most famous of which, called the Kellwasser event, occurred at the Frasnian-Famennian boundary; it is characterized by the widespread deposition of two organic-rich levels respectively called the Lower and Upper Kellwasser levels (Buggisch, 1991; Joachimski and Buggisch, 1993; House, 2002; Bond et al., 2004; Racki, 2005; Riquier et al., 2006; Becker et al., 2012). Though considered as one of the major extinction events in

Earth history, affecting both marine and continental faunas (see McGhee et al., 2013 for a recent review), the exact timing, causes and consequences of the Kellwasser event is still a matter of active research (Racki, 2005; Becker et al., 2016). The resultant massive organic carbon burial may have had a major impact on the global carbon cycle thereby enhancing a significant drop of the CO₂ atmospheric content and a global cooling trend in Late Devonian times (Algeo et al., 1995; Joachimski and Buggisch, 2002; Godderis and Joachimski, 2004; Averbuch et al., 2005; Riquier et al., 2010).

North Africa is characterized by the presence of several large to giant petroleum systems associated with Paleozoic source rocks (Boote et al., 1998; Craig et al., 2008). In well logging, the organic-rich intervals are recognizable by their high radioactivity in gamma ray logs, and are qualified as “radioactive shales” or “hot shales”. The dominant source rocks in North African petroleum systems are of Silurian age, in particular the Tanezzuft Formation in Algeria Tunisia and Libya, which base shows a characteristic hot shales interval (Boote et al., 1998; Craig et al., 2008). Organic-rich deposits of Late Devonian age are also widespread in North Africa and represent the second most important source rocks (Boote et al., 1998; Echikh, 1998; Lüning et al., 2003). From available stratigraphic correlations, the age of the Upper Devonian hot shales nevertheless varies from the early Frasnian to the early Famennian, from one basin to the other (Lüning et al., 2003). In the Ghadames Basin (Libya and Tunisia, Fig. 1), the Upper Devonian organic-rich level corresponds to a radioactive shales and limestones interval of late Frasnian age, sometimes referred to as the Cues limestones Horizon (Massa, 1988; Weyant and Massa, 1991). This interval well recognized in gamma ray logs (Weyant and Massa, 1991; Boote et al., 1998) could therefore be coeval to the Upper Kellwasser horizon (Joachimski and Buggisch, 1993; Bond et al., 2004). The two Kellwasser levels are well known in Morocco (Riquier et al., 2005), and Soua (2014) proposed that the two late Frasnian hot-shale intervals detected in South Tunisia were coeval to the Kellwasser levels.

This paper presents the results of a multidisciplinary study of a Frasnian-Famennian section in the Libyan part of the Ghadames Basin. The aim of the study was to determine the factors which favored OM deposition in this basin during the Late Devonian, and potentially relate it to the general context of the extensive Kellwasser environmental crisis.

2. Materials and methods

2.1 Studied interval

This study is based on samples from borehole D1-26 (30° 29' 21" N, 10° 46' 28" E) cored in 1960-61 by the Oasis Oil Company in Libya, in the central part of the Ghadames Basin (Fig. 1). Samples were taken from cores 6, 7 and 8 in the Aouinet Ouenine (or Awaynat Wanin) formations III and IV (AO III and AO IV, Fig. 2) and were labeled according to their depth in the core, originally given in feet. In the studied borehole, the AO III and AO IV formations mostly consist in grey to dark grey, finely laminated and micaceous shales to silty shales (Fig. 2). The upper part of the AO III Fm. is more calcareous and passes to alternating shales and fossiliferous limestones in the lower part of the AO IV Fm. (Fig. 2). This latter interval should correspond to the "Cues Limestone Horizon" and associated radioactive shales (Weyant and Massa, 1991). Sandy levels appear and become more abundant in the upper part of the AO IV Fm. (Fig. 2). Biostratigraphic constraints, based on palynomorphs and conodonts, and gamma ray correlations with other cores from the Ghadames Basin (Loboziak and Streel, 1989; Weyant and Massa, 1991; Spina et al., 2017), indicate that the samples from core 8 are of late Frasnian age, likely within the *Palmatolepis rhenana* conodont Zone (Streel, 2009), while those from core 6 are of late Famennian age (Fig. 2), likely within the *Siphonodella praesulcata* conodont Zone (Streel, 2009). Core 7 was collected in so-called radioactive shales (Massa, 2005, personal communication). This radioactive shale interval generally is considered of late Frasnian age (Massa, 1988; Boote et al., 1998). Nevertheless, Weyant and Massa (1991) analyzed the conodont content of two samples of unknown depth from core 7, and proposed an early Famennian age, in the Middle to Upper *Palmatolepis triangularis* Zone, or *Palmatolepis crepida* Zone.

2.2 Palynology

Eighteen core samples from the Aouinet-Ouenine III and IV formations (samples 7813 to 7081, 2381.4 m to 2158.3 m) were processed and analyzed. About 20-30 g of sample have been initially treated in hydrochloric acid (37 %), followed by digestion in hydrofluoric acid (50 %) and by treatment in 30% boiling hydrochloric acid to remove fluorosilicates. Residues were then subjected to density separation using zinc bromide solution in order to separate OM from heavy minerals. Finally, the organic-rich residues were sieved with acid resistant 50 and 15 µm sieves. No oxidation by nitric acid was performed. A minimum of three palynological

strew-slides were mounted for each sample using Eukitt as mounting medium. Palynological slides were examined using transmitted light microscopes, commonly with $\times 20$ (dry) and $\times 100$ (oil immersion) objectives. Palynological slides are stored in the collection of the UMR 8198 Evo-Eco-Paleo, Université de Lille-CNRS, Villeneuve d'Ascq (France). For palynofacies analysis, 300 particles were counted in each level, in order to estimate the proportions of the organic constituents. The organic constituents from D1-26 borehole were grouped according to Whitaker (1984) classification, modified to emphasize the organic components useful for palaeoenvironmental observations in the Devonian time span. The allochthonous fraction, of continental origin, includes the palynomaceral group (plant debris) and sporomorphs. The palynomaceral (PM) group is divided in PM1, orange to dark brown fragments, translucent, partially oxidized; PM2, cell-structured orange to dark-brown fragments, moderately oxidized; PM4, black opaque fragments, strongly oxidized. PM4 is the most stable palynomaceral and can be transported for a long distance before being degraded. Sporomorphs mostly correspond to cryptospores (Strother, 1991) and miospores (Steemans, 1999). The autochthonous fraction, of marine origin, comprises acritarchs, prasynophycean cysts, chitinozoans and amorphous organic matter (AOM). Acritarchs are organic-walled microfossils of uncertain biological affinity (Evitt, 1963), but mostly representing marine phytoplankton (Tappan, 1980; Martin, 1993; Colbath and Grenfell, 1995). In the Aouinet Ouenine sediments, prasynophycean cysts mainly correspond to *Tasmanites* (Tasmanaceae). This type of algal microfossil occurs in many marine facies from Cambrian to Miocene (Martín-Closas, 2003). Abundance of *Tasmanites* is often observed in marine (shelf and oceanic) organic-rich sediments (Tyson, 1995). Chitinozoans are organic-walled microfossil of uncertain affinity, but most probably representing marine microzooplankton (Paris and Nölvak, 1999). AOM is the degradation product of benthic and pelagic cyanobacteria and sulfur bacteria in oxygen deficient environments. Being the most dissolvable OM, it can be preserved only in oxygen-deprived (suboxic to anoxic) environments, where it is not destroyed and biodegraded by consumers and decomposers or by the oxidizing processes (Hart, 1986). The quantity of this latter was established on unfiltered palynological residues, counting only particles larger than 50 μm .

2.3 Bulk geochemistry

Twelve samples in the 2158.3 m to 2381.4 m interval (samples 7081 to 7813) were analyzed for bulk geochemistry. Rock-Eval analyses were performed on 100 mg of ground

bulk rock at the Institut Français du Pétrole-Energies Nouvelles using a Rock-Eval 6 apparatus and standard analytical conditions (Behar et al., 2001). The main parameters obtained are total organic carbon content (TOC, in weight %), Hydrogen index (HI, mg HC/g TOC), oxygen index (OI, mg CO₂/g TOC), that are proportional to the H/C and O/C ratio of the kerogen, respectively, and Tmax, an indicator of OM maturity. The sulfur content was determined by elemental analysis of the sediment with a FlashEA 1112 Elemental Analyser (Thermo). The analysis was performed on 1.5 to 2 mg of ground bulk rock added to approximately 5 mg of vanadium pentoxide, used as a combustion catalyst. 2,5-Bis(5-tert-butyl-benzoxazol-2-yl)thiophene (BBOT) was used as standard.

Major, minor, trace and rare earth elements (REE) concentrations were determined on ground bulk rock by inductively coupled plasma optical emission spectrometry (ICP-OES) and inductively coupled plasma mass spectrometry (ICP-MS), at Activation Laboratories Ltd. (Ancaster, Canada). Samples were mixed with a flux of lithium metaborate (LiBO₂) and lithium tetraborate (Li₂B₄O₇), and fused in an induction furnace. Molten sample was immediately poured into a solution of 5% nitric acid (HNO₃) containing an internal standard, and mixed continuously until completely dissolved. The analytical accuracy and precision are found to be better than 1–2% for major elements, 5% for REE and 5–10% for the other elements, as checked by international standards and analysis of replicate samples. The enrichment factors were calculated for trace elements often enriched biogenically or authigenically, as follows: $X_{EF} = [(X/Al)_{\text{sample}} / (X/Al)_{\text{PAAS}}]$, where X and Al represent the weight % concentrations of element X and Al, respectively. Samples were normalized using the Post Archean Average shale (PAAS) compositions (McLennan, 1989). An enrichment factor larger than 1 theoretically points to the enrichment of the element relative to the PAAS; in other words it indicates whether an element is biogenically or authigenically enriched compared to what its concentration would be, if it were of merely clastic origin. Practically, enrichment factors are taken into consideration when they out pass the value of 3. Full results are provided in Table S1. For calculation purpose, concentrations below the detection limit (Table S1) were taken as half the value of the detection limit.

2.4 Biomarker analysis

For biomarker analysis, rock fragments were extracted (24 h) with dichloromethane (DCM) in a refrigerator to remove possible contamination from the sample surface. The fragments were then crushed to extract the lipids from inside the rock. Ca. 30 g pulverized

sample were extracted using a soxhlet apparatus with DCM (24 h). Elemental sulfur was removed by addition of activated copper to the recovered extracts. The solvent was removed by means of rotary evaporation and the residue re-dissolved in cyclohexane. The cyclohexane-soluble fraction (maltenes) was further separated by means of column chromatography. A mixture of standard compounds (2,2,4,4,6,8,8-heptamethylnonane, Dr Ehrenstorfer-GmbH; anthracene D10, Dr Ehrenstorfer-GmbH; nonadecan-2-one, Sigma-Aldrich) was added to the maltenes prior to fractionation. The aliphatic fraction was recovered from the maltenes by elution with cyclohexane on an activated silica column. Elution with a mixture of cyclohexane/DCM (9/1, v/v) recovered the aromatic fraction, after which the polar fraction was recovered by elution with DCM/MeOH (1/1, v/v). The aliphatic and aromatic fractions were analyzed by way of gas chromatography-mass spectrometry (GC-MS) using either a Trace GC 2000 gas chromatograph equipped with a DB5ht column (30 m × 0.25 mm i.d.; 0.1 µm film thickness). The oven temperature program was: 100 °C (1 min) to 310 °C (held 16.5 min) at 4 °C/min. Helium was used as carrier gas. The mass spectrometer conditions were as follow: ionization energy 70 eV for electron ionization (EI) with a mass range m/z 50–700. Alternatively, samples were analyzed with a Perkin Elmer Clarus 680 GC equipped with a DB5MS ultra inert column (30 m × 0.25 mm i.d.; 0.25 µm film thickness). The oven temperature program was: 60 °C (1 min) to 100°C at 15°C/min, from 100°C to 200°C at 3°C/min, from 200 °C to 310 °C (held 20 min) at 4 °C/min. Helium was used as carrier gas. The mass spectrometer was operated at 70 eV for electron ionization (EI) with a mass range m/z 50–550 (full scan). Compounds were identified by comparison of mass spectra and elution order with published data.

3. Results and interpretation

3.1 Palynofacies

The palynofacies of the studied samples shows variable proportions of palynomorphs as well as AOM (Fig. 3). The palynofacies of the first levels from the AO III Fm. (base of core 8, samples 7813 to 7809, Fig. 3) shows high abundances of acritarchs and prasinophycean cysts (tasmanaceans) representing about 40% of the entire palynofacies. AOM is abundantly present in this interval, reaching up to ~40% of the palynofacies. Partially oxidized palynomacerals (e.g. PM1+PM2) and sporomorphs are scarcely present. Chitinozoans are relatively abundant (5-7 %). In the overlying levels of the AO III Fm. (samples 7802 to 7780,

Fig. 3), the palynofacies is dominated by the same marine elements but the proportion of acritarchs and tasmanaceans increases (up to 60 %) while the AOM content is reduced (0 to 10 %). At the base of AO IV Fm. (core 7, Fig.3), corresponding to the "radioactive shales", AOM in high proportion, acritarchs and prasynophycean cysts characterize almost all the entire palynofacies. Sporomorphs and chitinozoans are very rare to absent. The acritarch fauna is well diversified in the samples from cores 8 and 7. A distinctive change in the palynofacies composition marks the uppermost levels (core 6) of AO IV Fm. (samples 7100 to 7081, Fig. 3). Strongly oxidized palynomacerals (PM4) and partially oxidized palynomacerals (PM1 + PM2) are abundant as well as well-preserved sporomorphs. This latter group reaches abundance around 35% in the levels close to the boundary with the overlying Tahara Fm. Microplankton here mainly consists of acritarchs (10 to 15 %). *Tasmanites* are uncommon (≤ 10 %). AOM and chitinozoans are very rare or absent.

3.2 Bulk geochemistry

3.2.1 Organic matter and sulfur

Total organic carbon (TOC) contents of the analyzed samples vary between 0.8 and 2.8 % (average 1.75 %, Table 1). Tmax values range between 431 and 440 °C. Average Tmax value in the uppermost samples is 435 °C indicating the beginning of the oil window. The average Tmax value is slightly higher in the samples from core 8 (438 °C) indicating a slightly higher thermal maturity.

HI values are between 60 and 319 mg HC/g TOC (av. 132 mg HC/g TOC) while OI values range between 0 and 47 mg CO₂/g TOC. In a pseudo van Krevelen (HI-OI) as well as in the HI-Tmax diagram (Fig. 4A), the samples plot in the area of Type II-III OM. TOC/N atomic ratios vary between 14 and 44 (Table 1). In average, higher values are observed in the samples from core 6 (av. 38) than in samples from cores 7 and 8 (av. 18). Higher values of the C/N ratio in the samples from the top of AO IV Fm. (samples 7100 to 7081) suggest a higher contribution from woody organic matter (Meyers, 1997), which is consistent with palynofacies observations.

The sulfur content of the sediment is relatively high, with an average value of 2.8 %. In a S vs TOC plot (Fig. 4B), most samples plot above the "normal marine" line (Berner, 1982), having a slope of 2 for the Devonian (Berner and Raiswell, 1983). It can be argued that this feature could result from a TOC loss upon maturation and petroleum expulsion, since

Rock Eval production index (PI) values between 0.10 and 0.48 are compatible with oil expulsion (Table 1). Nevertheless, considering an initially doubled TOC values does not significantly change the S-TOC relationship as most samples would remain above this “normal marine” line (Fig. 4C). This S-TOC behavior is often considered as indicating deposition under strongly reducing to euxinic conditions (Leventhal, 1983).

3.2.2 Productivity (Ba, Ni, Cu) and redox proxies (Mo, U, V, Ni, Cu, Zn, Co, Cr, Ce*)

In order to be valid as productivity or redox proxies, elements must not be positively correlated with the Si (quartz) and/or Al (clay) content of the sediment. In the studied samples, Si and Al are negatively correlated, suggesting that the clay and silica content vary in opposite proportions. Most productivity and redox-sensitive elements in D1-26 samples are neither correlated with Si nor Al, except for Cr which is positively correlated with Al ($r = 0.818$, $n=12$) and Ba which is positively correlated with Si ($r = 0.933$, $n=12$). The latter correlation might indicate the presence of silica of biogenic origin. The average value of the Ge/Si ratio of $8.1 \cdot 10^{-6}$ however points for a mainly detrital origin of the silica in D1-26 samples, therefore corresponding to detrital quartz (Tribovillard, 2013). This indicates that Cr and Ba have a dominant detrital origin and cannot be considered as environmental proxies.

The average enrichment factors (EF) of the different elements which are not influenced by the detrital supply range between 0.5 and 3 (Fig. 5A), pointing for no significant authigenic enrichment in these sediments. In detail, most elements do not show a marked temporal trend except for molybdenum. The Mo_{EF} curve shows values generally above 1 in the lowermost samples and a maximum value (22) in sample 7712. Above this sample, Mo is below the detection limit (Table S1) so that Mo_{EF} values are below 1. The cerium anomaly (Ce^*), though potentially influenced by changes in the detrital flux and sea level variations (Wilde et al., 1996), is negative for all the samples (average -0.048). The temporal variations of Ce^* are somehow opposite to Mo variations, the lowest values being observed in the lowermost samples, while higher values are observed in the upper part of the studied interval (Fig. 5B). Overall, these data point for only moderately reducing conditions during deposition of the two formations. The most reducing conditions, as indicated by the highest enrichment in molybdenum, are observed in the sample 7712, corresponding to the base of the organic-rich, radioactive clays. The AO III Fm. overall appears deposited in slightly more reducing conditions than the AO IV Fm.

3.2.3 Detrital proxies

In addition to Si and Al, elements mostly associated with detrital minerals comprise K (feldspar and/or illitic clay minerals), Ti (rutile, titanite or ilmenite), Zr (zircon) and rare earth elements (REE, which can be associated with several heavy minerals; Mongelli, 1995; Cullers and Podkovyrov, 2002). When normalized to the Al content, the concentration of these elements is a good indicator of changes in the detrital flux.

The geochemical composition of the lowermost five samples is relatively constant (Fig. 6). Only REE ratios show slight variations in this part (Fig. 6B). From sample 7780 upwards the geochemical composition is more variable, with overall higher element/Al ratios for Ti, Hf, Th and Zr (Fig. 6A). The REE profiles also are more variable as indicated by the (La/Lu)_N and (Gd/Yb)_N ratios (Fig. 6B). For major and trace elements a maximum in the Al-normalized content is observed in the sample 7087 (Fig. 6A). This sample is also characterized by a PAAS-normalized REE pattern markedly different from that of the other samples (Fig. 7).

Increase and variability of Al-normalized concentration of the different elements considered here mainly indicate a change from an argillaceous AO III Fm. to a more silty or more sandy AO IV Fm. The relative stability of REE patterns in the studied samples—except for sample 7087—suggests that these changes are mainly due to grain sorting rather than from changes in the origin of the detrital grains. In the case of sample 7087, the REE pattern of which is markedly different, the higher Zr/Al and Hf/Al ratios indicate a higher proportion of heavy minerals in the sediment. Such characteristics are observed in sediments deposited by turbidity currents (e.g. Riboulleau et al., 2014) and can be related to the increased proportion of sandy layers in the upper part of the AO IV Fm., reflecting a relative sea-level fall during the deposition of this formation (Massa, 1988; Carr, 2002).

3.3 Biomarkers

3.3.1 Saturated fraction

The saturate fraction is dominated by a series of linear alkanes (*n*-alkanes). Hopanes, steranes, linear isoprenoids and cycloalkanes are present in lesser proportion.

n-Alkanes range from C₁₃ to C₃₇ with a maximum in C₁₆ or C₁₇ (Fig. 8A). The contribution from long chain *n*-alkanes is generally low, as indicated by the low value of the terrestrial to aquatic *n*-alkane ratio (TAR; Bourbonniere and Meyers, 1996; TAR values 0.06-

0.52). Only the sample 7085 shows a higher contribution of long-chain alkanes (TAR = 0.74). No marked even or odd predominance is observed, as indicated by the CPI values close to 1 (Bray and Evans, 1961). The highest CPI values (1.2) are observed in the two uppermost samples (7087 and 7085). Linear isoprenoids range from C₁₄ to C₂₃, maximum in C₁₉. The Pr/Ph ratio ranges from 1.0 to 2.2. This highest value is obtained for the two uppermost samples.

The distribution of bicyclic and tricyclic terpenoids is similar in all the samples. Bicyclic sesquiterpenoids range from C₁₄ to C₁₆ and are dominated by *ent*-8 β (H)-homodrimane followed by a C₁₅ compound (Fig. 8B). *Ent*-8 β (H)-drimane is also present in lesser amount. Three C₁₉ tricyclic terpanes are also present in low abundance in all the samples. From its mass spectrum, the first eluting compound corresponds to a 13 α (methyl)-tricyclic terpane previously as described by Wang and Simoneit (1995). No other element from this series is apparently present. The following compounds correspond to compounds B and C of Greenwood and George (1999). A series of regular tricyclic terpanes (cheilantanes) ranging from C₁₉ to C₂₆ (max C₂₃) is also observed in all the samples (Fig. 8C).

Hopanoids are present in all the samples but are relatively more abundant in the two uppermost samples. Hopanoids are dominated by a series of C₂₇ to C₃₄ regular $\alpha\beta$ hopanes (Fig. 8C). $\beta\alpha$ Moretanes ranging from C₂₉ to C₃₄ are present in low abundance. In addition, several series of rearranged hopanes are observed: C₂₄ and C₂₆ 17,21-secohopanes (Lu et al., 2009), C₂₉ and C₃₀ 17 α (H)-diahopanes, C₂₇ to C₃₀ 18 α (H)-neohopanes (Ts series) and C₂₉-C₃₂ early eluting series (Moldowan et al., 1991; Farrimond and Telnæs, 1996). Norgammacerane and gammacerane are also detected in low amounts, mostly in the two uppermost samples.

Steroids are present in all the samples, but in very low abundance. They are slightly more abundant in the two uppermost samples. The distribution of compounds is comparable in all the samples with a few variations, and comprises the dia- and regular series, dominated by the C₂₇ and C₂₉ compounds (Fig. 8D, Table 2). Diasteroids are relatively less abundant in the two uppermost samples (Table 2). Methylsteranes and dinosterane isomers were not detected. Short chain steroids C₂₁ to C₂₃ are also observed in all the samples (Fig. 8D). They are generally present in relatively low abundance compared to the longer C₂₇-C₂₉ compounds.

3.3.2 Aromatic fraction

This fraction is dominated by polycyclic aromatic hydrocarbons (PAHs) among which naphthalenes and phenanthrenes. Aromatic biomarkers mostly correspond to triaromatic steroids and arylisoprenoids.

Triaromatic steroids are observed in all the samples (not shown). They range from C₂₀ to C₂₈ and are dominated by the C₂₆ to C₂₈ compounds. Short chain compounds (C₂₀-C₂₂) are however more abundant in the lower samples. The distribution is dominated by the C₂₈ compounds for all the samples, however, the uppermost 2 samples show a higher contribution of the C₂₆ compounds. 3-Methyl and 4-methyltriaromatic steroids are also present in minor proportion in all the samples, however no triaromatic dinosteroids were identified.

Aromatic compounds derived from hopanoids were detected in very low proportions in the two uppermost samples. They mostly correspond to a C₂₄ ABCD-tetraaromatic hopanoid and des-E-D:C-friedo-25-norhopa-5,7,9-triene identified by Hauke et al. (1993). Cadalene is present in trace amounts in all the samples. Retene and the aromatic abietanoids were not detected.

A series of 2,3,6-trimethyl-arylisoprenoids ranging from C₁₃ to C₃₁ (max C₁₈) is observed in all the samples (Fig. 9). This series of compounds is well known as the product of degradation of isorenieratene (Requejo et al., 1992). A second series corresponding to 3,4,5-trimethyl-arylisoprenoids is also observed (Fig. 9). These compounds derived from the degradation of the aromatic carotenoid palaerenieratene (Brown et al., 2000) are present in all the samples except the two uppermost samples (7087 and 7085). Isorenieratane was detected in low amount in all the samples while palaerenieratane is observed in the four lowest samples (7812 to 7712). Several other diagenetic/catagenetic products of isorenieratene previously described by Koopmans et al. (1996) were also detected in all the samples, where they show a similar distribution: a series of diaromatic isoprenoids with a biphenyl head characterized by a m/z 237 fragment ranging from C₁₉ (max) to C₂₂, and C₂₁ to C₂₄ diaromatic compounds characterized by a m/z 133 fragment (compounds I to III, Fig. 9).

Alkylated di-, tri-, tetra-, and pentaaromatics are detected in abundance in all the samples. Naphthalene and methylated counterparts from C₁ to C₆ are present in all the samples. They are dominated by the C₂ or C₃ isomers. Phenanthrenes from C₀ to C₄ are observed in all the samples. Phenanthrene is the most abundant compound. Dibenzofurans are present in all the samples, but their proportion is higher in the uppermost two samples. They range from C₀ to C₂ and are dominated by the C₁ isomers. Dibenzothiophenes (DBT) from C₀ to C₃ are observed in all the samples. Their relative abundance is however lower in the two uppermost

samples. Their distribution is dominated by DBT or by 4-methyl-DBT. Fluoranthene, pyrene and methylated counterparts from C₁ to C₂ are detected in all the samples, however, their relative abundance is higher in the two uppermost samples. Triphenylene, chrysene their methylated counterparts from C₁ to C₄ are present in all the samples. The relative abundance of the C₀ counterparts and relative distribution of methylated homologs however differs between the lower four and upper two samples. In particular, a large dominance of the non-methylated counterparts is observed in the two uppermost samples. Isomer distribution ratios of selected PAHs are presented in Table 2.

Highly condensed PAH's such as benzo[e]pyrene, benzo[ghi]perylene or coronene are observed only in the uppermost two samples. They are not observed in the lowermost four samples, even by selective ion detection.

4 Discussion

4.1 Organic matter maturity

From the Rock-Eval Tmax values (Fig. 4A), the studied samples are located at the beginning of the oil window, with slightly higher maturity for samples from core 8 than for samples from core 6. This range of maturity is also indicated by the color and state of preservation of the miospores, acritarchs and *Tasmanites* in the samples (Spina et al., 2017). Using average correlation between biomarker isomerization ratios and thermal maturity (Peters et al., 2005), the ratio obtained in the studied samples (Table 2) tends to indicate a higher maturity than Rock-Eval Tmax, samples from core 8 having reached the peak of oil generation while samples from core 6 would be slightly less mature. Methylphenanthrene isomerization ratios indicate even more maturity with calculated vitrinite reflectance values higher than 0.7 % (Table 2). Hrouda (2004) previously observed a relatively poor correlation between molecular maturity indicators, vitrinite reflectance and Rock Eval in the Upper Devonian formations of the Ghadames Basin. Nevertheless, the relatively low maturity indicated by the Rock Eval data is supported by vitrinite reflectance measurements of 0.51 to 0.59 performed by Underdown et al. (2007) on the same depth interval. These maturities are also consistent with those obtained by Hrouda (2004) for Upper Devonian formations in nearby cores from the Ghadames Basin. This low maturity allows interpret further the biomarker distribution in the analyzed samples.

4.2 Organic matter source

Based on palynofacies analysis, TOC/N ratio, and biomarker content, the OM in the Aouinet Ouenine Fm. shows a mixed contribution from continental and marine fractions. The palynofacies content and TOC/N ratio allow to clearly distinguish the lower samples (AO III Fm. and radioactive shales) dominated by marine elements from the upper part of AO IV Fm., where the contribution from continental OM is more significant. Such distinction is less obvious basing on the biomarkers, since terrestrial plant biomarkers are almost absent from the record. Terrestrial plants in the Late Devonian were mostly restricted to bryophytes and early tracheophytes (Taylor et al., 2009). Earliest plants have not been so far related to an abundance of diversified biomarkers (Versteegh and Riboulleau, 2010), nevertheless a few higher plant terpenoids were previously identified in Devonian rocks, including in sediments of the Ghadames Basin (Kashirtsev et al., 2010; Romero-Sarmiento et al., 2011). Devonian plants already possessed a waxy coating made of long chain *n*-alkanes, typical of plant cuticles (Eglinton and Hamilton, 1967; Versteegh and Riboulleau, 2010). The dominance of short chain *n*-alkanes ($<C_{20}$) in the samples is typical of sedimentary OM dominated by an autochthonous algal input (Gelpi et al., 1970). Nevertheless, the increased proportion of long *n*-alkanes in samples from core 6, as indicated by the higher TAR values (Table 2), is consistent with an increased contribution of OM of terrestrial origin. Contribution from land-derived OM in the two samples from core 6 (7087 and 7085) is also indicated by the PAH distribution, showing a high proportion of dibenzofurans, which have been related to dehydration of plant cellulose and polysaccharides (Sephton et al., 1999) and the contribution of unsubstituted highly condensed PAH such as coronene, which have been related to wildfires (Marynowski and Simoneit, 2009).

The sterane distribution, dominated by the C_{29} and C_{27} isomer (Table 2), is typical of OM of marine origin. Methylated counterparts, in particular 4-methyl isomers are generally of algal origin. Dinosteroids and 4-methyl steroids are often assigned to dinoflagellates and/or acritarchs. Though dinosteroids are absent from the extracts of D1-26 samples, the methylated steroids can be related to the marine algal input. In particular, several studies of Paleozoic rocks and petroleum have related these compounds to acritarchs (Moldowan et al., 1996; Moldowan and Talyzina, 1998).

Hopanoids and their diagenesis products, as well as drimane and homodrimane are related to bacterial inputs (Ourisson et al., 1979; Alexander et al., 1983). The sterane/hopane ratio is often used as an indicator of the proportion of bacterial biomass compared to the eukaryotic one. The highest values of this ratio are observed in the samples from the AO IV

Fm. and therefore suggest higher contribution from eukaryotes in these samples, while a higher contribution from bacterial biomass is observed in the lowermost four samples.

Though their origin is still debated, tricyclic terpanes (cheilantanes) have been mostly related to the contribution from prasinophytes and in particular tasmanaceans (Aquino Neto et al., 1992; Dutta et al., 2006). This is consistent with the frequent observation of *Tasmanites* in the palynofacies of D1-26 samples (Fig. 3). High relative abundances of cheilantanes, indicated by the tricyclic/17 α hopane ratio are observed in the four lower samples (7812 to 7712, Table 2). The upper two samples (7087 and 7085) show lower relative abundance of cheilantanes (Table 2), which can be related to the lower relative abundance of *Tasmanites* in the palynofacies (Fig. 3). Abundance of prasinophytes in general or of *Tasmanites* in particular, has also been related to an increase of the C₂₈/C₂₉ sterane ratio (Schwark and Empt, 2006). This is for instance the case at the Frasnian-Famennian transition in the famous Kowala section in Poland, where prasinophytes represent 80 to 90% of palynomorphs (Filipiak, 2002; Schwark and Empt, 2006). No relationship is observed between the C₂₈/C₂₉ sterane ratio and the relative abundance of *Tasmanites* in the palynofacies of D1-26 samples (Table 2, Fig. 3), which could be related to the fact that prasinophytes rarely account for more than 40% of palynomorphs in these samples.

4.3 Redox conditions

Isorenieratene is a pigment synthesized by green sulfur bacteria (*Chlorobiaceae*), obligate anaerobes requiring light and free sulfides for living (Liaaen-Jensen, 1978; Overmann, 2008). In particular, *Chlorobiaceae* have been observed in the water column of the Black Sea (Overmann et al., 1992). For this reason, the observation of isorenieratene derivatives in sediment extracts is often used as an indicator of photic zone anoxia (Summons and Powell, 1986). The source of palaerenieratene is unknown but several arguments suggest this compound originates from extinct organisms related to *Chlorobiaceae* (Hartgers et al., 1994). A recent review of the occurrence of palaerenieratene in oils and sediments confirms this compound is almost restricted to the Paleozoic (French et al., 2015). When present, palaerenieratene is often more abundant than isorenieratene (Requejo et al., 1992; Behrens et al., 1998; Clifford et al., 1998; Joachimski et al., 2001; Armstrong et al., 2009; Bushnev, 2009; Maslen et al., 2009; Racka et al., 2010; Marynowski et al., 2012; Melendez et al., 2013; Tulipani et al., 2015). Changes in the proportion of palaerenieratene vs. isorenieratene derivatives in relation to paleoenvironments are rarely reported in the literature. Nevertheless, in the Middle Devonian Keg River Fm., Behrens et al. (1998) observed that while both

palaereniaratane and isorenieratane and their derivatives were present in transgressive facies, the regressive facies of this formation only showed the presence of isorenieratane and its derivatives.

The total concentration of isorenieratene and palaerenieratene derivatives in the extracts of D1-26 generally decreases up-section through the studied interval, with the maximum value observed in the lowermost sample and the minimum value in the uppermost sample (Fig. 10A). When plotted in the Pr/Ph ratio versus the aryl isoprenoid ratio (AIR*) modified from Schwark and Frimmel (2004), the lowermost four samples from D1-26 plot in a sector of relatively persistent photic-zone anoxia and reducing sediment, while the uppermost two plot in an area of oxygenated sediment with more episodic photic-zone anoxia (Fig. 10C). Location of D1-26 samples in this plot are also consistent with previous observations in the Posidonia black shale, regarding relative sea-level variations (Schwark and Frimmel, 2004): samples from AO III Fm. and radioactive shales, deposited in a rising sea level plot in the “intermediate sea level” sector, while samples from the higher part of AO IV Fm., deposited in a regressive context, plot in the “high sea level” and “low sea level” sectors. The ratio of the 3,4,5-trimethyl- to 2,3,6-trimethyl-aryl isoprenoids (palaerenieratene to isorenieratene derivatives ratio, PIR) slightly increases from the base to the middle part of the series and falls to 0 in the uppermost two samples (Fig. 10B). The disappearance of palaerenieratane and its derivatives in the regressive, highest part of AO IV Fm. is consistent with the previous observations of Behrens et al. (1998).

The abundance of prasinophytes in the palynofacies of D1-26 samples shows a good parallel with the relative abundance of isorenieratene and palaerenieratene and their derivatives in the extracts (Figs. 3, 10): present in significant proportion in AO III Fm. and the radioactive shales, they become minor in the upper part of AO IV Fm. This observation fully supports the proposition of Prauss (2007), who suggested that prasinophyte abundance in sediments is controlled by, and therefore is an indicator of, reducing conditions in the photic zone.

Based on redox-sensitive trace metal concentration, the studied Devonian sediments from borehole D1-26 appear deposited under mainly oxic conditions. This appears inconsistent with the occurrence of *Chlorobiaceae* biomarkers and of *Tasmanites*, in all the samples. The only sample showing enrichment in redox-sensitive trace metals is sample 7712, belonging to the very base of the radioactive shales. Its enrichment in uranium remains modest (EF = 2) while a more notable enrichment is observed for molybdenum (EF=21). The contrasted enrichments in Mo and U, respectively suggest molybdenum enrichment

associated with redox cycling of manganese or iron oxyhydroxides, at the sediment water-interface or within the water column (Algeo and Tribovillard, 2009). It means that the water column would have been regularly oxygenated but the redox-cline would have been lying at short distance beneath the sediment-water interface, thence an anoxic sediment. High enrichment in trace elements in the radioactive shales is reported by Weyant and Massa (1991), nevertheless these authors did not indicate from which boreholes or outcrops these data originated, moreover, no data are available for the rest of the Aouinet Ouenine group.

The discrepancy between molecular and inorganic redox proxies could result from a so-called “reservoir effect” (Algeo, 2004; Algeo and Lyons, 2006). When a basin is poorly connected with the open sea, scavenging of molybdenum and other redox-sensitive trace metals by the precipitation of authigenic minerals may overcome the inflow of these elements to the basin. This process results in the progressive depletion of these elements in the water column (Algeo, 2004; Algeo and Lyons, 2006). Sediments deposited below such depleted water column will not be enriched in trace elements, even if they are organic-rich and/or anoxic (Algeo, 2004; Tribovillard et al., 2008). During the Devonian, the North African domain was structured into several basins separated by topographic highs (Lüning et al., 2003; Guiraud et al., 2005; Wendt et al., 2006; Frizon de Lamotte et al., 2013). Poor connections of the bottom part of these basins with the open sea are suggested by the widespread deposition of organic-rich sediments in the various North African basins (Lüning et al., 2003; Wendt et al., 2006). Such configuration might therefore have favored progressive depletion of the water column in trace elements, leading to the poor enrichment observed in the studied samples from borehole D1-26.

4.4 Organic matter deposition and relationships with the F/F events

The studied samples from the Aouinet Ouenine Group where the TOC content was measured do not include the “radioactive shales” and are not very organic rich (Table 1). Nevertheless, the TOC values from 1 to 2% are consistent with previous studies of the Aouinet Ouenine Group in nearby boreholes (Hrouda, 2004).

The different methods used here indicate that in the Frasnian AO III Fm., OM mainly originates from an algo-bacterial biomass. The biomarker content and the significant contribution of *Tasmanites* in the palynofacies indicate that conditions in the sediment and water column were frequently reducing (Fig. 11A). The notable proportion of AOM in the palynofacies (Fig. 3) points to efficient OM preservation under reducing conditions. The moderate TOC values of the sediment despite recurrent anoxia, the notable proportion of

land-derived OM, and the abundant and well-diversified fauna of acritarchs in the palynofacies, nevertheless suggest that planktonic productivity was moderate (Fig. 11A). Consistent, Massa (1988) indicates that the facies in the AO III Fm. is confined and that the macrofauna are impoverished.

The palynofacies content and biomarkers in the radioactive shales from D1-26 borehole only moderately differ with those of the AO III Fm., indicating a dominantly algal bacterial OM and frequently anoxic conditions in the sediment and water column (Fig. 11B,C). The high TOC content documented in the radioactive shales (Massa, 1988) is however a major difference with the AO III Fm., which, added to the low proportion of terrestrial OM in the palynofacies, suggests increased planktonic productivity and eutrophic conditions during deposition (Fig. 11B,C). Chitinozoans have been recently shown to be produced by relatively shelfal epiplanktonic organisms (Vandenbroucke et al., 2010). Their decreased abundance in the palynofacies of the radioactive shales compared to AO III Fm. as well as the absence of sporomorphs are consistent with a more pelagic setting, and therefore an increased sea level (Fig. 11B,C).

Eutrophication leading to the deposition of the radioactive shales could have resulted from the input of land-derived nutrients in the basin (Algeo et al., 1995; Averbuch et al., 2005), however, the Late Devonian was dominantly a period of subsidence and not of uplift around the Ghadames Basin (Craig et al., 2008). Moreover, though the detrital elements suggest a change in the detrital flux between AO III and AO IV formations (Fig. 6), this change appears to be modest (Fig. 7) and is already observed in sample 7780, several meters below the radioactive shales. The present data therefore do not support an increased supply of land-derived nutrients in the Ghadames Basin. Since depositional conditions were already reducing in the Ghadames Basin during the deposition of AO III Fm. (Fig. 11A), nutrients could either have been recycled from the water-column by more frequent water-column mixing (Murphy et al., 2000), or originate from incursions of nutrient-rich water from the Paleotethys. On the one hand, we do not have arguments regarding more frequent water-column mixing. On the other hand, incursions of water from the open sea were likely favored by the sea level rise, and are supported by the Mo/Al ratio of the samples (Fig. 5). As previously discussed, the redox-sensitive trace elements suggest that the water column of the Ghadames Basin was relatively depleted in dissolved trace metals during deposition of AO III Fm. The enrichment in molybdenum observed in sample 7712 was therefore only possible because of Mo inputs, refuelled by the ingression of « new » sea water into the basin (Algeo, 2004; Algeo and Lyons, 2006).

The ambiguity on the age of samples from core 7 compels us to consider several scenarios relating the radioactive shales with the Frasnian-Famennian transition events. Massa (1988; pers. comm., 2005) suggested that the radioactive shales were of latest Frasnian age. If so, the radioactive shales of the Ghadames Basin might be equivalent to the Upper Kellwasser level. The two Kellwasser levels are well known in Morocco (Riquier et al., 2005), and the Frasnian radioactive shales have been related to the upper Kellwasser level in the Berkine area in Algeria (Lüning et al., 2004) and in southern Tunisia (Soua, 2014). The deposition of the upper Kellwasser level and of the radioactive shales from the Ghadames Basin were both related to a Late Frasnian second-order sea-level high-stand (Carr, 2002; Chen and Tucker, 2003; Dardour et al., 2004; Bond and Wignall, 2008). The eutrophication and possible development of more reducing conditions during the Upper Kellwasser event could result from the incursion of nutrient-rich and anoxic waters into the Ghadames Basin originating from the Paleotethys (Fig. 11B). This option would give support to the scenarios relating the deposition of the Kellwasser level to widespread oceanic eutrophication and development of anoxia in marginal basins (Joachimski and Buggisch, 1993; Bond et al., 2004; Averbuch et al., 2005; Riquier et al., 2006; Carmichael et al., 2014).

Alternately, if the radioactive shales were of early Famennian age (Weyant and Massa, 1991), their deposition could be related to the early Famennian sea-level rise documented in several basins (Sandberg et al., 2002; Haq and Schutter, 2008). Though sea-level variations around the Frasnian-Famennian are still a matter of discussion (Bond and Wignall, 2008), this could correspond to T-R cycle IIe of Johnson et al. (1985). Deposition of black shales in basinal settings during the early Famennian have been described in Poland, France, Morocco and Eastern North America (Wendt and Belka, 1991; Bond et al., 2004; Schieber and Lazar, 2004; Riquier et al., 2005) and may be related to this sea-level rise. For instance, in the Illinois Basin, the lowermost deposits of Famennian age in the New Albany shales show a more basinal facies and more eutrophic conditions than late Frasnian deposits (de la Rue et al., 2007). The radioactive shales would then correspond to a model of « transgressive black shales », where connection with the open sea allows the entrance of nutrient-rich water and eutrophication (Fig. 11C), as observed in the Cariaco Basin (Venezuela) tuned to Quaternary sea-level oscillations (Peterson et al., 2000).

In addition to its well known $\delta^{13}\text{C}$ excursion (Joachimski and Buggisch, 1993), recent studies indicate the Upper Kellwasser level could be characterized by molecular signatures: increase of the $\text{C}_{28}/\text{C}_{29}$ sterane ratio (Schwark and Empt, 2006) related to the prominence of prasinophytes and low values of the hopane/sterane ratio, indicating high relative input of

eukaryotes (Haddad et al., 2016). The radioactive shales of borehole D1-26 show a slight decrease of the C_{28}/C_{29} sterane ratio compared to the other samples (Table 2). The hopane/sterane ratio is lower in the radioactive shales than in the samples from the AO IV formation (Table 2). Nevertheless, similarly low hopane/sterane ratios are observed in the samples from AO III Fm. (Table 2). None of these parameters therefore allows clarify the ambiguity on the age of samples from core 7.

The highest part of AO IV Fm. is characterized by a change in sedimentary facies compared to underlying levels. The proportion of sand increases (Fig. 2, 6), indicating a more proximal depositional setting. This lithological evolution is related to a third- or second-order relative sea-level fall (Carr, 2002; Dardour et al., 2004; Fig. 11D). Consistent, the OM content changes and is dominated by land-derived material (Fig. 3). The lower sea-level and restricted conditions could have favored oligotrophic conditions in the Ghadames Basin (Fig. 11D). Palynofacies and biomarkers point to relatively oxidative conditions in the sediment (Fig. 11D), with only episodic photic zone anoxia. Though conditions were less prone to OM preservation, the highest part of the AO IV Fm. shows TOC values comparable to the AO III Fm. OM enrichment in this interval therefore appears to be related to an increased contribution of relatively resistant and less oil-prone terrestrial OM (Fig. 11D).

5. Conclusions

The multidisciplinary analysis of the Frasnian-Famennian Aouinet Ouenine III and IV formations in borehole D1-26, located in the Ghadames Basin (Libya) allowed to better understand the factors of OM enrichment in this important petroleum system. Our data indicate that during the Late Frasnian the Ghadames Basin was restricted and subject to frequent episodes of photic-zone anoxia. These conditions allowed a good preservation of OM derived from a moderate phytoplanktonic productivity. The beginning of OA IV Fm. is marked by the deposition of a radioactive shale interval, well known in core loggings of the Ghadames Basin, and corresponds to an episode of increased primary productivity and photic-zone anoxia. Though ambiguity remains on the exact age of this radioactive shale interval, our data suggest eutrophication was favored by the entrance of nutrient-rich waters into the Ghadames Basin. The upper part of OA IV Fm. is marked by a facies change associated with the Famennian regression. Photic zone anoxia was less frequent but the sediment is still organic-rich because of increased delivery of terrestrial OM.

The Late Devonian was a period of important changes in the carbon cycle marked by several episodes of oceanic anoxia and biotic crisis. While detailed and high-resolution studies of Late Devonian events in European, Asian and North-American successions are numerous, similar studies in North Gondwana remain rare and are mainly restricted to Morocco and Algeria, where the sediments present relatively proximal depositional facies. The present study demonstrates that the Ghadames Basin, in addition to its important petroleum interest, contains a rich sedimentary succession prone to future high-resolution studies of the Late Devonian events in North Gondwana.

Acknowledgments

This article is dedicated to the memory of Dominique Massa who provided the samples and supported research in the Paleozoic of North Africa. Jonhatan Redfern (University of Manchester) is acknowledged for providing detailed information on D1-26 borehole. This is a contribution to the project “The terrestrialization process: modeling complex interaction at the biosphere–geosphere interface” financially supported by the INSU ECLIPSE program, and the ANR TERRES project (ANR-10-BLAN-0607). Université de Lille is acknowledged for a 12-month postdoctoral grant awarded to A. Spina. Nicolas Visez and Sylvie Gosselin (PC2A, Université de Lille) are acknowledged for GC-MS injections. Thomas Algeo, Grzegorz Racki and an anonymous reviewer are acknowledged for constructive comments on the first version of this manuscript.

Literature

- Alexander, R., Kagi, R., Noble, R. (1983) Identification of the bicyclic sesquiterpenes drimane and eudesmane in petroleum. *Journal of the Chemical Society, Chemical Communications*, 5, 226–228.
- Algeo, T.J. (2004) Can marine anoxic events draw down the trace element inventory of seawater? *Geology*, **32**, 12, 1057.
- Algeo, T.J., Berner, R.A., Maynard, J.B., Scheckler, S.E. (1995) Late Devonian oceanic anoxic events and biotic crises: “rooted” in the evolution of vascular land plants? *GSA Today*, **5**, 3, 45, 64–66.

- Algeo, T.J., Lyons, T.W. (2006) Mo–total organic carbon covariation in modern anoxic marine environments: Implications for analysis of paleoredox and paleohydrographic conditions. *Paleoceanography*, **21**, 1, PA1016.
- Algeo, T.J., Scheckler, S.E. (1998) Terrestrial-marine teleconnections in the Devonian: links between the evolution of land plants, weathering processes, and marine anoxic events. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **353**, 1365, 113–130.
- Algeo, T.J., Tribovillard, N. (2009) Environmental analysis of paleoceanographic systems based on molybdenum–uranium covariation. *Chemical Geology*, **268**, 3–4, 211–225.
- Aquino Neto, F.R., Trigüis, J., Azevedo, D.A., Rodriques, R., Simoneit, B.R.T. (1992) Organic geochemistry of geographically unrelated tasmanites. *Organic Geochemistry*, **18**, 6, 791–803.
- Armstrong, H.A., Abbott, G.D., Turner, B.R., Makhoul, I.M., Muhammad, A.B., Pedentchouk, N., Peters, H. (2009) Black shale deposition in an Upper Ordovician–Silurian permanently stratified, peri-glacial basin, southern Jordan. *Organic-Carbon-Rich Sediments through the Phanerozoic: Processes, Progress, and Perspectives*, **273**, 3, 368–377.
- Averbuch, O., Tribovillard, N., Devleeschouwer, X., Riquier, L., Mistiaen, B., Van Vliet-Lanoe, B. (2005) Mountain building-enhanced continental weathering and organic carbon burial as major causes for climatic cooling at the Frasnian–Famennian boundary (c. 376 Ma)? *Terra Nova*, **17**, 1, 25–34.
- Becker, R.T., Gradstein, F.M., Hammer, O. (2012) Chapter 22 - The Devonian Period. *The Geologic Time Scale*, Elsevier, Boston, pp. 559–601.
- Becker, R.T., Königshof, P., Brett, C.E. (2016) *Devonian climate, sea level and evolutionary events*. Geological Society of London, 481 p.
- Behar, F., Beaumont, V., De B. Pentead, H.L. (2001) Technologie Rock-Eval 6 : performances et développements. *Oil & Gas Science and Technology - Rev. IFP*, **56**, 2, 111–134.
- Behrens, A., Wilkes, H., Schaeffer, P., Clegg, H., Albrecht, P. (1998) Molecular characterization of organic matter in sediments from the Keg River formation (Elk Point group), western Canada sedimentary basin. *Organic Geochemistry*, **29**, 8, 1905–1920.
- Berner, R.A. (1982) Burial of organic carbon and pyrite sulfur in the modern ocean: its geochemical and environmental significance. *American Journal of Science*, **282**, 451–473.
- Berner, R.A., Raiswell, R. (1983) Burial of organic carbon and pyrite sulfur in sediments over Phanerozoic time: a new theory. *Geochimica et Cosmochimica Acta*, **47**, 855–862.

- Bond, D., Wignall, P.B., Racki, G. (2004) Extent and duration of marine anoxia during the Frasnian–Famennian (Late Devonian) mass extinction in Poland, Germany, Austria and France. *Geological Magazine*, **141**, 2, 173–193.
- Bond, D.P.G., Wignall, P.B. (2008) The role of sea-level change and marine anoxia in the Frasnian–Famennian (Late Devonian) mass extinction. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **263**, 3, 107–118.
- Boote, D.R.D., Clark-Lowes, D.D., Traut, M.W. (1998) Palaeozoic petroleum systems of North Africa. *Geological Society, London, Special Publications*, **132**, 1, 7.
- Bourbonniere, R.A., Meyers, P.A. (1996) Sedimentary geolipid records of historical changes in the watersheds and productivities of Lakes Ontario and Erie. *Limnology and Oceanography*, **41**, 2, 352–359.
- Bray, E.E., Evans, E.D. (1961) Distribution of *n*-paraffins as a clue to recognition of source beds. *Geochimica et Cosmochimica Acta*, **22**, 2–15.
- Brown, E.T., Le Callonnec, L., German, C.R. (2000) Geochemical cycling of redox-sensitive metals in sediments from Lake Malawi: a diagnostic paleotracer for episodic changes in mixing depth. *Geochimica et Cosmochimica Acta*, **64**, 20, 3515–3523.
- Buggisch, W. (1991) The global Frasnian-Famennian »Kellwasser Event«. *Geologische Rundschau*, **80**, 1, 49–72.
- Bushnev, D. (2009) Organic matter of the Ukhta domanik. *Doklady Earth Sciences*, **426**, 4, 1531–8354.
- Canaple, J. (1963) *Synthèse régionale du sud-tunisien. Corrélations entre les forages BS1-OS1-SN1-DN1-MG1-DI.26*. Compagnie Franco-tunisienne des pétroles,.
- Carmichael, S.K., Waters, J.A., Suttner, T.J., Kido, E., DeReuil, A.A. (2014) A new model for the Kellwasser Anoxia Events (Late Devonian): Shallow water anoxia in an open oceanic setting in the Central Asian Orogenic Belt. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **399**, 394–403.
- Carr, I.D. (2002) Second-order sequence stratigraphy of the Palaeozoic of North Africa. *Journal of Petroleum Geology*, **25**, 3, 259–280.
- Chen, D., Tucker, M.E. (2003) The Frasnian–Famennian mass extinction: insights from high-resolution sequence stratigraphy and cyclostratigraphy in South China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **193**, 1, 87–111.
- Clifford, D.J., Clayton, J.L., Sinninghe Damsté, J.S. (1998) 2,3,6-/3,4,5-Trimethyl substituted diaryl carotenoid derivatives (Chlorobiaceae) in petroleum of the Belarussian Pripyat River Basin. *Organic Geochemistry*, **29**, 5–7, 1253–1267.

757 Colbath, G.K., Grenfell, H.R. (1995) Review of biological affinities of Paleozoic acid-
758 resistant, organic-walled eukaryotic algal microfossils (including “acritarchs”). *Review of*
759 *Palaeobotany and Palynology*, **86**, 3, 287–314.

760 Craig, J., Rizzi, C., Said, F., Thusu, B., Lüning, S., Asbali, A.I., Keeley, M.L., Bell, J.F.,
761 Durham, M.J., Eales, M.H., et al. (2008) Structural styles and prospectivity in the
762 Precambrian and Palaeozoic hydrocarbon systems of North Africa. In : Salem, M.J., Oun,
763 K.M., Essed, A.S., (eds.). *The Geology of East Libya, Vol. IV*, pp. 51–122.

764 Cullers, R.L., Podkovyrov, V.N. (2002) The source and origin of terrigenous sedimentary
765 rocks in the Mesoproterozoic Ui group, southeastern Russia. *Precambrian Research*, **117**, 3–
766 4, 157–183.

767 Dardour, A.M., Boote, D.R.D., Baird, A.W. (2004) Stratigraphic controls on Palaeozoic
768 petroleum systems, Ghadames Basin, Libya. *Journal of Petroleum Geology*, **27**, 2, 141–162.

769 Dutta, S., Greenwood, P.F., Brocke, R., Schaefer, R.G., Mann, U. (2006) New insights into
770 the relationship between *Tasmanites* and tricyclic terpenoids. *Organic Geochemistry*, **37**, 1,
771 117–127.

772 Echikh, K. (1998) Geology and hydrocarbon occurrences in the Ghadames Basin, Algeria,
773 Tunisia, Libya. *Geological Society, London, Special Publications*, **132**, 1, 109.

774 Eglinton, G., Hamilton, R.J. (1967) Leaf epicuticular waxes. *Science*, **156**, 3780, 1322–1335.

775 Evitt, W.R. (1963) A discussion and proposals concerning fossil dinoflagellates,
776 hystrichospheres, and acritarchs, I. *Proceedings of the National Academy of Sciences*, **49**, 2,
777 158–164.

778 Farrimond, P., Telnæs, N. (1996) Three series of rearranged hopanes in Toarcian sediments
779 (northern Italy). *Organic Geochemistry*, **25**, 3–4, 165–177.

780 Filipiak, P. (2002) Palynofacies around the Frasnian/Famennian boundary in the Holy Cross
781 Mountains, southern Poland. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **181**, 1–3,
782 313–324.

783 French, K.L., Rocher, D., Zumberge, J.E., Summons, R.E. (2015) Assessing the distribution
784 of sedimentary C₄₀ carotenoids through time. *Geobiology*, **13**, 2, 139–151.

785 Frizon de Lamotte, D., Tavakoli-Shirazi, S., Leturmy, P., Averbuch, O., Mouchot, N., Raulin,
786 C., Leparmentier, F., Blanpied, C., Ringenbach, J.-C. (2013) Evidence for Late Devonian
787 vertical movements and extensional deformation in northern Africa and Arabia: Integration in
788 the geodynamics of the Devonian world. *Tectonics*, **32**, 2, 107–122.

789 Gelpi, E., Schneider, H., Mann, J., Oro, J. (1970) Hydrocarbons of geochemical significance
790 in microscopic algae. *Phytochemistry*, **9**, 3, 603–612.

791 Godderis, Y., Joachimski, M.M. (2004) Global change in the Late Devonian: modelling the
792 Frasnian-Famennian short-term carbon isotope excursions. *Palaeogeography,*
793 *Palaeoclimatology, Palaeoecology*, **202**, 3–4, 309–329.

794 Greenwood, P.F., George, S.C. (1999) Mass spectral characteristics of C₁₉ and C₂₀ tricyclic
795 terpanes detected in Latrobe Tasmanite oil shale. *European Mass Spectrometry*, **5**, 3, 221–
796 230.

797 Guiraud, R., Bosworth, W., Thierry, J., Delplanque, A. (2005) Phanerozoic geological
798 evolution of Northern and Central Africa: An overview. *Phanerozoic Evolution of Africa*, **43**,
799 *I*, 83–143.

800 Haddad, E.E., Tuite, M.L., Martinez, A.M., Williford, K., Boyer, D.L., Droser, M.L., Love,
801 G.D. (2016) Lipid biomarker stratigraphic records through the Late Devonian
802 Frasnian/Famennian boundary: Comparison of high- and low-latitude epicontinental marine
803 settings. *Organic Geochemistry*, **98**, 38–53.

804 Haq, B.U., Schutter, S.R. (2008) A chronology of Paleozoic sea-level changes. *Science*, **322**,
805 5898, 64.

806 Hart, G.F. (1986) Origin and classification of organic matter in clastic systems. *Palynology*,
807 **10**, *I*, 1.

808 Hartgers, W.A., Sinninghe Damsté, J.S., Requejo, A.G., Allan, J., Hayes, J.M., Ling, Y., Xie,
809 T.-M., Primack, J., de Leeuw, J.W. (1994) A molecular and carbon isotopic study towards the
810 origin and diagenetic fate of diaromatic carotenoids. *Organic Geochemistry*, **22**, 3, 703–725.

811 Hauke, V., Wehrung, P., Hussler, G., Trendel, J.M., Albrecht, P., Riva, A., Connan, J. (1993)
812 Rearranged des-*E*-hopanoid hydrocarbons in sediments and petroleum. *Organic*
813 *Geochemistry*, **20**, 3, 415–423.

814 House, M.R. (2002) Strength, timing, setting and cause of mid-Palaeozoic extinctions.
815 *Palaeogeography, Palaeoclimatology, Palaeoecology*, **181**, 1–3, 5–25.

816 Hrouda, M. (2004) The hydrocarbon source potential of the Palaeozoic rocks of the
817 Ghadames Basin, NW Libya. Newcastle University.

818 Joachimski, M.M., Buggisch, W. (1993) Anoxic events in the late Frasnian—Causes of the
819 Frasnian-Famennian faunal crisis? *Geology*, **21**, 8, 675–678.

820 Joachimski, M.M., Buggisch, W. (2002) Conodont apatite $\delta^{18}\text{O}$ signatures indicate climatic
821 cooling as a trigger of the Late Devonian mass extinction. *Geology*, **30**, 8, 711.

822 Joachimski, M.M., Ostertag-Henning, C., Pancost, R.D., Strauss, H., Freeman, K.H., Littke,
823 R., Sinninghe Damsté, J.S., Racki, G. (2001) Water column anoxia, enhanced productivity

824 and concomitant changes in $\delta^{13}\text{C}$ and $\delta^{34}\text{S}$ across the Frasnian-Famennian boundary (Kowala
825 - Holy Cross Mountains/Poland). *Chemical Geology*, **175**, 1–2, 109–131.

826 Johnson, J.G., Klapper, G., Sandberg, C.A. (1985) Devonian eustatic fluctuations in
827 Euramerica. *GSA Bulletin*, **96**, 5, 567–587.

828 Kashirtsev, V.A., Moskvina, V.I., Fomin, A.N., Chalaya, O.N. (2010) Terpanes and steranes in
829 coals of different genetic types in Siberia. *Russian Geology and Geophysics*, **51**, 4, 404–411.

830 Koopmans, M.P., Köster, J., Van Kaam-Peters, H.M.E., Kenig, F., Schouten, S., Hartgers,
831 W.A., de Leeuw, J.W., Sinninghe Damsté, J.S. (1996) Diagenetic and catagenetic products of
832 isorenieratene: Molecular indicators for photic zone anoxia. *Geochimica et Cosmochimica*
833 *Acta*, **60**, 22, 4467–4496.

834 Leventhal, J.S. (1983) An interpretation of carbon and sulfur relationships in Black Sea
835 sediments as indicators of environments of deposition. *Geochimica et Cosmochimica Acta*,
836 **47**, 133–137.

837 Liaaen-Jensen, S. (1978) Marine carotenoids. In : Scheuer, P.J., (ed.). *Marine Natural*
838 *Products, Chemical and Biological Perspectives*, Academic Press, pp. 1–73.

839 Loboziak, S., Streel, M. (1989) Middle-upper Devonian miospores from the Ghadamis Basin
840 (Tunisia-Libya): Systematics and stratigraphy. *Review of Palaeobotany and Palynology*, **58**,
841 2–4, 173–196.

842 Lu, H., Chen, T., Grice, K., Greenwood, P., Peng, P., Sheng, G. (2009) Distribution and
843 significance of novel low molecular weight steranes in an immature evaporitic sediment from
844 the Jinxian Sag, North China. *Organic Geochemistry*, **40**, 8, 902–911.

845 Lüning, S., Adamson, K., Craig, J. (2003) Frasnian organic-rich shales in North Africa:
846 regional distribution and depositional model. *Geological Society, London, Special*
847 *Publications*, **207**, 1, 165.

848 Lüning, S., Wendt, J., Belka, Z., Kaufmann, B. (2004) Temporal–spatial reconstruction of the
849 early Frasnian (Late Devonian) anoxia in NW Africa: new field data from the Ahnet Basin
850 (Algeria). *Sedimentary Geology*, **163**, 3, 237–264.

851 Martin, F. (1993) Acritarchs: a review. *Biological Reviews*, **68**, 4, 475–537.

852 Martín-Closas, C. (2003) The fossil record and evolution of freshwater plants: A review.
853 *Geologica Acta*, **1**, 4, 315–338.

854 Marynowski, L., Simoneit, B.R.T. (2009) Widespread Upper Triassic to Lower Jurassic
855 wildfire records from Poland: evidence from charcoal and pyrolytic polycyclic aromatic
856 hydrocarbons. *Palaios*, **24**, 12, 785–798.

857 Marynowski, L., Zatoń, M., Rakociński, M., Filipiak, P., Kurkiewicz, S., Pearce, T.J. (2012)
 858 Deciphering the upper Famennian Hangenberg Black Shale depositional environments based
 859 on multi-proxy record. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **346**, 66–86.
 860 Maslen, E., Grice, K., Gale, J.D., Hallmann, C., Horsfield, B. (2009) Crocetane: A potential
 861 marker of photic zone euxinia in thermally mature sediments and crude oils of Devonian age.
 862 *Organic Geochemistry*, **40**, 1, 1–11.
 863 Massa, D. (1988) Paléozoïque de Libye occidentale : stratigraphie et paléogéographie.
 864 McGhee, G.R., Clapham, M.E., Sheehan, P.M., Bottjer, D.J., Droser, M.L. (2013) A new
 865 ecological-severity ranking of major Phanerozoic biodiversity crises. *Palaeogeography,*
 866 *Palaeoclimatology, Palaeoecology*, **370**, 260–270.
 867 McLennan, S.M. (1989) Rare earth elements in sedimentary rocks; influence of provenance
 868 and sedimentary processes. *Reviews in Mineralogy and Geochemistry*, **21**, 1, 169–200.
 869 Melendez, I., Grice, K., Trinajstić, K., Ladjavardi, M., Greenwood, P., Thompson, K. (2013)
 870 Biomarkers reveal the role of photic zone euxinia in exceptional fossil preservation: An
 871 organic geochemical perspective. *Geology*, **41**, 2, 123–126.
 872 Meyers, P. (1997) Organic geochemical proxies of paleoceanographic, paleolimnologic, and
 873 paleoclimatic processes. *Organic Geochemistry*, **27**, 5/6, 213–250.
 874 Moldowan, J.M., Dahl, J., Jacobson, S.R., Huizinga, B.J., Fago, F.J., Shetty, R., Watt, D.S.,
 875 Peters, K.E. (1996) Chemostratigraphic reconstruction of biofacies: molecular evidence
 876 linking cyst-forming dinoflagellates with pre-Triassic ancestors. *Geology*, **24**, 2, 159–162.
 877 Moldowan, J.M., Fago, F.J., Carlson, R.M.K., Young, D.C., an Duvne, G., Clardy, J., Schoell,
 878 M., Pillinger, C.T., Watt, D.S. (1991) Rearranged hopanes in sediments and petroleum.
 879 *Geochimica et Cosmochimica Acta*, **55**, 11, 3333–3353.
 880 Moldowan, J.M., Talyzina, N.M. (1998) Biogeochemical evidence for dinoflagellate
 881 ancestors in the Early Cambrian. *Science*, **281**, 1168–1170.
 882 Mongelli, G. (1995) Trace elements distribution and mineralogical composition in the < 2-μm
 883 size fraction of shales from the Southern Apennines, Italy. *Mineralogy and Petrology*, **53**, 1–
 884 3, 103–114.
 885 Murphy, A.E., Sageman, B.B., Hollander, D.J., Lyons, T.W., Brett, C.E. (2000) Black shale
 886 deposition and faunal overturn in the Devonian Appalachian Basin: Clastic starvation,
 887 seasonal water-column mixing, and efficient biolimiting nutrient recycling.
 888 *Paleoceanography*, **15**, 3, 280–291.
 889 Oasis oils Company (1961) D1-26 final log.

890 Ourisson, G., Albrech, P., Rohmer, M. (1979) The hopanoids - Paleochemistry and
891 biochemistry of a group of natural products. *Pure and Applied Chemistry*, **51**, 4, 709–729.

892 Overmann, J. (2008) Ecology of phototrophic sulfur bacteria. In : Hell, R., Dahl, C., Knaff,
893 D., Leustek, T., (eds.). *Sulfur Metabolism in Phototrophic Organisms*, Springer Netherlands,
894 Dordrecht, pp. 375–396.

895 Overmann, J., Cypionka, H., Pfennig, N. (1992) An extremely low-light adapted phototrophic
896 sulfur bacterium from the Black Sea. *Limnology and Oceanography*, **37**, 1, 150–155.

897 Paris, F., Nölvak, J. (1999) Biological interpretation and paleobiodiversity of a cryptic fossil
898 group: The “chitinozoan animal.” *Geobios*, **32**, 2, 315–324.

899 Peters, K.E., Walters, C.W., Moldowan, J.M. (2005) *The biomarker guide, second edition*.
900 Cambridge University Press, Cambridge, 1155 p.

901 Peterson, L.C., Haug, G.H., Murray, R.W., Yarincik, K.M., King, J.W., Bralower, T.J.,
902 Kameo, K., Rutherford, S.D., Pearce, R.B. (2000) Late Quaternary stratigraphy and
903 sedimentation at ODP Site 1002, Cariaco Basin (Venezuela). In : Leckie, R.M., Sigurdsson,
904 H., Acton, G.D., Draper, G., (eds.). *Proceedings of the ODP, Scientific Results*, pp. 85–99.

905 Prauss, M.L. (2007) Availability of reduced nitrogen chemospecies in photic-zone waters as
906 the ultimate cause for fossil prasinophyte prosperity. *Palaios*, **22**, 5, 489–499.

907 Racka, M., Marynowski, L., Filipiak, P., Sobstel, M., Piszczowska, A., Bond, D.P.G. (2010)
908 Anoxic Annulata Events in the Late Famennian of the Holy Cross Mountains (Southern
909 Poland): Geochemical and palaeontological record. *Palaeogeography, Palaeoclimatology,*
910 *Palaeoecology*, **297**, 3, 549–575.

911 Racki, G. (2005) Toward understanding Late Devonian global events: few answers, many
912 questions. In : Over, D.J., Morrow, J.R., Wignall, P.B., (eds.). *Understanding Late Devonian*
913 *and Permian-Triassic Biotic and Climatic Events: Towards an Integrated Approach*, Elsevier,
914 pp. 5–36.

915 Radke, M., Welte, D.H. (1983) The Methylphenanthrene Index (MPI): a maturity parameter
916 based on aromatic hydrocarbons. In : Bjorøy, P., Albrecht, P., Cornford, C., (eds.). *Advances*
917 *in Organic Geochemistry 1981*, Wiley, Chichester, pp. 504–512.

918 Radke, M., Willsch, H., Leythaeuser, D., Teichmüller, M. (1982) Aromatic components of
919 coal: relation of distribution pattern to rank. *Geochimica et Cosmochimica Acta*, **46**, 10,
920 1831–1848.

921 Requejo, A.G., Allan, J., Creaney, S., Gray, N.R., Cole, K.S. (1992) Aryl isoprenoids and
922 diaromatic carotenoids in Paleozoic source rocks and oils from the Western Canada and
923 Williston Basins. *Organic Geochemistry*, **19**, 1–3, 245–264.

924 Riboulleau, A., Bout-Roumazelles, V., Tribovillard, N., Guillot, F., Recourt, P. (2014)
 925 Testing provenance diagrams: Lessons from the well-constrained Cariaco Basin. *Chemical*
 926 *Geology*, **389**, 91–103.

927 Riquier, L., Averbuch, O., Devleeschouwer, X., Tribovillard, N. (2010) Diagenetic versus
 928 detrital origin of the magnetic susceptibility variations in some carbonate Frasnian–
 929 Famennian boundary sections from Northern Africa and Western Europe: implications for
 930 paleoenvironmental reconstructions. *International Journal of Earth Sciences*, **99**, 1, 57–73.

931 Riquier, L., Tribovillard, N., Averbuch, O., Devleeschouwer, X., Riboulleau, A. (2006) The
 932 Late Frasnian Kellwasser horizons of the Harz Mountains (Germany): Two oxygen-deficient
 933 periods resulting from different mechanisms. *Chemical Geology*, **233**, 1–2, 137–155.

934 Riquier, L., Tribovillard, N., Averbuch, O., Joachimski, M.M., Racki, G., Devleeschouwer,
 935 X., El alban, A., Riboulleau, A. (2005) Productivity and bottom water redox conditions at the
 936 Frasnian-Famennian boundary on both sides of the Eovariscan Belt: constraints from trace-
 937 element geochemistry. *Understanding Late Devonian And Permian-Triassic Biotic and*
 938 *Climatic Events Towards an Integrated Approach*, Elsevier, pp. 199–224.

939 Romero-Sarmiento, M.-F., Riboulleau, A., Vecoli, M., Versteegh, G.J.-M. (2011) Aliphatic
 940 and aromatic biomarkers from Gondwanan sediments of Late Ordovician to Early Devonian
 941 age: An early terrestrialization approach. *Organic Geochemistry*, **42**, 6, 605–617.

942 de la Rue, S.R., Rowe, H.D., Rimmer, S.M. (2007) Palynological and bulk geochemical
 943 constraints on the paleoceanographic conditions across the Frasnian–Famennian boundary,
 944 New Albany Shale, Indiana. *International Journal of Coal Geology*, **71**, 1, 72–84.

945 Sandberg, C.A., Morrow, J.R., Ziegler, W. (2002) Late Devonian sea-level changes,
 946 catastrophic events, and mass extinctions. In : Koeberl, C., MacLeod, K.G., (eds.).
 947 *Catastrophic Events and Mass Extinctions: Impacts and Beyond*, Geological Society of
 948 America, p.

949 Schieber, J., Lazar, R.O. (2004) *Devonian black shales of the eastern U.S.: New insights into*
 950 *sedimentology and stratigraphy from the subsurface and outcrops in the Illinois and*
 951 *Appalachian Basins*. Indiana Geological Survey, Bloomington, Ind., 90 p.

952 Schwark, L., Empt, P. (2006) Sterane biomarkers as indicators of palaeozoic algal evolution
 953 and extinction events. *Evolution of the System Earth in the Late Palaeozoic: Clues from*
 954 *Sedimentary Geochemistry*, **240**, 1, 225–236.

955 Schwark, L., Frimmel, A. (2004) Chemostratigraphy of the Posidonia Black Shale, SW-
 956 Germany: II. Assessment of extent and persistence of photic-zone anoxia using aryl
 957 isoprenoid distributions. *Chemical Geology*, **206**, 3–4, 231–248.

958 Sephton, M.A., Looy, C.V., Veeffkind, R.J., Visscher, H., Brinkhuis, H., de Leeuw, J.W.
 959 (1999) Cyclic diaryl ethers in a Late Permian sediment. *Organic Geochemistry*, **30**, 4, 267–
 960 273.

961 Soua, M. (2014) Paleozoic oil/gas shale reservoirs in southern Tunisia: An overview. *Journal*
 962 *of African Earth Sciences*, **100**, 450–492.

963 Spina, A., Vecoli, M., Riboulleau, A., Clayton, G., Cirilli, S., Di Michele, A., Marcoguiseppe,
 964 A., Rettori, R., Sassi, P., Servais, T., et al. (2017) *Assessment of thermal maturity of organic*
 965 *matter of Lower Palaeozoic rocks (Ghadamis Basin, North Africa): application of*
 966 *Palynomorph Darkness Index (PDI) and other thermal maturity indicators*. Perugia-Italy,.

967 Steemans, P. (1999) Paléodiversification des spores et des cryptospores de l'Ordovicien au
 968 Dévonien inférieur. *Geobios*, **32**, 2, 341–352.

969 Streel, M. (2009) Upper Devonian miospore and conodont zone correlation in western
 970 Europe. *Geological Society, London, Special Publications*, **314**, 1, 163.

971 Strother, P.K. (1991) A classification schema for the cryptospores. *Palynology*, **15**, 1, 219–
 972 236.

973 Summons, R.E., Powell, T.G. (1986) Chlorobiaceae in Palaeozoic seas revealed by biological
 974 markers, isotopes and geology. *Nature*, **319**, 6056, 763–765.

975 Tappan, H.N. (1980) *The paleobiology of plant protists*. W. H. Freeman,.

976 Taylor, T.N., Taylor, E.L., Krings, M. (2009) *Paleobotany: The Biology and Evolution of*
 977 *Fossil Plants*. Academic Press, 1230 p.

978 Tribovillard, N. (2013) The Ge/Si ratio as a tool to recognize biogenic silica in chert. *Comptes*
 979 *Rendus Geoscience*, **345**, 3, 160–165.

980 Tribovillard, N., Averbuch, O., Devleeschouwer, X., Racki, G., Riboulleau, A. (2004) Deep-
 981 water anoxia over the Frasnian–Famennian boundary (La Serre, France): a tectonically
 982 induced oceanic anoxic event? *Terra Nova*, **16**, 5, 288–295.

983 Tribovillard, N., Bout-Roumazeilles, V., Algeo, T., Lyons, T.W., Sionneau, T., Montero-
 984 Serrano, J.C., Riboulleau, A., Baudin, F. (2008) Paleodepositional conditions in the Orca
 985 Basin as inferred from organic matter and trace metal contents. *Marine Geology*, **254**, 1–2,
 986 62–72.

987 Tulipani, S., Grice, K., Greenwood, P.F., Haines, P.W., Sauer, P.E., Schimmelmann, A.,
 988 Summons, R.E., Foster, C.B., Böttcher, M.E., Playton, T., et al. (2015) Changes of
 989 palaeoenvironmental conditions recorded in Late Devonian reef systems from the Canning
 990 Basin, Western Australia: A biomarker and stable isotope approach. *Gondwana Research*, **28**,
 991 4, 1500–1515.

- Tyson, R.V. (1995) *Sedimentary organic matter*. Chapman & Hall, London., 615 p.
- Ulmishek, G.F., Klemme, H.D. (1990) *Depositional controls, distribution, and effectiveness of world petroleum source rocks*.
- Underdown, R., Redfern, J., Lisker, F. (2007) Constraining the burial history of the Ghadames Basin, North Africa: an integrated analysis using sonic velocities, vitrinite reflectance data and apatite fission track ages. *Basin Research*, **19**, 4, 557–578.
- Vandenbroucke, T.R.A., Armstrong, H.A., Williams, M., Paris, F., Sabbe, K., Zalasiewicz, J.A., Nölvak, J., Verniers, J. (2010) Epipelagic chitinozoan biotopes map a steep latitudinal temperature gradient for earliest Late Ordovician seas: Implications for a cooling Late Ordovician climate. *Early Palaeozoic Palaeoenvironments*, **294**, 3, 202–219.
- Versteegh, G.J.M., Riboulleau, A. (2010) An organic geochemical perspective on terrestrialization. *Geological Society, London, Special Publications*, **339**, 1, 11–36.
- Wang, T.-G., Simoneit, B.R.T. (1995) Tricyclic terpanes in Precambrian bituminous sandstone from the eastern Yanshan region, North China. *Chemical Geology*, **120**, 1, 155–170.
- Wendt, J., Belka, Z. (1991) Age and depositional environment of upper devonian (early Frasnian to early famennian) black shales and limestones (Kellwasser facies) in the eastern Anti-Atlas, Morocco. *Facies*, **25**, 1, 51–89.
- Wendt, J., Kaufmann, B., Belka, Z., Klug, C., Lubeseder, S. (2006) Sedimentary evolution of a Palaeozoic basin and ridge system: the Middle and Upper Devonian of the Ahnet and Mouydir (Algerian Sahara). *Geological Magazine*, **143**, 3, 269.
- Weyant, M., Massa, D. (1991) Contribution of conodonts to the Devonian biostratigraphy of Western Libya. In : Salem, M.J., Hammuda, O.S., Eliagoubi, B.A., (eds.). *The Geology of Libya*, Elsevier, Amsterdam, pp. 1297–1322.
- Whitaker, M.F. (1984) The usage of palynology in definition of Troll Field geology. In: Reduction of uncertainties in innovative reservoir geomodelling. *6th Offshore Northern Seas Conference and Exhibition*, Norsk petroleum forening, Stavanger, pp. 1–50.
- Wilde, P., Quinby-Hunt, M.S., Erdtmann, B.-D. (1996) The whole-rock cerium anomaly: a potential indicator of eustatic sea-level changes in shales of the anoxic facies. *Sedimentary Geology*, **101**, 1–2, 43–53.

Figure captions

Figure 1: Location map of the Ghadames Basin showing the location of borehole D1-26.

Figure 2: Lithological and stratigraphic description of the Upper Devonian formations in borehole D1-26, with sonic and resistivity (ILD) curves (unavailable scale). Compilation of data from Oasis oils Company (1961), Canaple (1963) Loboziak and Streel (1989), Weyant and Massa (1991) Spina et al. (2017). The peaks in the ILD curve (grey area) underline an alternance of shale and limestones corresponding to the radioactive shales and Cues Limestone horizon.

Figure 3: Palynofacies observations of the samples from borehole D1-26. A) Particle counting; B,C) Prasinophycean cysts, AOM and acritarchs (B: slide 7802; C: slide 7780); D) Prasinophycean cysts and AOM (slide 7704); E) miospores, Prasinophycean cysts and palynomacerals (slide 7083). Photographs show the >50µm palynological residue. Legend of the log as in fig. 2.

Figure 4: Rock-Eval and bulk geochemistry of samples from borehole D1-26. A) Hydrogen index (HI) *versus* Tmax plot showing the type and maturity of the organic matter. B) Total sulfur *versus* total organic carbon (TOC) of the sediment. The dotted line corresponds to the relationship observed in “normal marine” shales (Berner, 1982; Berner and Raiswell, 1983). C) Same as B after doubling the TOC content in order to account for thermal loss of organic matter.

Figure 5: Mineral redox proxies in the samples from borehole D1-26. A) Enrichment factor (EF) of the main redox-sensitive trace elements. B) Cerium anomaly. The grey interval corresponds to the Cues Limestone horizon and radioactive shales.

Figure 6: Detrital proxies in the samples from borehole D1-26. A) major and trace element content normalized to the aluminum content. B) Rare earth element ratios and europium anomaly. The grey interval corresponds to the Cues Limestone horizon and radioactive shales.

Figure 7: Rare earth element content in the samples from borehole D1-26 normalized to the Post Archean Average Shale (PAAS, McLennan, 1989).

Figure 8: Representative fragmentograms showing the biomarker distribution in the aliphatic fraction of the extracts from borehole D1-26 samples. A) *n*-alkanes and linear isoprenoids (m/z 57; sample 7087). B) Di- and tri-cyclic terpenoids (m/z 123; sample 7087). C) Cheilantanes and hopanoids (m/z 191; sample 7085). D) Steroids and diasteroids (m/z 217; sample 7780). Numbers above symbols indicate carbon number.

Figure 9: Example of distribution of isorenieratene and palaerenieratene derivatives in the aromatic fraction of the extracts from borehole D1-26 samples (m/z 133+134; sample 7810).

Figure 10: Molecular indicators of redox conditions in the extracts from D1-26 samples. A) Total concentration of isorenieratene and palaerenieratene derivatives in the extracts. B) C₁₆-C₃₁ palaerenieratene to isorenieratene derivatives ratio (PIR). The grey interval corresponds to the Cues Limestone horizon and radioactive shales. C) Pristane to phytane ratio (Pr/Ph) versus the modified aryl isoprenoid ratio (AIR*). Scheme from Schwark and Frimmel (2004)

Figure 11: Depositional model for organic matter in the Ghadames Basin during the Frasnian to upper Famennian interval. See text for explanations.

Supplementary material

Table S1: major, trace and rare earth element content in samples from borehole D1-26.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Tables

Table 1: Rock Eval and elemental analysis results of the samples from borehole D1-26.

Sample	S1 ¹	S2 ¹	Tmax °C	TOC %	PI ²	HI ³	OI ⁴	C _{min} (%)	S (%)	N (%)	TOC/N
7081	0.19	1.08	432	1.8	0.15	60	0	0.1	7.23	0.06	35
7083	0.24	1.78	434	2.68	0.12	66	0	0.14	3.73	0.08	39
7085	0.36	3.53	436	2.78	0.09	127	8	0.12	1.79	0.08	41
7087	0.15	1.01	439	0.86	0.13	117	48	0.96	1.27	0.03	33
7100	0.2	1.76	438	1.87	0.1	94	22	0.86	1.63	0.05	44
7712	0.85	2.8	434	2.21	0.23	127	5	0.16	3.23	0.12	21
7780	1.55	8.7	442	2.73	0.15	319	2	0.14	0.73	0.13	25
7800	0.7	0.77	431	1.06	0.48	73	27	0.13	2.55	0.07	18
7809	0.65	1.19	435	1.06	0.35	112	0	0.05	2.32	0.09	14
7810	0.78	2.2	441	1.29	0.26	171	12	0.09	1.95	0.1	15
7812	0.86	2.05	441	1.26	0.29	163	0	0.1	2.33	0.09	16
7813	0.73	2.2	440	1.38	0.25	159	4	0.12	2.17	0.08	20

¹ mgHC/g

² Production index: $PI = S1 / (S1 + S2)$

³ Hydrogen index in mgHC/gTOC

⁴ Oxygen index in mgCO₂/gTOC

Table 2: Biomarker ratios in the extracts of samples from borehole D1-26.

Sample	7085	7087	7712	7780	7810	7812
Source/environment						
Pr/Ph	2.07	2.23	1.29	1.01	1.30	1.42
Pr/C ₁₇	0.82	1.03	0.65	0.76	0.77	0.74
Ph/C ₁₈	0.43	0.53	0.51	0.69	0.66	0.57
CPI ¹	1.19	1.22	0.97	1.05	1.07	0.93
TAR ²	0.96	0.41	0.07	0.39	0.2	0.52
tricyclic/17 α hopanes ³	0.11	0	0.43	0.3	0.5	0.46
Hop/ster ⁴	3.6	3.24	1.41	1.32	1.27	1.22
%C ₂₇ ⁵	26.6	27.8	34.6	30.9	22.5	23.6
%C ₂₈ ⁵	23.5	17.3	14.2	17.5	22.9	30.2
%C ₂₉ ⁵	49.9	54.9	51.2	51.6	54.6	46.2
AIR* ⁶	2.22	1.02	0.84	0.4	0.86	1.26
Maturity						
Diasterane S/(S+R) ⁷	0.58	0.59	0.55	0.59	0.6	0.62
Sterane C ₂₉ S/(S+R) ⁸	0.43	0.45	0.48	0.5	0.48	0.33
Sterane C ₂₉ $\beta\beta$ /($\beta\beta$ + $\alpha\alpha$) ⁹	0.34	0.41	0.61	0.61	0.66	0.67
Hopane C ₃₂ 22S/(22S+22R)	0.6	0.59	0.62	0.56	0.6	0.56
Hopane C ₃₀ $\beta\alpha$ /($\beta\alpha$ + $\alpha\beta$)	0.17	0.15	0	0.13	0.15	0.16
Ts/(Ts+Tm)	0.35	0.36	-	0.72	0.81	0.82
MPI ¹⁰	0.62	0.73	0.61	0.66	0.73	0.81
MPR ¹¹	1.36	1.48	0.81	0.88	0.96	0.94
Vitrinite reflectivity Rc ¹²	0.77	0.84	0.77	0.79	0.84	0.89

¹ *n*-Alkanes Carbon Preference Index:

$$CPI = \{[(C_{25}+C_{27}+C_{29}+C_{31}+C_{33})/(C_{26}+C_{28}+C_{30}+C_{32}+C_{34})] + [(C_{25}+C_{27}+C_{29}+C_{31}+C_{33})/(C_{26}+C_{28}+C_{30}+C_{32}+C_{34})]\} / 2$$

(Bray and Evans, 1961).

² *n*-Alkanes Terrestrial to aquatic ratio: $TAR = (C_{27}+C_{29}+C_{31})/(C_{15}+C_{17}+C_{19})$ (Bourbonniere and Meyers, 1996).

³ $\Sigma(C_{21}-C_{29})$ cheilantanes/ $\Sigma(C_{29}-C_{35})$ $\alpha\beta$ hopanes

⁴ Calculated using the sum of dia- and regular steroids and the sum of all hopanoids (diahopanes, TS series, hopanes and moretanes).

⁵ Relative percentage of C₂₇, C₂₈ and C₂₉ regular steranes, determined on the $\alpha\alpha\alpha$ R isomers.

⁶ Modified arylisoprenoid ratio: $AIR^* = (C_{13}-C_{17})/(C_{18}-C_{22})$ arylisoprenoids using both the 2,3,6- and 3,4,5-trimethyl-aryl isoprenoids on m/z 133+134 fragmentogram.

⁷ Calculated using C₂₇ $\beta\alpha$ diasteranes.

⁸ Calculated using C₂₉ $\alpha\alpha$ steranes.

⁹ Calculated using C₂₉ R and S stereoisomers.

¹⁰ Methylphenanthrene index $MPI = 1.5[3-MP+2-MP]/[P+9-MP+1-MP]$ (Radke and Welte, 1983).

¹¹ Methylphenanthrene ratio $MPR = [2-MP]/[1-MP]$ (Radke et al., 1982).

¹² $Rc = 0.6 * MPI + 0.4$ (Radke and Welte, 1983)

ACCEPTED MANUSCRIPT