



HAL
open science

Evidence for the implication of the histone code in building the genome structure

Kirti Prakash, David Fournier

► **To cite this version:**

Kirti Prakash, David Fournier. Evidence for the implication of the histone code in building the genome structure. *BioSystems*, 2018, 164, pp.49 - 59. 10.1016/j.biosystems.2017.11.005 . hal-01787125

HAL Id: hal-01787125

<https://hal.sorbonne-universite.fr/hal-01787125>

Submitted on 7 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evidence for the implication of the histone code in building the genome structure

Kirti Prakash^{a,b,c,*} and David Fournier^{d,*}

^aPhysico-Chimie Curie, Institut Curie, CNRS UMR 168, 75005 Paris, France; ^bOxford Nanoimaging Ltd, OX1 1JD, Oxford, UK; ^cMicron Advanced Bioimaging Unit, Department of Biochemistry, University of Oxford, Oxford, UK; ^dFaculty of Biology and Center for Computational Sciences, Johannes Gutenberg University Mainz, 55128 Mainz, Germany; *Correspondence: kirtiprakash1.61@gmail.com, dfournie@uni-mainz.de

Histones are punctuated with small chemical modifications that alter their interaction with DNA. One attractive hypothesis stipulates that certain combinations of these histone modifications may function, alone or together, as a part of a predictive histone code to provide ground rules for chromatin folding. We consider four features that relate histone modifications to chromatin folding: charge neutralisation, molecular specificity, robustness and evolvability. Next, we present evidence for the association among different histone modifications at various levels of chromatin organisation and show how these relationships relate to function such as transcription, replication and cell division. Finally, we propose a model where the histone code can set critical checkpoints for chromatin to fold reversibly between different orders of the organisation in response to a biological stimulus.

DNA | nucleosomes | histone modifications | chromatin domains | chromosomes | histone code | chromatin folding | genome structure

Introduction

The genetic information within chromosomes of eukaryotes is packaged into chromatin, a long and folded polymer of double-stranded DNA, histones and other structural and non-structural proteins. The repeating units of the polymer, the nucleosomes, are 147 base-pairs (1.75 turn) of DNA wrapped around an octamer of 4 histone proteins [1, 2]. Nucleosomes are thought to be further compacted into a higher order 30 nm chromatin fibre by linker histone H1 [3]. The structure of nucleosomes can be altered post-translationally by the small chemical modifications of histone protein [4, 5]. Subsequently, one can characterise the organisation of chromatin into three interrelated categories: (1) the basic building blocks, (2) the functional structure of chromatin and (3) the higher order spatial arrangement of chromatin.

The two classical building blocks (Figure 1A, first column): beads-on-a-string and 30 nm chromatin fibre have been extensively studied previously [1, 6–8]. Regarding the intermediary level of compaction, chromatin can display several configurations (active, repressed, inactive) depending upon enrichment of a particular histone mark (Figure 1A, second column). At a higher order (Figure 1A, third column), chromatin can be either described as a bimodal heterochromatin/euchromatin model (condensed and open regions, respectively), chromosome territories [9] or as a very condensed structure in the




Fig. 1. Chromatin spatial organization and relationship with the histone code. (A) The spatial organization of chromatin can be studied at three levels: at the lowest orders, which include the beads-on-a-string model and the chromatin fibre (left column), at the level of functional chromatin domains (middle column) and high order chromatin patterns (right column). The organization of chromatin domains can be modelled using various post-translational histone modifications. Depending upon the kind of histone modification a nucleosome is enriched with, chromatin can be either highly condensed, in an open conformation or switch between these two extreme forms. The higher order chromatin patterns can either be viewed from: the chromosome territory model of DNA organization, the over-simplified bimodal classification of condensed and open chromatin as hetero- and euchromatin respectively, and the highly condensed configuration of metaphase chromosomes. (B) The components of the histone code. On the left, the input system A, made of combinations of histone modifications (coloured pins). One unique combination can be found at three different positions 1,2,3, in the genome. The combinations of the input system A is translated into components of the output system B, with a black and white colour-scale. The translation happens via adaptors such as transcription factors or epigenetic-regulators that recognise specific histone modifications to change chromatin state.

case of the metaphase chromosome. Chromatin fibres indeed present a variety of sizes [10] and of shapes ([11] shows a

solenoid model of chromatin), while recent studies attempt to challenge their existence [12], hinting that the hierarchy beads-on-a-string/fibres/domains might be much more complicated and diverse than we currently think.

On the functional side, it has been shown that biochemical changes made to specific histones tails are associated with different condensation levels of chromatin. For instance, trimethylaton of lysine 9 on histone 3 (H3K9me3) is usually associated with condensed chromatin and centromeric regions [15, 16]. On the other hand, trimethylaton of lysine 4 on histone 3 (H3K4me3) is strongly enriched at promoter regions of active genes where chromatin exists in an open conformation [17, 18]. Chromatin can also exist in a switchable state between these two extreme forms if histone H3 is trimethylated at lysine 27 (H3K27me3) [19]. H3K27me3 is a histone mark characteristic of repressed genes.

The number of ways histone proteins can be modified is quite extensive and can explain the vast majority of possible combinations that can lead to various functional outcomes. The relationship between these combinations and the function they perform is referred to as the "histone code" [20–22]. Previously, biological codes have been thoroughly described [23–25]; they comprise of input system made of signs with no biological function that are translated into organic output functions via adaptor molecules. In the example of the genetic code, the inputs are codons, the adaptor is the translation machinery

and the outputs are amino acids. In the case of the histone code, combinations of histone modifications at a given position on the genome constitute the input system, the adaptors are the epigenetic regulators (for instance, the enzymatic complex Ezh2) that bind the modifications and outputs are chromatin features such as the level of chromatin compaction or gene expression (Figure 1B).

Here, we present supportive evidence for a combinatorial occurrence of histone modifications and its consequence for higher-order chromatin folding. We start with a brief historical introduction to chromatin biology. We then show the correlation among histone modifications at the level of nucleosomes, regulatory regions, TAD and chromosomes. Finally, we propose a model showing that these distinct chromatin domains can co-exist on a single chromosome.

Histone modifications, a robust but flexible matrix for evolution

Histones were first identified as a fundamental component of the nucleus in the early days of molecular biology [26, 27] and were described as circular structures responsible for compacting DNA [1, 6, 28]. They were soon associated with DNA periodicity and compaction [6, 29–32], where DNA coils around an octamer of four core histone proteins (H2A/H2B dimer and H3/H4 tetramer, Figure 2A) [1, 33]. The histone core with DNA was termed nucleosome and is known to be the first level of chromatin compaction. Early experiments showed that histones, via their N-terminal tails, can experience modifications such as acetylation and methylation, that can further alter the compaction of chromatin [4]. The nomenclature of histone modification were named the following way: in the case of H3K9me3, H3 refers to the core histone protein, K refers to the amino acid, the number 9 indicates the position of lysine residue from N-terminal end of the amino acid tail of histone protein and me3 refers to the type of modification on the lysine residue (Figure 2B).

Over the course of the last decade, amino acids have been shown to experience several modifications, of at least twelve types: acetylation (lysine), methylation (lysine and arginine), phosphorylation (serine and threonine), sumoylation (lysine), ubiquitylation (lysine), ADP ribosylation, butyrylation, citrullination, crotonylation, formylation, proline isomerization, propionylation [34–39].

At a fundamental level, most of these modifications are neutralizing the net charge between DNA and histone proteins, and much of the gene regulation is a by product of effective net charge in the environment [44, 45]. Hence a combination of marks must work in synergy to provide sufficient proof reading so that no gene is accidentally turned on or off [46–

Outlook

The urge to discover secrets is deeply ingrained in human nature; even the least curious mind is roused by the promise of sharing knowledge withheld from others. Most of us are driven to sublimate this urge by the solving of artificial puzzles devised for our entertainment. Detective stories or crossword puzzles cater for the majority; the solution of secret codes may be the hobby of the few.

J. Chadwick, The Decipherment of Linear B [13].

Citation inspired from The Code Book by Simon Singh [14].

From the day/night dichotomy to the genetic code, nature is full of symmetric, antagonistic exemplars and patterns. One such example is the organisation of structurally distinct chromatin states (active, inactive) on a single chromosome. In this article, we try to show how simple combinations of essential elements such as histone modifications can participate in sophisticated cellular features such as the structure of the genome. Here a code is identified, where an input system (histone modifications) is translated into an output system (chromatin states) via adaptors (epigenetic regulators or transcription factors). Such a code has a distinct importance in gene regulation and consequently for the cellular phenotype.




Fig. 2. Histone modifications are major biochemical features of chromatin. Histone can experience various post-translational modifications in their protruding N-terminal tails but also within the C-terminal regions. These changes affect not only the overall compaction of chromatin but also gene expression. (A) The principal modifications on the four core histones: H2A, H2B, H3, H4. Two patterns worth noting here are the arrangement of lysines (K4, K9, K14, K18, K23, K27) and occurrence of Lysine (K), Serine (S) and Arginine (A) at (9, 10, 11) and (27, 28, 29) position on the amino acid tail of histone H3. Modified from [40] with permission. (B) Nomenclature of histones post-transcriptional modifications. The different features are the type of histone (H3 in our example), the amino acid and its position (Lysine at position 27) and the chemical modification, here a trimethylation, or triple methylation of the lysine 27 from the N-terminal end of the amino acid tail. (C) Different enzymes lead to writing, erasing and reading of chemical groups on the tails of histone proteins. Writers are enzymes that introduce various chemical modifications to histones. There are about 100 different enzymes that can write acetyl, methyl and phosphoryl groups (circles) to individual amino acid residues. Erasers are the enzymes that remove histone modifications and perform the opposite task of writers. Readers are enzymes that recognise specific chemical groups on the amino acid residues. For instance, bromodomains engage with acetylated lysines (modified from [41] with permission). Ac, acetylation; Me, methylation; P, phosphorylation; Ub, ubiquitination.

[48]. Lastly, the combination of histone modifications are stably maintained across many species and protein motifs have evolved to recognise histone modifications [49].

We consider four measures which characterize the function of histone modifications

1. charge neutralisation
2. molecular specificity
3. robustness
4. evolvability

Charge neutralisation: Histone modifications are usually modified via enzymes that can neutralise the excess of charge on the DNA [50, 51]. A nucleosome core particle has an overall residual charge of $-150e$ (DNA contributes $-294e$,

and histones contribute $+144e$) and is therefore electrically not neutral [52]. Tails can attract negatively remote DNA molecules (the negatively charged due to phosphate oxide groups) to induce regional DNA compaction. Subsequently, the folding of DNA is also highly dependent on the positive counterions in the environment [53, 54].

Early experiments showed that acetylation (CH₃-CH₂-) or methylation (CH₃-) of nucleosomes occur respectively in 50-60% and 40% of histones and can lead to increase and decrease of polymerase activity, respectively [44]. Methyl groups are produced during the metabolism of methionine, a subpart of the B12 vitamin circuit while acetylation is generated from the acetyl-Coenzyme A [55, 56]. Histone acetylation impairs the affinity of histones to DNA via charge antagonism, reduction of compaction and subsequent recruitment of certain factors such as SIR at telomeres or TAF1 at promoters, inducing activation of transcription [57–59]. This property is accompanied by the higher mobility of histones along DNA [60]. Differently, methylation is associated to compaction of chromatin and so to reduced transcription, with reduced histone mobility [44, 60, 61].

Molecular specificity: The histone modification sequence is orchestrated by the enzymes that either deposit, read or remove the marks, and serve as the adaptors molecules essential to biological codes [23]. Enzymes that deposit marks are diverse and reflect the large spectrum of modifications available in nature (Figure 2C). Well-known examples are of histone acetyl-transferases (HAT) which transfer the acetyl group from acetyl-Coenzyme A to the tail of histones, and histone deacetylases (HDAC) which remove the acetyl groups, most notably upon chromatin compaction. Other complexes involved in histone regulation include the Trithorax complex (Trx) which is involved in methylation of histones and Polycomb (PcG) with module PRC2 involved in the deposition of H3K27me3 mark via methyltransferase Ezh2 [62, 63].

Additionally, the H3K27me3 mark is bound by the PRC1 complex that can mono-ubiquitinate histone H2A on lysine 119 (H2AK119Ub1). This process is known to silence genes and is an excellent example of a combination of marks found at a genomic site with an impact on the function. In contrast, transcription factors with bromodomains are known to bind acetylated lysines [64] and induce gene activation via acetyl-transferase activity [65].

Robustness: Robustness is the capacity of living systems to preserve a given function during evolutionary times. Many important core features of organisms are robust and found in a wide variety of species; for instance, the skeleton is found in all vertebrates and can be considered a robust feature. Small or big structural variations are present in all groups, but the




Fig. 3. A binary combinatorial histone code at the level of nucleosomes. (A). Different clusters of histone modifications are revealed by hierarchical clustering. Correlation between modifications are displayed with a color scheme, red to blue from negative to positive correlation coefficient. Clusters are identified by black boxes (figure from [42], courtesy Kernel Press, Mainz). (B) Different cis and trans patterns of histone crosstalk (figure from [43], reprinted with permission). (C) The impact of three primary histone modifications on chromatin compaction. H3K4me3-rich nucleosomes are associated with active chromatin in an open conformation, H3K27me3-rich nucleosomes are associated with repressed chromatin, while H3K9me3-rich nucleosomes are associated with highly-compacted, inactive chromatin.

template remains the same. Robustness seems to imply two features: cooperation and redundancy; cooperation of the different basic molecular elements of the system, including quality control checkpoints [66] and a high redundancy of the molecular elements, in order to resist to small changes. In the example of the skeleton, many different genes are involved, each of them participating to build the scaffold (cooperation). Nevertheless, many genes perform the same function so if one gene fails or fluctuates, several other ones can take over to perform the same function (redundancy).

Overall, robustness is a requirement for all fundamental systems of life, including all the core biological codes such as the genetic code, which can produce proteins despite occasional changes [67]. Similarly, the histone code seems to feature properties favouring robustness. As we will see later, very similar histone modifications such as acetylations or methylations act in concert to form the histone code (cooperation), which potentially means that the loss of one mark does not necessarily affect the output of the code very strongly (redundancy). For instance, for a given genomic region, active promoters, different combinations are possible, with small variations (Figure 4). Such a redundancy stems in the high similarity of nucleosome structure across kingdoms archaea and eukaryotes [68] and the fact that most well-known histone modifications are found across eukaryotes, including human, mouse, fly and *C.elegans* [69]. For instance, H3K9me3 is found to be present

in vertebrates, invertebrates and yeast [70, 71]. Functions of enzymes associated with histone modifications remodelling seem to be very conserved across eukaryotes [72, 73]. Subsequently, the histone code provides a robust system that is not prone to mistakes and delivers a reliable answer to stimuli, and therefore seems to fill all requirements of a robust system.

Evolvability: A consequence of robustness is that small variations are usually harmless, a property that opens doors to small changes through evolution, with various outcomes. Evolvability of the histone code indeed does not operate at the level of histone sequence itself, as histone sequences are found to be very conserved. Nevertheless, minute changes in histone associations are found between relatively close species such as chimpanzee and human, or primates and rodents [69, 74]. Other marks such as H3K4me3 seem to have different patterns in complex organisms compared to rather simple distribution in yeast.

As a result of these different features, the histone code can be viewed as a stable system (notion of robustness) with enough flexibility that provides a basic ground to generate new functions by accumulation of slight changes of its elements (notion of evolvability).

Histone code dictates combinatorial associations between histone modifications at the nucleosome level

Many histone modifications have been proven to function in a concerted manner with other marks. Theory by Jenuwein and Allis hypothesizes that these combinations serve as elementary signs which are translated into instructions by chromatin protein complexes to regulate the genome [21]. Genomic approaches such as ChIP-Seq have identified the precise localization of proteins on DNA, and even more strikingly, have been able to distinguish between different modifications due to highly specific antibodies against mono- di- or trimethylation, among others (see [75], a pioneering example). The findings in these studies show that histone modifications of different classes are found simultaneously at the same genomic positions, *id est* in the same nucleosome. This is the result of an amplification process, for example, H4K20me1 can be recognized by enzyme Suv4-20h1 which triggers a second methylation event on the same lysine, generating a H4K20me2 mark [76, 77]. Another example is the PcG complex involved in gene silencing, which is made of the association of complexes PRC1 and PRC2. PRC2 is specialized in depositing the H3K27me3 mark [62, 63], while PRC1 recognizes H3K27me3 [78] and subsequently ubiquitinates lysine 119 of histone H2A via protein Ring 1B E3 ubiquitin ligase [79], amplifying the signal.

More recent studies have demonstrated that variants of the PRC1 complex can come in first to deposit H2A ubiquitination, which is thereafter recognised by PRC2 to finally deposit H3K27me3 [80, 81]. Histone modifications do not need to be exactly on the same histone protein to act synergistically or on neighbor nucleosomes, providing that the different modifications acting synergistically are close together in 3D space [82]. The combinations of histone marks affect transcription and so indirectly cellular events which are important for cell and organism identity. For instance, during interphase, phosphorylation at H3S10 associates with H3K9 and H3K14 acetylation and induces chromatin relaxation [83]. During mitosis, DNA needs to be packed and H3S10 phosphorylation heavily triggers compaction. Two combinations with the same mark participate to two different cellular outcomes.

The amount of information carried by the overall collection of modifications can be probed by checking the correlation of marks between individual histones, using data from ChIP-Seq [42]. The data reveals a global redundancy factor of about 1:7 with five different clusters using 39 different marks (Figure 3A). As a result, the number of possible functions that can be directed by the system, even if the number of marks is high, is relatively low, which is confirmed by genomic studies [84]. An insightful model has also shown that a simple two-marks histone code can already provide a wide panel of

function, hinting that few core elements can lead to a large combination of genomic functions [85, 86].

Direct experimental evidence for associations of histone modifications was brought by studies using co-marking of histone modifications [87] and genomic experiments associated to microscopy (co-occurring iChIP in [88]). The single-nucleosome experiments confirm very well-known histone code rule to happen on isolated nucleosomes, such as the bivalent state (H3K27me3/H3K4me3), which is typical of embryonic stem cells. An example of a *cis* and *trans* crosstalk within a nucleosome is shown in Figure 3B.

Advance microscopy techniques [89–92] have been powerful at seeing the distribution of histone modifications in the nucleus [93–95]. In particular, single molecule localisation microscopy can discriminate between regions painted with H3K4me3 and H3K27me3 [96, 97]. These marks are highly associated with the general compaction level of chromatin (Figure 3C). Overall, one usually distinguishes between euchromatin and heterochromatin, the first being opened, in an "active" state prompt for genes to be transcribed, while the latter is in a repressed state [98–100].

Histone code associations with genome function and structure

Historically, the association of histone modifications with transcription is known since long [44, 103]. Regarding modification associations, studies have revealed that an increase in the complexity of the histone make-up results in progressive changes in the gene expression [104], with acetylation to be globally associated to increase in transcription and deacetylation with decrease in transcription. To our knowledge, H3K36 methylation by the SET2 complex (co-transcriptionally binding to elongating Pol II) targets Rpd3S deacetylases to suppress cryptic transcription start sites within genes [105, 106].

ChIP-seq experiments have helped to draw a more refined picture of the relationships between histone modifications and gene function [101]. Hidden Markov models were employed to guess where a histone modification combination (or "chromatin state") starts and ends. After testing many combinations of parameters, the model was found to compromise 51 states, which is a surprisingly large number. Most of these states are found to be associated with specific genomic regions. 11 out of the 51 states fall in promoter regions, which is a significant enrichment when considering the low amount of promoter regions in the genome compared to non-coding regions (Figure 4A). Promoter regions are associated with acetylated marks, all of which show a similar bimodal profile around transcription start sites (TSS) (Figure 4B). Transcribed regions are associated with particular kinds of methylation, entirely different from




Fig. 4. The histone code at the level of genomic features. (A) The combinations of histone modifications and their associations to various genomic states (figure modified from [101], reprinted with permission). Here, the genome is partitioned into five broad regions: promoter states, transcribed states, active intergenic, repressive and repetitive state. A combination of histone marks is associated with each genomic state. The relative degree of association is indicated by a colour scale spanning frequency values between 0 and 1. (B) Profile of histone modifications around the transcription start site (TSS). On top panel, profile for acetylation marks; below panel: methylations. X-axes: Distance to TSS. The negative values are upstream from the gene and the positive values downstream, irrespective of the strandness of the gene. Y-axes: Normalized read counts. Profiles were generated by us with a R script [102].

the ones observed in repeated regions, namely methylations at lysine 4 of histone 3 (H3K4me1 and H3K4me2) along with methylation on lysine 79 of histone 3. Finally, the table shows that repeated regions display a unique combination of acetylation and methylation which is quite predictable. The rules illustrated in Figure 3A have since been extensively described in a broad range of tissues studied in the large consortium devoted to epigenetic studies called ENCODE [71].

Histone modification is indeed only meaningful in synergy with other genomic features, for instance, methylation of cytosines at CpG sites. DNA methylation is known to be associated with gene silencing. For this to happen, histone methyltransferase HP1 recognises methylation events on lysine 9 of histone 3. During DNA replication and after binding, HP1 recruits the DNA methyltransferase DNMT1 that in-

duces hemi-methylation of neighbour cytosines, inducing gene promoter inactivation [108]. Differently, DNA methylation is known to stimulate the deposition of repressive mark H3K9me3 and induce DNA compaction at promoters [109], via binding of methylated cytosine by MeCP2 and deposition of the mark by coupled enzyme Suv39h1 [110].

The impact of histone modifications on transcription and other cellular functions is largely based on the influence of chromatin structure. At a nanoscale, methylation or deacetylation of histone tails ensure the stability of nucleosomes [50]. At a gene scale, the makeup is thought to be locally uniform, guiding folding properties, for instance in the region of Hox genes [67]. Areas with similar epigenetic make-up may fold together while repelling other parts of different make-up, thus contributing to the formation of locally coherent topological chromatin




Fig. 5. The histone code predicts the degree of folding of chromatin domains. (A) A section of chromosome 19 shows different subcompartments based on histone modifications. Here, chromatin domains with high contact frequencies correlate with histone marks associated with inactive chromatin such as H3K9me3 and H4K20me3. Such relation between H3K9me3 and H4K20me3 was also found at the nucleosomes level based on the entirely different method and cell line. This shows that relationship between histone modifications holds at various length scales. Figure reprinted from [84], with permission. (B) Imaging of histone modifications further reveals the distribution of domains associated to different histone modifications. The different histone modifications have distinct molecular signatures, probably due to different folding rules and roles in gene regulatory mechanisms. While H3K9me3 seems to be mostly enriched at chromocenters as expected, H3K27me3 is distributed more in speckle-like patterns. The acetylation patterns (H3K14ac) vary in density and are more diverse in distribution. Figure adapted from [107]. Scale bar: 1 μm . (C) A model portraying three stages of a Hox gene activation, with progressive suppression of a silent chromatin domain made up of H3K27me3- and H3K9me3-rich histones toward the installation of an active domain rich with H3K4me3. The change of domain happens literally at the same place, with inhibitory marks being progressively evicted from the region to be replaced by the active domain. Note the region of inhibitory marks remain at the bottom, to keep the two extremities of the loop in a condensed fashion and leave the loose active region in a relatively consistent open shape.

domains in the size range of 0.5-1 Mb, or TAD described by recent chromatin conformation capturing approaches [84, 111]. Indeed, we have recently shown that histone modifications cluster at genome scale using localisation microscopy [112]. Previously, it was shown with the confocal microscope, that H3K4me3-rich, H3K27me3-rich and H3K9me3-rich regions partially segregate (for instance in [113]). On our localisation images, interphase and M-phase chromatin could also be roughly divided into H3K4me3-rich, H3K27me3-rich and H3K9me3-rich regions of highly different patterns. These marks are known to be respectively associated with active genes, inactive genes and deeply repressed chromatin [114].

Along with microscopy, genomics has been the most powerful at characterising the relationship between modifications of histones and structure. The HiC method has helped to identify regions of structurally complex DNA folding, whether loose or compacted, that are thought to be functional units [111]. ChIP-seq of histone modifications has shown that chromatin domains described with the HiC method can be associated with combinations of histones (Figure 5A). In mammals, domains related to active chromatin usually show H3K4me3 and acetylated histones H3 and H4 at the promoters of genes, while

inactive domains display genes silenced and mark H3K27me3. Differently, H3K27me3-rich domains have the potential to be activated on demand, depending on the tissue, and so are more open than H3K9me3-rich domains [115]. Finally, domains associated to H3K9me2/H3K9me3 are very compact and constitute a significant portion of chromatin that is never active and is structurally very condensed. Figure 5B shows localisation microscopy images with the clusters of active, inactive and repressed chromatin, painted with H3K14ac, H3K9me3 and H3K27me3 respectively [10]. From these different findings, we propose a model for chromatin domain formation (Figure 5C).

Histone code across the cell cycle

During the cell cycle, the genome structure is heavily modified via epigenetic mechanisms. Nevertheless, activity and regulation of genes have to continue to a certain extent. Keeping the epigenetic makeup of chromatin during DNA condensation at the end of metaphase to switch to mitosis is relatively costly in terms of energy, so some optimal regulatory mechanisms have to happen to "memorise" the epigenetic status of the dividing cell. So far, we have considered cells in G1 or G0 phase, but the picture is rather different during mitosis, es-




Fig. 6. Histone code segregates functional compartments in the meiotic chromosomes. (A) Super-resolution microscopy reveals higher-order clusters of chromatin patterns along the pachytene chromosome. The DNA map of meiotic chromosomes at pachytene stage shows different condensation levels. These levels were found to be constrained by anti-correlating clusters of histone modifications (pink) along the synaptonemal complex proteins 3 (SYCP3) (green). Histone modifications associated with transcription (H3K4me3) emanate radially in loop-like structures while the histone modifications associated with repressive chromatin (H3K27me3) are confined to axial regions of the SYCP3. The histone modifications associated with centromeric chromatin (H3K9me3) are found at one end of the SYCP3 (figure and caption modified from [42, 114]). (B) A model for co-existence of different chromatin domains on a single chromosome. The two strands of the SYCP3 are displayed in black. In purple, active chromatin marked by H3K4me3 covers the entire surface of chromosomes (shown in blue). Symmetrical and periodic clusters of repressed chromatin H3K27me3 are shown with yellow spots. Inactive polar chromatin marked by H3K9me3 appears as an orange spiral pattern. The plot in the middle shows the distance distribution of DNA, active (H3K4me3) and inactive (H3K27me3, H3K9me3) regions of the genome from the central axis of SYCP3.

pecially with respect to histone modifications. Immediately following replication, histone proteins start to accumulate on the newly-synthesised DNA to form nucleosomes, involving a specific variant of histone H3, H3.1. The late cell cycle phase has two kinds of epigenetic regulations [116], which we review briefly below.

Firstly, some marks are involved in the mechanism of cell cycle progression itself. Accumulated H4K20me1 mark at gene bodies is read and leads to recruitment of factors participating in the initiation of DNA synthesis at origins of replication such as L3MBT1 and condensin II [117, 118]. This recognition mechanism leads to amplified events of methylation, and accumulation of marks H4K20me2 and H4K20me3 via methyltransferases SUV4-20H1 and SUV4-20H2 to switch from S- to M-phase [119–121]. Another mark involved in the process is H3S10 phosphorylation (H3S10ph). Along with methylation at position H4K20, it is the only histone site which varies through the cell cycle. This simple two-mark regulation is enough to induce cascades of factor expression involved in further triggering chromatin compaction during mitosis [122].

Secondly, besides the epigenetic mechanisms that accompany replication and mitosis, another kind of epigenetic regulation is devoted to preserving the transcriptional status of genes so that activity can be restored in daughter cells at the end of mitosis. The view gets even more complex when considering symmetric and asymmetric division [123, 124]. The two

marks which participate the most in this kind of remodelling are H3K27me3 and H3K9me3 [116], driving gene silencing and high chromatin compaction, respectively. A fundamental biological question is whether the epigenetic marks and so the histone code rules are preserved through replication, a mechanism named epigenetic inheritance [125–127]. In order replication to happen, the DNA becomes naked, and the nucleosomes need to be reassembled so chromatin keeps its functions in later stages. Many mechanisms are known, some involving a de novo “random” refolding which is independent of the DNA status before S-phase, and one mechanism dependent on the previous status, involving an epigenetic memory. In one model, all histone modifications are preserved and memorised on each DNA strand of the two double helices after replication. Differently, histone modifications can be partially transmitted to the daughter DNA molecules, but via copy mechanisms involving histone modifications, readers and writers recruited locally. In this case, all gaps can be filled to restore an epigenetic state very close to the one of the parent DNA molecule.

Another important question is the recycling of nucleosomes and histones. Are nucleosomes de-assembled and reconstructed in the newly synthesised DNA molecules or are they mostly transmitted in an intact form, with missing nucleosomes being de novo assembled? The answer is that in most cases, each nucleosome is transmitted intact to one of the two daughter strands, so the histone code is preserved [123, 128]. The

equivalent position on the other strand is possibly filled with information coming from adjacent redundant nucleosomes via enzymes CAF-1 and ASF1 [129].

Despite costly at first sight, epigenetic inheritance may be relatively cheap to maintain with the recruitment of proper readers and writers in the vicinity of replication sites to stochastically recreate the “parental” epigenetic make-up in daughter strands. In the future, integration of modelling, genomic and microscopy features will be key to understanding the mechanisms of epigenetic inheritance.

The association of histone code with cell identity and dynamics

Overall, genomic studies have shown that there is a substantial reshuffling of histone modifications during the cell cycle [84]. To show these associations, careful choice of FISH probes associated with H3K4me3/H3K27me3/H3K9me3 were found to demonstrate that chromatin domain stained with any of these marks anti-localise (Figure 6), a fact confirmed in *Drosophila* more recently, with overlaps [115]. The model of domains that we described previously is static, and in reality, chromatin domains experience a large change in embryonic development and differentiation processes of adult tissues (Figure 7). According to microscopy and genomic data, during early stages of development, a heavy remodelling of chromatin structure happens. Most of the chromatin is relatively loose at first, and becomes more compacted in later stages, with associated silencing of many genes. In embryonic stem cells, a relatively high number of genes are expressed, including genes for maintenance of stem cell status, such as *cMyc* or *Sox2* [130]. These genes are usually painted with H3K4me3 and H3K27ac marks at their promoters, prone to recruit transcription factors such as *Pou5f1*, *Sox2*, *cMyc* and *Klf4* [131]. Many specialised genes are on the other hand showing poised promoters. These promoters have an ambivalent epigenetic status, comprising both active and inactive marks H3K4me3 and H3K27me3, respectively, and have the potential to either be activated or repressed during differentiation.

Another key event of cell differentiation is the increase of the proportion of densely repressed chromatin, usually associated with telomeres and nucleoli. These regions are associated with H3K9me3 and eventually H3K27me3 (Figure 6B). Cells become slowly more specialised during development, which means that their specialised metabolism focuses on expressing of only a small subset of genes with a majority of genes being put in a repressed state. A lot of genomic locations indeed become repressed, histone combination usually turns to a deacetylated state with repressed marks, such as H3K27me3. Some tissues experience bigger changes than others, for instance, T




Fig. 7. Histone code during developmental processes. The model depicts the rules of the histone code associated to major developmental and regulatory processes happening in the human body. The different colours of circle feature different kinds of histone modifications combinations with the most significant modifications displayed. Blue: Euchromatin (H3K4me2/3 and H3K27ac, see Hawkins et al. 2010 [132]). Brown: repressed chromatin (H3K9me2/3, *lbid*). Red: Inactive chromatin (H3K27me3, *lbid*). Purple: Poised transcription (H3K4me3 and H3K27me3, see Gapp et al. 2014 [133]). Black: Quiescent chromatin (H4K20me2/3, see Everitts et al. 2013 and Onodera and Nakayama 2015 [134, 135]). Pathways from one genome status to another are figured by pale grey lines associated with names of chromatin modifiers or readers involved.

lymphocytes present an enrichment for the H3K27me3 mark compared to B lymphocytes [135].

A tight orchestration of gene regulation happens during the developmental processes associated with *Hox* genes, which are key in defining general body plans [136, 137]. They form clusters and are either transcribed together or sequentially during different developmental phases. In *Drosophila*, *Hox* genes are repressed at the beginning of development, under the influence of endogenous PcG complexes of maternal source [138, 139]. With time, *Hox* genes are de-repressed and activated by methylation of lysine 4 of histone three while H3K27me3 is dissociated. Progression is also spatial, with *Hox* genes being activated from one end of the cluster to the other [140]. Furthermore, both “active” and “inactive” parts of the *Hox* genes cluster form separate chromatin domains (Figure 5C). Differently, in tissues with no *Hox* gene expression, such as brain, most of clusters are silenced and fold according to cluster positions, instead of active/inactive status, showing a deep connection between histone code, dynamics of transcriptional regulation, high-order DNA folding and body-wide phenotype.

Conclusion

We have presented arguments for the existence of a spectrum of epigenetic features related to chromatin structure, epitomized by the term histone code. A consequence of the intrinsic redundancy of the code is robustness to small changes and therefore the message it delivers will not be very much altered if, for instance, an enzyme depositing one mark is not fully functional. Nevertheless, the position themselves where the histone modifications occur, are not prone to variations. As an example, the alteration of lysine H3K27 into a methionine is found in a certain type of glioma [141]. We can only hypothesize that similar mutations in other important modifications may have other dramatic effects for cellular phenotype and life span of organisms with either cancer or developmental defects. However, most importantly, redundancy of the code is a way to increase the probability for factors to be brought to the nucleosomes, for instance in a case of a protein complex where one protein binds to one histone modification, and another protein binds to a different histone modification. This is shown by the fact that factors from the same complex usually bind similar marks or associated marks either on the same histone or close positions. The free energy of the compound/nucleosome system will increase, increasing the probability of binding [142].

Redundancy of information provided by the histone code is an essential feature of robustness of cellular processes. As the number of histone modifications defining a given chromatin state increases, so does the probability to recruit specific factors that will change or modulate the state of chromatin. Overall, the complexity of histone modulation helps to ensure that events will happen at the right time in the right place. Though not fully understood yet, these processes are certainly a key point to understand the mechanisms that connect gene regulation and chromatin folding.

ACKNOWLEDGMENTS. D.F. thanks the Center for Computational Sciences in Mainz for the financial support. K.P. thanks Wioleta Dudka for proofreading the manuscript.

References

1. Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ (1997) Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature* 389(6648):251–260.
2. Andrews AJ, Luger K (2011) Nucleosome structure (s) and stability: variations on a theme. *Annual review of biophysics* 40:99–117.
3. Wong J et al. (1998) Distinct requirements for chromatin assembly in transcriptional repression by thyroid hormone receptor and histone deacetylase. *The EMBO Journal* 17(2):520–534.
4. Grunstein M (1997) Histone acetylation in chromatin structure and transcription. *Nature* 389(6649):349.
5. Lanctôt C, Cheutin T, Cremer M, Cavalli G, Cremer T (2007) Dynamic genome architecture in the nuclear space: regulation of gene expression in three dimensions. *Nature Reviews Genetics* 8(2):104–115.
6. Kornberg RD, Thonmas JO (1974) Chromatin structure: oligomers of the histones. *Science* 184(4139):865–868.

7. Richmond T, Finch J, Rushton B, Rhodes D, Klug A (1983) Structure of the nucleosome core particle at 7 Å resolution. *Nature* 311(5986):532–537.
8. Kornberg RD, Lorch Y (1999) Twenty-five years of the nucleosome, fundamental particle of the eukaryote chromosome. *Cell* 98(3):285–294.
9. Markaki Y et al. (2011) Functional nuclear organization of transcription and dna replication a topographical marriage between chromatin domains and the interchromatin compartment in *Cold Spring Harbor symposia on quantitative biology*. (Cold Spring Harbor Laboratory Press).
10. Prakash K (2017) Structure, function and dynamics of chromatin in *Chromatin Architecture*. (Springer International Publishing), pp. 63–103.
11. Dubochet J, Adrian M, Schultz P, Oudet P (1986) Cryo-electron microscopy of vitrified sv40 minichromosomes: the liquid drop model. *The EMBO journal* 5(3):519.
12. Fussner E, Ching RW, Bazett-Jones DP (2011) Living without 30nm chromatin fibers. *Trends in biochemical sciences* 36(1):1–6.
13. Chadwick J (1990) *The decipherment of Linear B*. (Cambridge University Press).
14. Singh S (2000) *The code book: the secret history of codes and code-breaking*. (Fourth Estate).
15. Fischle W et al. (2005) Regulation of hp1-chromatin binding by histone h3 methylation and phosphorylation. *Nature* 438(7071):1116.
16. Ekwall K (2007) Epigenetic control of centromere behavior. *Annu. Rev. Genet.* 41:63–81.
17. Santos-Rosa H et al. (2002) Active genes are tri-methylated at k4 of histone h3. *Nature* 419(6905):407–411.
18. Ruthenburg AJ, Allis CD, Wysocka J (2007) Methylation of lysine 4 on histone h3: intricacy of writing and reading a single epigenetic mark. *Molecular cell* 25(1):15–30.
19. Chandraratna T et al. (2012) Independence of repressive histone marks and chromatin compaction during senescent heterochromatic layer formation. *Molecular cell* 47(2):203–214.
20. Strahl BD, Allis CD (2000) The language of covalent histone modifications. *Nature* 403(6765):41–45.
21. Jenuwein T, Allis CD (2001) Translating the histone code. *Science* 293(5532):1074–1080.
22. Turner BM (2002) Cellular memory and the histone code. *Cell* 111(3):285–291.
23. Barbieri M (2003) *The organic codes: an introduction to semantic biology*. (Cambridge University Press).
24. Barbieri M (2008) Biosemiotics: a new understanding of life. *Naturwissenschaften* 95(7):577–599.
25. Kühn S, Hofmeyr JHS (2014) Is the “histone code” an organic code? *Biosemiotics* 7(2):203–222.
26. Stedman E, Stedman E (1951) The basic proteins of cell nuclei. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 235(630):565–595.
27. Wilkins M (1956) Physical studies of the molecular structure of deoxyribose nucleic acid and nucleoprotein in *Cold Spring Harbor symposia on quantitative biology*. (Cold Spring Harbor Laboratory Press), Vol. 21, pp. 75–90.
28. Finch J, Klug A (1976) Solenoidal model for superstructure in chromatin. *Proceedings of the National Academy of Sciences* 73(6):1897–1901.
29. Busch H (2012) *The cell nucleus*. (Elsevier) Vol. 3.
30. Langridge R et al. (1957) Molecular structure of deoxyribonucleic acid (dna). *The Journal of biophysical and biochemical cytology* 3(5):767.
31. Wilkins M, Zubay G, Wilson H (1959) X-ray diffraction studies of the molecular structure of nucleohistone and chromosomes. *Journal of Molecular Biology* 1(2):179IN7–185IN10.
32. Pardon J, Richards B, Cotter R (1974) X-ray diffraction studies on oriented nucleohistone gels in *Cold Spring Harbor symposia on quantitative biology*. (Cold Spring Harbor Laboratory Press), Vol. 38, pp. 75–81.
33. Oudet P, Gross-Bellard M, Chambon P (1975) Electron microscopic and biochemical evidence that chromatin structure is a repeating unit. *Cell* 4(4):281–300.
34. DeLange RJ, Hooper JA, Smith EL (1973) Histone iii. sequence studies on the cynaogen bromide peptides; complete amino acid sequence of calf thymus histone iii. *Journal of Biological Chemistry* 248(9):3261–3274.
35. Adamietz P, Rudolph A (1984) Adp-ribosylation of nuclear proteins in vivo. identification of histone h2b as a major acceptor for mono-and poly (adp-ribose) in dimethyl sulfate-treated hepatoma ah 7974 cells. *Journal of Biological Chemistry* 259(11):6841–6846.
36. Liebich H et al. (1993) Non-enzymatic glycation of histones. *Biological mass spectrometry* 22(2):121–123.
37. Hymes J, Fleischhauer K, Wolf B (1995) Biotinylation of histones by human serum biotinidase: assessment of biotinyl-transferase activity in sera from normal individuals and children with biotinidase deficiency. *Biochemical and molecular medicine* 56(1):76–83.
38. WONDRAK GT, CERVANTES-LAUREAN D, JACOBSON EL, JACOBSON MK (2000) Histone carbonylation in vivo and in vitro. *Biochemical Journal* 351(3):769–777.
39. Tan M et al. (2011) Identification of 67 histone marks and histone lysine crotonylation as a

- new type of histone modification. *Cell* 146(6):1016–1028.
40. Rodríguez-Paredes M, Esteller M (2011) Cancer epigenetics reaches mainstream oncology. *Nature medicine* pp. 330–339.
 41. Tarakhovskiy A (2010) Tools and landscapes of epigenetics. *Nature immunology* 11(7):565–568.
 42. Prakash K, Fournier D (2017) Histone code and higher-order chromatin folding: A hypothesis. *Genomics and Computational Biology* 3(2):41.
 43. Kouzarides T (2007) Chromatin modifications and their function. *Cell* 128(4):693–705.
 44. Allfrey V, Faulkner R, Mirsky A (1964) Acetylation and methylation of histones and their possible role in the regulation of rna synthesis. *Proceedings of the National Academy of Sciences* 51(5):786–794.
 45. Wade PA, Pruss D, Wolffe AP (1997) Histone acetylation: chromatin in action. *Trends in biochemical sciences* 22(4):128–132.
 46. Henikoff S (2005) Histone modifications: combinatorial complexity or cumulative simplicity? *Proceedings of the National Academy of Sciences of the United States of America* 102(15):5308–5309.
 47. Sneppen K, Dodd IB (2012) A simple histone code opens many paths to epigenetics. *PLoS computational biology* 8(8):e1002643.
 48. Cepeda-Humerez SA, Rieckh G, Tkačik G (2015) Stochastic proofreading mechanism alleviates crosstalk in transcriptional regulation. *Physical review letters* 115(24):248101.
 49. Winston F, Allis CD (1999) The bromodomain: a chromatin-targeting module? *Nature Structural & Molecular Biology* 6(7):601–604.
 50. Biswas M, Langowski J, Bishop TC (2013) Atomistic simulations of nucleosomes. *Wiley Interdisciplinary Reviews: Computational Molecular Science* 3(4):378–392.
 51. Norton VG, Imai BS, Yau P, Bradbury EM (1989) Histone acetylation reduces nucleosome core particle linking number change. *Cell* 57(3):449–457.
 52. Cortini R et al. (2016) The physics of epigenetics. *Reviews of Modern Physics* 88(2):025002.
 53. Bertin A, Mangelot S, Renouard M, Durand D, Livolant F (2007) Structure and phase diagram of nucleosome core particles aggregated by multivalent cations. *Biophysical journal* 93(10):3652–3663.
 54. Yang Z, Hayes JJ (2011) The divalent cations ca²⁺ and mg²⁺ play specific roles in stabilizing histone–dna interactions within nucleosomes that are partially redundant with the core histone tail domains. *Biochemistry* 50(46):9973–9981.
 55. Kuo MH, Allis CD (1998) Roles of histone acetyltransferases and deacetylases in gene regulation. *Bioessays* 20(8):615–626.
 56. Wang J et al. (2012) Nutrition, epigenetics, and metabolic syndrome. *Antioxidants & redox signaling* 17(2):282–301.
 57. Mathis DJ, Oudet P, Wasyluk B, Chambon P (1978) Effect of histone acetylation on structure and in vitro transcription of chromatin. *Nucleic acids research* 5(10):3523–3548.
 58. Hong L, Schroth G, Matthews H, Yau P, Bradbury E (1993) Studies of the dna binding properties of histone h4 amino terminus. thermal denaturation studies reveal that acetylation markedly reduces the binding constant of the h4" tail" to dna. *Journal of Biological Chemistry* 268(1):305–314.
 59. Morinière J et al. (2009) Cooperative binding of two acetylation marks on a histone tail by a single bromodomain. *Nature* 461(7264):664.
 60. Cosgrove MS, Boeke JD, Wolberger C (2004) Regulated nucleosome mobility and the histone code. *Nature structural & molecular biology* 11(11):1037–1043.
 61. Prakash K (2017) A condensed history of chromatin research in *Chromatin Architecture*. (Springer International Publishing), pp. 1–24.
 62. Cao R et al. (2002) Role of histone h3 lysine 27 methylation in polycomb-group silencing. *Science* 298(5595):1039–1043.
 63. Czermin B et al. (2002) Drosophila enhancer of zeste/esc complexes have a histone h3 methyltransferase activity that marks chromosomal polycomb sites. *Cell* 111(2):185–196.
 64. Dhalluin C, Carlson JE, Zeng L, He C, et al. (1999) Structure and ligand of a histone acetyltransferase bromodomain. *Nature* 399(6735):491.
 65. Santillan DA et al. (2006) Bromodomain and histone acetyltransferase domain specificities control mixed lineage leukemia phenotype. *Cancer research* 66(20):10032–10039.
 66. Wagner A (2008) Robustness and evolvability: a paradox resolved. *Proceedings of the Royal Society of London B: Biological Sciences* 275(1630):91–100.
 67. Barbieri M et al. (2017) Active and poised promoter states drive folding of the extended hoxb locus in mouse embryonic stem cells. *Nature structural & molecular biology* 24(6):515–524.
 68. Mattioli F et al. (2017) Structure of histone-based chromatin in archaea. *Science* 357(6351):609–612.
 69. Ho JW et al. (2014) Comparative analysis of metazoan chromatin organization. *Nature* 512(7515):449–452.
 70. Consortium EP, et al. (2004) The encode (encyclopedia of dna elements) project. *Science* 306(5696):636–640.
 71. Consortium EP, et al. (2012) An integrated encyclopedia of dna elements in the human genome. *Nature* 489(7414):57.
 72. Laible G et al. (1997) Mammalian homologues of the polycomb-group gene enhancer of zeste mediate gene silencing in drosophila heterochromatin and at s. cerevisiae telomeres. *The EMBO journal* 16(11):3219–3232.
 73. Luense LJ et al. (2016) Comprehensive analysis of histone post-translational modifications in mouse and human male germ cells. *Epigenetics & chromatin* 9(1):24.
 74. Shulha HP et al. (2012) Human-specific histone methylation signatures at transcription start sites in prefrontal neurons. *PLoS biology* 10(11):e1001427.
 75. Barski A et al. (2007) High-resolution profiling of histone methylations in the human genome. *Cell* 129(4):823–837.
 76. Pei H et al. (2011) Mms19 regulates histone h4k20 methylation and 53bp1 accumulation at dna damage sites. *Nature* 470(7332):124.
 77. Kuo AJ et al. (2012) Orc1 bah domain links h4k20me2 to dna replication licensing and meier-gorlin syndrome. *Nature* 484(7392):115.
 78. Müller J et al. (2002) Histone methyltransferase activity of a drosophila polycomb group repressor complex. *Cell* 111(2):197–208.
 79. de Napoles M et al. (2004) Polycomb group proteins ring1a/b link ubiquitylation of histone h2a to heritable gene silencing and x inactivation. *Developmental cell* 7(5):663–676.
 80. Cooper S et al. (2014) Targeting polycomb to pericentric heterochromatin in embryonic stem cells reveals a role for h2ak119u1 in prc2 recruitment. *Cell reports* 7(5):1456–1470.
 81. Blackledge NP et al. (2014) Variant prc1 complex-dependent h2a ubiquitylation drives prc2 recruitment and polycomb domain formation. *Cell* 157(6):1445–1459.
 82. Fraser P, Bickmore W (2007) Nuclear organization of the genome and the potential for gene regulation. *Nature* 447(7143):413.
 83. Prigent C, Dimitrov S (2003) Phosphorylation of serine 10 in histone h3, what for? *Journal of cell science* 116(18):3677–3685.
 84. Rao SS et al. (2014) A 3d map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 159(7):1665–1680.
 85. Dodd IB, Micheelsen MA, Sneppen K, Thon G (2007) Theoretical analysis of epigenetic cell memory by nucleosome modification. *Cell* 129(4):813–822.
 86. Dodd IB, Sneppen K (2011) Barriers and silencers: a theoretical toolkit for control and containment of nucleosome-based epigenetic states. *Journal of molecular biology* 414(4):624–637.
 87. Bernstein BE et al. (2006) A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* 125(2):315–326.
 88. Shema E et al. (2016) Single-molecule decoding of combinatorially modified nucleosomes. *Science* 352(6286):717–721.
 89. Betzig E et al. (2006) Imaging intracellular fluorescent proteins at nanometer resolution. *Science* 313(5793):1642–1645.
 90. Rust MJ, Bates M, Zhuang X (2006) Stochastic optical reconstruction microscopy (storm) provides sub-diffraction-limit image resolution. *Nature methods* 3(10):793.
 91. Schermelleh L et al. (2008) Subdiffraction multicolor imaging of the nuclear periphery with 3d structured illumination microscopy. *Science* 320(5881):1332–1336.
 92. Prakash K (2017) High-density superresolution microscopy with an incoherent light source and a conventional epifluorescence. *bioRxiv* p. 121061.
 93. Żurek-Biesiada D et al. (2015) Localization microscopy of dna in situ using vybrant® dye-cycle violet fluorescent probe: A new approach to study nuclear nanostructure at single molecule resolution. *Experimental cell research*.
 94. Żurek-Biesiada D et al. (2016) Quantitative super-resolution localization microscopy of dna in situ using vybrant® dye-cycle violet fluorescent probe. *Data in Brief* 7:157–171.
 95. Szczurek AT et al. (2014) Single molecule localization microscopy of the distribution of chromatin using hoechst and dapi fluorescent probes. *Nucleus* 5(4):331–340.
 96. Prakash K et al. (2015) Superresolution imaging reveals structurally distinct periodic patterns of chromatin along pachytene chromosomes. *Proceedings of the National Academy of Sciences* 112(47):14635–14640.
 97. Kirmes I et al. (2015) A transient ischemic environment induces reversible compaction of chromatin. *Genome biology* 16(1):1–19.
 98. Popken J et al. (2015) Remodeling of the nuclear envelope and lamina during bovine preimplantation development and its functional implications. *PLoS One*.
 99. Zinner R, Teller K, Versteeg R, Cremer T, Cremer M (2007) Biochemistry meets nuclear architecture: multicolor immuno-fish for co-localization analysis of chromosome segments and differentially expressed gene loci with various histone methylations. *Advances in enzyme regulation* 47(1):223–241.
 100. Garcia BA et al. (2007) Chemical derivatization of histones for facilitated analysis by mass spectrometry. *Nature protocols* 2(4):933.
 101. Ernst J, Kellis M (2010) Discovery and characterization of chromatin states for systematic

- annotation of the human genome. *Nature biotechnology* 28(8):817–825.
102. Prakash K (2012) A binary combinatorial histone code. *Master thesis*. Aalto University.
 103. Tse C, Sera T, Wolffe AP, Hansen JC (1998) Disruption of higher-order folding by core histone acetylation dramatically enhances transcription of nucleosomal arrays by rna polymerase iii. *Molecular and cellular biology* 18(8):4629–4638.
 104. Zentner GE, Henikoff S (2013) Regulation of nucleosome dynamics by histone modifications. *Nature structural & molecular biology* 20(3):259–266.
 105. Carrozza MJ et al. (2005) Histone h3 methylation by set2 directs deacetylation of coding regions by rpd3s to suppress spurious intragenic transcription. *Cell* 123(4):581–592.
 106. Krogan NJ et al. (2003) Methylation of histone h3 by set2 in *saccharomyces cerevisiae* is linked to transcriptional elongation by rna polymerase ii. *Molecular and cellular biology* 23(12):4207–4218.
 107. Prakash K (2017) Chromatin architecture: Advances from high-resolution single molecule dna imaging.
 108. Cedar H, Bergman Y (2009) Linking dna methylation and histone modification: patterns and paradigms. *Nature reviews. Genetics* 10(5):295.
 109. Lunyak VV et al. (2002) Corepressor-dependent silencing of chromosomal regions encoding neuronal genes. *Science* 298(5599):1747–1752.
 110. Fuks F et al. (2003) The methyl-cpg-binding protein mecp2 links dna methylation to histone methylation. *Journal of Biological Chemistry* 278(6):4035–4040.
 111. Dixon JR et al. (2012) Topological domains in mammalian genomes identified by analysis of chromatin interactions. *Nature* 485(7398):376–380.
 112. Prakash K (2017) Periodic and symmetric organisation of meiotic chromosomes in *Chromatin Architecture*. (Springer International Publishing), pp. 105–133.
 113. Smeets D et al. (2014) Three-dimensional super-resolution microscopy of the inactive x chromosome territory reveals a collapse of its active nuclear compartment harboring distinct xist rna foci. *Epigenetics & chromatin* 7(8).
 114. Prakash K (2016) The periodic and dynamic structure of chromatin. *PhD thesis*. Heidelberg University.
 115. Boettiger AN et al. (2016) Super-resolution imaging reveals distinct chromatin folding for different epigenetic states. *Nature* 529(7586):418–422.
 116. Zee BM, Britton LMP, Wolle D, Haberman DM, Garcia BA (2012) Origins and formation of histone methylation across the human cell cycle. *Molecular and cellular biology* 32(13):2503–2514.
 117. Liu W et al. (2010) Phf8 mediates histone h4 lysine 20 demethylation events involved in cell cycle progression. *Nature* 466(7305):508.
 118. Trojer P et al. (2007) L3mbtl1, a histone-methylation-dependent chromatin lock. *Cell* 129(5):915–928.
 119. Jørgensen S et al. (2007) The histone methyltransferase set8 is required for s-phase progression. *J Cell Biol* 179(7):1337–1345.
 120. Rice JC et al. (2002) Mitotic-specific methylation of histone h4 lys 20 follows increased pr-set7 expression and its localization to mitotic chromosomes. *Genes & development* 16(17):2225–2230.
 121. Schotta G et al. (2008) A chromatin-wide transition to h4k20 monomethylation impairs genome integrity and programmed dna rearrangements in the mouse. *Genes & development* 22(15):2048–2061.
 122. Wilkins BJ et al. (2014) A cascade of histone modifications induces chromatin condensation in mitosis. *Science* 343(6166):77–80.
 123. Xu M et al. (2010) Partitioning of histone h3-h4 tetramers during dna replication-dependent chromatin assembly. *Science* 328(5974):94–98.
 124. Tran V, Lim C, Xie J, Chen X (2012) Asymmetric division of drosophila male germline stem cell shows asymmetric histone distribution. *Science* 338(6107):679–682.
 125. Henikoff S, Furuyama T, Ahmad K (2004) Histone variants, nucleosome assembly and epigenetic inheritance. *Trends in Genetics* 20(7):320–326.
 126. Martin C, Zhang Y (2007) Mechanisms of epigenetic inheritance. *Current opinion in cell biology* 19(3):266–272.
 127. Probst AV, Dunleavy E, Almouzni G (2009) Epigenetic inheritance during the cell cycle. *Nature reviews Molecular cell biology* 10(3):192–206.
 128. Jackson V (1990) In vivo studies on the dynamics of histone-dna interaction: evidence for nucleosome dissolution during replication and transcription and a low level of dissolution independent of both. *Biochemistry* 29(3):719–731.
 129. Budhavarapu VN, Chavez M, Tyler JK (2013) How is epigenetic information maintained through dna replication? *Epigenetics & chromatin* 6(1):32.
 130. Chen T, Dent SY (2014) Chromatin modifiers: regulators of cellular differentiation. *Nature reviews. Genetics* 15(2):93.
 131. Takahashi K, Yamanaka S (2006) Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *cell* 126(4):663–676.
 132. Hawkins RD et al. (2010) Distinct epigenomic landscapes of pluripotent and lineage-committed human cells. *Cell stem cell* 6(5):479–491.
 133. Gapp K et al. (2014) Implication of sperm rnas in transgenerational inheritance of the effects of early trauma in mice. *Nature neuroscience* 17(5):667–669.
 134. Everitts AG et al. (2013) H4k20 methylation regulates quiescence and chromatin compaction. *Molecular biology of the cell* 24(19):3025–3037.
 135. Onodera A, Nakayama T (2015) Epigenetics of t cells regulated by polycomb/trithorax molecules. *Trends in molecular medicine* 21(5):330–340.
 136. Bernstein BE et al. (2005) Genomic maps and comparative analysis of histone modifications in human and mouse. *Cell* 120(2):169–181.
 137. Chi P, Allis CD, Wang GG (2010) Covalent histone modifications—miswritten, misinterpreted and mis-erased in human cancers. *Nature reviews cancer* 10(7):457–469.
 138. Beuchle D, Struhl G, Muller J (2001) Polycomb group proteins and heritable silencing of drosophila hox genes. *Development* 128(6):993–1004.
 139. Spemann A, van Lohuizen M (2006) Polycomb silencers control cell fate, development and cancer. *Nature Reviews Cancer* 6(11):846–856.
 140. Montavon T, Duboule D (2013) Chromatin organization and global regulation of hox gene clusters. *Phil. Trans. R. Soc. B* 368(1620):20120367.
 141. Chan KM et al. (2013) The histone h3. 3k27m mutation in pediatric glioma reprograms h3k27 methylation and gene expression. *Genes & development* 27(9):985–990.
 142. Ruthenburg AJ, Li H, Patel DJ, Allis CD (2007) Multivalent engagement of chromatin modifications by linked binding modules. *Nature reviews. Molecular cell biology* 8(12):983.