



HAL
open science

Tidal variability of CO₂ and CH₄ emissions from the water column within a *Rhizophora* mangrove forest (New Caledonia)

Adrien Jacotot, Cyril Marchand, Michel Allenbach

► To cite this version:

Adrien Jacotot, Cyril Marchand, Michel Allenbach. Tidal variability of CO₂ and CH₄ emissions from the water column within a *Rhizophora* mangrove forest (New Caledonia). *Science of the Total Environment*, 2018, 631-632, pp.334-340. 10.1016/j.scitotenv.2018.03.006 . hal-01826186

HAL Id: hal-01826186

<https://hal.sorbonne-universite.fr/hal-01826186v1>

Submitted on 29 Jun 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Short Communication

Tidal variability of CO₂ and CH₄ emissions from the water column within a *Rhizophora* mangrove forest (New Caledonia)☆

Adrien Jacotot^{a,b,*}, Cyril Marchand^a, Michel Allenbach^b


^a IMPMC, Institut de Recherche pour le Développement (IRD), UPMC, CNRS, MNHN, Noumea, New Caledonia, France

^b Université de la Nouvelle-Calédonie, ISEA, EA 7484, BPR4, 98851 Noumea, New Caledonia

HIGHLIGHTS

- CO₂ and CH₄ fluxes from the water column were studied within a mangrove forest.
- Mean emissions were 3.35 mmolC m⁻² h⁻¹ for CO₂, and 18.30 μmolC m⁻² h⁻¹ for CH₄.
- δ¹³CO₂ ranged from -26.9‰ to -8.6‰, suggesting a mixing between different sources.
- CO₂ and CH₄ emissions were 1.9 and 5.5 times higher during ebb than during flow.
- Spring tides induced higher CO₂ and CH₄ emissions than neap tides.

GRAPHICAL ABSTRACT



ABSTRACT

We performed a preliminary study to quantify CO₂ and CH₄ emissions from the water column within a *Rhizophora* spp. mangrove forest. Mean CO₂ and CH₄ emissions during the studied period were 3.35 ± 3.62 mmolC m⁻² h⁻¹ and 18.30 ± 27.72 μmolC m⁻² h⁻¹, respectively. CO₂ and CH₄ emissions were highly variable and mainly driven by tides (flow/ebb, water column thickness, neap/spring). Indeed, an inverse relationship between the magnitude of the emissions and the thickness of the water column above the mangrove soil was observed. δ¹³CO₂ values ranged from -26.88‰ to -8.6‰, suggesting a mixing between CO₂-enriched pore waters and lagoon incoming waters. In addition, CO₂ and CH₄ emissions were significantly higher during ebb tides, mainly due to the progressive enrichment of the water column by diffusive fluxes as its residence time over the forest floor increased. Eventually, we observed higher CO₂ and CH₄ emissions during spring tides than during neap tides, combined to depleted δ¹³CO₂ values, suggesting a higher contribution of soil-produced gases to the emissions. These higher emissions may result from higher renewable of the electron acceptor and enhanced exchange surface between the soil and the water column. This study shows that CO₂ and CH₄ emissions from the water column were not negligible and must be considered in future carbon budgets in mangroves.

Keywords:

Water to atmosphere fluxes
Greenhouse gas
Blue carbon
Carbon dioxide isotope
Semi-arid mangrove forest

1. Introduction

Mangroves are considered as major ecosystems in the carbon cycle along tropical and subtropical coastlines, being among the most efficient blue carbon sinks (Kauffman et al., 2011). Due to their high primary productivity, estimated at 218 ± 72 TgC year⁻¹ (Bouillon et al.,

☆ Declarations of interest: none.

* Corresponding author at: Institut de Recherche pour le Développement, BPA5, 98848 Nouméa, New Caledonia.

E-mail address: adrien.jacotot@protonmail.com (A. Jacotot).

2008), and their long-term carbon storage capacities (Donato et al., 2011), mangroves have been recognized as having a key role in climate change mitigation (Howard et al., 2017; Mcleod et al., 2011). However, part of mangroves primary productivity is mineralized in their soils, producing greenhouse gas (GHG) that can be subsequently emitted towards the atmosphere (Chen et al., 2014). Among GHG, methane may be of major concern due to its global warming potential, 34 times higher than CO₂ using a 100 year time frame and climate carbon feedback (Stocker et al., 2013). In anoxic mangrove soils, when electron acceptors such free oxygen, metal oxides, nitrates, and sulfates have been exhausted, methanogenesis can occur. Recently, it was demonstrated that sulfate reducing and methanogens microorganisms can coexist in mangrove soils and it was thus suggested that methane emissions have been underestimated (Chauhan et al., 2015; Lyimo et al., 2009).

In mangrove soils, organic matter (OM) decay processes depend on numerous factors, including waterlogging (Kristensen et al., 2008a). In fact, mangroves are subjected to tidal cycles that influence the redox characteristics of their soils, modifying the rate of GHG production. In addition, tides can also influence GHG emissions towards the atmosphere since gas diffusion differs when the soil is immersed or unsaturated. Recently, many studies focused on CO₂ or CH₄ emissions at the soil-air interface at low tide (Bulmer et al., 2015; Chanda et al., 2013; Chauhan et al., 2015; Chen et al., 2016; Chen et al., 2010; Leopold et al., 2013, 2015; Livesley and Andrusiak, 2012; Wang et al., 2016), or at the water-air interface in adjacent tidal creeks (Borges et al., 2003; Bouillon et al., 2003b; Call et al., 2015; Maher et al., 2015). However, to our knowledge, no study focused on water to atmosphere GHG emissions when the forest is inundated, which may frequently occur, depending on the tidal range and the position of the forest in the intertidal zone.

Stable isotope of carbon ($\delta^{13}\text{C}$) is a widely used tool for studying carbon dynamic in natural environments (Gonneea et al., 2004; Graham et al., 2001; Lu et al., 2016; Yamamuro, 2000), including the partitioning of the different sources (Midwood and Millard, 2011; Millard et al., 2010; Paterson et al., 2009). In mangrove ecosystems, mangrove litter, benthic microalgae, phytoplankton and seagrass detritus are the main OM sources (Kristensen et al., 2008a), and the latter may be distinguishable thanks to their specific $\delta^{13}\text{C}$ values. Marine OM has typical value of $\delta^{13}\text{C}$ ranging from -30 to -16% , whereas mangroves plants are C3 photosynthetic pathways, and therefore produce OM that have a $\delta^{13}\text{C}$ value ranging from -32 to -21% (Lamb et al., 2006). In addition, due to a low carbon fractionation during respiration processes (Lin and Ehleringer, 1997; Maher et al., 2015), the $\delta^{13}\text{C}$ of the CO₂ produced is close to the $\delta^{13}\text{C}$ value of its source. Recent development of advanced technologies such as cavity ring-down spectroscopy (CRDS) allows high resolution in situ measurements of $\delta^{13}\text{CO}_2$. Therefore, these new analytical means could help to identify the origins of the CO₂ emitted from mangrove ecosystems.

Within this context, the present study aimed to (i) quantify the CO₂ and CH₄ emissions at the water-air interface within a *Rhizophora* spp. mangrove forest, (ii) evaluate the variability of these emissions along tidal cycles, and (iii) identify the origin of the CO₂ fluxes measured. To reach our goals, we measured CO₂ and CH₄ fluxes, as well as $\delta^{13}\text{CO}_2$ values, in the field, using a dark floating chamber connected to a cavity ring-down spectrometer analyzer (CRDS), along different tidal cycles from neap to spring tides.

2. Material and methods

2.1. Study site

The present study was conducted in the mangrove of Ouemo (22°16'50"S, 166°28'16"E), in New Caledonia, a French overseas archipelago located in the South Pacific (21°21'S, 165°27'E). The studied mangrove was dominated by three *Rhizophora* mangrove species: *R. stylosa*, *R. samoensis* and *R. selala*. Climate in the region is semi-arid

and strongly influenced by the inter-tropical convergence zone and by the El Niño Southern Oscillation (ENSO). Average air temperature varied between 20.5 and 26.6 °C, with a mean annual precipitation of 1070 mm (data from meteofrance.com). The tidal regime is semi-diurnal, with a tidal range ranging from 1.10 to 1.70 m.

2.2. Gas fluxes measurements

CO₂, $\delta^{13}\text{CO}_2$ and CH₄ measurements were performed using a dark custom-built floating chamber (466 cm²; 5050 cm³) connected to a G2131-*i* CRDS analyzer (Picarro Inc., Santa Clara, CA, USA) that measures gas concentrations at a frequency of 1 Hz. Guaranteed precision of the analyzer are 200 ppb + 0.05% of reading, 50 ppb + 0.05% of reading and 0.1‰, for CO₂ and CH₄, and $\delta^{13}\text{CO}_2$, respectively. Accuracy of the CRDS analyzer was periodically checked using certified N₂ (0 ppm CO₂ and CH₄), CO₂ (503 ppm) and CH₄ (100 ppm) gas standard samples (Calgaz, Air Liquide, USA).

We are aware that floating chambers may induce a bias in flux measurements during windy conditions or with high current velocity that induce artificial turbulences and, as a result, increase fluxes (Kremer et al., 2003; Yang et al., 2014). However, in low turbulence environment, like inside the mangrove forest where the wind is almost null, and the water flow is slow, the floating chamber technique can be a powerful method (Lorke et al., 2015; Vachon et al., 2010). Thus, we feel confident about the validity of our measurements. In addition, floating chambers have the capability to capture ebullition events that may account for a large proportion of the gas transferred to the atmosphere, particularly CH₄ (Chuang et al., 2017).

Measurements were performed every three weeks, from December 2016 to September 2017, at a single sampling station. Measurements were done during sunny days, with the slack high tide around noon, and included different tidal regimes. During neap tides, the tidal range varied from 1.10 to 1.25 m, which corresponded to a water column thickness above the mangrove sediment of 15 to 30 cm at the maximum of the high tide. During spring tides, the tidal range varied from 1.25 to 1.55 m, and, therefore, the water column thickness at the maximum of the high tide ranged from 30 cm to 60 cm. Measurements were performed all along the high tide, from the beginning of the flow to the end of the ebb. For each incubation, an integrating period of 3 to 6 min was chosen, depending on the linearity of the signal. Therefore, 19 to 40 measurements were performed during each campaign, resulting in a total of 284 flux measurements.

In addition, before each incubation, water level was measured thanks to a water gauge, and air temperature was recorded thanks to a handheld Skymate SM-19 thermometer.

2.3. Flux calculation

Water to atmosphere fluxes of CO₂ and CH₄ were integrated as a function of time, and calculated using the following formula:

$$F_{(\text{CO}_2, \text{CH}_4)} = (d(\text{CO}_2, \text{CH}_4)/dt) * V / (R * S * T) * 3.6$$

where F is the water to atmosphere fluxes of CO₂ or CH₄ (mmolC m⁻² h⁻¹); d(CO₂, CH₄) / dt is the variation in CO₂ or CH₄ as a function of time (ppm s⁻¹); V is the total volume of the system (m³); R is the ideal gas constant of 8.205746 10⁻⁵ (atm m³ K⁻¹ mol⁻¹); T is the absolute air temperature (K); and S is the area of the bottom of the incubation chamber (m²).

2.4. $\delta^{13}\text{CO}_2$

To measure the isotopic value of the CO₂ ($\delta^{13}\text{CO}_2$) released from the water column, we used a Keeling plot approach (Keeling, 1958, 1961; Pataki et al., 2003). By plotting the $\delta^{13}\text{CO}_2$ value CO₂ as a function of the inverse of the CO₂ concentration ($\delta^{13}\text{CO}_2 = f(1/\text{CO}_2)$) during each incubation, the intercept of a linear regression with the y-axis is equivalent to the $\delta^{13}\text{CO}_2$ value of the flux.

In addition, three supplementary incubations were realized at high tide outside the mangrove forest to evaluate the $\delta^{13}\text{CO}_2$ value of the CO_2 produced within the water column in the lagoon.

2.5. Statistical analyses

Student's *t*-tests were used to test the significant differences ($p < 0.05$) in the emissions between flow and ebb tides as well as between spring and neap tides. All statistical analyses were performed under R software version 3.3.1 (R Development Core Team, 2008).

3. Results and discussion

CO_2 and CH_4 emissions from the water column to the atmosphere varied from 0.13 to 18.41 $\text{mmolC m}^{-2} \text{h}^{-1}$ and from 0.18 to 172.07 $\mu\text{molC m}^{-2} \text{h}^{-1}$, respectively (Fig. 1). Thus, when converted into CO_2 -equivalent and considering its 100-year global warming potential (Stocker et al., 2013), methane represented 18% of the total emissions measured ($\text{CO}_2 + \text{CH}_4$). Gas transfer to the atmosphere may occur by upward molecular diffusion (Kristensen et al., 2008b) but also through gas bubble ebullition, mainly for methane due to its lower solubility (Barnes et al., 2006; Komiya et al., 2015; Stamp et al., 2013). However, in the present study, considering the linear evolution of CO_2 and CH_4 concentrations with time during each flux measurement, we suggest that only the diffusive fluxes were measured. As a consequence, the emissions reported herein may have been under-estimated, particularly methane.

To our knowledge, this study was the first one interested in measuring GHG emissions at the water-air interface within a mangrove forest. Therefore, we decided to compare our results with fluxes from other interfaces (*i.e.* from mangrove soils and creek waters to the atmosphere). CO_2 emissions were in the same order of magnitude and even higher than from mangrove soils at low tide or from mangrove tidal creeks (Table 1). It was demonstrated that soil surface biofilm may limit GHG emissions from mangrove soils, notably by acting as a physical barrier but also by using the CO_2 for primary production through photosynthesis (Bulmer et al., 2015; Leopold et al., 2015; Leopold et al., 2013). As a result, some authors reported negative fluxes at the soil surface (*e.g.* Chen et al., 2014; Leopold et al., 2015), which was never the case from the water column in the studied *Rhizophora* forest. Closed canopy and elevated water turbidity probably limited the development of




Fig. 1. Mean CO_2 ($\text{mmolC m}^{-2} \text{h}^{-1}$) and CH_4 ($\mu\text{molC m}^{-2} \text{h}^{-1}$) (+SD) emissions during the complete tidal cycles ($n = 284$), the flow tides ($n = 141$) and the ebb tides ($n = 143$) in the mangrove of Ouemo for the period December 2016 to September 2017. Different letters indicate significant differences (two-samples *t*-test, CO_2 : $t_{(205.66)} = 5.11$, $p < 0.001$, CH_4 : $t_{(146.3)} = 7.96$, $p < 0.001$).

microalgae that might have used the CO_2 produced in the soil. In addition, low tide fluxes measurements are strictly representative of the surface that is covered by the incubation chamber, and, usually, roots and crab burrows are excluded to avoid overestimations of the fluxes. At the opposite, water-air fluxes integrate the entire soil surface that is flooded by the water, and, therefore, roots and burrows are an important factor to take into account as they represent preferential diffusion pathways for the gases to escape from the sediment. The density of *Rhizophora* roots and crab burrows could not be measured in our study site, however, the density of crab burrows was reported for other mangrove in New Caledonia and was evaluated to 83 burrows m^{-2} for the fiddler crab *Uca* spp. (Nielsen et al., 2003), and from 8.9 to 29.1 crabs ha^{-1} for the mud crab *Scylla serrata* (Dumas et al., 2012).

However, regarding CH_4 , some studies reported much higher values from mangrove soils but lower from tidal creeks (Table 1). When CH_4 diffuses from anoxic mangrove soils into oxic lagoon water, it may be subject to aerobic oxidation by methane-oxidizing bacteria or to photo-oxidation (Morana et al., 2015; Smith et al., 2003), therefore limiting emissions at the water-air interface both in mangrove forests and in tidal creeks. We thus suggest that further measurements of GHG emissions must be performed in mangroves at high tide with different physiographic conditions. We also suggest including these results in future mangrove carbon budgets to make them more accurate, this ecosystem being flooded part of the time.

Our results demonstrate that CO_2 and CH_4 fluxes were highly variable with tides. As the thickness of the water column above the mangrove soil increased, CO_2 and CH_4 emissions decreased (Fig. 2a, b, c and d). The $\delta^{13}\text{CO}_2$ value of the CO_2 fluxes measured at the lowest tidal level was -26.88% (Fig. 2e), which was close to the $\delta^{13}\text{C}$ values of *Rhizophora* roots, ranging from -32 to -25% (Weiss et al., 2016), and organic matter, ranging from -30 to -20% (Bouillon et al., 2003a; Saintilan et al., 2013; Weiss et al., 2016). Oppositely, the $\delta^{13}\text{CO}_2$ value measured at the highest tidal level that was -8.6% (Fig. 2e and f), which was similar to the values of $\delta^{13}\text{CO}_2$ measured in the lagoon outside the mangrove. Consequently, we suggest that (i) the variability of the CO_2 fluxes measured resulted from a mixing between these two sources, *i.e.* the CO_2 produced within the soil and the one produced within the water column, and that (ii) emissions decreases with the increasing water column thickness resulted from the dilution of pore waters, enriched in CO_2 , by lagoon waters. We also suggest that few biogeochemical processes affected CO_2 dynamic within the water column due to (i) its low thickness that implied a low residence time of the gas, and (ii) limited photosynthetic processes resulting from dense canopy and water turbidity, as explained earlier. Regarding CH_4 , water mixing and aerobic oxidation, within the water column, as it was observed in other studies (Abril et al., 2007; Bouillon et al., 2007; Dutta and Mukhopadhyay, 2016) may partly explain the results obtained. However, measurements of both $\delta^{13}\text{CH}_4$ and dissolved oxygen are needed to comfort these hypotheses.

In addition, significant higher emissions were measured during the ebb than during the flow (two-samples *t*-test, CO_2 : $t_{(205.66)} = 5.11$, $p < 0.001$, CH_4 : $t_{(146.3)} = 7.96$, $p < 0.001$), but also during spring tides than during neap tides (CO_2 , $t_{(180.56)} = -4.60$, $p < 0.001$, and CH_4 , $t_{(134.1)} = -5.32$, $p < 0.001$) (Fig. 2a, b, c and d). We suggest that these differences may result from a combination of different factors including: residence times of the water in the soil, volume of lagoon water entering the mangrove, area of submerged mangrove, and electron acceptor renewal.

During neap tides, CO_2 emissions reached up to 9.59 $\text{mmolC m}^{-2} \text{h}^{-1}$ at the beginning of the rising tide, and up to 14.42 $\text{mmolC m}^{-2} \text{h}^{-1}$ at the end of the ebb tide (Fig. 2a and b). Meanwhile, as water thickness decreased during ebb, the $\delta^{13}\text{CO}_2$ values showed clearly an increased contribution of the CO_2 originated from mangrove soils, with more depleted values at the end of the ebb tides (Fig. 2e and f). Several forcing factors may enhance the exchanges of gaseous compounds between the soil and the overlying water, including: (i) flushing of the upper centimeters

Table 1

Various dark CO₂ (mmolC m⁻² h⁻¹) and CH₄ (μmolC m⁻² h⁻¹) emissions from mangrove forests during emersion periods, and from mangrove creeks and estuaries.

Location	Climate	CO ₂ (mmolC m ⁻² h ⁻¹)	CH ₄ (μmolC m ⁻² h ⁻¹)	Reference
Mangrove waters New Caledonia	Semi-arid	0.13 to 18.41	0.18 to 172.07	This study
Mangrove soils New Zealand	Temperate	7.02 ± 1.91	–	Bulmer et al. (2015)
New Caledonia	Semi-arid	–0.71 to 2.41	–	Leopold et al. (2015)
China	Subtropical	6.92 to 20.56	34.24 to 5168.60	Chen et al. (2010)
China	Subtropical	–0.19 to 4.62	21.56 to 1919.68	Wang et al. (2016)
Indonesia	Tropical	–1.34 to 3.88	–6.05 to 13.14	Chen et al. (2014)
Tanzania	Tropical	1.50 to 3.67	0 to 3.65	Kristensen et al. (2008b)
Mangrove creeks and estuaries Brazil	Semi-arid	0.62 ± 0.30	0.0000093	Nóbrega et al. (2016)
Australia	Subtropical	0.38 to 26.21	0.54 to 26.38	Call et al. (2015)
Florida	Subtropical	–	–2.60 to 9.61	Cabezas et al. (2017)
Australia	Subtropical	0.79 to 2.92	0.29 to 2.13	Maher et al. (2015)
India	Tropical	–	0.08 to 5.61	Biswas et al. (2007)
India	Tropical	–	0.37	Dutta et al. (2015)




Fig. 2. Emissions of CO₂ and CH₄ fluxes, and δ¹³C-CO₂ values (‰) as a function of the water level above the sediment: (a, c and e) during flow tides, and (e, d and f) during the ebb tides.

of soil driven by bottom currents inducing pressure gradients at the soil surface, concept of “skin circulation” introduced by Billerbeck et al. (2006); (ii) macrofaunal burrowing activities during inundation, causing an increase in soil permeability (Santos et al., 2012); (iii) flushing of crab burrows, the later significantly enhancing the effective surface area of mangrove soils (up to 7-fold), therefore increasing the diffusive fluxes towards the water column (Stieglitz et al., 2000, 2013; Heron and Ridd, 2008). We suggest that the increased residence time of the water during inundation resulted in enhanced exchanges across the soil-water interface due to the different processes listed above, inducing its enrichment in gaseous compounds, and explaining thus the higher emissions measured during the ebb than during the flow. Additional processes may be involved for CH₄ since the difference between the beginning of the flow and the end of the ebb was much higher than for CO₂ (Fig. 2); CH₄ concentrations was <21 μmolC m⁻² h⁻¹ during the flow and reached >170 μmolC m⁻² h⁻¹ during the ebb when the water levels were minimum (Fig. 2c and d). During low tide, atmospheric air can penetrate deeper in the soil, limiting its anoxic character and thus CH₄ production (Kristensen et al., 2008a). However, although its production can occur deeper in the soil, CH₄ may be oxidized during its transit to the atmosphere within the soil, resulting in the low emissions measured at the beginning of the flow. Conversely, at the end of the ebb, most of electron acceptors may have been exhausted due to the installation of an anoxic environment during the high tide, and thus, methanogenesis may occur, resulting in higher CH₄ emissions.

During spring tides, emissions intensity differed from neap tides. If fluxes were not significantly higher during the flow, they were during the ebb, with CO₂ and CH₄ emissions reaching up to 18.41 mmolC m⁻² h⁻¹ and 172.07 μmolC m⁻² h⁻¹, respectively (Fig. 1 and Fig. 2a, b, c and d). One possible explanation is a higher renewable of the electron acceptors pool within the soil during spring tide, which enhanced the microbial activity and consequently greenhouse gas production. Then, at low tide, the water level within the soil may be lower during spring tide, resulting in higher contribution of the soil deep layers to CH₄ emissions, as gas diffusion is higher in air than in water. Additionally, higher water level during spring tides may have flooded the upper intertidal zones within the mangrove forest, increasing therefore the exchanges between the surface pore waters and the overlying water column. The flushing of surface pore waters and burrows of this upper intertidal zone, due to changes in hydrostatic pressure along the topographic gradient during ebb, may be involved in the enhancement of the emissions, analogically to the concept of “mangrove tidal pumping”, which has been described to significantly enrich tidal creeks in nutrients, gaseous and dissolved compounds (Call et al., 2015; Dittmar and Lara, 2001; Maher et al., 2013, 2015). In addition, the more depleted δ¹³CO₂ values during ebb of spring tides, in comparison to ebb of neap tides (Fig. 2f), and except at the lowest water column thickness, support the hypothesis of a higher contribution of CO₂ produced within mangrove soils to the water column. Surprisingly, the lowest δ¹³CO₂ values were measured at the beginning of the flow (Fig. 2e), which were close to the δ¹³C values of mangrove soil organic matter or mangrove roots as explain earlier in this study, suggesting that mangrove pore waters strongly dominated the water column composition at this period of the tide, which is clearly different from neap tides (Fig. 2e). This result may be related to the tidal wave spread when entering the mangrove, inducing higher pore water advection during the spring tides notably through crab burrows. However, although higher mean CO₂ and CH₄ emissions were observed for the flow during spring tides, the maximal value reached by the emissions was not higher than during neap tides (Fig. 2a). One possible explanation is that CO₂ and CH₄ did not accumulate within the soil at low tide due to the more efficient diffusion of GHG in air than in water, and thus were rapidly emitted to the atmosphere. When the tide increased, the deep soil pore waters, depleted in δ¹³CO₂, were transported to the surface, resulting in low δ¹³CO₂

values emitted at the water surface. This study thus highlighted that tides (flow/ebb, water column thickness, neap/spring) is a key factor controlling CO₂ and CH₄ emissions from the water column during mangrove forests immersion periods. In a future research effort, the influence of other environmental parameters such as seasonal temperature variations should be studied. Although the sampling period in this study encompassed the two main seasons in New Caledonia, they were not entirely covered, and considering the high tidal variability, our data set did not allow us to conclude on a possible seasonal effect.

4. Conclusion

This preliminary study demonstrates that tide characteristics (flow/ebb, water column thickness, neap/spring) drove CO₂ and CH₄ emissions at the water-air interface within the studied *Rhizophora* spp. mangrove forest. Simultaneous measurements of CO₂, CH₄ and δ¹³CO₂ values highlighted that:

- the water column above the forest floor was a mix between soil pore waters, enriched in gaseous compounds originated from OM decomposition and root respiration, and incoming lagoon water, poor in CO₂ and CH₄ and characterized by enriched δ¹³C values, resulting in lower fluxes when the water column thickness increased,
- the progressive enrichment of the water column in CO₂ and CH₄ by diffusive fluxes, as its residence time over the forest floor increased, induced higher fluxes during the ebb tide than during the flow,
- CO₂ and CH₄ emissions were higher during spring tides than during neap tides possibly due to (i) higher renewable of the electron acceptor pool within the soil that enhanced microorganism's activity, and (ii) flooding of the upper intertidal zones that induced a higher exchange surface between the soil and the water column.

These results also showed that CO₂ and CH₄ emissions were not negligible and therefore, we suggest that these fluxes should be integrated in future carbon budget to make them more accurate. Further studies should now examine the variability of these emissions across mangroves that differ by their physiographic conditions to assess their spatial variability.

Acknowledgments

This work was funded by the Province Sud of New Caledonia, the City of Mont Dore, KNS Koniambo Nickel SAS, Vale NC, and the GOPS. We thank the Air Liquide Foundation for funding the CRDS analyzer. We are grateful to Inès Gayral for her support in field work.

Declaration of interest

Authors have no conflict of interest to declare.

References

- Abril, G., Commarieu, M.-V., Guérin, F., 2007. Enhanced methane oxidation in an estuarine turbidity maximum. *Limnol. Oceanogr.* 52:470–475. <https://doi.org/10.4319/lo.2007.52.1.0470>.
- Barnes, J., Ramesh, R., Purvaja, R., Nirmal Rajkumar, A., Senthil Kumar, B., Kritchika, K., Ravichandran, K., Uher, G., Upstill-Goddard, R., 2006. Tidal dynamics and rainfall control N₂O and CH₄ emissions from a pristine mangrove creek. *Geophys. Res. Lett.* 33 (L15405). <https://doi.org/10.1029/2006GL026829>.
- Billerbeck, M., Werner, U., Polerecky, L., Walpersdorf, E., DeBeer, D., Huettel, M., 2006. Surficial and deep pore water circulation governs spatial and temporal scales of nutrient recycling in intertidal sand flat sediment. *Mar. Ecol. Prog. Ser.* 326, 61–76.
- Biswas, H., Mukhopadhyay, S.K., Sen, S., Jana, T.K., 2007. Spatial and temporal patterns of methane dynamics in the tropical mangrove dominated estuary, NE coast of Bay of Bengal, India. *J. Mar. Syst.* 68:55–64. <https://doi.org/10.1016/j.jmarsys.2006.11.001>.
- Borges, A.V., Djenidi, D., Lacroix, G., Théate, J., Delille, B., Frankignoulle, M., 2003. Atmospheric CO₂ flux from mangrove surrounding waters. *Geophys. Res. Lett.* 30. <https://doi.org/10.1029/2003GL017143>.

- Bouillon, S., Dahdouh-Guebas, F., Rao, A., Koedam, N., Dehairs, F., 2003a. Sources of organic carbon in mangrove sediments: variability and possible ecological implications. *Hydrobiologia* 495, 33–39.
- Bouillon, S., Frankignoulle, M., Dehairs, F., Velimirov, B., Eiler, A., Abril, G., Etcheber, H., Borges, A.V., 2003b. Inorganic and organic carbon biogeochemistry in the Gautami Godavari estuary (Andhra Pradesh, India) during pre-monsoon: the local impact of extensive mangrove forests. *Glob. Biogeochem. Cycles* 17 n/a–n/a. <https://doi.org/10.1029/2002GB002026>.
- Bouillon, S., Middelburg, J.J., Dehairs, F., Borges, A.V., Abril, G., Flindt, M.R., Ulomi, S., Kristensen, E., 2007. Importance of intertidal sediment processes and porewater exchange on the water column biogeochemistry in a pristine mangrove creek (Ras Dege, Tanzania). *Biogeoosci. Discuss.* 4, 317–348.
- Bouillon, S., Connolly, R.M., Lee, S.Y., 2008. Organic matter exchange and cycling in mangrove ecosystems: recent insights from stable isotope studies. *J. Sea Res.* 59:44–58. <https://doi.org/10.1016/j.seares.2007.05.001>.
- Bulmer, R.H., Lundquist, C.J., Schwendenmann, L., 2015. Sediment properties and CO₂ efflux from intact and cleared temperate mangrove forests. *Biogeosciences* 12: 6169–6180. <https://doi.org/10.5194/bg-12-6169-2015>.
- Cabezas, A., Mitsch, W.J., MacDonnell, C., Zhang, L., Bydątek, F., Lasso, A., 2017. Methane emissions from mangrove soils in hydrologically disturbed and reference mangrove tidal creeks in southwest Florida. *Ecol. Eng.* <https://doi.org/10.1016/j.ecoleng.2017.08.041>.
- Call, M., Maher, D.T., Santos, I.R., Ruiz-Halpern, S., Mangion, P., Sanders, C.J., Erler, D.V., Oakes, J.M., Rosentreter, J., Murray, R., Eyre, B.D., 2015. Spatial and temporal variability of carbon dioxide and methane fluxes over semi-diurnal and spring-neap-spring timescales in a mangrove creek. *Geochim. Cosmochim. Acta* 150:211–225. <https://doi.org/10.1016/j.gca.2014.11.023>.
- Chanda, A., Akhand, A., Manna, S., Dutta, S., Das, I., Hazra, S., Rao, K.H., Dadhwal, V.K., 2013. Measuring daytime CO₂ fluxes from the inter-tidal mangrove soils of Indian Sundarbans. *Environ. Earth Sci.* 72:417–427. <https://doi.org/10.1007/s12665-013-2962-2>.
- Chauhan, R., Datta, A., Ramanathan, A., Adhya, T.K., 2015. Factors influencing spatio-temporal variation of methane and nitrous oxide emission from a tropical mangrove of eastern coast of India. *Atmos. Environ.* 107:95–106. <https://doi.org/10.1016/j.atmosenv.2015.02.006>.
- Chen, G.C., Tam, N.F.Y., Ye, Y., 2010. Summer fluxes of atmospheric greenhouse gases N₂O, CH₄ and CO₂ from mangrove soil in South China. *Sci. Total Environ.* 408:2761–2767. <https://doi.org/10.1016/j.scitotenv.2010.03.007>.
- Chen, G.C., Ulumuddin, Y.I., Pramudji, S., Chen, S.Y., Chen, B., Ye, Y., Ou, D.Y., Ma, Z.Y., Huang, H., Wang, J.K., 2014. Rich soil carbon and nitrogen but low atmospheric greenhouse gas fluxes from North Sulawesian mangrove swamps in Indonesia. *Sci. Total Environ.* 487:91–96. <https://doi.org/10.1016/j.scitotenv.2014.03.140>.
- Chen, G., Chen, B., Yu, D., Tam, N.F.Y., Ye, Y., Chen, S., 2016. Soil greenhouse gas emissions reduce the contribution of mangrove plants to the atmospheric cooling effect. *Environ. Res. Lett.* 11, 124019. <https://doi.org/10.1088/1748-9326/11/12/124019>.
- Chuang, P.-C., Young, M.B., Dale, A.W., Miller, L.G., Herrera-Silveira, J.A., Paytan, A., 2017. Methane fluxes from tropical coastal lagoons surrounded by mangroves, Yucatán, Mexico: methane fluxes from coastal lagoons. *J. Geophys. Res. Biogeosci.* 122: 1156–1174. <https://doi.org/10.1002/2017JG003761>.
- Dittmar, T., Lara, R.J., 2001. Molecular evidence for lignin degradation in sulfate-reducing mangrove sediments (Amazonia, Brazil). *Geochim. Cosmochim. Acta* 65, 1417–1428.
- Donato, D.C., Kauffman, J.B., Mordiyarso, D., Kurnianto, S., Stidham, M., Kanninen, M., 2011. Mangroves among the most carbon-rich forests in the tropics. *Nat. Geosci.* 4: 293–297. <https://doi.org/10.1038/ngeo1123>.
- Dumas, P., Léopold, M., Frotté, L., Peignon, C., 2012. Mud crab ecology encourages site-specific approaches to fishery management. *J. Sea Res.* 67:1–9. <https://doi.org/10.1016/j.seares.2011.08.003>.
- Dutta, M. Kumar, Mukhopadhyay, S.K., 2016. Reviews and syntheses: methane biogeochemistry in Sundarbans mangrove ecosystem, NE coast of India; a box modeling approach. *Biogeoosci. Discuss.* <https://doi.org/10.5194/bg-2016-58>.
- Dutta, M.K., Mukherjee, R., Jana, T.K., Mukhopadhyay, S.K., 2015. Biogeochemical dynamics of exogenous methane in an estuary associated to a mangrove biosphere; the Sundarbans, NE coast of India. *Mar. Chem.* 170:1–10. <https://doi.org/10.1016/j.marchem.2014.12.006>.
- Gonneea, M.E., Paytan, A., Herrera-Silveira, J.A., 2004. Tracing organic matter sources and carbon burial in mangrove sediments over the past 160 years. *Estuar. Coast. Shelf Sci.* 61:211–227. <https://doi.org/10.1016/j.ecss.2004.04.015>.
- Graham, M.C., Eaves, M.A., Farmer, J.G., Dobson, J., Fallick, A.E., 2001. A Study of Carbon and Nitrogen Stable Isotope and Elemental Ratios as Potential Indicators of Source and Fate of Organic Matter in Sediments of the Forth Estuary. *Scotland. Estuar. Coast. Shelf Sci.* 52:375–380. <https://doi.org/10.1006/ecss.2000.0742>.
- Heron, S.F., Ridd, P.V., 2008. The tidal flushing of multiple-loop animal burrows. *Estuar. Coast. Shelf Sci.* 78:135–144. <https://doi.org/10.1016/j.ecss.2007.11.018>.
- Howard, J., Sutton-Grier, A., Herr, D., Kleypas, J., Landis, E., Mcleod, E., Pidgeon, E., Simpson, S., 2017. Clarifying the role of coastal and marine systems in climate mitigation. *Front. Ecol. Environ.* 15:42–50. <https://doi.org/10.1002/fee.1451>.
- Kauffman, J.B., Heider, C., Cole, T.G., Dwire, K.A., Donato, D.C., 2011. Ecosystem carbon stocks of Miocenesian mangrove forests. *Wetlands* 31:343–352. <https://doi.org/10.1007/s13157-011-0148-9>.
- Keeling, C.D., 1958. The concentration and isotopic abundances of atmospheric carbon dioxide in rural areas. *Geochim. Cosmochim. Acta* 13, 322–334.
- Keeling, C.D., 1961. The concentration and isotopic abundances of carbon dioxide in rural and marine air. *Geochim. Cosmochim. Acta* 24:277–298. [https://doi.org/10.1016/0016-7037\(61\)90023-0](https://doi.org/10.1016/0016-7037(61)90023-0).
- Komiya, S., Noborio, K., Katano, K., Pakoktom, T., Siangliw, M., Toojinda, T., 2015. Contribution of ebullition to methane and carbon dioxide emission from water between plant rows in a tropical rice paddy field. *Int. Sch. Res. Not.* 2015. <https://doi.org/10.1155/2015/623901>.
- Kremer, J.N., Nixon, S.W., Buckley, B., Roques, P., 2003. Technical note: conditions for using the floating chamber method to estimate air-water gas exchange. *Estuaries* 26, 985–990.
- Kristensen, E., Bouillon, S., Dittmar, T., Marchand, C., 2008a. Organic carbon dynamics in mangrove ecosystems: a review. *Aquat. Bot.* 89:201–219. <https://doi.org/10.1016/j.aquabot.2007.12.005>.
- Kristensen, E., Flindt, M.R., Ulomi, S., Borges, A., Abril, G., Bouillon, S., 2008b. Emission of CO₂ and CH₄ to the atmosphere by sediments and open waters in two Tanzanian mangrove forests. *Mar. Ecol. Prog. Ser.* <https://doi.org/10.3354/meps07642>.
- Lamb, A.L., Wilson, G.P., Leng, M.J., 2006. A review of coastal palaeoclimate and relative sea-level reconstructions using $\delta^{13}\text{C}$ and C/N ratios in organic material. *Earth-Sci. Rev.* 75:29–57. <https://doi.org/10.1016/j.earscirev.2005.10.003>.
- Leopold, A., Marchand, C., Deborde, J., Chaduteau, C., Allenbach, M., 2013. Influence of mangrove zonation on CO₂ fluxes at the sediment-air interface (New Caledonia). *Geoderma* 202, 62–70.
- Leopold, A., Marchand, C., Deborde, J., Allenbach, M., 2015. Temporal variability of CO₂ fluxes at the sediment-air interface in mangroves (New Caledonia). *Sci. Total Environ.* 502, 617–626.
- Lin, G., Ehleringer, J.R., 1997. Carbon Isotopic Fractionation Does Not Occur during Dark Respiration in C₃ and C₄ Plants. *Plant Physiol.* 114:391–394. <https://doi.org/10.1104/pp.114.1.391>.
- Livesley, S.J., Andrusiak, S.M., 2012. Temperate mangrove and salt marsh sediments are a small methane and nitrous oxide source but important carbon store. *Estuar. Coast. Shelf Sci.* 97:19–27. <https://doi.org/10.1016/j.ecss.2011.11.002>.
- Lorke, A., Bodmer, P., Noss, C., Alshboul, Z., Koschorreck, M., Somlai-Haase, C., Bastviken, D., Flury, S., McGinnis, D.F., Maec, A., Müller, D., Premke, K., 2015. Technical Note: Drifting Versus Anchored Flux Chambers for Measuring Greenhouse Gas Emissions from Running Waters. <https://doi.org/10.5194/cg-12-7013-2015>.
- Lu, L., Cheng, H., Pu, X., Wang, J., Cheng, Q., Liu, X., 2016. Identifying organic matter sources using isotopic ratios in a watershed impacted by intensive agricultural activities in Northeast China. *Agric. Ecosyst. Environ.* 222:48–59. <https://doi.org/10.1016/j.agee.2015.12.033>.
- Lyimo, T.J., Pol, A., Jetten, M.S., Op den Camp, H.J., 2009. Diversity of methanogenic archaea in a mangrove sediment and isolation of a new Methanococoides strain. *FEMS Microbiol. Lett.* 291, 247–253.
- Maher, D.T., Santos, I.R., Golsby-Smith, L., Gleeson, J., Eyre, B.D., 2013. Groundwater-derived dissolved inorganic and organic carbon exports from a mangrove tidal creek: the missing mangrove carbon sink? *Limnol. Oceanogr.* 58:475–488. <https://doi.org/10.4319/lo.2013.58.2.0475>.
- Maher, D.T., Cowley, K., Santos, I.R., Macklin, P., Eyre, B.D., 2015. Methane and carbon dioxide dynamics in a subtropical estuary over a diel cycle: insights from automated in situ radioactive and stable isotope measurements. *Mar. Chem.* 168:69–79. <https://doi.org/10.1016/j.marchem.2014.10.017>.
- McLeod, E., Chmura, G.L., Bouillon, S., Salm, R., Björk, M., Duarte, C.M., Lovelock, C.E., Schlesinger, W.H., Silliman, B.R., 2011. A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO₂. *Front. Ecol. Environ.* 9:552–560. <https://doi.org/10.1890/110004>.
- Midwood, A.J., Millard, P., 2011. Challenges in measuring the $\delta^{13}\text{C}$ of the soil surface CO₂ efflux. *Rapid Commun. Mass Spectrom.* 25:232–242. <https://doi.org/10.1002/rcm.4857>.
- Millard, P., Midwood, A.J., Hunt, J.E., Barbour, M.M., Whitehead, D., 2010. Quantifying the contribution of soil organic matter turnover to forest soil respiration, using natural abundance $\delta^{13}\text{C}$. *Soil Biol. Biochem.* 42:935–943. <https://doi.org/10.1016/j.soilbio.2010.02.010>.
- Morana, C., Borges, A.V., Roland, F.A.E., Darchambeau, F., Descy, J.-P., Bouillon, S., 2015. Methanotrophy within the water column of a large meromictic tropical lake (Lake Kivu, East Africa). *Biogeosciences* 12:2077–2088. <https://doi.org/10.5194/bg-12-2077-2015>.
- Nielsen, O.I., Kristensen, E., Macintosh, D.J., 2003. Impact of fiddler crabs (*Uca* spp.) on rates and pathways of benthic mineralization in deposited mangrove shrimp pond waste. *J. Exp. Mar. Biol. Ecol.* 289:59–81. [https://doi.org/10.1016/S0022-0981\(03\)00041-8](https://doi.org/10.1016/S0022-0981(03)00041-8).
- Nóbrega, G.N., Ferreira, T.O., Siqueira Neto, M., Queiroz, H.M., Artur, A.G., Mendonça, E.D.S., Silva, E.D.O., Otero, X.L., 2016. Edaphic factors controlling summer (rainy season) greenhouse gas emissions (CO₂ and CH₄) from semiarid mangrove soils (NE-Brazil). *Sci. Total Environ.* 542:685–693. <https://doi.org/10.1016/j.scitotenv.2015.10.108>.
- Pataki, D.E., Ehleringer, J.R., Flanagan, L.B., Yakir, D., Bowling, D.R., Still, C.J., Buchmann, N., Kaplan, J.O., Berry, J.A., 2003. The application and interpretation of Keeling plots in terrestrial carbon cycle research. *Glob. Biogeochem. Cycles* 17.
- Paterson, E., Midwood, A.J., Millard, P., 2009. Through the eye of the needle: a review of isotope approaches to quantify microbial processes mediating soil carbon balance. *Tansley review. New Phytol.* 184:19–33. <https://doi.org/10.1111/j.1469-8137.2009.03001.x>.
- R Development Core Team, 2008. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Saintilan, N., Rogers, K., Mazumder, D., Woodroffe, C., 2013. Allochthonous and autochthonous contributions to carbon accumulation and carbon store in southeastern Australian coastal wetlands. *Estuar. Coast. Shelf Sci.* 128:84–92. <https://doi.org/10.1016/j.ecss.2013.05.010>.
- Santos, I.R., Eyre, B.D., Huettel, M., 2012. The driving forces of porewater and groundwater flow in permeable coastal sediments: a review. *Estuar. Coast. Shelf Sci.* 98:1–15. <https://doi.org/10.1016/j.ecss.2011.10.024>.
- Smith, K.A., Ball, T., Conen, F., Dobbie, K.E., Massheder, J., Rey, A., 2003. Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and

- biological processes. *Eur. J. Soil Sci.* 54:779–791. <https://doi.org/10.1046/j.1351-0754.2003.0567.x>.
- Stamp, I., Baird, A.J., Heppell, C.M., 2013. The importance of ebullition as a mechanism of methane (CH₄) loss to the atmosphere in a northern peatland. *Geophys. Res. Lett.* 40: 2087–2090. <https://doi.org/10.1002/grl.50501>.
- Stieglitz, T., Ridd, P., Müller, P., 2000. Passive irrigation and functional morphology of crustacean burrows in a tropical mangrove swamp. *Hydrobiologia* 421:69–76. <https://doi.org/10.1023/A:1003925502665>.
- Stieglitz, T.C., Clark, J.F., Hancock, G.J., 2013. The mangrove pump: the tidal flushing of animal burrows in a tropical mangrove forest determined from radionuclide budgets. *Geochim. Cosmochim. Acta* 102:12–22. <https://doi.org/10.1016/j.gca.2012.10.033>.
- Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, B., Midgley, B.M., 2013. *IPCC, 2013: climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge University Press.
- Vachon, D., Prairie, Y.T., Cole, J.J., 2010. The relationship between near-surface turbulence and gas transfer velocity in freshwater systems and its implications for floating chamber measurements of gas exchange. *Limnol. Oceanogr.* 55:1723–1732. <https://doi.org/10.4319/lo.2010.55.4.1723>.
- Wang, H., Liao, G., D'Souza, M., Yu, X., Yang, J., Yang, X., Zheng, T., 2016. Temporal and spatial variations of greenhouse gas fluxes from a tidal mangrove wetland in Southeast China. *Environ. Sci. Pollut. Res.* 23:1873–1885. <https://doi.org/10.1007/s11356-015-5440-4>.
- Weiss, C., Weiss, J., Boy, J., Iskandar, I., Mikutta, R., Guggenberger, G., 2016. Soil organic carbon stocks in estuarine and marine mangrove ecosystems are driven by nutrient colimitation of P and N. *Ecol. Evol.* 6:5043–5056. <https://doi.org/10.1002/ece3.2258>.
- Yamamuro, M., 2000. Chemical tracers of sediment organic matter origins in two coastal lagoons. *J. Mar. Syst.* 26:127–134. [https://doi.org/10.1016/S0924-7963\(00\)00049-X](https://doi.org/10.1016/S0924-7963(00)00049-X).
- Yang, L., Lu, F., Zhou, X., Wang, X., Duan, X., Sun, B., 2014. Progress in the studies on the greenhouse gas emissions from reservoirs. *Acta Ecol. Sin.* 34:204–212. <https://doi.org/10.1016/j.chnaes.2013.05.011>.