



**HAL**  
open science

# Challenging exploration of troubled waters: a decade of surveys of the giant freshwater pearl mussel *Margaritifera auricularia* in Europe

Vincent Prié, Joaquin Soler, Rafael Araújo, Xavier Cucherat, Laurent Philippe, Nicolas Patry, Benjamin Adam, Nicolas Legrand, Philippe Jugé, Nina Richard, et al.

## ► To cite this version:

Vincent Prié, Joaquin Soler, Rafael Araújo, Xavier Cucherat, Laurent Philippe, et al.. Challenging exploration of troubled waters: a decade of surveys of the giant freshwater pearl mussel *Margaritifera auricularia* in Europe. *Hydrobiologia*, 2018, 810 (1), pp.157-175. 10.1007/s10750-017-3456-0 . hal-01826265

**HAL Id: hal-01826265**

<https://hal.sorbonne-universite.fr/hal-01826265v1>

Submitted on 29 Jun 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Challenging exploration of troubled waters: ten years' surveys of the giant freshwater pearl  
2 mussel *Margaritifera auricularia* in Europe

3

4 Vincent Prié<sup>1, 2</sup>

5 Joaquin Soler<sup>3</sup>

6 Rafael Araujo<sup>4</sup>

7 Xavier Cucherat<sup>5</sup>

8 Laurent Philippe<sup>1</sup>

9 Nicolas Patry<sup>1</sup>

10 Benjamin Adam<sup>1</sup>

11 Nicolas Legrand<sup>1</sup>

12 Philippe Jugé<sup>6</sup>

13 Nina Richard<sup>6</sup>

14 Karl M. Wantzen<sup>3, 7</sup>

15

16

17 <sup>1</sup> Biotope, SIDI, 22 Bd Maréchal Foch, 34140 Mèze, France

18 <sup>2</sup> Muséum national d'Histoire naturelle, Département Systématique et Evolution, ISyEB (UMR 7205  
19 CNRS/UPMC/MNHN/EPHE), 43, Rue Cuvier, 75231 Paris, France.

20 <sup>3</sup> Université François Rabelais, UMR 7324 – CITERES, 33 Allée Ferdinand de Lesseps, 37204 Tours cedex 03,  
21 France

22 <sup>4</sup> Museo Nacional de Ciencias Naturales - C.S.I.C. c/ José Gutiérrez Abascal 2, 28006 Madrid, Spain

23 <sup>5</sup> 10 rue Louis Aragon, 59147 Gondecourt, France

24 <sup>6</sup> Université François-Rabelais, CETU Elmis Ingénieries, 11 Quai Danton, 37500 Chinon, France

25 <sup>7</sup> UNESCO River Culture – Fleuves et Patrimoines Chair, Université François Rabelais, UMR 7324 – CITERES,  
26 33 Allée Ferdinand de Lesseps, 37204 Tours cedex 03, France

27

28 Corresponding author:

29 Vincent Prié; [vprie@biotope.fr](mailto:vprie@biotope.fr); +33 7 71 08 93 46

30

31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

Acknowledgments:

This work was conducted within the scope of the LIFE project “Life13BIOFR001162 Conservation of the Giant Pearl Mussel in Europe”. We thank Dominique Tesseyre from the Adour-Garonne Water Agency; Julie Marcinkowsky and Gérard Tardivo from the DREAL Centre as well as the DREAL Picardie for providing financial support for large-scale surveys of *M. auricularia* in France; Elodie Hugues, Guillaume Métayer (Conseil Général de Charente Maritime), David Bécart (Voies Navigables de France) and Amandine Szurpicki (COSEA), Frédérique Moinot and Olivier Guerri (EPIDOR) for financing focus surveys in the Charente, Seine, Vienne and Garonne Rivers; and for Spain FMC Forest S.A., Enagas, Gas Natural, INYPSA, Hidroeléctrica La Zaida, Edison Mission Energy and EID Consultores.

Abstract

The critically endangered Giant Freshwater Pearl Mussel *Margaritifera auricularia* was presumed extinct before its re-discovery in Spain in 1985 and France in 2000. Since then, numerous surveys have been set up to search for living populations in France and Spain. This article presents an up-to-date distribution of the species based on available data, i.e. literature, Museum collections and recent field surveys; and provides unpublished molecular data for France. The Giant Freshwater Pearl Mussel is still living as three populations in the Ebro River in Spain, and eight populations in France (two in the Loire drainage, one in the Charente drainage, two in the Garonne drainage and three in the Adour drainage). The biggest population lives in the Charente River with an estimated 100.000 individuals. Recruitment is very scarce in all populations but living specimens estimated to be less than 10 years old have been found in the Ebro in Spain and in the Vienne, Charente, Dronne and Adour rivers in France. Recent populations rediscovery in France were mainly a result of intensive dedicated surveys including scuba-diving. Subsequent advances in knowledge show how large rivers and downstream ecosystems remain a *terra incognita* for the hydrobiologist.

Distribution; museum collections; historical data; scuba diving surveys; large rivers; conservation

60 Introduction

61

62 Freshwater ecosystems are the most threatened ecosystems worldwide (Dudgeon et al. 2006) and freshwater  
63 bivalves rank amongst the most threatened animals in the world (Lydear et al. 2004; Lopes-Lima et al. 2016).  
64 One of them, the Giant Freshwater Pearl Mussel *Margaritifera auricularia* (Spengler, 1793), ranges amongst the  
65 most imperilled bivalve species. Although it was considered widespread in most of the western Europe rivers at  
66 the beginning of the 20th century, it is now considered as critically endangered by the IUCN (Araujo & Ramos,  
67 2001; Prié 2010). The Giant Freshwater Pearl Mussel had become so rare during the 20<sup>th</sup> century that it was not  
68 even considered when the European Habitat Directive species lists have been established. Indeed, the Giant  
69 Freshwater Pearl Mussel is difficult to observe: it lives downstream in large rivers, a habitat that is difficult to  
70 survey due to deepness, turbidity, current and often navigation. Hence, not surprisingly, it has been overlooked  
71 by malacologists of the 20<sup>th</sup> century. However, it nowadays still survives as a few populations in south-west  
72 France and eastern Spain.

73 The species was first rediscovered in Spain in 1985 (Altaba, 1990) and in France in 2000 (Cochet, 2001). Since  
74 1998, the biology, distribution and lifecycle of the Giant Freshwater Pearl Mussel in Spain were described  
75 (Araujo & Ramos, 1998a, b; Araujo & Ramos, 2000 a, b; Araujo & Ramos, 2000; Grande *et al.*, 2001; Araujo *et*  
76 *al.*, 2001; 2002; 2003; Gómez & Araujo, 2008). Since then, some few news of the species in Spain have been  
77 released in national and international congresses (i. e. Nakamura et al., 2015; Online Resource 1), but, apart from  
78 Araujo & Álvarez-Cobelas (2016) there are no new scientific results published since 2008. In France, focused  
79 surveys have led to the rediscovery of many populations since 2007, but most of these results are unpublished  
80 (but see Prié et al., 2007; Prié et al., 2008; Prié et al., 2010) or available only as grey literature (Online Resource  
81 1).

82 An extensive review of all available data on *Margaritifera auricularia*'s distribution is provided here for the first  
83 time, together with new data from museum collections and recent field surveys. This article clarifies the past and  
84 present distribution of the species, presents the results of the last ten years' surveys in France and Spain and  
85 discusses conservation perspectives.

86

87 Material and Methods

88

89 Bibliography review

90 The bibliography since 1793 (species description date) has been extensively reviewed. Local publication and  
91 grey literature were also consulted when available. Bibliographic data was generally imprecise, but allowed  
92 figuring a broad image of the original distribution and ecology of *M. auricularia* (Fig. 1). Bibliography review  
93 thus provided the first indications for where to look for this species.

94

95 Museum collections

96 A first review of museum collections had been performed by Araujo & Ramos (2000a) at a global scale. This  
97 review mostly aimed at large national museum collections and included also *Margaritifera marocana* (Pallary,  
98 1918), a species living only in Morocco (Araujo et al., 2009a). We then inventoried all the regional museum and  
99 Universities collections in France. Fifty-eight local natural history collections were identified. Each of them was  
100 contacted and questioned about the presence of malacological collections, freshwater bivalves and eventually *M.*  
101 *auricularia* specimens. When *M. auricularia* specimens were recorded in the inventories or discovered in the  
102 collection by the curator, pictures were sent to us to confirm identification. Eventually, some of the most  
103 important collections (Musée des Confluences in Lyon, Museum d'Histoire Naturelle in Bordeaux, Museum  
104 d'Histoire Naturelle in Toulouse, Museum national d'Histoire naturelle in Paris, Museum d'Histoire Naturelle in  
105 Lille, Museum d'Histoire Naturelle in Nantes, Museum d'Histoire naturelle in Orleans, University of Rennes,  
106 University of Montpellier) were visited by one of us.

107 Specimens collected since 2000, year of the re-discovery of the species in France, were not included in the  
108 results presented here.

109

110 Field surveys and population sizes

111

112 Numerous field surveys aiming at freshwater mussels have been performed in France and Spain (Fig. 1, Table  
113 1).

114 These dedicated surveys aimed at places most likely to host the species, i.e. places identified by literature data,  
115 museum collection data or, for France, species habitats modelling (Prié et al., 2014). Moreover, some surveys  
116 took place into the frame of impact studies. These impact studies were triggered when *M. auricularia* was living

117 - or when available data suggested that it could still be living - in an area impacted by a development project.  
118 The results of these impact studies are generally not published, consisting only in various cryptic reports (but see  
119 Prié et al., 2007; Prié et al., 2008; Araujo & Alvarez-Cobelas, 2016). We here summarize for the first time all the  
120 grey literature related to *M. auricularia* in France and Spain (Online Resource 1).

121 *M. auricularia* mainly lives in downstream ecosystems. Surveying this habitat is challenging because it is often  
122 deep, turbid, strongly flowing and navigable. In the Ebro historic channels, sampling depends on the hydraulic  
123 works made by the Confederación Hidrográfica del Ebro; it is necessary to decrease the water level in order to  
124 wade the channel bottom to find the specimens (Gómez & Araujo, 2008). In France, some populations are  
125 readily accessible, living in the banks (Vienne River) or in shallow waters (Creuse, Luy or Arros River). For  
126 those populations, snorkelling or wading with viewing glasses allowed efficient surveys. However, cumbersome  
127 methods based on a team of scuba-divers were needed in most cases. For some surveys, a boat was used to  
128 shuttle the divers from a place to another. For others, divers dove from the river banks and sampling plans were  
129 then constrained by river accessibility.

130 Population sizes given here were estimated based on exhaustive counts of observed living individuals (Luy,  
131 Creuse and Vienne Rivers); statistical analyses (Ebro, Arros and Charente Rivers), or in the worst case, by a  
132 subjective appreciation based on the density of specimens observed (Dronne, Adour and Save Rivers).

133 The Seine (downstream) and Eure Rivers could only be surveyed by dredging. The dredger used had an aperture  
134 of 50 cm, a 25-mm mesh, weighted 11 kg and was propelled by a 30-horsepower engine Zodiac by means of a  
135 30 m long rope. In the Eure River, different biotopes and flow facies were aimed at (mud, sand, stones, riffles,  
136 vegetation). In the Seine River, water was up to 6 meters deep and too troubled for operators to see the river bed.  
137 Catches were then randomly positioned. Catches were 8 to 10 m long in the Eure River, and up to 40-50 m long  
138 in the Seine River. Sediment collected by the dredger was pulled up and sorted out on the boat. Wading surveys  
139 were adopted upstream the Seine River.

140 In the Somme River, a boat was used to shuttle divers and 82 bank to bank transects were sampled on a 26 km  
141 long river stretch. In the Oise River, the divers were also transported by boat from a spot to another, but diving  
142 plans were constrained by river condition (from very strong current to muddy bottoms). Areas with very strong  
143 current were sampled combining scuba-diving and climbing technics, with a 100 m long static rope secured on a  
144 tree on the bank. The diver used a climbing harness and caving equipment in addition to scuba diving gear to  
145 progress on the rope. Fins were used to go from side to side in the current, allowing to cover a ca.90 m long  
146 cone-shaped surface on the river bottom. Altogether, 115 dives have been carried out on a 35 km long stretch of

147 the river, from the confluence with the Aisne River downstream to a few kilometres upstream the town of  
148 Sempigny. Upstream this stretch of river, surveys were carried out wading randomly in suitable habitats.

149 In the Charente River, the population was estimated based on scuba-diving transects surveys. A boat was used to  
150 shuttle the divers from a transect to another. A 20 m long line was settled down on the bottom of the river and  
151 scuba divers counted every living specimen left and right of the line at a distance of 2 m. Each sample then  
152 covered 80 m<sup>2</sup>. Transects were repeated every kilometer in the river stretch where mussels were present, and  
153 then every three kilometers downstream and upstream the population's distribution limits. A total of 43 transects  
154 were repeated on a stretch of 60 km. Detection probability has been estimated at 75% using iterated observations  
155 analysed with the software MARK (White & Burnham, 1999). Geographical statistics (Anselin, 1996) were  
156 performed using GeoDa software (Anselin et al., 2006). Suitable habitat length in the whole river was delimited  
157 downstream by the limit of the mud cover due to the influence of the Saint-Savinien's impoundment, upstream  
158 by the limit of the living population. Between these limits, the substrate and general ecological quality of the  
159 river was very homogenous. In this stretch of favourable habitat, live specimens have been observed wherever  
160 we have dived between 2010 and 2016, thus confirming that the population is uniformly distributed.

161 Fourteen sampling surveys were undertaken between September 2000 and June 2006 in the Ebro River,  
162 totalising 25 km, wading in shallow waters and with a team of divers in the deeper parts of the river. Divers used  
163 submerged ropes to perform bank to bank or longitudinal transects (survey methods were detailed and reported  
164 in Gómez & Araujo, 2008; Araujo & Álvarez-Cobelas, 2016).

165 In the Dronne and Isle Rivers, about 100 km stretch of each river upstream their confluence was surveyed, both  
166 by wading and scuba-diving from the banks. The estimation of the population size was based on author's  
167 appreciation only, and is likely underestimated: over 50 specimens have been observed during the surveys, with  
168 a subpopulation of 30 specimens in the lower location (exhaustive count). We estimate that about half of the  
169 living individuals have been observed during surveys, which is unlikely given the detection probability in this  
170 large river.

171 About 60 km of the Save River was surveyed by wading and scuba-diving, aiming at an exhaustive count of the  
172 few remaining specimens which were found only in the lower section of the river. Most of the sampling in the  
173 Adour River was undertaken by wading and snorkelling, with scuba divers requested only for a few deeper  
174 places. As for the Dronne River, few specimens were found in isolated places, with biggest subpopulation  
175 numbering about ten specimens. Population size is estimated based on experts' appreciation only. The Arros  
176 River is highly impacted by agriculture practices. The remaining favourable habitats were found isolated

177 between the numerous impoundments' influences. A first survey was conducted by scuba divers, but the deepest  
178 places did not have suitable habitats. A more intensive survey was then organized by a team wading with  
179 viewing glasses. The total length of river stretches having suitable habitats was 54 km. Within these 54 km,  
180 sixteen sites were sampled. On each sampled site, stretches of 100 m to 1 km were exhaustively surveyed.  
181 Population size was estimated based on average densities observed during surveys, multiplied by favourable  
182 habitat's surface. In the Luy River, divers explored the deepest pools while most of the river can be explored by  
183 wading. The main population is found in a very shallow place, and exhaustive counts were performed three  
184 times (years 2010, 2011, 2012) by five persons wading in a line, about one meter apart, ensuring efficient  
185 scanning of every single place of the river bed. However, detection probability is never 100%. Some specimens  
186 may spend some time completely buried in the sediment and are overlooked (see below the results for the Luy  
187 River). The results of these assumed exhaustive counts are therefore likely underestimated.

188 The most intensive surveys took place in the Vienne and Creuse Rivers. The surveys aimed at providing  
189 exhaustive counts of all living specimens. Observers with viewing glasses and divers (depending on the depth)  
190 were lined one meter apart and moved forward upstream, ensuring efficient scanning of every single place of the  
191 river bed. Sampling was reiterated several time between 2009 and 2016 using the same methods.

192 In this study, when shells only have been collected, we considered "ancient shells" those that were worn and  
193 uncomplete, without periostracum nor ligament remains. "Recent shells" include shells with at least  
194 periostracum and ligament remains. We consider as "juveniles" specimens with shell length lower than 11 cm,  
195 "subadults" specimens from 11 to 14 cm. Occasionally, some adult specimens had very short shells, especially in  
196 the Charente River, but these were obviously very old given the growth lines density and shell wear.

197

#### 198 Genetic analyses

199 Tissue samples have been collected from ten specimens from the Ebro River in Spain, and ten specimens from  
200 the Vienne River (Loire coastal drainage), two specimens from the Luy River (Adour River coastal drainage),  
201 two specimens from the Charente River and one specimen from the Save River (Garonne River coastal drainage)  
202 in France. Foot tissue samples were snipped in the field and preserved in 90° ethanol for molecular analysis.

203 For Spanish specimens, DNA was extracted using CTAB protocol: tissue samples, preserved in ethanol or  
204 frozen, were ground to a powder in liquid nitrogen before adding 600  $\mu$ L of CTAB lysis buffer (2% CTAB, 1.4  
205 M NaCl, 0.2%  $\beta$ -mercaptoethanol, 20 mM EDTA, 0.1 M TRIS [pH = 8]) and subsequently digested with  
206 proteinase K (100  $\text{mg}\cdot\text{ml}^{-1}$ ) for 2–5 h at 60° C. Total DNA was extracted according to standard



207 phenol/chloroform procedures (Sambrook & Maniatis, 1989). For French specimen DNA was extracted using  
208 the Nucleospin Tissue Kit (marketed by Macherey–Nagel), following the manufacturer’s protocol. Extractions,  
209 amplifications and sequencing were performed by Genoscreen (France).

210 To test genetic variability between populations, we examined fragments of two mitochondrial genes, COI and  
211 16S, used previously by Huff et al. (2004); these showed the greatest phylogenetic resolution power for  
212 relationships among margaritiferids. 28S nuclear gene fragments were also amplified, but different fragments  
213 were targeted for French and Spanish specimens. The COI, 16S and 28S gene were amplified by polymerase  
214 chain reaction (PCR) using the protocol described by Prié & Puillandre, 2014 for French specimens, and  
215 described by Machordom et al. (2003) and Araujo et al. (2016) for Spanish specimens. The amplified fragments  
216 were purified by ethanol precipitation prior to sequencing both strands using BigDye Terminator kits (Applied  
217 Biosystems, ABI). Products were electrophoresed on an ABI 3730 genetic Analyser (Applied Biosystems). The  
218 forward and reverse DNA sequences obtained for each specimen were aligned and checked using the Sequencer  
219 program (Gene Code Corporation) after removing primer regions. Sequences were automatically aligned using  
220 ClustalW multiple alignments implemented in BioEdit 7.0.5.3 (Hall, 1999). The accuracy of automatic  
221 alignments was confirmed by eye. Genebank accession numbers are provided in Table 2.

222

223 Results

224 Bibliography

225 Available literature provided valuable data, although generally without precise location nor date. Nevertheless, a  
226 first historical distribution map could be drawn from ancient literature data. *Margaritifera auricularia* is known  
227 from the Netherland, England and Germany from fossil records only. However, some shells collected in the  
228 Unstrut River in Germany are very well preserved and probably date back to historical times, at least until the  
229 early Middle Ages (Bössneck et al., 2006). Fossil data in Spain includes a Mediterranean Quaternary river in  
230 Yecla (Murcia) with 129.000-140.000 years old specimens (Andrés & Ortuño, 2014) and many other Atlantic  
231 rivers with 5.000 years old specimens (Araujo & Moreno, 1999). In France, fossil data near Marseille (coming  
232 from archaeological excavation) and in Massif Central (found amongst fossils collected in a cave) were  
233 presumably a result of human transportation.

234 According to historical data collected, *Margaritifera auricularia* was only found in large rivers, in a calcareous  
235 substrate, in France, Spain and Italy. In France, historical data mainly comes from the Atlantic and Channel sea  
236 coastal drainages, with only one occurrence in the Mediterranean coastal drainages, in the Saône River (Rhône

237 tributary). In Italy and Spain, the species is historically known from two Mediterranean coastal drainages, the Po  
238 and Ebro Rivers (Araujo & Ramos, 2000a). In Spain *M. auricularia* lived in two historic channels from the Ebro  
239 River, the Canal Imperial and the Canal de Tauste, where there were about 5000 live specimens. The more  
240 recent data published about these Spanish populations were recorded in Araujo & Ramos (2000b), Gómez &  
241 Araujo (2008) and Araujo & Álvarez-Cobelas (2016).

242

#### 243 Museum collections

244 The Museum collections have been examined first by Araujo & Ramos (2001) at a wide scale, focusing mainly  
245 on national museums worldwide. Prié et al. (unpublished data, Online Resource 1) have focused on French  
246 regional collections only. Out of the 58 collections identified, 25 had at least one specimen of *M. auricularia*  
247 (Fig. 2A): Musée du Château in Annecy, Musée des Confluences in Lyon, Museum of Perpignan, Musée  
248 zoologique of Strasbourg, Muséum - Aquarium of Nancy, Museum of Auxerre, Muséum d'histoire naturelle in  
249 Bordeaux, Muséum d'histoire naturelle in Bourges, Muséum d'histoire naturelle in Grenoble, Museum d'Histoire  
250 Naturelle in Nantes, Muséum d'Histoire Naturelle in Toulouse, Museum d'Histoire Naturelle Victor Brun in  
251 Montauban, Museum d'Histoires Naturelles in Colmar, Muséum of Orléans, Muséum of Dijon, Muséum Lecoq  
252 in Clermont-Ferrand, Muséum national d'Histoire naturelle in Paris, Muséum national d'histoire naturelle in  
253 Lille, Paraclat center of ONEMA in Boves, Pôle muséal of Troyes, Université of Bourgogne in Dijon, Université  
254 of Montpellier I, Université of Rennes I, Museum d'histoire naturelle in la Rochelle, Museum of Cherbourg-  
255 Octeville. Part of the data from Museum collections were fossil specimens. A total of 400 non-fossil specimens  
256 were found in Museum collections, including the 37 specimens already found by Araujo & Ramos (2001).  
257 Among them, 332 were localized at a river drainage scale. A third of the specimens came from the Garonne  
258 drainage, 19 % from the Saône River (half of them coming from a single batch collected by Coutagne in 1879)  
259 and 17% from the Ebro River (Fig. 2B). Other drainages represent less than 30% of the Museum collections  
260 specimens. About 80% of the specimens dated were collected before the beginning of the 20<sup>th</sup> century.

261

#### 262 Field surveys and populations sizes

263 A total of 2.500 km of rivers has been surveyed for *M. auricularia* in France and Spain during the last ten years  
264 (see bibliography and Online Resource 1 for details). These surveys covered most of the river stretches for which  
265 literature or museum collections data was available. Eleven populations could be identified, eight in France and  
266 three in Spain, plus a single individual found recently in the Ebro River (pers. comm. from R. Álvarez-Halcón to

267 R. Araujo) upstream Zaragoza (Fig. 3, Table 3). In Spain, the main population, with 5.000 live specimens, live at  
268 the Canal Imperial in Aragón. Although there have been some recent mortalities, some young specimens  
269 probably under ten years old have been observed during the last years (pers. comm. from J. Guerrero to R.  
270 Araujo). The other two Spanish populations, on the Canal de Tauste and the lower Ebro River, are today  
271 practically testimonials (pers. com. of the Generalitat of Catalonia to R. Araujo). See Gómez & Araujo (2008),  
272 Araujo (2012) and Araujo & Álvarez-Cobelas (2016) for more information.

273 In France, field surveys allowed finding ancient shells in the Seine, in the Vesle and in the Aisne Rivers; in the  
274 Saône River (Rhône drainage) near Pontailleur-sur-Saône and in the Garonne River near Agen, findings which  
275 corroborate historical data. We believe the species was extirpated long time ago in those rivers. In the Oise River  
276 (Seine drainage), very recent shells have been found in 2007 and 2008, some of them still embedded in their  
277 natural position, suggesting that the species became extirpated very little time before the surveys took place.

278 The populations of the Creuse and Vienne Rivers (Loire drainage) are the most studied in France. They live in  
279 shallow and clear water, allowing regular surveys using viewing glasses or snorkelling. Although these  
280 populations are rather small (about 250 specimens altogether), over 40 juveniles were found in the Vienne and  
281 Creuse Rivers, which represent about 15% of the population.

282 Three sites with a few tens of live specimens were discovered in the Dronne River, including one juvenile of  
283 about ten cm. Additionally, some isolated individuals were also observed, suggesting the population is scarce but  
284 relatively widespread. In the Save River, only 5 live specimens were observed. Sampling conditions are difficult,  
285 with variable depth and current strength, and very low visibility. We can therefore suppose that our detection  
286 probability is low. But based on survey results, we estimate that the population should not exceed a few tens of  
287 living individuals. It is likely rapidly declining given the bad condition of the river and the large number of  
288 recent shells collected compared to the very few living specimens observed. The Adour drainage rivers were  
289 known to host *M. auricularia* from both literature and Museum collections data. In the Adour mainstream, the  
290 population is now highly fragmented, with only three sites where live specimens could be found. One of them,  
291 the most upstream, is now extirpated (Prié et al., 2010). The total population is estimated to be about 300  
292 specimens in the total length of the Adour mainstream, but we still need a better estimation based on an  
293 appropriate sampling protocol. On the Luy tributary, a population of about 150 specimens is found in a very  
294 small stretch of river. Interestingly, although this River is very shallow (from 30 cm to 1,5 m), clear and easy to  
295 survey (hence detection probability is optimal), successive counts of 2010, 2011 and 2012 lead to respectively  
296 110, 96 and 145 specimens. We suppose that a significant part of the population lives buried in the sediment,

297 which biases the results of the counts. The Arros River had been overlooked by literature review and field  
298 surveys up to 2016. Following the findings in Museum collections, dedicated field surveys were conducted in  
299 2016, allowing the rediscovery of a living population. This population's size was estimated to about 200  
300 individuals on the 54 km of favourable habitat. The Charente River was known from the ancient literature to host  
301 an important population of *M. auricularia* (Bonnemère, 1901). Shell fragments and very few live specimens had  
302 been found by naturalists since 2003 (Nienhuis, 2003; P. Jourde pers. com.). Intensive field surveys performed in  
303 2007, 2010, 2016 led to the discovery of the largest population worldwide. Geographical statistics based on  
304 scuba-diving transects showed that the population was not aggregated. Hence the total population size could be  
305 estimated by multiplying the average density by the total surface of suitable habitat in the stretch of river  
306 inhabited by *M. auricularia*. The population size in the Charente River was estimated to be about 100.000  
307 (80.000 – 120.000) individuals, between the towns of Cognac upstream to Port-d'Envaux downstream.

308

#### 309 Genetic diversity

310 *Margaritifera auricularia* is genetically remarkably homogenous. The specimens from France and Spain all  
311 shared the same 16S and COI haplotype, but two specimens from Spain: specimens vouchered FW1238-14 and  
312 FW1238-12, with for COI T-> A in position 37, T->A in position 50 and G->C in position 73; and for 16S T-> C  
313 in position 176. The French and Spanish specimens could not be compared for 28S as different gene fragments  
314 were amplified. But within France, all specimens shared the same haplotype and within Spain, all specimens  
315 shared the same haplotype.

316

#### 317 Discussion

##### 318 Historical and actual data

319 The number of specimens found in the various regional museum collections was unexpected. *Margaritifera*  
320 *auricularia* is a large species that retained collector's attention. Most data from museum collections  
321 corresponded to the literature data, excepted those from the Arros and Vezere Rivers in France. Surprisingly,  
322 most French specimens came from the Garonne and Saône Rivers, where the species is now believed to be  
323 extirpated or very rare. In contrast, very few specimens came from the Charente River, where the largest  
324 population is found nowadays, and where industrial fisheries were established to make nacre shirt buttons  
325 (Bonnemère, 1901). Similarly, museum collections host no specimen from the Vienne or Creuse Rivers, where

326 healthy populations live in shallow and clear waters. In the Seine drainage, most shells came from upstream and  
327 the Aisne tributary, while the Oise tributary seems to have host the last population.

328 The historical review confirmed that *M. auricularia* was once present as far as the Thames in England and  
329 Netherlands and Germany where fossil specimens have been found and studied (Araujo & Ramos, 2001). On  
330 historical times, we found museum records (recent shells) from the Rhine in France or Germany (precise  
331 location being unknown), the Seine and the Rhône in France, the Pô in Italy and the Tajo in Spain, where the  
332 species is now believed to be extirpated (Araujo & Ramos, 2001). Today, *Margaritifera auricularia* is  
333 considered restricted to five coastal drainages: from north to south the Loire drainage (two close populations in  
334 the Vienne and Creuse Rivers), the Charente drainage, the Garonne drainage (two very isolated populations, in  
335 the Dronne and Save Rivers), the Adour drainage (at least three isolated populations, one in the Adour itself, one  
336 in the Luy and one in the Arros) and the Ebro River (three populations, two in channels and a small one  
337 remaining in the Ebro itself). As has been previously estimated (Prié et al., 2014), *Margaritifera auricularia*'s  
338 range contraction has probably reached about 90% in the last two centuries.

339

340 Surveying downstream ecosystems

341 Large rivers are amongst the most difficult ecosystems to sample. Deepness, turbidity and water current are  
342 challenging conditions. In addition, large rivers are subject to navigation, which makes scuba-diving potentially  
343 hazardous. Nevertheless, scuba diving appears to be the most efficient way to produce data for species such as  
344 *M. auricularia*: despite malacological surveys undertaken with canoes and dredging, only a few shell fragments  
345 had been collected in the Charente River before scuba diving sampling had been set up. Scuba divers met  
346 hundreds of shells and living specimens there. Similarly, scuba-divers collected the few living specimens, that  
347 today are probably dead by now, in the main Ebro River in Spain (Araujo & Álvarez-Cobelas, 2016). In the Oise  
348 River, a few ancient shell fragments had been collected on the banks by amateur malacologists, but scuba-diving  
349 allowed finding numerous shells in most of the river stretches investigated. In the Garonne River mainstream, a  
350 malacologist spent about 20 days wading and searching for shells on the gravelled banks. In two days, a team of  
351 three divers found four shell fragments.

352 While bivalve surveys have been conducted in the Saône River (ex. Mouthon & Daufresne, 2006), no shell  
353 fragments had ever been collected before 2016's scuba-diving prospections. The advances in the distribution  
354 knowledge of *M. auricularia* in France and Spain are directly linked to new investigation methods and scuba  
355 diving is so far the most efficient mean of survey for this species.

356 Conservation and further perspectives

357 Main threats

358 While overfishing may have contributed to the species decline in the past (Bonnemère, 1901; Prié et al., 2011;  
359 Araujo & Álvarez-Cobelas, 2016), it is obviously river management and agriculture impacts that nowadays  
360 cause the most important threats to the Giant Freshwater Pearl Mussel. Both causes are linked together, at least  
361 in the southern part of the species distribution area: river management aims at providing freshwater for corn  
362 culture, especially in summer. Hence, numerous dams are built, even in small rivers, to maintain pools for  
363 pumping water in the dry season. These dams produce lotic and silty conditions unsuitable for the Giant  
364 Freshwater Pearl Mussel. Altogether, these small dams can affect about than 70% of a given rivers stretch. In the  
365 Dronne, Arros and Save Rivers in France for example, the Giant Pearl Freshwater Mussel populations survive in  
366 the form of dashed lines, only in riffles (shallow parts of streams where water flows brokenly) with gravel or  
367 stony bottoms, between long portions of lotic conditions. Moreover, these dams constitute obstacles for potential  
368 fish hosts. The presumed natural host fish of the Giant Freshwater Pearl Mussel, the European Sturgeon  
369 *Acipenser sturio*, has been extirpated from almost all European rivers mainly because of dams (Lepage &  
370 Rochard, 1995; Gesner et al., 2010). River management has been an important threat in Spain too. Water  
371 regulation and the replacement of natural bottoms with concrete have been responsible for a massive death of  
372 Naiads. Recently, there has been an unusual high mortality of adults in the Imperial Channel (pers. com. from  
373 the Diputación General de Aragón to R. Araujo), but the causes are unknown.

374 Although a probable cause of recruitment failure, moderate levels of pollution and eutrophication have not  
375 demonstrated to be a significant threat to adult specimens. Some population survive in highly human-impacted  
376 waters. For example, one of the highest Giant Freshwater Pearl Mussel densities spot lies just downstream the  
377 Saintes sewage system in the Charente River. The same kind of conditions occurs at the Canal Imperial in  
378 Aragón with the water coming from the Ebro River, which is highly polluted. Overall, the species survives in  
379 rivers that are highly impacted by agriculture and domestic effluents. But we still don't know how these  
380 eutrophic and polluted waters may impact juvenile survival (Augspurger et al., 2007; Strayer & Malcom, 2012;  
381 Archambault et al., 2014).

382 Invasive species probably add to the threats *M. auricularia* is facing. Widespread invasive species such as  
383 *Corbicula fluminea* probably affect the freshwater mussels of Europe like it has been demonstrated for other  
384 species in North America (eg. Soussa et al., 2014). However, no clear impacts have been described for *M.*  
385 *auricularia*, and the healthiest populations survive in rivers largely colonized by *Corbicula*. The zebra mussel

386 *Dreissena polymorpha* attaches to the valves of *M. auricularia* in the Ebro, probably affecting filtration  
387 efficiency. This phenomenon has not been observed in France, where the zebra mussel remains at low densities  
388 in the rivers of the Atlantic coast.

#### 389 Habitat management

390 Contrarily to the Freshwater Pearl Mussel *M. margaritifera*, for which experiments of habitat managements have  
391 proved to be successful (Altmüller & Dettmer, 2006), the Giant Freshwater Pearl Mussel lives in downstream  
392 ecosystems. Attempts to implement broad scale drainage system management are therefore unrealistic. However,  
393 some realistic management objectives can be achieved to improve the habitat quality locally, in a short or middle  
394 term. The deconstruction of the numerous impoundments (many of them being disused) seems the most efficient  
395 way to restore suitable riverbed conditions for the Giant Freshwater Pearl Mussel. Although the negative impact  
396 of pollution and eutrophication are not clearly known, they are for sure not needed for the species survival.  
397 Improving water quality through reasonable agricultural practices, with buffer strips or grass strips along  
398 waterways, should be a middle-term objective.

#### 399 Farming projects

400 Breeding farms have been established for many endangered mussel species. In Europe, there is an abundant  
401 literature dealing with *M. margaritifera* breeding farms. Some trials are also ongoing for *U. crassus* and for  
402 various *Unio* species in Spain (Araujo et al., 2015). Regarding the Giant Freshwater Pearl Mussel, attempts of  
403 ex-situ breeding have been performed in Spain (Nakamura et al., 2015), and a LIFE project is ongoing in France  
404 to artificially breed the species in controlled conditions. Juveniles have been successfully produced (Nakamura  
405 et al., 2015), but we still face obstacles in the rearing of these juveniles (although some juveniles are still alive,  
406 Nakamura com. pers.).

407

#### 408 Genetic diversity

409 The very low genetic diversity for the mitochondrial genes studied was unexpected as (i) the Giant Freshwater  
410 Pearl Mussel populations are geographically isolated for a long time; and (ii) strong morphological differences  
411 are found between populations (Fig. 4). (i) The populations from France belong to the Atlantic drainage and the  
412 population from Spain to the Mediterranean drainage, two geographically isolated bioregions. Strong genetic  
413 divergences are observed for other freshwater mussel species from the Iberian Peninsula: *U. delphinus* from the  
414 “*pictorum*” lineage and *U. tumidiformis* from the “*crassus*” lineage were recently considered as distinct species  
415 based on molecular divergences (Reis & Araujo, 2009; Araujo et al., 2009b). But on the other hand, some

416 species do not show significant genetic divergences (ex. *U. mancus*, Prié et al., 2012; *Potomida littoralis*, Araujo  
417 et al., 2016; Froufe et al., 2016). (ii) The different populations known today have obvious morphological  
418 differences in shell size and shape (Fig. 4). The specimens from the Charente River population have a peculiarly  
419 small and conspicuously ear-like shell shape, contrasting to the Vienne and Dordogne Rivers populations, which  
420 are larger and more elongated; and to the Arros and Save Rivers populations, which are remarkable with their  
421 huge sizes. Some populations live in deep coastal rivers (ex. Ebro, Vienne and Charente populations) while  
422 others seem to be confined to shallow riffle sections of the upstream rivers (ex Save and Adour populations), but  
423 these ecological traits are not linked to shell morphological differences.

424 Margaritiferidae are known to have very low mitochondrial DNA evolution rates (Araujo et al., 2016; Bolotov et  
425 al., 2016). Population genetics based on microsatellites allowed to differentiate evolutionary units within the  
426 related species *Margaritifera margaritifera* (Geist et al., 2010; Stoekle et al., 2016) and *M. marocana* (Soussa et  
427 al., submitted). But first studies using microsatellites based on *M. margaritifera* primers have failed to reveal any  
428 population structure in France (Prié, unpublished data). If the ex-situ breeding projects are successful, the  
429 population genetics question will become unavoidable.

430

431 The fish host issue

432 The known host fish of *Margaritifera auricularia* are sturgeon species *Acipenser sturio*, *A. nacari* and *A. baeri*,  
433 the River Blenny *Salaria fluviatilis* and the Eastern Mosquitofish *Gambusia holbrooki* (Araujo & Ramos, 1998b;  
434 Araujo et al., 2000; Araujo et al., 2001; Altaba & Lopez, 2001; Lopez & Altaba, 2005; Lopez et al., 2007).

435 The only native *Acipenser* species in the area of occurrence of *Margaritifera auricularia* is the European  
436 sturgeon *A. sturio*. This species became extirpated from most European Rivers during the 20th century.  
437 Nowadays, it is almost extinct, with last documented natural reproduction dating back to 1994 in the Garonne  
438 River. The River Blenny is a Mediterranean species whose range does not overlap with the French populations  
439 of *M. auricularia*. The Eastern Mosquitofish, an introduced species, lives in shallow and standing to slow-  
440 flowing waters. It is not usually found in places favoured by *Margaritifera auricularia*. Reported success as host  
441 fish for *M. auricularia* glochidia was questionable. Experiments with other common fish species that occur  
442 within the distribution range of *M. auricularia* (*Anguilla Anguilla*, *Barbus graellsii*, *Barbus haasi*,  
443 *Parachondrostoma toxostoma*, *Cobitis paludicola*, *Liza aurata*, *Mugil cephalus*, *Alburnus alburnus*, *Carassius*  
444 *auratus*, *Cyprinus carpio*, *Gobio gobio*, *Scardinius erythrophthalmus* and *Tinca tinca*) failed to produce juveniles  
445 (Araujo et al., 2001; Lopez & Altaba, 2005).



446 The actual knowledge on *M. auricularia* host fish cannot explain the recruitment observed recently in the  
447 Atlantic drainage rivers. We therefore suspect an overlooked host fish species. For example, *Alosa* species,  
448 which are migratory fish and still breed in the drainages where *M. auricularia* produces juveniles, are good  
449 candidates (Llorente et al., 2015). But there must be another fish host to explain reproduction in the Dronne and  
450 Charente Rivers, which are isolated from the sea by impoundments; or in very upstream populations such as  
451 those of the Arros or Aisne Rivers, where migratory fishes do not breed. The other hypothesis could be that  
452 reproduction occurred in France periodically taking advantage of accidental releases of *A. baeri*, a common  
453 species in French fish farms (but not in Spain). We have recently succeeded in completing the full cycle on the  
454 three-spined stickleback *Gasterosteus aculeatus* in controlled conditions (Soler et al., in prep.). As this species is  
455 widespread within the range of *M. auricularia* and tolerant to brackish waters, it could also be a good candidate  
456 as a natural fish host. To find the natural host fish species of *M. auricularia* in France is now vital for the  
457 survival and conservation of this freshwater bivalve.

458

459 Conclusion

460 *Margaritifera auricularia* has become very rare in the last century, with an estimated range contraction of 90%.  
461 Only three populations were known worldwide before 2007. Intensive surveys in the last decade allowed re-  
462 discovering nine more. Given the magnitude of the efforts allocated to surveying the species in its historical  
463 range, we now believe that there are very little chances to rediscover unnoticed populations (excepted maybe in  
464 north-east France).

465

466 Although some juveniles were found recently, they remain very scarce and most extant populations seem to live  
467 on a borrowed time. Within the time lapse of this study, some populations already became extirpated in the Ebro  
468 and Adour Rivers. The status of the species therefore remains worrying. Priority populations for conservation are  
469 the Charente River's population, because it is by far the largest worldwide; the Vienne and Creuse population,  
470 because it has the higher level of natural recruitment; the Adour drainage populations, because they form an  
471 important and unique metapopulation; and the Ebro population because it is now the only remaining one in the  
472 Mediterranean drainage. Conservation challenges for the next years are (i) an appropriate management of the  
473 rivers which host the priority populations; (ii) the development of farming projects, in order to reinforce existing  
474 populations; (iii) research on fish hosts, for a better comprehension of the species' threats; ecological  
475 requirements, to understand which are the habitat factors driving the species' recruitment success; population

476 genetics to plan conservation efforts according to the genetic diversity of the remaining populations; (iv) a wide  
477 scale development of modern survey methods such as scuba diving and environmental DNA in order to discover  
478 the potentially remaining unnoticed populations.

479 Despite these efforts, we may fail to save the Giant Freshwater Pearl Mussel from extinction. However, current  
480 researches help to shed light on the obscure river downstream ecosystems' ecological functions and threats, as  
481 well as to develop exploring methods for this challenging environment.

482

483

#### 484 References

485

486 Andrés, D. & A. Ortuño, 2014. *Margaritifera auricularia* (Spengler, 1793): primeras evidencias sobre el antiguo  
487 cuaternario de Yecla (Murcia). *Athene* 24: 21-37.

488 Anselin, L., 1996. The Moran Scatterplot as an ESDA Tool to Assess Local Instability in Spatial Association. in:  
489 Fischer, M., Scholten, H. & D. Unwin [Eds]. *Spatial Analytical Perspectives on GIS in Environmental and*  
490 *Socio-Economic Sciences*. Taylor and Francis, London.

491 Anselin, L., I. Syabri & Y. Kho, 2006. GeoDa: An introduction to spatial data analysis. *Geographical Analysis*  
492 38: 5-22.

493 Altaba, C. R., 1990. The last known population of the freshwater mussel *Margaritifera auricularia* (Bivalvia,  
494 Unionoidea): A conservation priority. *Biological conservation* 52: 271-286.

495 Altaba, C. R. & M. A. Lopez, 2001. Experimental demonstration of viability for the endangered giant  
496 pearl mussel *Margaritifera auricularia* (Bivalvia: Unionoidea) in its natural habitat. *Bolleti de la Societat*  
497 *d'Historia Natural de les Balears* 44: 15-21.

498 Altmüller, R. & R. Dettmer, 2006. Erfolgreiche Artenschutzmaßnahmen für die Flussperlmuschel *Margaritifera*  
499 *margaritifera* L. durch Reduzierung von unnatürlichen Feinsedimentfrachten - Erfahrungen im Rahmen des  
500 Lutterprojekts. *Informationsdienst Naturschutz Niedersachsen* 26: 192 -204.

501 Araujo, R. & M. A. Ramos, 1998a. *Margaritifera auricularia* (Unionoidea, Margaritiferidae), the giant  
502 freshwater pearl mussel rediscovered in Spain. *Graellsia* 54: 129-130.

503 Araujo, R. & M. A. Ramos, 1998b. Description of the glochidium of *Margaritifera auricularia* (Spengler, 1793)  
504 (Bivalvia, Unionidae). *Philosophical Transactions of The Royal Society of London B* 353: 1553-1559.

505 Araujo, R. & R. Moreno, 1999. Former Iberian distribution of *Margaritifera auricularia* (Spengler) (Bivalvia:

506 Margaritiferidae). *Iberus* 17: 127-136.

507 Araujo, R. & M. A. Ramos, 2000a. A critical revision of the historical distribution of the endangered  
508 *Margaritifera auricularia* (Spengler, 1782) (Mollusca: Margaritiferidae) based on Museum specimens.  
509 *Journal of Conchology* 37: 49-59.

510 Araujo, R. & M. A. Ramos, 2000b. Status and conservation of the relict giant European freshwater pearl mussel  
511 *Margaritifera auricularia* (Spengler, 1793). *Biological Conservation* 96: 233-239.

512 Araujo, R., D. Bragado & M. A. Ramos, 2000. Occurrence of glochidia of the endangered *Margaritifera*  
513 *auricularia* (Spengler, 1793) and other mussel species (Bivalvia: Unionoida) in drift and on fishes in an  
514 ancient channel of the Ebro River, Spain. *Archiv für Hydrobiologie* 148: 147-160.

515 Araujo, R. & Ramos, M. A. 2001. Action Plan for *Margaritifera auricularia*. Convention on the Conservation of  
516 European Wildlife and Natural Habitats (Bern Convention). *Council of Europe Publishing*. Nature and  
517 environment, No. 117. Strasbourg.

518 Araujo, R., D. Bragado & M. A. Ramos, 2001. Identification of the river blenny, *Salaria fluviatilis*, as a host to  
519 the glochidia of *Margaritifera auricularia*. *Journal of Molluscan Studies* 67: 128-129.

520 Araujo, R., N. Cámara & M. A. Ramos, 2002. Glochidium metamorphosis in the endangered freshwater mussel  
521 *Margaritifera auricularia* (Spengler, 1793): A histological and scanning electron microscopy study. *Journal*  
522 *of Morphology* 254: 259-265.

523 Araujo, R., M. Quirós & M. A. Ramos, 2003. Laboratory propagation and culturing of juveniles of the  
524 endangered freshwater mussel *Margaritifera auricularia* (Spengler, 1793). *Journal of Conchology* 38: 53-60.

525 Araujo, R., C. Toledo, D. Van Damme, M. Ghamizi & A. Machordom, 2009a. *Margaritifera marocana* (Pallary,  
526 1918) a valid current species inhabiting the Moroccan rivers. *Journal of Molluscan Studies* 75: 95-101.

527 Araujo, R., J. Reis, A. Machordom, C. Toledo, M. J. Madeira, I. Gomez, J. C. Velasco, J. Morales, J. M. Barea,  
528 P. Ondina & I. Ayala, 2009b. Las Nayades de la península Ibérica. *Iberus* 27: 7-72.

529 Araujo, R. 2012. *Margaritifera auricularia*. In: VV.AA., Bases ecológicas preliminares para la conservación de  
530 las especies de interés comunitario en España: Invertebrados. Ministerio de Agricultura, Alimentación y  
531 Medio Ambiente. Madrid. 54 pp.

532 Araujo, R., C. Feo, Q. Pou & M. Campos, 2015. Conservation of two endangered European freshwater mussels.  
533 A three-year semi-natural breeding experiment. *The Nautilus* 129: 126-135.

534 Araujo, R. & M. Álvarez-Cobelas, 2016. Influence of flow diversions on Giant freshwater pearl mussel  
535 populations in the Ebro River, Spain. *Aquatic Conservation: Marine and Freshwater Ecosystems*, DOI:  
536 10.1002/aqc.2622

537 Araujo, R., S. Schneider, K. J. Roe, D. Erpenbeck & A. Machordom, 2016. The origin and phylogeny of  
538 Margaritiferidae (Bivalvia, Unionoida): a synthesis of molecular and fossil data. *Zoologica scripta* DOI  
539 10.1111/zsc.12217

540 Araujo, R, D. Buckley, K.-O. Nagel & A. Machordom, 2016. *Potomida littoralis* (Bivalvia, Unionidae)  
541 evolutionary history: slow evolution or recent speciation? *Zoological Journal of the Linnean Society* DOI:  
542 10.1111/zoj.12470

543 Archambault, J.-M., W. G. Cope, & T. J. Kwak, 2014. Survival and behaviour of juvenile unionid mussels  
544 exposed to thermal stress and dewatering in the presence of a sediment temperature gradient. *Freshwater*  
545 *Biology* 59: 601-613.

546 Augspurger, T., F. J. Dwyer, C. G. Ingersoll & C. M. Kane, 2007. Advances and opportunities in assessing  
547 contaminant sensitivity of freshwater mussel (Unionidae) early life stages. *Environmental Toxicology and*  
548 *Chemistry* 26: 2025-2028.

549 Bolotov, I. N., I. V. Vikhrev, Y. V. Bespalaya, M. Y. Gofarov, A. V. Kondakov, E. S. Konopleva, N. N. Bolotov  
550 & A. A. Lyubas, 2016. Multi-locus fossil-calibrated phylogeny, biogeography and a subgeneric revision of  
551 the Margaritiferidae (Mollusca: Bivalvia: Unionoida). *Molecular Phylogenetics and Evolution* 103: 104-121

552 Bonnemère, L., 1901. Les mollusques des eaux douces de France et leurs perles. Institut international de  
553 bibliographie scientifique, Paris.

554 Bössneck, U., R. Kleemann & L. Buttstedt, 2006. Die Große Flussperlmuschel (*Pseudunio auricularius*  
555 Spengler, 1793) in Mitteleuropa: Neue Befunde zur historischen und fossilen Verbreitung (Bivalvia:  
556 Margaritiferidae). *Malakologische Abhandlungen* 24: 141-156.

557 Cochet, G., 2001. Redécouverte d'une population vivante de la Grande Mulette, *Margaritifera auricularia*, sur la  
558 Vienne et la Creuse. *Recherches Naturalistes en Région Centre* 10: 3-16.

559 Dudgeon, D., A. H. Arthington, M. O. Gessner, Z.-I. Kawabata, D. J. Knowler, C. Lévêque, R. J. N. Aiman, A.-  
560 H. Prieur-Richard, D. Soto, M. L. J. Stiassny & C. A. Sullivan, 2006. Freshwater biodiversity: importance,  
561 threats, status and conservation challenges. *Biological Reviews* 81: 163–182.

562 Froufe, E., V. Prié, J. Faria, M. Ghamizi, D. V. Gonçalves, M. Emre Gürlek, I. Karaouzas, Ü. Kebapçı, H.  
563 Sereflişan, C. Sobral, R. Sousa, A. Teixeira, S. Varandas, S. Zogaris & M. Lopes-Lima, 2016. Phylogeny,

564 phylogeography, and evolution in the Mediterranean region: News from a freshwater mussel (*Potomida*,  
565 Unionida). *Molecular Phylogenetics and Evolution* 100: 322-332.

566 Geist, J., H. Söderberg, A. Karlberg & R. Kuehn, 2010. Drainage-independent genetic structure and high genetic  
567 diversity of endangered freshwater pearl mussels (*Margaritifera margaritifera*) in northern Europe.  
568 *Conservation Genetics* 11:1339-1350.

569 Gesner, J., P. Williot, E. Rochard, J. Freyhof & M. Kottelat, 2010. *Acipenser sturio*. The IUCN Red List of  
570 Threatened Species 2010: e.T230A13040963. <http://dx.doi.org/10.2305/IUCN.UK.2010-1.RLTS.T230A13040963.en>. Downloaded on 19 December 2016.

571

572 Gómez, I. & R. Araujo, 2008. Channels and ditches as the last shelter for freshwater mussels. The case of *M.*  
573 *auricularia* and other naiads at the mid Ebro River basin, Spain. *Aquatic Conservation: Marine and*  
574 *Freshwater Ecosystems* 18: 658-670.

575 Grande, C., R. Araujo & M. A. Ramos, 2001. The gonads of *Margaritifera auricularia* (Spengler, 1793) and  
576 *Margaritifera margaritifera* (L. 1758) (Bivalvia: Unionoidea). *Journal of Molluscan Studies* 67: 27-35.

577 Hall T.A., 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for  
578 Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.

579 Lepage, M. & E. Rochard, 1995. Threatened fishes of the world: *Acipenser sturio* Linnaeus, 1758  
580 (Acipenseridae). *Environmental Biology of Fishes* 43: 28.

581 Llorente, L., E. Roselló, A. Morales, C. Liesau, A. Daza & R. Araujo, 2015. Las náyades (Mollusca, Unionoidea)  
582 del Calcolítico de Camino de las Yeseras (San Fernando de Henares, Madrid). *La Investigación*  
583 *Arqueomalacológica en la Península Ibérica: nuevas aportaciones. Actas de la IV Reunión de*  
584 *Arqueomalacología de la Península Ibérica*. Santander. Ed: Gutiérrez, I., D. Cuenca & M. R González: 125-  
585 134.

586 Lopes-Lima M., R. Sousa, J. Geist, D. C. Aldridge, R. Araujo, J. Bergengren, Y. Bernal, E. Bódis, L.  
587 Burlakova, D. Van Damme, K. Douda, E. Froufe, D. Georgiev, C. Gumpinger, A. Karatayev, Ü. Kebapçı, I.  
588 Killeen, J. Lajtner, B. M. Larsen, R. Lauceri, A. Legakis, S. Lois, S. Lundberg, E. Moorkens, G. Motte, K.-  
589 O. Nagel, P. Ondina, A. Outeiro, M. Paunovic, V. Prié, T. von Proschwitz, N. Riccardi, M. Rudzīte, M.  
590 Rudzītis, C. Scheder, M. Seddon, H. Şereflişan, V. Simić, S. Sokolova, K. Stoeckl, J. Taskinen, A. Teixeira,  
591 F. Thielen, T. Trichkova, S. Varandas, H. Vicentini, K. Zajac, T. Zajac & S. Zogaris, 2016. Conservation  
592 status of freshwater mussels in Europe: state of the art and future challenges. *Biological Reviews*  
593 doi:10.1111/brv.12244

594 López, M. A. & C. R. Altaba, 2005. Fish host determination for *Margaritifera auricularia* (Bivalvia:  
595 Unionoida): results and implications. *Bolletino Malacologico* 41: 89-98.

596 Lopez, M. A., C. R. Altaba, T. Rouault & E. Gisbert, 2007. The European sturgeon *Acipenser sturio* is a suitable  
597 host for the glochidia of the freshwater pearl mussel *Margaritifera auricularia*. *Journal of Molluscan Studies*  
598 73: 207-209.

599 Lydeard C., R.H. Cowie, W.F. Ponder, A.E. Bogan, P. Bouchet, S.A. Clark, K.S. Cumming, T.J. Frest, O.  
600 Gargominy, D.G. Herbert, R. Hershler, K.E. Perez, B. Roth, M. Seddon, E.E. Strong & F.G. Thompson,  
601 2004. The global decline of nonmarine mollusks. *BioScience* 54: 321–330.

602 Machordom, A., R. Araujo, D. Erpenbeck & M.-A. Ramos, 2013. Phylogeography and conservation genetics of  
603 endangered European Margaritiferidae (Bivalvia: Unionoidea). *Biological Journal of the Linnean Society* 78:  
604 235–252.

605 Mouthon, J. & M. Daufresne, 2006. Effects of the 2003 heatwave and climatic warming on mollusc communities  
606 of the Saône: a large lowland river and its two main tributaries (France). *Global Change Biology*, 12 : 441-  
607 449.

608 Nakamura, K., M. A. Yanguas, I. Ruiz, I. de Blas, E. Elbaile & C. Salinas, 2015. The first year of *Margaritifera*  
609 *auricularia* (Spengler, 1793). Breeding in captivity in Aragon, Spain. 2<sup>nd</sup> International seminar on the rearing  
610 of unionoid mussels. Tuesday 24<sup>th</sup> November – Thursday 26<sup>th</sup> November 2015, Clervaux (Luxembourg).

611 Nakamura, K., L. Cucala, A. Mestre, F. Mesquita-Joanes, E. Elbaile, C. Salinas & M. Ángel Muñoz-Yanguas, *in*  
612 *press*. Modelling growth in the critically endangered freshwater mussel *Margaritifera auricularia* (Spengler,  
613 1793) in the Ebro basin. *Hydrobiologia*.

614 Nienhuis, J. A. J. H., 2003. The rediscovery of Spengler’s freshwater pearl mussel *Pseudunio auricularius*  
615 (Spengler, 1793) (Bivalvia, Unionoidea, Margaritiferidae) in two river systems in France, with an analysis of  
616 some factors causing its decline. *Basteria* 67: 67-86.

617 Prié, V., L. Philippe & G. Cochet, 2007. Evaluation de l’impact d’un projet de canal sur les naïades de l’Oise  
618 (France) et découverte de valves récentes de *Margaritifera auricularia* (Spengler, 1793) (Bivalvia :  
619 Margaritiferidae). *MalaCo* 4: 178-182.

620 Prié, V., G. Cochet, & L. Philippe, 2008. Une population majeure de la très rare Grande Mulette *Margaritifera*  
621 *auricularia* (Bivalvia ; Margaritiferidae) dans la Charente (France). *MalaCo* 5: 230-239.

622 Prié, V., 2010. *Margaritifera auricularia*. The IUCN Red List of Threatened Species 2010: e.T12798A3381899.  
623 <http://dx.doi.org/10.2305/IUCN.UK.2010-4.RLTS.T12798A3381899.en>. Downloaded on 30 September  
624 2016.

625 Prié, V., P. Bousquet, A. Serena, E. Tabacchi, P. Jourde, B. Adam, T. Deschamps, M. Charneau, T. Tico, M.  
626 Bramard & G. Cochet, 2010. Nouvelles populations de Grande Mulette *Margaritifera auricularia* (Spengler,  
627 1793) (Bivalvia, Margaritiferidae) découvertes dans le sud-ouest de la France. *MalaCo* 6: 294-297.

628 Prié, V., L. Philippe & G. Cochet, 2011. Plan National d'Actions en faveur de la Grande Mulette *Margaritifera*  
629 *auricularia* – 2012-2017. Ministère de l'écologie, du développement durable, des transports et du logement.

630 Prié, V., N. Puillandre, & P. Bouchet 2012. Bad taxonomy can kill: molecular reevaluation of *Unio mancus*  
631 Lamarck, 1819 (Bivalvia: Unionidae) and its accepted subspecies. *Knowledge and Management of Aquatic*  
632 *Ecosystems* 405 (8): DOI <http://dx.doi.org/10.1051/kmae/2012014>

633 Prié, V., Q. Molina & B. Gamboa 2014. French Naiad (Bivalvia: Margaritiferidae, Unionidae) species  
634 Distribution Models: prediction maps as tools for conservation. *Hydrobiologia* 735: 81-94.

635 Prié, V. & N. Puillandre, 2014. Molecular phylogeny, taxonomy and distribution of French *Unio* species  
636 (Bivalvia, Unionidae). *Hydrobiologia* 735: 95-110.

637 Reis, J. & R. Araujo, 2009. Redescription of *Unio tumidiformis* Castro, 1885 (Bivalvia, Unionidae), an  
638 endemism from the south-western Iberian Peninsula. *Journal of Natural History* 43: 1929-1945.

639 Soussa, R., A. Novais, R. Costa & D. L. Strayer, 2014. Invasive bivalves in fresh waters: impacts from  
640 individuals to ecosystems and possible control strategies. *Hydrobiologia* 735: 233-255.


641 Stoeckle, B. C., R. Araujo, J. Geist, R. Kuehn, C. Toledo & A. Machordom, 2016. Strong genetic differentiation  
642 and low genetic diversity of the freshwater pearl mussel (*Margaritifera margaritifera* L.) in the southwestern  
643 European distribution range. *Conservation Genetics*. DOI: 10.1007/s10592-016-0889-3

644 Strayer, D. L. & H. M. Malcom, 2012. Causes of recruitment failure in freshwater mussel populations in  
645 southeastern New York. *Ecological applications* 22: 1780-1790.

646 Villasante, F., J. Abad, R. Araujo & J. Basset, 2016. Aportación al conocimiento de la presencia de  
647 *Margaritifera auricularia* (Spengler, 1793) en el cauce del río Tajo (España). *Iberus*, 34: 79-82.

648 White, G.C. & K. P. Burnham, 1999. Program MARK: Survival estimation from populations of marked  
649 animals. *Bird Study* 46: 120-138.

650



652



653 Figure 1: fossil (black crosses) and historical data (white dots for precise locations, blue lines for rivers names

654 only) collected from the literature and Museum collections; and subsequent intensive field surveys locations


655 (polygons).

656






657  
 658 Figure 2: A: location of main museum collection investigated (dot size according to number of *M. auricularia*  
 659 specimens). B: number of specimens held in Museum collections per main coastal drainages (the Saône River is  
 660 actually a tributary of the Rhône, but all the specimens are located in the Saône and none elsewhere in the  
 661 Rhône).



662

663 Figure 3: results of last ten years' field surveys and known past and actual distribution of *M. auricularia*. Fossil  
 664 data (black crosses), historical data (white dots), shells collected in the last 10 years (orange dots) and still living  
 665 populations (red dots).

666



667

668 Figure 4: morphological variability of *Margaritifera auricularia*. A: Vienne River; B-C: Charente River; D:

669 Dronne River; E: Save River; F: Luy River; G: Arros River, H: Ebro River.

670  
671  
672

Table 1: summary of the methods used in France for surveying *M. auricularia*. For Spain, survey methods were detailed and reported in Gómez & Araujo, 2008 and Araujo & Álvarez-Cobelas, 2016.

Coastal drainage	River	Year	Use of a boat	Scuba-diving	Other methods	Authors involved
Somme	Somme River	2011	X	4 divers, 5 days		XC, VP
	Seine River (downstream)	2011	X		Dredging (2 persons for 4 days)	XC
	Seine River (upstream)	2015			Snorkeling (2 persons for one days)	XC
	Oise River	2007-2008	X	3 divers, 20 days		VP, LP, XC
Seine	Aisne River	2011		2-4 divers, 5 days		XC, VP
Rhône	Saône and Doubs Rivers	2016		4 divers, 5 days	Wading with viewing glasses (2 persons, 5 days)	VP, LP, NL, NP, BA
Loire	Loire, Indre and Cher Rivers	2010-2011		2-3 divers, about 5 days	Wading with viewing glasses (2-4 persons, estimated to about 10 days altogether)	LP, VP
	Vienne and Creuse Rivers	2009-2016		2 to 6 divers, over 20 days altogether	Wading with viewing glasses and snorkeling (2 to 6 persons, estimated to over 30 days altogether)	VP, XC, LP
Charente	Charente River	2007-2016	X	2 to 3 divers, about 20 days altogether		VP
	Dronne and Isle Rivers	2012-2014		2 divers, about 10 days altogether	Wading with viewing glasses (1-2 persons, about 5-10 days altogether)	VP
	Dordogne River	2016	X	2 divers, 3 days altogether	Wading with viewing glasses (1 person, about 5 days)	VP
	Vézère River	2016		3 divers, 3 days	Wading with viewing glasses and snorkeling (2 persons 3 days)	VP, LP, NP, BA, NL
	Garonne River	2016		3 divers, 2 days	Wading with viewing glass (1 person, 30 days)	VP, NL
Garonne	Save River	2009		2 divers, 1 day	Wading with viewing glass and snorkeling (1 person, 5 days)	VP, BA
Adour	Adour River	2012-2014		2 divers approx. 9 days	Wading with viewing glasses and snorkeling (2 to 6 persons, 10 days)	VP, BA, NL
	Arros River	2016		3 divers, 3 days	Wading with viewing glasses (6 persons, 4 days)	VP, BA, NL

673  
674

675 Table 2: genes and Genbank accession numbers of French specimens used for DNA analyses.

676

Coastal drainage	River	Specimen voucher number	Genbank accession number		
			COI	16S	28S
Charente	Charente	MNHN-IM-2009-12596	MF494673	MF494681	MF494677
		MNHN-IM-2009-12597	MF494674	MF494682	MF494678
Garonne	Save	MNHN-IM-2009-12601	MF494675	MF494683	MF494679
Adour	Luy	MNHN-IM-2009-12662	MF494671	MF494696	MF494676
		MNHN-IM-2009-12663	MF494672	MF494697	
Loire	Vienne	Maur91	MF494670	MF494695	MF494680
		MNHN-IM-2009-12611	MF494661	MF494684	
		MNHN-IM-2009-12615	MF494662	MF494685	
		Maur70	MF494663	MF494686	
		Maur72	MF494664	MF494687	
		Maur74	MF494665	MF494688	
		Maur76	MF494666	MF494689	
		Maur77	MF494667	MF494690	
		Maur78	MF494668	MF494691	
		Maur85	MF494669	MF494693	
		Maur79		MF494692	
Maur88		MF494694			

677

678

680 Table 3: summary of literature data, museum collections (fossil data are not considered) and field surveys.  
681

Country	Coastal drainage	Literature	Museum collections (number of specimens)	Dedicated field surveys (references with an * refer to grey literature, summarized in Online Ressource 1)	Recent surveys results	Estimated population size	
France	Charente	Charente	X	9	Prié et al., 2007*; Prié, 2010; Prié & Mouton, 2016*	Live specimens and juveniles	100.000
	Garonne	Garonne (mainstream)	X	125	Prié et al., 2016*	Ancien shells	
		Isle	X	0	Prié, 2012*	Recent shells	
		Dronne	X	0	Prié, 2012*; Prié, 2013*	Live specimens and juveniles	> 100
		Save		0	Prié, 2012*	Few live specimens, declining population	< 30
	Adour	Adour (mainstream)	X	9	Prié, 2012*	Live specimens and juveniles	> 300
		Arros		6	Prié & Néri, 2016*	Live specimens and subadult	200
		Luy		0	Prié, 2012*	Live specimens and subadults	150
	Loire	Loire (mainstream)	X	1		Recent shells	
		Vienne		0	Cochet, 2006*; Philippe et al., 2009*; Philippe et al., 2010*, 2011*, 2012*	Live specimens and juveniles	>100
		Indre		0	Dohogne, 2008*; Philippe et al., 2009*	Recent shells	
		Creuse		0	Philippe et al., 2012*; Philippe et al., 2013*, 2014*, 2015*, 2016*	Live specimens and juveniles	>150
		Cher		1	Prié et al., 2011*; Prié et al., 2016*	Nothing	
	Seine	Seine (mainstream)	X	12	Cucherat et al., 2011*	Ancien shells	
		Oise		0	Prié et al., 2007*	Recent shells	
		Aube	X	1	Cucherat et al., 2011*	Ancien shells	

		Aisne	X	6	Philippe et al., 2009*; Cucherat et al., 2011*	Ancien shells	
	Escaut	Escaut	X	2		No survey	
	Somme	Somme	X	1	Cucherat & Prié, 2011*	Nothing	
	Rhône	Saône	X	65	Prié et al., 2016*	Ancien shells	
Italy	Po	Po	X	15		No survey	
Spain	Ebro	Upper Ebro	X		Araujo et al., 2009b*; Araujo & Álvarez-Cobelas, 2016 ; pers. comm. from R. Álvarez-Halcón to R. Araujo	Live specimens and juveniles (at Gallur)	1
		Ribera alta			Nakamura & Guerrero, 2008 ; Araujo & Álvarez-Cobelas, 2016	Used to be 38-40 live specimens, today likely extirpated	
		Canal Imperial de Aragon	X	55	Gomez & Araujo, 2008; Araujo et al., 2009; Araujo & Alvarez-Corbela, 2016 ; pers. comm. from J. Guerrero to R. Araujo	Live specimens and juveniles	4.000
		Canal de Tauste	X		Araujo et al., 2009b*; pers. comm. from J. Guerrero to R. Araujo	Live specimens and juveniles	200
		Quinto ditch			Gomez & Araujo, 2008; Nakamura et al., 2017	Recent shells	
	Lower Ebro	X		Araujo et al., 2009b*; Araujo & Alvarez-Corbela, 2016	Used to be 70 live specimens, today extirpated		
	Tajo	Tajo	X	1	Villasante et al., 2016	Nothing	
<i>Fossil data</i>							
U.K.	Thames	Thames		17		No survey	
Germany	Rhine	Rhine	X	3		No survey	
Netherlands	Rhine	Rhine	X			No survey	

683 **Online resource 1: grey literature of the last decade. References of the various reports**  
684 **and impact studies aiming specifically at *M. auricularia*.**

685  
686  
687

---

688 2016

---

689

690 Philippe, L., N. Patry & C. Mathieu 2016. Suivi des populations de Grande Mulette déplacée  
691 dans le cadre de la LGV SEA. LISEA – Biotope

692 Prié, V. & F. Néri 2016. Recherche de la Grande Mulette *Margaritifera auricularia* sur  
693 l'Arros (Gers). Conservatoire d'Espaces Naturels Midi-Pyrénées – Biotope.

694 Prié, V., B. Adam, X. Cucherat, N. Legrand, N. Patry & L. Philippe 2016. Etude  
695 biogéographique de la Grande Mulette *Margaritifera auricularia* en France. Prospections  
696 dans le Cher, la Saône, la Vézère, l'Arros et la Garonne. Université François-Rabelais de  
697 Tours - Biotope

698 Prié, V. & N. Mouton 2016. Bathymétrie de la Charente – Caractérisation de l'habitat  
699 favorable à la Grande Mulette à partir de relevés bathymétriques. Biotope - INGEO.

700 Rubio Millán, C., Gloria Muñoz-Camarillo, I. Sanz Bayón, R. M. Álvarez Halcón & A. Calvo  
701 Tomás. 2016. Estado de conservación de las náyades en el Ebro medio. *Naturaleza*  
702 *Aragonesa*, n° 33.

703

---

704 2015

---

705

706 Philippe, L., N. Patry & C. Mathieu 2015. Suivi des populations de Grande Mulette déplacée  
707 dans le cadre de la LGV SEA. LISEA - Biotope

708 Prié, V., B. Adam, X. Cucherat, N. Legrand, N. Patry & L. Philippe 2015. Etude  
709 biogéographique de la Grande Mulette *Margaritifera auricularia* en France. Synthèse  
710 bibliographique et recherches muséographiques. Université François-Rabelais de Tours -  
711 Biotope

712

---

713 2014

---

714

715 Le Bloch, F., R. Henry, G. Dicev, N. Flamant, S. Sibley, A. Vacher, E. Loufti, E. Monnier, T.  
716 Sauzon, Q. Vanel, M. Camus, S. Montagne, J. Loiseau, P. Clevenot, D. Mollard, C.  
717 Gibeau, H. Bouyon, S. Vrignaud, D. Genoud & J. Mouthon 2014. Projet de mise à grand  
718 gabarit de la Seine entre Bray-sur-Seine (77) et Nogent-sur-Seine (10). Voies Navigables  
719 de France – Écosphère – Hydrosphère

720 Philippe, L., N. Patry & C. Mathieu 2014. Suivi des populations de Grande Mulette déplacée  
721 dans le cadre de la LGV SEA. COSEA – Biotope

722

---

723 2013

---

724

725 Philippe, L., N. Patry, C. Mathieu & S. Walter 2013. Suivi des populations de Grande Mulette  
726 déplacée dans le cadre de la LGV SEA. COSEA – Biotope

727 Prié, V. 2013. Recherche et caractérisation des populations de Grande Mulette *Margaritifera*  
728 *auricularia* dans la Dronne. EPIDOR – Biotope

729

---

730 2012

---

731



- 732 Nakamura Antonacci, K., E. Elbaile Pérez, M.A. Muñoz Yanguas, C. Catalá Roca & C.  
733 Salinas Yuste 2012. Captive breeding of the endangered pearl mussel *Margaritifera*  
734 *auricularia* (Spengler, 1793). Large scale laboratory production of juveniles. Poster.  
735 International Meeting on Biology and Conservation of Freshwater Bivalves, 4-7 September  
736 2012, Bragança, Portugal
- 737 Nakamura Antonacci, K., E. Elbaile Pérez, M.A. Muñoz Yanguas, C. Catalá Roca & C.  
738 Salinas Yuste 2012. Size-specific growth pattern of fresh water mussel *Margaritifera*  
739 *auricularia* in the Ebro river channels (Spain). Poster. International Meeting on Biology  
740 and Conservation of Freshwater Bivalves, 4-7 September 2012, Bragança, Portugal
- 741 Philippe, L., V. Prié, X. Cucherat, N. Patry, D. Froment, M. Jamier, G. Fahrner, B. Faure, C.  
742 Mathieu, S. Walter, Q. Molina & N. Tapko 2012. Etude d'impact du franchissement de la  
743 Vienne par la LGV SEA – Déplacement de la population des stations de Grande Mulette et  
744 de Mulette épaisse impactées. COSEA - Biotope
- 745 Philippe, L., V. Prié, X. Cucherat, N. Patry, D. Froment, G. Fahrner, S. Walter, Q. Molina &  
746 N. Tapko 2012. Etude d'impact du franchissement de la Vienne par la LGV SEA –  
747 Déplacement de la population des stations de Grande Mulette et de Mulette épaisse  
748 impactées. COSEA - Biotope
- 749 Prié, V. 2012. Inventaire de la Grande mulette *Margaritifera auricularia* dans le Sud-Ouest  
750 de la France (III). Agence de l'Eau Adour-Garonne – Biotope
- 
- 751
- 752 2011
- 
- 753
- 754 Cucherat, X. & V. Prié 2011. Etude d'impact du projet de curage du canal de la Somme entre  
755 Sailly-Laurette et Sormont (Département de la Somme) - Recherches spécifiques de la  
756 Grande Mulette et de la Mulette épaisse. Conseil Général de la Somme - Biotope
- 757 Cucherat, X., V. Prié & N. Tapko 2011. Étude d'impact du projet de curage du canal de la  
758 Somme entre Sailly-Laurette et Sormont (Département de la Somme) - Recherches  
759 spécifiques de la Grande Mulette et de la Mulette épaisse. Conseil Général de la Somme –  
760 Biotope
- 761 Cucherat, X., M. Jamier, L. Philippe, V. Prié, L. Simon & N. Tapko 2011. Mise en place d'un  
762 plan de conservation pour les mollusques inscrits aux annexes de la Directive « Habitats-  
763 Faune-Flore » et protégés au titre de l'arrêté du 23 avril 2007 – Phase II - DREAL Picardie  
764 - Biotope
- 765 Muñoz-Yanguas, MA, C. Catalá, E. Elbaile, C. Salinas & K. Nakamura 2011. First  
766 assessment of the population effects of the maintenance channel works in the naiads of the  
767 Canal Imperial de Aragón. EUROMALACOL. 18 al 23 de Julio 2011. Vitoria, Spain.
- 768 Prié V., L. Philippe & N. Patry 2011. Recherche de la présence d'espèces patrimoniales de  
769 mollusques aquatiques dans la rivière le Cher. DREAL Centre - Biotope
- 770 Philippe, L., V. Prié & N. Patry 2011. Etude d'impact du franchissement de la Vienne par la  
771 LGV SEA – Test de marquage et de déplacement sur la population de Grande Mulette  
772 impactée. COSEA – Biotope
- 773 Prié, V. 2011. Inventaire de la Grande mulette *Margaritifera auricularia* dans le Sud-Ouest  
774 de la France (II). Agence de l'Eau Adour-Garonne - Biotope
- 
- 775
- 776 2010
- 
- 777
- 778 Philippe, L., V. Prié, X. Cucherat, N. Tapko & P. Bousquet 2010. Etude d'impact du  
779 franchissement de la Vienne par la LGV SEA - Suivi de la population des stations de  
780 Grande Mulette et de Mulette épaisse potentiellement impactées. COSEA – Biotope

- 781 Prié, V. 2010. Inventaire de la Grande moulette *Margaritifera auricularia* dans le Sud-Ouest  
782 de la France (I). Agence de l'Eau Adour-Garonne - Biotope
- 783 Prié, V. 2010. La Grande Mulette *Margaritifera auricularia* sur la Charente, complément  
784 d'inventaire sur le site Natura 2000 FR 5400-472. Ligue pour la Protection des Oiseaux -  
785 Biotope
- 
- 787 2009
- 
- 789 Cucherat, X. 2009. Mise en place d'un plan de conservation pour les mollusques inscrits aux  
790 annexes de la Directive « Habitats-Faune-Flore » et protégés au titre de l'arrêté du 23 avril  
791 2007 en Picardie – Phase I- DREAL Picardie - Biotope
- 792 Philippe, L., V. Prié, X. Cucherat & P. Bousquet 2009. Inventaires des mollusques protégés  
793 sur l'Aisne et la Meuse dans le cadre du projet de réhabilitation de 29 barrages à aiguilles.  
794 Voies Navigables de France – Biotope.
- 795 Philippe, L., V. Prié & G. Cochet 2009. Etude préliminaire au franchissement de la Vienne  
796 par la LGV SEA – Inventaire des moulettes protégées sur 59 cours d'eau entre Bordeaux et  
797 Tours et suivi de la population des stations de Grande Mulette de l'A10. RFF – Biotope
- 798 Araujo, R., M.J. Madeira & I. Ayala 2009. Estudio del estado actual de conservación de  
799 *Margaritifera auricularia* en las aguas del río Ebro. Informe final. CSIC-ACEBI. Madrid.
- 
- 801 2008
- 
- 803 Dohogne, R. 2008.- Recherche de la Moule perlière (*Margaritifera margaritifera*) et de la  
804 Grande moulette (*Margaritifera auricularia*) dans le département de l'Indre. Rapport  
805 d'étude, Indre nature : 13 pp.
- 806 Nakamura, K. & J. Guerrero 2008. *Margaritifera auricularia*, un difícil reto de conservación.  
807 *Quercus* 265: 20-25.
- 
- 809 2007
- 
- 811 Araujo, R., M.J. Madeira & I. Ayala 2007. Estudio del estado actual de *Margaritifera*  
812 *auricularia* en las aguas del Rio Ebro. Confereracion Hidrografica del Ebro
- 813 Prié, V., L. Philippe & G. Cochet 2007. Etude d'impact du projet de curage de la Charente  
814 sur les bivalves et recherche de la Grande Mulette *Margaritifera auricularia* Etablissement  
815 Public Territorial de Bassin du fleuve Charente - Biotope
- 816 Prié, V., L. Philippe & G. Cochet 2007. Etude d'impact du projet de canal Seine-Nord-Europe  
817 sur les Nâïades – rivière Oise. Voies Navigables de France – Biotope
- 
- 819 2006
- 
- 821 Cochet, G. 2006. La Grande Mulette (*Margaritifera auricularia*) dans la Vienne au niveau du  
822 tracé de la LGV SEA. Réseau Ferré de France, 35 pp.
- 823 Zapater Galve M., R. Araujo Armero, R. Álvarez Halcón, K. Nakamura Antonacci & M.  
824 Alcántara de la Fuente (coord.) 2006. Las almejas de agua dulce en Aragón : *Margaritifera*  
825 *auricularia* y otros bivalvos. Serie Especie. Consejo de Protección de la Naturaleza de  
826 Aragón, D.L. 68 pp.
- 827
- 828
- 829