The variation of invariant graphs in forced systems

Bastien Fernandez, Anthony Quas

► To cite this version:

Bastien Fernandez, Anthony Quas. The variation of invariant graphs in forced systems. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 2018, 28 (8), pp.083101. 10.1063/1.5026551. hal-01879947

HAL Id: hal-01879947

<https://hal.sorbonne-universite.fr/hal-01879947>

Submitted on 24 Sep 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

The variation of invariant graphs in forced systems

Bastien Fernandez, and Anthony Quas

Citation: *Chaos* **28**, 083101 (2018); doi: 10.1063/1.5026551

View online: <https://doi.org/10.1063/1.5026551>

View Table of Contents: <http://aip.scitation.org/toc/cha/28/8>

Published by the [American Institute of Physics](#)

Articles you may be interested in

[Asymmetry in initial cluster size favors symmetry in a network of oscillators](#)

Chaos: An Interdisciplinary Journal of Nonlinear Science **28**, 081101 (2018); 10.1063/1.5043588

[Epidemic spreading on metapopulation networks including migration and demographics](#)

Chaos: An Interdisciplinary Journal of Nonlinear Science **28**, 083102 (2018); 10.1063/1.5021167

[The dynamics of knowledge acquisition via self-learning in complex networks](#)

Chaos: An Interdisciplinary Journal of Nonlinear Science **28**, 083106 (2018); 10.1063/1.5027007

[Chaos in Kuramoto oscillator networks](#)

Chaos: An Interdisciplinary Journal of Nonlinear Science **28**, 071102 (2018); 10.1063/1.5041444

[Chaos in homeostatically regulated neural systems](#)

Chaos: An Interdisciplinary Journal of Nonlinear Science **28**, 083104 (2018); 10.1063/1.5026489

[Overcoming network resilience to synchronization through non-fast stochastic broadcasting](#)

Chaos: An Interdisciplinary Journal of Nonlinear Science **28**, 071104 (2018); 10.1063/1.5044420

The variation of invariant graphs in forced systems

Bastien Fernandez¹ and Anthony Quas²

¹Laboratoire de Probabilités, Statistique et Modélisation, CNRS—Université Paris 7 Denis Diderot—Sorbonne Université, 75205, Paris Cedex 13, France

²Department of Mathematics and Statistics, University of Victoria, Victoria, British Columbia V8W 3R4, Canada

(Received 21 February 2018; accepted 24 July 2018; published online 24 August 2018)

In skew-product systems with contractive factors, all orbits asymptotically approach the graph of the so-called sync function; hence, the corresponding regularity properties primarily matter. In the literature, sync function Lipschitz continuity and differentiability have been proved to hold depending on the derivative of the base reciprocal, if not on its Lyapunov exponent. However, forcing topological features can also impact the sync function regularity. Here, we estimate the total variation of sync functions generated by one-dimensional Markov maps. A sharp condition for bounded variation is obtained depending on parameters, which involves the Markov map topological entropy. The results are illustrated with examples. *Published by AIP Publishing.* <https://doi.org/10.1063/1.5026551>

To describe the properties of the long term response to a deterministic stimulus is a ubiquitous issue in Chaotic Dynamics. In the context of dissipative factors driven by autonomous systems, this question boils down to evaluating the characteristics of the so-called synchrony function. So far, focus has been made on sync function features that depend on the forcing derivative (in a broad sense). However, discontinuous examples in applications suggest the need to evaluate other basic features, such as the total variation. Here, we prove in simple examples that the sync function is, or is not, of bounded variation, depending only on the forcing topological entropy. Therefore, basic properties of invariant graphs also depend on topological features of the corresponding forcing systems.

I. INTRODUCTION

An important issue in the theory of forced systems is to evaluate the regularity of their synchronization graph. Suppose that an autonomous (discrete time) dynamical system

$$x^{t+1} = f(x^t), \quad t \in \mathbb{N} \cup \{0\},$$

is given together with a dissipative factor

$$y^{t+1} = g(x^t, y^t).$$

Here, f is an invertible map with invariant subset U in a Banach space X with norm $\|\cdot\|_X$ and $g : X \times Y \rightarrow Y$ (Y is a Banach space with norm $\|\cdot\|_Y$) such that

$$\sup_{x \in U} \|g(x, y) - g(x, y')\|_Y \leq \gamma \|y - y'\|_Y, \quad \forall y, y' \in Y,$$

for some $0 < \gamma < 1$. Then, the orbits $\{(x^t, y^t)\}$ of the skew-product system (f, g) are attracted by the graph of the corresponding sync function.^{2,17,19,22} This sync function, say, $\phi : U \rightarrow Y$, can be defined by the conjugacy equation

$$g[x, \phi(x)] = \phi \circ f(x), \quad \forall x \in U,$$

which ensures invariance of the corresponding graph $\{[x, \phi(x)]\}_{x \in U}$. In this context, regularity properties of this

function matter because they determine those dynamical characteristics of the drive system that carry over to the factor. For instance, Lipschitz continuity implies control of dimension estimates. Applications range from filtering of chaotic signals^{3,6,8} to damage detection in material science.¹⁸

The study of sync functions can be regarded as part of the analysis of inertial manifolds in dynamical systems.^{12,13} Besides existence and continuity, a standard result in this theory is the proof of differentiability under the condition

$$K\gamma < 1,$$

where $K = \sup_{x \in U} \|Df^{-1}(x)\|_X$, f is a diffeomorphism, and g is continuously differentiable.^{7,14,23,24} (If f^{-1} and g are merely Lipschitz continuous, then the same inequality—where K now stands for the Lipschitz constant of f^{-1} —implies that the function ϕ itself is Lipschitz continuous.) When $K\gamma \geq 1$, ϕ may not be differentiable/Lipschitz continuous;^{4,16} however, it is certainly Hölder continuous.^{1,7,23–25}

The analysis has subsequently been extended to accommodate non-uniformly hyperbolic effects. In particular, ϕ has been proved to be differentiable in the Whitney sense (i.e., given an invariant measure μ for f , ϕ is uniformly differentiable on sets of measure $1 - \epsilon$ for every $\epsilon > 0$) provided that^{7,24}

$$e^{\text{Lyap}(f^{-1})} \gamma < 1,$$

where $\text{Lyap}(f^{-1})$ is the maximum of the largest Lyapunov exponent of f^{-1} over a set of full measure μ and satisfies $e^{\text{Lyap}(f^{-1})} \leq K$.

Investigations have been pursued beyond the homeomorphic case, either when f is non-invertible^{5,21,25} or when it has discontinuities, more precisely, when f^{-1} is a discontinuous map of the interval.^{1,4,6} In this case, discontinuities transfer to the sync function, where they typically form dense subsets. A natural quantifier in this setting is the total variation which, roughly speaking, measures the length of the corresponding graph, see, e.g., Ref. 15 for a definition. The sync function has been shown to be of bounded variation under

FIG. 1. Schematic diagram of the properties of $\phi_\gamma = \sum_{k=0}^{+\infty} \gamma^k T^{k+1}$ for $0 < \gamma < 1$, when T is a continuous piecewise affine and expanding, and transitive Markov map of the interval.

the condition

$$e^{h_{\text{top}}(f^{-1})} \gamma < 1,$$

where $h_{\text{top}}(f^{-1})$ is the topological entropy of f^{-1} . On the other hand, the total variation of ϕ becomes infinite when γ is sufficiently large.¹⁰ A sync function of bounded variation makes it easier to show that the factor statistics inherit absolute continuity of the forcing statistics.¹¹ In any case, the topological feature of f^{-1} can also affect regularity properties of ϕ .

II. MAIN RESULTS

This letter aims to provide similar total variation estimates in more general cases, when the forcing is not necessarily discontinuous. For simplicity, we shall work in the framework of (one-dimensional) linear filters and skew-product inverse forcing (although certain results extend to more general cases without additional conceptual difficulties).^{3,6,9,25} More precisely, the set U can be written as $U = [0, 1] \times U_2$ (where U_2 will be irrelevant), and letting $x = (x, x_2)$ where $x \in [0, 1]$, the inverse forcing is given by

$$f^{-1}(x, x_2) = [T(x), f_2(x, x_2)],$$

where $T : [0, 1] \rightarrow [0, 1]$ (and, again, the mapping f_2 will play no role here). For a systematic way of defining f such that f^{-1} is as here, see Ref. 10. Assuming in addition that the filter satisfies

$$g(x, y) = x + \gamma y, \quad \forall y \in \mathbb{R} = Y, \quad x = (x, x_2) \in [0, 1] \times U_2,$$

the sync function then takes the following compact expression:

$$\phi_\gamma = \sum_{k=0}^{+\infty} \gamma^k T^{k+1}. \quad (1)$$

FIG. 2. Examples of maps satisfying the assumptions of Proposition 2.

A Markov map of the interval²⁰ is said to be transitive if the corresponding transition matrix is irreducible. The main result of this paper states that, for T a piecewise affine expanding and transitive Markov map, the total variation of ϕ_γ shows a sharp transition at $\gamma = e^{-h_{\text{top}}(T)}$.

Theorem 1. *Assume that T is a piecewise affine and expanding, and transitive Markov map. Then, the total variation of ϕ_γ defined by (1) on $[0, 1]$*

- *is finite when $e^{h_{\text{top}}(T)} \gamma < 1$ and*
- *is infinite when $e^{h_{\text{top}}(T)} \gamma > 1$, except maybe for at most $2N - 1$ values of γ , where N is the number of atoms of T .*

For maps T that are also continuous, this result can be combined with the previously mentioned ones to obtain the schematic diagram of the sync function properties presented in Fig. 1.

As the proof will show, the first part of the statement actually holds for every piecewise continuous and monotone map T of the interval and also for more general filters such as

$$g(x, y) = p(x) + \gamma y,$$

where p is assumed to be smooth. An example of such extension is $p(x) = \cos(2\pi x)$ and $T(x) = \beta x - \lfloor \beta x \rfloor$, where the sync function essentially boils down to the Weierstrass function¹⁶ (NB: in this case we have $K = e^{h_{\text{top}}(T)} = e^{L_{\text{yap}}(T)} = \beta$).

On the other hand, the proof of the second part strongly relies on the assumptions on T , although we believe that infinite variation occurs for sync functions associated with more general maps. Furthermore, we also believe that ϕ_γ has infinite variation for all $\gamma \geq e^{-h_{\text{top}}(T)}$, i.e., there are no exceptional values. This can be proved in some special cases (see examples in Fig. 2).

FIG. 3. Examples of the function ϕ_γ for the tent map with maximum at $x = a$ (left picture in Fig. 2) and $a = \frac{1}{4}$ here. Top left: $\gamma = 0.2$ is such that $K\gamma < 1$. Top right: $\gamma = 0.45$ (viz., $K\gamma < 1 < e^{h_{\text{top}}(T)}\gamma$). Bottom left: $\gamma = 0.55$ ($e^{h_{\text{top}}(T)}\gamma < 1 < e^{\text{Lyap}(T)}\gamma$). Bottom right: $\gamma = 0.75$ ($e^{\text{Lyap}(T)}\gamma > 1$).

Proposition 2. Assume that T is a piecewise affine and expanding, and transitive Markov map with a full branch and such that

- either $T(0+0) = 0$ or $T(1-0) = 1$,
- or there exist $\underline{a} < a < \bar{a}$ such that $T(\bar{a}) = T(\underline{a}) = a$ and either $T(a) = 0$ and $T(0) > 0$ or $T(a) = 1$ and $T(1) < 1$.

Then, the total variation of ϕ_γ on $[0, 1]$ is infinite when $e^{h_{\text{top}}(T)}\gamma \geq 1$.

To illustrate the results, examples of graphs of the sync function ϕ_γ , obtained with the tent map (left picture in Fig. 2), are presented in Fig. 3.

III. PROOFS

Proof of Theorem 1. In order to prove the first assertion, consider an arbitrary subdivision $0 = x_1 < x_2 < \dots < x_n = 1$ of $[0, 1]$. Uniform convergence in the definition of ϕ_γ implies

$$\sum_{i=1}^{n-1} |\phi_\gamma(x_{i+1}) - \phi_\gamma(x_i)| \leq \sum_{k=0}^{+\infty} \gamma^k \sum_{i=1}^{n-1} |T^{k+1}(x_{i+1}) - T^{k+1}(x_i)|.$$

Using that $\sum_{i=1}^{n-1} |T^{k+1}(x_{i+1}) - T^{k+1}(x_i)| \leq \text{Var}_{[0,1]} T^{k+1}$ and taking the supremum over all subdivisions then yield

$$\text{Var}_{[0,1]} \phi_\gamma \leq \sum_{k=0}^{+\infty} \gamma^k \text{Var}_{[0,1]} T^{k+1}.$$

Now, given $k \geq 0$, let n_k be the number of affine branches of T^k . Since each branch has variation at most 1 and the jump between branches is at most 1, we have $\text{Var}_{[0,1]} T^k \leq 2n_k$ and the first assertion follows from the fact that for a Markov map, we have $h_{\text{top}}(T) = \lim_{k \rightarrow +\infty} \frac{\log n_k}{k}$.²⁰

For the second assertion, let $\{I_i\}_{i=1}^N$ be the intervals on which T is affine and let s_i be the slope on I_i . Let X be

the subshift of finite type consisting of admissible sequences of elements of $\mathcal{A} = \{1, 2, \dots, N\}$ and let M be its transition matrix.

Given a real function f defined on an interval I , define the reduced variation of f on I by

$$\text{RV}_I(f) = \inf_a \text{Var}_I[f(x) - ax].$$

The reduced variation satisfies the following properties:

- $\text{RV}_I(f) = \text{RV}_I(f + g)$ for any affine function g on I ;
- $\text{RV}_I(\alpha f) = \alpha \text{RV}_I(f)$ for every $\alpha \in \mathbb{R}^+$;
- $\text{RV}_{[a,c]}(f) \geq \text{RV}_{[a,b]}(f) + \text{RV}_{[b,c]}(f)$ for every $a < b < c$;
- $\text{RV}_I(f) = 0$ if and only if f is affine on I .

Lemma 3. Assume that T is a piecewise affine and transitive Markov map and let $\gamma > e^{-h_{\text{top}}(T)}$. If $\text{Var}_{[0,1]} \phi_\gamma < +\infty$, then ϕ_γ must be affine on each I_i .

Proof. Let $v_i = \text{RV}_{I_i}(\phi_\gamma)$. The relation $\phi_\gamma = T + \gamma \phi_\gamma \circ T$, the above properties, the Markov property, and the fact that the branches are linear imply

$$\begin{aligned} v_i &= \text{RV}_{I_i}(T + \gamma \phi_\gamma \circ T) = \gamma \text{RV}_{I_i}(\phi_\gamma \circ T) \\ &\geq \gamma \sum_{i \rightarrow j} \text{RV}_{I_i \cap T^{-1}(I_j)}(\phi_\gamma \circ T) = \gamma \sum_{i \rightarrow j} v_j. \end{aligned}$$

That is, $v \geq \gamma Mv$ and $v = \{v_i\}_{i=1}^N$ is a non-negative vector. Since order is preserved, we can iterate this inequality giving $v \geq (\gamma M)^n v$ for all $n \in \mathbb{N}$. The assumption $\gamma > e^{-h_{\text{top}}(T)}$ implies that γM is a non-negative irreducible matrix with spectral radius greater than 1. According to the Perron-Frobenius Theorem, the only non-negative v satisfying $v \geq \gamma Mv$ is $v = 0$, viz., ϕ_γ must be affine on each I_i . $\square$

It remains to show that there are at most finitely many values of γ such that ϕ_γ is affine on each interval I_i . To that goal, we separate the proof into two cases:

- (1) there exist $j \in \{1, \dots, N\}$, $x, y \in I_j$, and $n \in \mathbb{N}$ such that $T'[T^n(x)] \neq T'[T^n(y)]$,
- (2) there exists $n \in \mathbb{N}$ such that T^n contains a discontinuity inside some I_i .

One of these conditions must always hold. Indeed, if all iterates T^n have constant derivatives on each I_i , then some iterates T^n must be discontinuous inside some I_i (otherwise, T being piecewise expanding, the length $|T^n(I_i)|$ of a sufficiently large iterate would be larger than 1, which is impossible).

Case (1). We may wlog assume that n is the smallest integer such that $T'[T^n(x)] \neq T'[T^n(y)]$. Since T is piecewise affine, there must exist $z \in (x, y) \cap T^{-n}(a)$, where a is an atom boundary point for which the left and right derivatives $T'(a-0) \neq T'(a+0)$ differ. The Markov assumption implies that the orbits of boundary points must be eventually periodic. Moreover, we may also assume that T is continuous at all iterates of a ; otherwise, case (2) applies. Therefore, there exist $p, q \geq 0$ such that $p + q \leq N + 1$ and an atom boundary point b so that

$$T^q(a) = b \quad \text{and} \quad T^p(b) = b.$$

Applying repeatedly the relation $\phi_\gamma = T + \gamma\phi_\gamma \circ T$ to successive iterates, one gets

$$\begin{aligned} \phi_\gamma(z) &= \sum_{k=0}^{n-2} \gamma^k T^{k+1}(z) \\ &+ \gamma^{n-1} \left[\sum_{k=0}^{q-1} \gamma^k T^k(a) + \frac{\gamma^q}{1-\gamma^p} \sum_{k=0}^{p-1} \gamma^k T^k(b) \right]. \end{aligned}$$

All iterates T^k involved in the RHS are piecewise affine; hence, the left and right derivatives $(T^k)'(\cdot - 0)$ and $(T^k)'(\cdot + 0)$ are well-defined at every point in the interior of $(0, 1)$. These derivatives can be extended by continuity to $[0, 1]$ [viz., $T'(0 - 0) := T'(0 + 0)$ and $T'(1 + 0) := T'(1 - 0)$]. Assuming that ϕ_γ is affine on the Markov partition, we have $(\phi_\gamma)' = T' + \gamma(\phi_\gamma)' \circ T$ giving a recurrence relation for the values of the derivative on the intervals of the Markov partition. Since $z \pm 0$ are pre-periodic, we can iterate and solve linear equations for $(\phi_\gamma)'(z \pm 0)$, yielding rational expressions for the left and right derivatives $(\phi_\gamma)'(z - 0)$ and $(\phi_\gamma)'(z + 0)$ that are similar to that for $\phi_\gamma(z)$ above.

That n is the smallest integer implies $(T^k)'(z - 0) = (T^k)'(z + 0)$ for $k = 1, \dots, n - 1$. It follows that the equation $(\phi_\gamma)'(z - 0) = (\phi_\gamma)'(z + 0)$ amounts to a polynomial one, with degree at most $2N - 1$, and which is non-degenerate since $T'(a - 0) \neq T'(a + 0)$. Therefore, except for at most $2N - 1$ values of γ , we have $(\phi_\gamma)'(z - 0) \neq (\phi_\gamma)'(z + 0)$ and then ϕ_γ cannot be affine on I_j , as desired.

Case (2). The argument is similar to the previous case. Let x in the interior of some I_i and a discontinuity a of T be such that $T^n(x) = a$ for some $n \in \mathbb{N}$. Wlog, we can assume that T is continuous at each $T^k(x)$ for $k = 1, \dots, n - 1$. Then we have

$$|\phi_\gamma(x - 0) - \phi_\gamma(x + 0)| = \gamma^n |\phi_\gamma(a - 0) - \phi_\gamma(a + 0)|.$$

Let $a_- = T(a - 0)$ and $a_+ = T(a + 0)$. Again, the orbits of a_-, a_+ must be eventually periodic. More precisely, there exist $q_-, p_- \geq 0$, $q_- + p_- \leq 2N$, $\{a_{-,k}\}_{k=1}^{q_- - 1}$, and $\{b_{-,k}\}_{k=0}^{p_- - 1}$ such that

$$\phi_\gamma(a - 0) = a_- + \sum_{k=1}^{q_- - 1} a_{-,k} \gamma^k + \frac{\gamma^{q_-}}{1 - \gamma^{p_-}} \sum_{k=0}^{p_- - 1} b_{-,k} \gamma^k$$

and similarly

$$\phi_\gamma(a + 0) = a_+ + \sum_{k=1}^{q_+ - 1} a_{+,k} \gamma^k + \frac{\gamma^{q_+}}{1 - \gamma^{p_+}} \sum_{k=0}^{p_+ - 1} b_{+,k} \gamma^k.$$

Similar to before, the equation $\phi_\gamma(a - 0) = \phi_\gamma(a + 0)$ amounts to a non-degenerate polynomial one since $a_- \neq a_+$. The polynomial degree is at most $\max\{q_-, q_+\} + p_- + p_+ - 1 \leq 2N - 1$ if $p_- \neq p_+$ (resp. $\max\{q_-, q_+\} + p_+ - 1 \leq 2N - 1$ if $p_- = p_+ = p$). Hence, except for at most $2N - 1$ values of γ , ϕ_γ cannot be affine on I_i . The proof of Theorem 1 is complete. $\square$

Proof of Proposition 2. Assume first that $T(0 + 0) = 0$. By assumption, there exists $a > 0$ such that T is continuous and expanding on $(0, a)$, $T : (0, a) \mapsto [0, T(a)]$, and $T(x) > x$ for $x \in (0, a)$. Therefore, any $x \in [0, T(a)]$ has a pre-image $y \in T^{-1}(x)$ such that $y < x$.

By transitivity, the periodic orbits of T are dense in $[0, 1]$. Given $p > 1$, let $(x_i)_{i=1}^p$ be one such orbit for which $x_1 = \min_i x_i \in [0, T(a)]$. Labelling in $(x_i)_{i=1}^p$ has been chosen so that $x_{i+1} = T(x_i)$ for $i = 1, \dots, p - 1$ and $T(x_p) = x_1$. Let $x_0 < x_1$ be the pre-image of x_1 in $(0, a)$.

Let $n \in \mathbb{N}$ and consider the pre-images $x_{0,i,n} \in T^{-n}(x_0)$, $x_{1,i,n} \in T^{-n}(x_1)$, and $x_{p,i,n} \in T^{-n}(x_p)$ by the same affine branch of T^n . Transitivity and the fact that T has a full branch ensure their existence for every branch of T^n , provided that n is large enough.

For $k = 1, \dots, n - 1$, the iterates $T^{k+1}(x_{0,i,n}), T^{k+1}(x_{1,i,n}), T^{k+1}(x_{p,i,n})$ lie in the same atoms. Hence, there exist $c_{i,n} \in \mathbb{R}$ such that we have

$$\begin{aligned} \phi_\gamma(x_{0,i,n}) - \phi_\gamma(x_{1,i,n}) &= \sum_{k=0}^{n-1} \gamma^k [T^{k+1}(x_{0,i,n}) - T^{k+1}(x_{1,i,n})] \\ &+ \gamma^n \sum_{k=0}^{+\infty} \gamma^k [T^{k+1}(x_0) - T^{k+1}(x_1)] \\ &= c_{i,n}(x_{0,i,n} - x_{1,i,n}) + \gamma^n [\phi_\gamma(x_0) - \phi_\gamma(x_1)], \end{aligned}$$

and $T(x_0) = x_1$ implies $\phi_\gamma(x_0) - \phi_\gamma(x_1) = x_1 - (1 - \gamma)\phi_\gamma(x_1)$. Similarly, we have

$$\begin{aligned} \phi_\gamma(x_{p,i,n}) - \phi_\gamma(x_{1,i,n}) &= c_{i,n}(x_{p,i,n} - x_{1,i,n}) \\ &+ \gamma^n [x_1 - (1 - \gamma)\phi_\gamma(x_1)]. \end{aligned}$$

Since $x_0 < x_1 < x_p$, the point $x_{1,i,n}$ must lie inside the interval delimited by $x_{0,i,n}$ and $x_{p,i,n}$. Hence, the first terms in these expressions of $\phi_\gamma(x_{0,i,n}) - \phi_\gamma(x_{1,i,n})$ and $\phi_\gamma(x_{p,i,n}) - \phi_\gamma(x_{1,i,n})$ must be of the opposite sign. This implies that

$$\max\{|\phi_\gamma(x_{0,i,n}) - \phi_\gamma(x_{1,i,n})|, |\phi_\gamma(x_{p,i,n}) - \phi_\gamma(x_{1,i,n})|\} \geq \gamma^n |x_1 - (1 - \gamma)\phi_\gamma(x_1)|.$$

Now, there are $N_n \simeq e^{nh_{\text{top}}(T)}$ triplets $x_{0,i,n}, x_{1,i,n}, x_{p,i,n}$, one for each cylinder of length $n + 1$, i.e., each element of $\bigcap_{k=0}^n T^{-k}([0, 1])$. Therefore, we have

$$\text{Var} \phi_\gamma|_{[0,1]} \geq N_n \gamma^n |x_1 - (1 - \gamma)\phi_\gamma(x_1)|,$$

and to ensure the claim, it suffices to show that $x_1 \neq (1 - \gamma)\phi_\gamma(x_1)$.

Using that $(x_i)_{i=1}^p$ is p -periodic, one gets

$$\phi_\gamma(x_1) = \frac{x_2 + \gamma x_3 + \dots + \gamma^{p-1} x_p + \gamma^p x_1}{1 - \gamma^p}$$

and the assumptions $x_1 = \min_i x_i$ and $p > 1$ imply

$$x_2 + \gamma x_3 + \dots + \gamma^{p-1} x_p + \gamma^p x_1 > \sum_{k=0}^p \gamma^k x_1$$

from where it follows that $(1 - \gamma)\phi_\gamma(x_1) > x_1$. This concludes the proof in the case $T(0 + 0) = 0$. The case $T(1 - 0) = 1$ can be treated identically.

Finally, in the case where there exist $\underline{a} < a < \bar{a}$ such that $T(\underline{a}) = T(\bar{a}) = a$ and either $T(a) = 0$ and $T(0) > 0$ or $T(a) = 1$ and $T(1) < 1$, a reasoning similar to the previous proof can be applied. The final condition for infinite variation becomes $0 - (1 - \gamma)\phi_\gamma(0) \neq 0$ [resp. $1 - (1 - \gamma)\phi_\gamma(1) \neq 0$].

0], which certainly holds because $\phi_\gamma(0) \geq T(0) > 0$ [resp. $\phi_\gamma(1) \leq T(1) + \frac{\gamma}{1-\gamma} < 1 + \frac{\gamma}{1-\gamma}$]. $\square$

- ¹V. S. Afraimovich, J.-R. Chazottes, and A. Cordonet, "Synchronization in directionally coupled systems: some rigorous results," *Discrete Cont. Dyn. Syst. B* **1**, 421–442 (2001).
- ²V. S. Afraimovich, N. N. Verichev, and M. I. Rabinovich, "Stochastic synchronization of oscillations in dissipative systems," *Quantum Electron.* **29**, 795–803 (1986).
- ³R. Badii, G. Broggi, B. Derighetti, M. Ravani, S. Ciliberto, A. Politi, and M. A. Rubio, "Dimension increase in filtered chaotic signals," *Phys. Rev. E* **60**, 979–982 (1988).
- ⁴R. Badii and A. Politi, "Dimension function and phase transition-like behavior in strange attractors," *Phys. Scr.* **35**, 243–246 (1987).
- ⁵E. Barreto, K. Josić, C. J. Morales, E. Sanders, and P. So, "The geometry of chaos synchronization," *Chaos* **13**, 151–164 (2003).
- ⁶D. S. Broomhead, J. P. Huke, and M. R. Muldoon, "Linear filters and nonlinear systems," *J. R. Stat. Soc. B* **53**, 373–382 (1992).
- ⁷K. M. Campbell and M. E. Davies, "The existence of inertial functions in skew product systems," *Nonlinearity* **9**, 801–817 (1996).
- ⁸T. L. Carroll, "Detecting recursive and nonrecursive filters using chaos," *Chaos* **20**, 013123 (2010).
- ⁹M. E. Davies and K. M. Campbell, "Linear recursive filters and nonlinear dynamics," *Nonlinearity* **9**, 487–499 (1996).
- ¹⁰B. Fernandez, "Discontinuous generalized synchronization of chaos," *Dyn. Syst.* **27**, 105–116 (2012).
- ¹¹B. Fernandez and A. Quas, "Statistical properties of invariant graphs in piecewise affine discontinuous forced systems," *Nonlinearity* **24**, 2477–2488 (2011).
- ¹²M. Hirsch and C. Pugh, "Stable manifolds and hyperbolic sets in Global Analysis," *Am. Math. Soc. Proc. Symp. Pure Math.* **14**, 133–163 (1970).
- ¹³M. Hirsch, C. Pugh, and M. Shub, *Invariant Manifolds*, Lecture Notes in Mathematics Vol. 583 (Springer, 1977).
- ¹⁴B. Hunt, E. Ott, and J. A. Yorke, "Differentiable generalized synchronization of chaos," *Phys. Rev. E* **55**, 4029–4034 (1997).
- ¹⁵A. N. Kolmogorov and S. V. Fomin, *Elements of the Theory of Functions and Functional Analysis* (Dover, 1999).
- ¹⁶J. A. Kaplan, J. Mallet-Paret, and J. A. Yorke, "The Lyapunov dimension of a nowhere differentiable attracting torus," *Ergod. Theory Dyn. Syst.* **4**, 261–281 (1984).
- ¹⁷L. Kocarev and U. Parlitz, "Generalized synchronization, predictability and equivalence in unidirectionally coupled dynamical systems," *Phys. Rev. Lett.* **76**, 1816 (1996).
- ¹⁸J. M. Nichols, S. T. Trickey, M. Seaver, and L. Moniz, "Use of fiber-optic strain sensors and Hölder exponents for detecting and localizing damage in an experimental plate structure," *J. Intell. Mater. Syst. Struct.* **18**, 51–67 (2007).
- ¹⁹L. M. Pecora and T. L. Carroll, "Synchronization in chaotic systems," *Phys. Rev. Lett.* **64**, 821 (1990).
- ²⁰M. Pollicott and M. Yuri, *Dynamical Systems and Ergodic Theory* (Cambridge University Press, 1998).
- ²¹N. F. Rulkov and V. S. Afraimovich, "Detectability of nondifferentiable generalized synchrony," *Phys. Rev. E* **67**, 066218 (2003).
- ²²N. F. Rulkov, M. M. Sushchik, L. S. Tsimring, and H. D. I. Abarbanel, "Generalized synchronization of chaos in directionally coupled chaotic systems," *Phys. Rev. E* **51**, 980–994 (1996).
- ²³J. Stark, "Invariant graphs for forced systems," *Physica D* **109**, 163–179 (1997).
- ²⁴J. Stark, "Regularity of invariant graphs for forced systems," *Ergod. Theory Dyn. Syst.* **19**, 155–199 (1999).
- ²⁵J. Urías, "Filters display inverse limit spaces," *Chaos* **14**, 963–968 (2004).