Tracer diffusion in crowded narrow channels

O. Bénichou, P. Illien, G. Oshanin, A. Sarracino, R. Voituriez

► To cite this version:

O. Bénichou, P. Illien, G. Oshanin, A. Sarracino, R. Voituriez. Tracer diffusion in crowded narrow channels. *Journal of Physics: Condensed Matter*, 2018, 30 (44), pp.443001. 10.1088/1361-648X/aae13a . hal-01908770

HAL Id: hal-01908770

<https://hal.sorbonne-universite.fr/hal-01908770>

Submitted on 30 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

TOPICAL REVIEW

Tracer diffusion in crowded narrow channels

To cite this article: O Bénichou *et al* 2018 *J. Phys.: Condens. Matter* **30** 443001

View the [article online](#) for updates and enhancements.

IOP | ebooks™

Bringing you innovative digital publishing with leading voices
to create your essential collection of books in STEM research.

Start exploring the **collection** - download the first chapter of
every title for free.

Topical Review

Tracer diffusion in crowded narrow channels

O Bénichou¹, P Illien², G Oshanin¹ , A Sarracino^{3,4} and R Voituriez^{1,5}¹ Sorbonne Université, CNRS, Laboratoire de Physique Théorique de la Matière Condensée (UMR 7600),

4 Place Jussieu, 75252 Paris Cedex 05, France

² ESPCI Paris, PSL Research University, UMR Gulliver 7083, 10 rue Vauquelin, 75005 Paris, France³ ISC-CNR and Dipartimento di Fisica Università Sapienza Roma, Piazzale Aldo Moro 2, 00185 Rome, Italy⁴ Dipartimento di Ingegneria, Università della Campania 'Luigi Vanvitelli', 81031 Aversa (CE), Italy⁵ Sorbonne Université, CNRS, Laboratoire Jean Perrin (UMR 8237), 4 Place Jussieu, 75005 Paris, FranceE-mail: benichou@lptmc.jussieu.fr, pierre.illien@espci.fr, oshanin@lptmc.jussieu.fr, ale.sarracino@gmail.com and voituriez@lptmc.jussieu.fr

Received 19 May 2018, revised 3 September 2018

Accepted for publication 13 September 2018

Published 10 October 2018

Abstract

We summarise different results on the diffusion of a tracer particle in lattice gases of hard-core particles with stochastic dynamics, which are confined to narrow channels—single-files, comb-like structures and quasi-one-dimensional channels with the width equal to several particle diameters. We show that in such geometries a surprisingly rich, sometimes even counter-intuitive, behaviour emerges, which is absent in unbounded systems. This is well-documented for the anomalous diffusion in single-files. Less known is the anomalous dynamics of a tracer particle in crowded branching single-files—comb-like structures, where several kinds of anomalous regimes take place. In narrow channels, which are broader than single-files, one encounters a wealth of anomalous behaviours in the case where the tracer particle is subject to a regular external bias: here, one observes an anomaly in the temporal evolution of the tracer particle velocity, super-diffusive at transient stages, and ultimately a giant diffusive broadening of fluctuations in the position of the tracer particle, as well as spectacular multi-tracer effects of self-clogging of narrow channels. Interactions between a biased tracer particle and a confined crowded environment also produce peculiar patterns in the out-of-equilibrium distribution of the environment particles, very different from the ones appearing in unbounded systems. For moderately dense systems, a surprising effect of a negative differential mobility takes place, such that the velocity of a biased tracer particle can be a non-monotonic function of the force. In some parameter ranges, both the velocity and the diffusion coefficient of a biased tracer particle can be non-monotonic functions of the density. We also survey different results obtained for a tracer particle diffusion in unbounded systems, which will permit a reader to have an exhaustively broad picture of the tracer diffusion in crowded environments.

Keywords: lattice gases, stochastic dynamics, tracer diffusion, narrow channels

(Some figures may appear in colour only in the online journal)

Contents

1. Introduction	2
2. Tracer diffusion in single-files, on infinite combs and on infinite (unbounded) lattices of spatial dimension $d \geq 2$..	3
2.1. Unbiased tracer diffusion.....	3
2.1.1. Single-files.....	3
2.1.2. Comb-like structures.....	4
2.1.3. Unbounded lattices with $d \geq 2$	5
2.2. Biased tracer diffusion.....	6
2.2.1. Single-files.....	6
2.2.2. Density profiles of the environment particles.....	8
2.2.3. Mean velocity of the tracer particle.....	8
2.2.4. Variance of the tracer particle displacement.....	9
3. Biased tracer diffusion in narrow channels.....	10
3.1. Density profiles of the environment particles.....	10
3.2. Mean velocity of a biased tracer particle.....	11
3.2.1. Transient mean velocity.....	11
3.2.2. Terminal mean velocity.....	11
3.2.3. General force-velocity relation.....	12
3.3. Variance of the biased tracer particle displacement: from super-diffusion to giant diffusion.....	12
4. Summary and outlook.....	14
Acknowledgments.....	14
References.....	14

1. Introduction

Transport of molecules and small particles in pores, channels, or other quasi-one-dimensional systems has attracted a great deal of attention within several last decades [1]. On the one hand, this interest stems from the relevance of the problem to a variety of realistic physical, biophysical and chemical systems, as well as important applications in nanotechnology and nanomedicine, e.g. in the creation of artificial molecular nanofilters. Few stray examples include, e.g. transport in porins [2–4], through the nuclear pores in eukaryotic cells [5–7], or along the microtubules [8] and dendritic spines [9], transport of microswimmers in narrow pores [10–12], translocation of polymers in pores [13–18] and their sequencing in nanopore-based devices [19], ionic currents across nanopores [20, 21], ionic liquids in supercapacitors [22–29], nano- and microfluidics [30–33] and transport of inertial particles advected by laminar flows [34–36]. On the other hand, the diversity and the significance of the problems emerging in this field represent a challenging area of research for theoreticians [37].

Molecular crowding has been recognised as an important factor (see, e.g. [38–53] and references therein), which strongly affects the behaviour of currents across the narrow channels, as well as the dynamics of individual molecules or probes, which move either due to thermal activation only, or are also subject to external constant forces. This latter case, i.e. dynamics

Figure 1. Tracer particle, subject to a regular constant force F pointing along the channel, in a confined bath of crowding particles.

of *biased* particles in crowded narrow channels is precisely the chief subject of this topical review. More specifically, we present here an overview of analytical and numerical work which deals with the behaviour observed in narrow channels—single-files, ramified single-files (comb-like structures) and quasi-one-dimensional channels with the width equal to several particle diameters—with rigid, structureless hard-walls, densely populated by a quiescent mixture of identical environment particles with purely repulsive short-ranged interactions. The environment particles undergo stochastic dynamics due to thermal noise resulting, e.g. from the interactions with the solvent present in the channel. In addition, there is a special tracer particle, which has the same size and the same interaction potential as the environment particles, but is also subject to a regular constant force F pointing along the channel (see figure 1). As one may notice, these settings mimic the standard experimental set-up of the so-called *active* micro-rheology [31, 54–61] and hence, the tracer particle can be thought of as the probe designed to measure, at a molecular scale, the response of the confined passive molecular crowding environment.

The channel geometry on which we concentrate here is given by two particular examples (see figure 2)—2D ideal strips, such as ones printed on a substrate and used in microfluidic devices [31], and 3D ideal capillaries; both types of systems are macroscopically large in the direction of the applied bias, and in the perpendicular to F direction have dimensions which are *comparable* to the diameters of the environment particles. The walls of the channels are perfect and do not contain any impurities or geometric corrugations, and the cross-section is constant across the channel. We note that a large amount of an available theoretical work concerns lattice systems with stochastic dynamics and interactions between the tracer particle and the environment particles, as well as between the environment particles themselves, being a mere hard-core exclusion. In realistic systems, of course, the particles evolve in continuous-space and the interactions may extend beyond the nearest-neighbouring particles, for example because of hydrodynamic or electrostatic interactions. This simplification, however, permits to unveil some universal features of the dynamical behaviour beyond the linear in the force F regime. Whenever possible, we also compare the theoretical predictions against the results of numerical Monte Carlo or molecular dynamics simulations.

The purpose of this review is three-fold. First, we propose a complete and comprehensive picture of essentially

Figure 2. Two-dimensional strip-like lattice with width $L = 3$ and a 3D, ‘capillary’-like simple cubic lattice with side $L = 3$.

non-equilibrium, cooperative phenomena emerging in crowded narrow channels in response to the passage of a biased intruder. We proceed to show that there is a large number of spectacular effects and some of them take place beyond the regime of the linear response. Strikingly, the behaviour appears to be distinctly different in extremely narrow channels—the so-called single-files and in somewhat broader channels, in which the particles can bypass each other. Second, we note that physical situations involving a single particle pulled by a constant force through a host medium composed of interacting mobile particles represents the typical settings for studying the linear-response Einstein relation (see, e.g. recent [62–64]), which links the diffusion coefficient of this particle in the absence of the force, and the mobility in the presence of the latter. We comment on the validity of the Einstein relation throughout the review and show that, curiously enough, it may be associated with some subdominant contributions to the dynamics. Lastly, as it has been already understood for narrow channels in which all the particles experience a constant bias—the so-called driven diffusive systems (see, e.g. [38–41]), the explicit treatment of the multiple degrees of freedom of the environment is of an essential importance. Here, as well, we proceed to show that many interesting phenomena take place due to the interplay between these degrees of freedom and the dynamics of the tracer particle. These effects are overlooked once one represents the environment in an integrated, effective form and studies a Markovian Langevin dynamics of the tracer particle in an effective environment.

2. Tracer diffusion in single-files, on infinite combs and on infinite (unbounded) lattices of spatial dimension $d \geq 2$

To set up the scene, we recall here some most prominent results on the tracer particle (TP) diffusion in interacting particles systems. We do not consider as our goal to provide an exhaustive review and to pay a due tribute to all the outstanding contributors to this field—this would require too much space. Instead, we merely list below some very significant achievements relevant to the subject of our review, which were spread across many different disciplines and journals within the last five decades. This will permit the reader to have a somewhat broader picture. In this way, as well, many remarkable and novel phenomena specific to the tracer diffusion in narrow channels will be made more apparent.

2.1. Unbiased tracer diffusion

It was understood for a long time that diffusion of a TP in a dynamical background formed by other interacting and

mobile (randomly moving) particles is coupled in a non-trivial way to the evolution of the environment itself (see, e.g. [65–67]). Clearly, in a sufficiently dense system, when a TP displaces in one unit of time over one unit of length away from its location occupied at the previous unit of time, it leaves a void of a clear space—a vacancy—behind it. Then, it is often more probable that the TP will return back to this location, than keep on going away from it where its motion will be hindered by other particles. This ‘anti-persistence’ of the TP motion and the circumstance whether it will be ‘permitted’ by the environment particles to return back depends essentially on how fast the environment can rearrange itself and close the void. In turn, this depends on plenty of physical factors—the density of the environment, the particle-particle interactions, the temperature and the viscosity of the embedding solvent, if any. Hence, the TP diffusion coefficient is expected to acquire some dependence on all the aforementioned parameters.

2.1.1. Single-files. The single-file concept was introduced first in biophysical literature [68] to describe diffusion through pores in membranes, which are so narrow such that the initial order of particles is preserved at all times. In mid 60 s it was realised that in single-files the effect of interactions with the environment can be even stronger than a mere renormalisation of the diffusion coefficient—it was shown that the single-file constraint can change the very temporal evolution of the TP mean-squared displacement $\overline{X_t^2}$. Analysing the TP dynamics in a single-file of interacting diffusions, it was discovered in [69] that $\overline{X_t^2}$ obeys, in the asymptotic limit $t \rightarrow \infty$, a slower than diffusive law, $\overline{X_t^2} \sim \sqrt{t}$, i.e. $\overline{X_t^2}$ exhibits not a linear Stokes–Einstein but an anomalous sub-diffusive growth with time. The point is that in single-files—an extreme case of narrow channels—due to the constraint that the environment particles cannot bypass each other, in order to explore a distance X_t the TP has to involve in a cooperative motion $\sim \rho X_t$ particles of the environment, with ρ being their mean density⁶. In consequence, the effective frictional force exerted by the medium on the TP also scales with the distance travelled by the TP, which entails a sub-diffusive motion⁷. In a way, this resembles (and in fact, it is linked on the level of the underlying mathematics) dynamics of a tagged bead in an infinitely long Rouse polymer chain; in order to explore progressively

⁶ See, e.g. the discussion after equation (2.22) in [70] for more details.

⁷ The situation is, of course, different in ‘single-files’ in contact with a vapour phase such that the environment particles can desorb from and re-adsorb back to the channel. In this case the order is not preserved and the TP undergoes a standard diffusive motion with, however, the diffusion coefficient dependent on all the system’s parameters [71, 72].

longer and longer spatial scales the tagged bead has to involve in motion more and more other beads of the chain [73].

This remarkable result has been subsequently re-derived in different settings by using a variety of analytical techniques [74–82] (see also recent [83, 84] for a more extensive review) to show that the exact leading asymptotic behaviour of $\overline{X_t^2}$ obeys

$$\overline{X_t^2} \underset{t \rightarrow \infty}{\sim} \frac{(1-\rho)a^2}{\rho} \sqrt{\frac{2}{\pi} \frac{t}{\tau}}, \quad (1)$$

where a is the lattice spacing and τ is the mean waiting time between jumps. When $\rho \rightarrow 1$, (i.e. the space available for diffusion shrinks), the right-hand-side of equation (1) vanishes, since the system becomes completely blocked. In turn, in the limit $\rho \rightarrow 0$ the right-hand-side of equation (1) diverges, meaning that a faster growth has to take place—in this limit, of course, one expects the standard Brownian motion result $\overline{X_t^2} \sim t$ to hold. For arbitrarily small, but finite ρ , the result in equation (1) will describe the ultimate long-time behaviour, while the diffusive law will appear as a transient.

The time dependence of equation (1) was also observed experimentally (see, e.g. [85–88] and also [83, 84]), which is not surprising given its universal nature. The TP dynamics in single-files still represents a challenging playground for testing different analytical techniques and probing other properties: convergence of the distribution of X_t to a Gaussian [78], explicit form of the distribution of X_t at arbitrary, not necessarily large times in dense systems [89], the TP dynamics in the presence of a slower-than-diffusive environment [90], large deviations properties [91–95] and emerging correlations [70, 96–100]. Surprisingly, some unexpected features of this seemingly exhaustively well-studied process keep on being revealed: it was shown recently [101] (see also [92, 102, 103]) that the preservation of the initial order implies, in fact, an *infinite* memory of the process on the specific initial conditions—in systems, in which the particles of the environment are initially out-of-equilibrium (also called a ‘quenched’ initial condition), the growth of the mean-squared displacement of the TP proceed a factor of $\sqrt{2}$ slower, than in systems with an equilibrium (also called ‘annealed’ or uniform) initial distribution of the environment particles⁸. In [92] such a dichotomy in dynamics under annealed and quenched initial conditions has been also shown to persist for higher-order cumulants of the TP position, for which it becomes even more striking.

2.1.2. Comb-like structures. Before we pass from single-files to the behaviour on unbounded regular lattices of dimension d higher than 1, it seems relevant to focus on some geometrically ‘intermediate’ situation, in which a variety of anomalous diffusions emerge. Recently [105] studied the dynamics of a TP in the presence of hard-core environment particles on a comb-like lattice, i.e. a 1D backbone (see figure 3) connected at each site to a tooth—an infinite 1D lattice, with passages of particles between the teeth being allowed only along the backbone. Dynamics of a single isolated particle in such a

Figure 3. Unbiased tracer particle (red filled circle) on a comb-like lattice populated by hard-core environment particles (blue filled circles). The x axis is a backbone, whereas the orthogonal lines are the infinitely long teeth of the comb.

system has been widely studied in the past as a toy model of dynamics in a geometrically disordered system [106]. It is well known that the mean-squared displacement of a single TP (in the absence of the environment particles) obeys, at sufficiently long times, $\overline{X_t^2} \sim \sqrt{t}$ [106, 107], i.e. the TP moves sub-diffusively along the x -axis due to progressively longer and longer excursions on the teeth. [105] thus analysed a natural extension of this model by adding stochastically moving environment particles and considering two situations: (a) the environment particles can move everywhere, i.e. both along the backbone and along the teeth, while the TP can go along the backbone only, and (b) both the TP and the environment particles are permitted to go everywhere.

At the first glance, the case (a) seems to be very close to the situation discussed in [71, 72], i.e. a 1D lattice attached to a reservoir of particles. Hence, one may expect essentially the same ‘diffusive’ behaviour, i.e. $\overline{X_t^2} \sim Dt$, where the diffusion coefficient D captures the combined effect of the geometry and dynamics. In reality, the behaviour appears to be more complicated. Bénichou *et al* [105] focused on densely populated systems and analysed the TP mean-squared displacement in the leading in the density $\rho_0 = 1 - \rho$ of vacancies order, in the limit $\rho_0 \rightarrow 0$. It was shown that, when the vacancies are initially uniformly spread across the comb with mean density ρ_0 , the mean-squared displacement of the TP along the x -axis obeys, for times t sufficiently large but less than a certain large cross-over time $t_1 \sim 1/\rho_0^4 \ln^8(1/\rho_0)$:

$$\lim_{\rho_0 \rightarrow 0} \frac{\overline{X_t^2}}{\rho_0} = \frac{1}{2^{5/4} \Gamma(7/4)} t^{3/4}, \quad (2)$$

where $\Gamma(x)$ is the gamma-function, which crosses over for $t > t_1$ to the diffusive behaviour of the form

$$\lim_{t \rightarrow \infty} \frac{\overline{X_t^2}}{t} = \rho_0^2 \left(\ln \frac{1}{\rho_0} \right)^2. \quad (3)$$

The results in equations (2) and (3) reveal two interesting features: first, it is the existence of a transient regime with a sub-diffusive behaviour, which is more extended in time the denser the system is. Second, in the ultimate diffusive regime,

⁸Note that this result has been obtained earlier in [104], but was not explicitly mentioned in this work.

the diffusion coefficient of the TP exhibits a rather unusual non-analytic dependence on the density of vacancies.

Bénichou *et al* [105] described an interesting out-of-equilibrium situation when the vacancies are all initially placed on the backbone only, with a linear density $\rho_0^{(\text{lin})} \ll 1$. Interestingly enough, in this case the mean-squared displacement of the TP grows sub-linearly

$$\overline{X_t^2} = \frac{\rho_0^{(\text{lin})}}{2^{7/4}\Gamma(5/4)} t^{1/4}, \quad (4)$$

for any (sufficiently large) time t , without any crossover to an ultimate diffusive regime. We note, as well, that all *even* cumulants of the TP displacement follow the dependences in equations (2) and (3) in case of a uniform distribution of vacancies on the comb, and the dependence in equation (4) in case of their placement on the backbone only, which signals that the distribution of the TP position is a Skellam distribution (see, [105] for more details). This means that the distribution converges to a Gaussian at long times, with an appropriately rescaled TP position.

Lastly, in the case (b) when both the environment particles and the TP can go along the teeth, the following dynamical scenario has been predicted and confirmed through an extensive numerical analysis [105]. For an extended time interval $0 \ll t \ll t^{(1)} \sim 1/\rho_0^2$, one has $\overline{X_t^2} \sim t^{3/4}$, in which regime the TP has not had enough time to explore any given tooth because its excursions were hindered by the environment particles. This regime is followed, on the time interval $t^{(1)} \ll t \ll t^{(2)}$ (precise form of $t^{(2)}$ was not provided in [105]), by a rather unusual dependence $\overline{X_t^2} \sim t^{9/16}$, associated with the large- t tail of the corresponding distribution of the time spent by the TP on a given tooth. Ultimately, for $t \gg t^{(2)}$, one finds again $\overline{X_t^2} \sim t^{3/4}$. Note that the coincidence of the dynamical exponent 3/4 characterising the initial and the final stages is occasional; the underlying physics is completely different. Note, as well, that at all stages the growth of the TP mean-squared displacement proceeds faster than in case of a single isolated TP on an empty comb ($\overline{X_t^2} \sim t^{1/2}$). This is a consequence of interactions with the environment particles, which do not permit the TP to enter too often into the teeth and also to travel too far within each tooth.

2.1.3. Unbounded lattices with $d \geq 2$. On lattices of spatial dimension $d \geq 2$, the ultimate long-time dynamics of the TP is diffusive⁹, such that $\overline{X_t^2} = 2dDt$. As we have already remarked, here the main issue is the calculation of the effective diffusion coefficient which embodies the full dependence on both the rates of the TP and of the environment particles, their density and the type of the lattice (if any), on which the evolution takes place. It was fairly well understood that calculation of the diffusion coefficient is a genuine many-body problem, which is unsolvable in the general case and one has to resort to either density expansions or some other approximations, verified by numerics. Numerous approaches

have been developed and a large number of different results, both analytical and numerical, have been obtained (see, e.g. [111–121] for a few stray examples, [122] for some early review and [123] for a more recent summary). We focus below particularly on two approaches, proposed in [112] and [120, 121], respectively, which will be used in what follows for the analysis of the TP dynamics in narrow channels.

In [112], the self-diffusion constant of a tracer on regular lattices partially populated by identical hard-core particles has been analysed using an approximate approach, based on a perturbative expansion in powers of $\rho(1-\rho)$, which is exact at the two extrema of the volume fraction, $\rho = 0$ and $\rho = 1$. This approach gives, in particular, for an unbounded d -dimensional hyper-cubic lattice the following expression for the TP diffusion coefficient

$$D = \frac{(1-\rho)a^2}{2d\tau} f(\rho, \omega), \quad (5)$$

where $1/\tau$ is the jump frequency of the TP, a is the lattice spacing, and $\omega = \tau^*/\tau$ with $1/\tau^*$ being the characteristic jump frequency of the environment particles. Lastly, $f(\rho, \omega)$ is the correlation factor which is explicitly given by

$$f(\rho, \omega) = \frac{(\omega(1-\rho) + 1)(1-\alpha)}{\omega(1-\rho) + 1 - \alpha(1 + \omega(1-3\rho))}, \quad (6)$$

with α being one of the Watson integrals [124]

$$\alpha = \frac{1}{(2\pi)^d} \int_{-\pi}^{\pi} dk_1 \dots \int_{-\pi}^{\pi} dk_d \frac{\sin^2(k_1)}{\sum_{j=1}^d (1 - \cos(k_j))}. \quad (7)$$

This latter factor relates D to the propagator of a special type of random walk on a d -dimensional hyper-cubic lattice. For $d = 1$, one has $\alpha \equiv 1$ and hence, D vanishes, as it should, since here we enter into the realm of a single-file diffusion. On comparing the analytical predictions against available at that time results of Monte Carlo simulations in $d = 3$ systems, the authors demonstrated that their approximation gives a good interpolation formula in between of the two extrema. In [125, 126] and [127], the expression in equation (5) has been generalised, using a similar approach based on a decoupling of the TP-particle-particle correlation function into a product of two TP-particle correlation functions, for situations when the density ρ of the environment particles is not explicitly conserved (unlike, e.g. the case of a monolayer in a slit-pore, confined between two solid surfaces) but the particles undergo continuous exchanges with a reservoir maintained at a constant chemical potential. References [125, 126] focused specifically on such a situation in two dimensions, i.e. a monolayer on top of a solid surface exposed to a vapour phase, while [127] presented an explicit expression for the TP diffusion coefficient in three-dimensions, which case represents some generalised model of a dynamic percolation. The resulting expressions for the diffusion coefficient are quite cumbersome and we do not present them here, addressing an interested reader to the original papers [125–127]. We note parenthetically that in case when exchanges with the vapour are permitted, dynamics becomes diffusive and D is well-defined [71, 72] even in single-files, since the initial order is destroyed.

⁹ See, however, [108–110] for the non-diffusive behaviour in case of an inhomogeneous crowding.

A different line of thought has been put forth in [120, 121], which studied the TP dynamics in a very densely populated system of hard-core particles, assuming that a continuous ‘reshuffling’ of the environment occurs due to the dynamics of the ‘vacancies’ (i.e. vacant sites, present on the lattice at mean density $\rho_0 = 1 - \rho$), rather than of the particles themselves. Focussing on such a ‘vacancy-assisted’ dynamical model on a 2D square lattice of a unit lattice spacing, the authors studied first the case of just a single vacancy, which moves in the discrete time $n, n = 0, 1, 2, \dots$, by exchanging its position, at each tick of the clock, with one of the neighbouring particles (including the TP, once it appears on the neighbouring site), chosen at random. In such a 2D model, any particle including the TP performs concerted random excursions and can, in principle, travel to infinity from its initial position. By counting all possible paths which bring the TP to position $\mathbf{X}$ at time moment n , the authors were able to calculate exactly the full probability distribution $P_n(\mathbf{X})$ of finding the TP at this very site at time moment n . Surprisingly, this distribution appears to be non-Gaussian even in the limit $n \rightarrow \infty$ and the mean-squared displacement $\overline{\mathbf{X}_n^2}$ shows a striking sub-diffusive behaviour in the leading in n order [120]

$$\overline{\mathbf{X}_n^2} = \frac{\ln(n)}{\pi(\pi - 1)}, \quad (8)$$

with a very non-trivial numerical amplitude (which depends, of course, on the lattice structure).

Brummelhuis and Hilhorst [121], focused on the case of a very small density of vacancies. Discarding the events when any two vacancies appear at the adjacent sites (whose probability is of order of ρ_0^2), an analogous expression for $P_n(\mathbf{X})$ has been derived, which does converge to a Gaussian in the limit $n \rightarrow \infty$. The TP dynamics in this case appears to be diffusive, since now the TP displacement is a sum of many independent events. In consequence, the mean-squared displacement, at sufficiently large n , obeys

$$\overline{\mathbf{X}_n^2} = \frac{\rho_0}{\pi - 1} n, \quad (9)$$

with, again, a very non-trivial numerical amplitude, which depends on the lattice structure. We emphasise that this result is only valid in the linear in the density of vacancies ρ_0 order. There exist, of course, ρ_0 -dependent corrections to the diffusion coefficient. Expanding the expression in equation (5) in Taylor series in powers of ρ_0 and retaining only the leading term in this expansion, one may straightforwardly verify that it coincides with the expression in equation (9).

The result in equation (9) has been subsequently generalised in [128] for the situations where the TP interacts with the environment particles in a different way, than the particles interact between themselves, which situation mimics, e.g. dynamics of an intruder (e.g. an Indium atom) in the close-packed upper most layer of atoms in a metal (e.g. Cu), in the presence of one or a few naturally existing defects of packing—the vacancies [129, 130]. The expression derived in [128] revealed an interesting non-monotonic dependence of the self-diffusion coefficient on the strengths of these interactions. Vacancy-assisted dynamics on different types of lattices has been amply discussed in [131–136].

By assuming the validity of the Einstein relation, a generalisation of the expression in equation (9) for a hyper-cubic lattice of an arbitrary dimension d can be obtained from the expression for the TP mobility (see the supplementary information to [137]). This gives the following expression

$$D = \frac{\rho_0}{2d} \frac{1 - \alpha}{1 + \alpha}, \quad (10)$$

where α is defined in equation (7), $\tau^* = \tau = 1$ and $a = 1$. We note that equation (10) coincides with the leading term in the expansion in Taylor series in powers of ρ_0 of the expression in equation (5).

2.2. Biased tracer diffusion

We turn to the situations when the particles of the environment still experience no other regular force except for random thermal ones, but the TP is subject to some regular constant force F pointing along the X -axis. It should be emphasised that the presence of a constant bias exerted on the TP only does not merely add another dimension to the parameter space, but crucially changes the dynamical behaviour in the system under study. Within the context of the aforementioned experimental technique—micro-rheology [31, 54–60]—the situations described in section 2.1 correspond to the so-called *passive* micro-rheology, when the TP, as well as the environment particles move solely due to thermal forces and the system is in thermal equilibrium. On the contrary, when the TP is biased by an external force, the situation corresponds to the typical settings of an *active* micro-rheology. Here, the TP entrains in motion the environment particles appearing in its vicinity, which produces micro-structural changes in the host medium and brings it out of equilibrium. This means that the system becomes characterised by highly asymmetric stationary density profiles of the environment particles, in the frame of reference moving with the TP. Except for the single-files, which still exhibit a very particular dynamics (i.e. the TP creeps, instead of moving ballistically, $\overline{X_t} \sim \sqrt{t}$, and the density profiles around it do not converge to a stationary form), in higher dimensions the TP attains a constant velocity V which results from an interplay between the jamming of the medium, produced by the TP, and diffusive re-arrangements of the environment. Various aspects of this problem have been studied experimentally (see, e.g. [138–140]), using a combination of numerical simulations and theoretical approaches (see, e.g. [141, 142]) and by extensive numerical simulations (see, e.g. [143–145]). An inverse problem in which the TP is kept fixed (e.g. by an optical tweezer), while the environment particles are subject to a constant force has also been rather extensively studied [146–149]. Below we present some most important results for single-files and unbounded, infinite in all direction systems, obtained for the models of lattice gases with stochastic dynamics.

2.2.1. Single-files. A model with a biased TP travelling in a lattice gas of particles with symmetric hopping probabilities has been studied in [150] in the particular limit when the force is infinitely large, such that the TP performs a totally directed

random walk in the direction of the applied bias, constrained by hard-core interactions with the environment particles. It was shown in [150] that when the TP jumps instantaneously to the neighbouring lattice site on its right, as soon as it becomes vacant, and does not jump in the opposite direction, its mean displacement obeys at sufficiently long times

$$\bar{X}_t = \gamma \sqrt{2D_0 t}, \quad (11)$$

where D_0 is the Stokes–Einstein diffusion coefficient a single isolated environment particle and the amplitude γ is defined implicitly as the solution of the following transcendental equation

$$\gamma I_+(\gamma) = \rho_0, \quad I_+(\gamma) = \sqrt{\frac{\pi}{2}} \exp\left(\frac{\gamma^2}{2}\right) \left(1 - \operatorname{erf}\left(\frac{\gamma}{\sqrt{2}}\right)\right), \quad (12)$$

with $\operatorname{erf}(x)$ being the error function. Remarkably, the TP does not move ballistically, i.e. does not have a constant velocity, but rather creeps. This happens because the displacement of the TP becomes effectively hindered by the environment particles; at each step to the right, the TP ‘pumps’ a vacancy to the left, such that the density of the environment particles in the phase in front of the TP effectively grows. In turn, there appears a sort of a traffic jam in front of the TP, which also grows in size in proportion to the distance X_t travelled by the TP. This means that the system is out of equilibrium (in contrast to the situations described in section 2.1) at any time, the density profile of the environment is a step-like function propagating to the right away of the TP such that more and more of the environment particles are entrained by the TP in a directed motion. Viewing X_t as a random process described by a Langevin equation, one may argue [150] that in this situation there is an effective frictional force exerted on the TP due to the appearance of the propagating traffic jam in front of it, which grows in proportion to X_t .

In [151, 152] a more complicated situation has been analysed when the force F exerted on the TP is finite, such that it may perform jumps in both directions¹⁰. It was shown in [151, 152] that the TP mean displacement still has a form in equation (11), but now the amplitude γ is determined by a more complicated transcendental equation

$$\left(\gamma I_+(\gamma) + \frac{e^{-\beta F} - \rho_0}{1 - e^{-\beta F}}\right) \left(\gamma I_-(\gamma) + \frac{1 - \rho_0 e^{-\beta F}}{1 - e^{-\beta F}}\right) = \frac{\rho_0^2 e^{-\beta F}}{(1 - e^{-\beta F})^2}, \quad (13)$$

where β is the reciprocal temperature and

$$I_-(\gamma) = \sqrt{\frac{\pi}{2}} \exp\left(\frac{\gamma^2}{2}\right) \left(1 + \operatorname{erf}\left(\frac{\gamma}{\sqrt{2}}\right)\right). \quad (14)$$

References [151, 152] have also demonstrated that the density profiles of environment particles in front of and past the TP attain a stationary form in variable $x = (X - \bar{X}_t)/\bar{X}_t$ characterised by a dense region in front of the TP and a depleted region in its wake. In consequence, in the laboratory frame, the environment particles progressively accumulate in front of

the TP, and the size of the depleted region past the TP, which is devoid of particles, also grows progressively in time. Overall, the biased TP entrains in a directed motion the entire system, which is quiescent in the absence of the external bias. This is quite evident for the particles in front of the TP, but less trivial for the particles in its wake. The explanation is, however, rather simple and relies on one of the results presented in [78]; namely, it was shown that in case of an asymmetric layout of the environment particles around an unbiased TP, such that they are initially present past the TP and are absent in front of it, the mean displacement of the TP follows $\bar{X}_t = \sqrt{t \ln(t)}$, i.e. an unbiased TP moves faster than $\sqrt{t}$ due to an additional logarithmic factor, which stems from the pressure exerted on the TP by the particle phase behind it. This means that in case of a symmetric placement of the environment particles around the biased TP, once the TP goes away from the phase behind it, the closest to the TP particle sees the void space in front of it and starts to move as $\sqrt{t \ln(t)}$ catching eventually the TP, which moves as $\sqrt{t}$.

The results in equations (11) and (13) permit to make the following, conceptually important conclusion: define a generalised, time-dependent mobility μ_t of the TP in the presence of the force as

$$\mu_t = \lim_{F \rightarrow 0} \frac{\bar{X}_t}{F t}, \quad (15)$$

and the time-dependent diffusivity D_t in the absence of the external force as $D_t = \bar{X}_t^2/2t$. Our aim is now to check if the Einstein relation $\mu_t = \beta D_t$ holds. Note that, first, for standardly defined mobility and the diffusion coefficient, which requires going to the limit $t \rightarrow \infty$, the Einstein relation trivially holds since $0 = 0$. A legitimate and non-trivial question is if it holds as well for time-dependent μ_t and D_t . Calculating D_t from equation (1) and μ_t —from equations (11) and (13), one realises that the Einstein relation holds exactly in the case when the initial distribution of the environment particles is random [151, 152], and does not hold when the latter is regular [101], which is a non-trivial result. The validity of the Einstein relation in single-file systems has been also investigated via molecular dynamics simulations in models of granular (inelastic) particles in [158, 159].

Within the last decade many other characteristic features of biased TP diffusion in single-file systems have been studied. In particular, the mean displacement of a biased TP has been analysed in case when the initial distribution of the environment particles has a shock-like profile: a higher density of the environment particles from the left from the TP than from the right of the TP, and the TP is subject to a constant force F pointing towards the higher density phase, (which mimics an effective attraction towards this phase) [160]. It was shown that $\bar{X}_t$ obeys equation (11) in which the amplitude γ is determined implicitly as the solution of a quite complicated transcendental equation (see, [160]). Interestingly enough, the TP not always moves in the direction of the force but only when the latter exceeds some threshold value F_c . For $F < F_c$, the TP moves against the force and the high density phase expands. When the force is exactly equal to this threshold value, $\gamma = 0$ such that the mean displacement vanishes; in this particular

¹⁰ See also [153] for some applications in biophysics, [154–156] for a relevant analysis within the harmonisation approximation and [157] for the solution in a similar mathematical set-up.

case, the mean-squared displacement is not zero, $\overline{X_t^2} \sim \sqrt{t}$ with a prefactor dependent on the densities in both phases (see, [160]).

In recent [161] dynamics of a pair of biased TPs in a single-file has been analysed in case when they experience action of the forces pointing in the opposite directions, such that the forces tend to separate the TPs. It was shown that in this situation, however, the TPs do indeed travel in the opposite directions only when the forces exceed some critical value; otherwise, the pair of TPs remains bounded.

The full distribution of the TP position as well as its cumulants have been calculated exactly for an arbitrary time in the leading order in the density of vacancies ρ_0 [89], upon an appropriate generalisation of the approach developed in [121]. The TP position distribution has been also determined in case of an arbitrary particles density within a certain decoupling approximation [162, 163], for the situation when exchanges of particles with a reservoir are permitted. It was shown that in the asymptotic limit $t \rightarrow \infty$ the distribution converges to a Gaussian. Interestingly enough, it was realised that in the presence of a bias the diffusion coefficient can be a non-monotonic function of the environment particles density ρ . Lastly, the emerging correlations in such systems have been analysed in [164, 165].

We close this subsection by noting that the models with a biased TP evolving in a single-file of particles with symmetric hopping probabilities appear also in different contexts (seemingly unrelated to the TP diffusion). In particular, they describe the dynamics of the front of a propagating precursor film which emanates from a spreading liquid droplet [166–170] or a time evolution of a boundary of a monolayer on a solid surface [171].

2.2.2. Density profiles of the environment particles. On unbounded lattices of spatial dimension $d \geq 2$, the biased TP attains ultimately a constant velocity V along the direction of the force, whose dependence on system's parameters is rather non-trivial and will be discussed below. At the same time, the TP alters the environment, involving in a directional motion some of the environment particles and hence, drives the environment out of equilibrium—it is no longer homogeneous and some density profiles emerge. In the frame of reference moving with the TP, these density profiles attain a steady-state form meaning that, (in contrast to the case of single-files), here the entrainment is only partial, in the sense that upon an encounter with the TP an environment particle travels alongside the TP for some random time, leaving it afterwards and being replaced by another particle. The total amount of the entrained environment particles stays constant, on average, in time.

The emerging steady-state density profiles have been studied in detail in [125–127] for square and simple cubic lattices. It was realised that, in general, in front of the TP and in the direction perpendicular to the force, the approach of the perturbed (local) density to its average value is exponential, with the characteristic length dependent on the coordinates, such that they are asymmetric. In the wake of the TP, the form of

these profiles depends essentially on whether one deals with the situation in which the number of the environment particles is explicitly conserved, or the system is exposed to some vapour phase and the environment particle may desorb from and re-adsorb into the system—in this latter case the particles number fluctuates in time around some average value. Then, the local density past the TP approaches the unperturbed value exponentially fast, but the characteristic length protrudes over much larger scales than in front of the TP. The situation is very different in the conserved number case. Here, the local density approaches its unperturbed value as a power-law, meaning that the environment remembers the passage of the TP over very large temporal and spatial scales. The approach of the local density $\rho(\lambda)$ to the mean density ρ obeys [125–127]

$$\rho - \rho(\lambda) \sim \frac{C}{\lambda^{(d+1)/2}}, \quad (16)$$

where λ is the distance past the TP. The amplitude C in this decay law has been also calculated in [125–127].

We close this subsection with the following two remarks:

First, we note that such a spectacular behaviour of the density of the environment particles in the wake of a biased TP has been also seen in various continuous space settings (see, e.g. [138, 142, 143, 145, 172]), in granular media [173] and also in colloidal experiments (see, e.g. [139, 147]).

Second, we note that an interesting phenomenon may take place in situations where there is not a single biased TP, but there are a few or even a concentration of them. The point is that an out-of-equilibrium environment may mediate long-ranged mutual interactions between intruders, which are non-reciprocal and violate Newton's third law (see, e.g. [174, 175]). When several TPs are present, each of them will perturb the distribution of the environment particles in its vicinity and in order to minimise the micro-structural changes in the environment and to reduce the dissipation, they will start to move collectively. Stochastic pairing of two biased TPs has been observed in simulations in a model of a 2D lattice gas [176], in experiments in colloidal suspensions [177, 178], and also theoretically predicted for the relative motion of two TPs in a nearly-critical fluid mixture, due to emerging critical Casimir forces [179]. Formation of string-like clusters of TPs, or 'trains' of TPs, which reveals an emerging effective attraction between them has been evidenced for situations with a small concentration of TPs [180–182]. Lastly, in case when the concentration of the TPs is comparable to that of the environment particles, the TPs arrange themselves in lanes [183–187]. This self-organisation of the TPs resembles spontaneous lane formation in binary mixtures of oppositely charged colloids (see, e.g. [188]), in dusty complex plasmas [189] and also for such seemingly unrelated systems as pedestrian counter flows [190].

2.2.3. Mean velocity of the tracer particle. The functional dependence of the TP ultimate velocity V on physical parameters has been studied in details within the last two decades. In the most general case, these parameters are the magnitude of the external force, the density of the environment, the rates of the particles exchanges with a vapour phase, and the jump

rates of both the TP and of the environment particles. Several analytical approaches have been used in this analysis—an analytical technique based on a decoupling of correlations between the tracer particle and two given environment particles (see, [125–127]), and also a different approach based on an appropriate generalisation of the vacancy-assisted dynamical model of [120, 121], in which a bias exerted on the TP has been taken into account explicitly [191].

References [125–127] studied the functional dependence of V on all physical parameters, including the exchange rates of the environment particles with the vapour phase (which define the value of the density ρ), in the general case of unequal jump rates of the TP and of the environment particles. The resulting expressions appear very cumbersome, and we address the reader to the original papers. Here we merely mention some features of the expressions obtained in [125–127]:

First, in the limit of very small forces, V attains a physically meaningful form

$$V = \xi^{-1}F, \quad (17)$$

which can be thought of as an analogue of the Stokes formula with ξ being the friction coefficient. This signifies that the frictional force exerted on the TP by the environment is *viscous* at small values of F . In turn, the friction coefficient can be decomposed into two contributions [125–127]: a mean-field one, proportional to an effective density of voids, and the second one which has a more complicated form and stems from the cooperative behaviour emerging in such a system—de-homogenisation of the environment by the TP and formation of some asymmetric density profiles of the environment particles around the TP, which we will discuss later on.

Second, it was observed in [125–127] that in the systems with an explicitly conserved particle density, the mobility of the TP exactly equals βD with D defined in equation (5) meaning that the Einstein relation holds.

Third, we note that the expressions obtained in [125–127] have been re-examined recently in [192] and it was realised that for the environments which evolve on time scales larger than the one which defines the TP jump rates, the terminal velocity V is a *non-monotonic*¹¹ function of the force F . The velocity first increases with an increase of F , as dictated by the linear response theory, reaches a maximal value and then, upon a further gradual increase of F , starts to decrease. In consequence, the differential mobility of the TP, defined beyond the linear response regime, becomes negative. In [192], a full phase chart has been presented indicating the region in the parameter space in which such a non-monotonic behaviour emerges. We note that the phenomenon of the so-called negative differential mobility is a direct consequence of an out of equilibrium dynamics. Various facets of this intriguing phenomenon have been studied in the past and it has been also observed in many realistic physical systems [34, 35, 193–203].

A different approach to calculation of the terminal velocity of a biased TP in very densely crowded environments has been pursued in [191], which generalised the analytical technique

originally developed in [120, 121], over the case when the TP is subject to an external constant force F . Bénichou and Oshanin [191] first focused on the case of a single vacancy-mediated dynamics evolving in discrete time n and calculated exactly the mean displacement of a biased TP on an infinite square lattice to get

$$\bar{X}_n = \frac{1}{\pi} \frac{\sinh(\beta F/2)}{(2\pi - 3) \cosh(\beta F/2) + 1} \ln(n), \quad (18)$$

i.e. the TP displaces anomalously slowly (logarithmically) with time. Accordingly, the time-dependent mobility μ_n (see equation (15)) of the biased TP obeys

$$\mu_n = \frac{\beta}{4\pi(\pi - 1)} \frac{\ln(n)}{n}. \quad (19)$$

Determining from equation (8) the effective diffusivity of the TP in the absence of the force, $D_n = \bar{X}_n^2/4n$, one arrives at a surprising conclusion [191]: the Einstein relation $\mu_n = \beta D_n$ holds exactly even for such an anomalously confined diffusion! The full distribution function of the TP position has been also determined in [191] and it was shown that it *does not* converge to a Gaussian as $n \rightarrow \infty$, similarly to the case without an external bias [120].

Bénichou and Oshanin [191] extended this analysis over the case when vacancies are present on a square lattice at very small density ρ_0 , to find that in the lowest order in the density of vacancies the mean displacement of the biased TP obeys

$$\bar{X}_n = \frac{\sinh(\beta F/2)}{(2\pi - 3) \cosh(\beta F/2) + 1} \rho_0 n, \quad (20)$$

such that the TP now moves ballistically with a finite velocity $V = \bar{X}_n/n$. Inspecting the mobility of the TP and the diffusion coefficient of the TP in equation (9), one realises that the Einstein relation holds [191]. The distribution of X_n has also been determined in [191] and it was shown that the latter converges to a Gaussian as $n \rightarrow \infty$.

In more recent [137], the result in equation (20) has been generalised, using essentially the same technique as in [191], to calculate the mean velocity of a biased TP on a d -dimensional hypercubic lattice (including the simple cubic one, $d = 3$) in the presence of a small density of vacancies. The result of [137] (see the supplementary materials to this paper) reads

$$V = a_0 \rho_0, \quad a_0 = \frac{(1 - \alpha) \sinh(\beta F/2)}{(d(1 + \alpha) - 1 + \alpha) \cosh(\beta F/2) + 1 - \alpha}, \quad (21)$$

where the parameter α is defined in equation (7). For $d = 2$, equation (21), multiplied by n , reduces to the expression in equation (20). For $d = 1$, the parameter α becomes equal to 1, such that V vanishes, as it should.

2.2.4. Variance of the tracer particle displacement. We concentrate on the time-evolution and the density dependence of the variance σ_x^2 of a biased TP displacement in the direction of the force F , $\sigma^2 = \bar{X}_t^2 - \bar{X}_t^2$, on a 2D square and a 3D simple cubic lattices. This question has been raised only recently and several rather surprising results have been obtained. Note that

¹¹ See also section 3.2.3 below for the analogous behaviour of the velocity versus force in strip-like geometries.

in the absence of the force F , the variance simply becomes $\sigma_x^2 = 2dDt$, where the forms of the diffusion coefficient have been discussed above.

We start with the case of a square-lattice, densely populated by the environment particles, and a biased TP, and focus on the vacancy-assisted dynamics in discrete time n , which was put forth in [120, 121]. The first observation of a ‘strange’ behaviour of the variance σ_x^2 has been made in [204], in which it was predicted analytically and verified numerically that $\sigma_x^2 \sim n \ln(n)$. Surprisingly, it appeared that in a dense system the spread of fluctuations in the TP position is (weakly) super-diffusive. This question has been further inspected in [205] and the physical mechanism responsible for such a super-diffusive growth of fluctuations has been explained—it was shown that it emerges due to correlations between the successive jumps of the TP, which originate from interactions with a single vacancy returning to the TP position many times and carrying it in the direction of the force.

Deeper analysis of this model has been presented in [137], and it was shown that this super-diffusive regime is only a transient one which prevails up to times of order of $1/\rho_0^2$. Not only the leading term but also the correction terms to the leading behaviour have been calculated, to provide the following exact expression

$$\lim_{\rho_0 \rightarrow 0} \frac{\sigma_x^2}{\rho_0} \sim a_0 n \left[\frac{2a_0}{\pi} \left(\ln(n) + \ln 8 + C - 1 + \frac{\pi^2(5-2\pi)}{(2\pi-4)} \right) + \coth\left(\frac{\beta F}{2}\right) \right], \quad (22)$$

where C is the Euler–Mascheroni constant, the parameter a_0 is defined in equation (21) and the omitted terms do not depend on n , i.e. are unimportant for sufficiently large n . Recalling that $a_0 \sim F$ for small forces, equation (22) implies that the coefficient in front of the leading super-diffusive term is proportional to F^2 , meaning that super-diffusion emerges beyond the linear response. Note that using equation (21) at small F and equation (22) at $F = 0$, the Einstein relation is shown to be satisfied.

For times greater than $1/\rho_0^2$, the weakly super-diffusive regime crosses over to a standard diffusive growth of the variance, which emerges due to renewal events—arrivals of other vacancies to the location of the TP—which fade out the memory about interactions with a single vacancy and make successive jumps of the TP uncorrelated. In this ultimate regime, one finds that in the leading in ρ_0 order the variance obeys [137]

$$\lim_{n \rightarrow \infty} \frac{\sigma_x^2}{n} \sim a_0 \rho_0 \left[\coth\left(\frac{\beta F}{2}\right) + 2a_0 \left(\frac{2}{\pi} \ln\left(\frac{1}{a_0 \rho_0}\right) + \frac{1}{\pi} \ln 8 + \frac{\pi(5-2\pi)}{2\pi-4} \right) \right]. \quad (23)$$

Here, again, one observes that the effective diffusion coefficient acquires some additional terms which are quadratic with the force in the limit of small forces. Note that for moderate forces, however, these terms can become large due to the

factor $\ln(1/\rho_0)$. The validity of the Einstein relation in this ultimate regime is again ensured by the term $a_0 n \coth(\beta F/2)$.

For the TP dynamics on a simple cubic lattice, densely populated by the environment particles, the growth of the variance of the TP displacement obeys at all time scales [137]

$$\lim_{\rho_0 \rightarrow 0} \frac{\sigma_x^2}{\rho_0} \sim a_0 n \left[\coth\left(\frac{\beta F}{2}\right) + A a_0 \right], \quad (24)$$

where A is a numerical constant. Hence, the spread of fluctuations in three- (and actually, higher-) dimensions is diffusive but the effective diffusion coefficient contains, as well, quadratic with the force terms in the limit of small applied forces. The validity of the Einstein relation is again ensured by the term $a_0 n \coth(\beta F/2)$.

Lastly, we discuss the observations made in [206], which analysed the density-dependence of the variance of the TP displacement in a system of the environment particles evolving on a discrete lattice of an arbitrary dimension in continuous time t . The approach of [206] was based on a decoupling of the TP-particle-particle correlation function, similar to the one employed in [125–127], and the results were verified numerically and in the high- and low-density of particles limits, in which exact solutions are available [137, 207–209]. In particular, it was shown in [206] that in the ultimate regime ($t \rightarrow \infty$) the effective diffusion coefficient can be a non-monotonic function of the density, for forces which exceed a certain threshold value, well above the linear response regime, and is a monotonic function of ρ for F below the threshold. Moreover, in the case when the jump rate of the environment particles is substantially lower than the one characterising the jumps of the TP, this effective diffusion coefficient can be also a non-monotonic function of the force F —a behaviour we have already observed for the velocity of a biased TP.

3. Biased tracer diffusion in narrow channels

We turn eventually to the dynamics of a biased TP in narrow channels—strip-like or capillary-like lattices depicted in figure 2. These lattices have width L , (with L being an integer), in case of a strip, and a cross-section $L \times L$ in case of a capillary, and are of an infinite extent in the direction of the force F acting on the TP only. As above, we will focus on the density profiles of the environment particles around a steadily moving TP, its velocity V , and the variance σ_x^2 of the TP displacement at time t . We proceed to show that in such confined geometries the time-evolution of the system exhibits many remarkable features, which are absent on infinite lattices with $d \geq 2$.

3.1. Density profiles of the environment particles

The density profiles of the environment particles around a steadily moving TP in strip-like and capillary-like geometries have been studied analytically, using a decoupling of the TP-particle-particle correlation functions into a product of two TP-particle correlation functions, and also numerically in recent [210, 211]. It was realised that, similarly to the case of lattices which have an infinite extent in all directions, these

density profiles attain a stationary form in the frame of reference moving with the TP. These profiles are asymmetric and are characterised by a dense, traffic jam-like region in front of the TP and a depleted by the environment particles region in the wake of the TP. The traffic jam-like region is more pronounced than in an infinite space [211], which is not counter-intuitive. This means that the frictional force exerted by the host medium on the TP is bigger and hence, the velocity is smaller in strips and capillaries than on infinite square and simple cubic lattices. The density, as a function of the distance from the TP in the direction perpendicular to the force, is described by an exponential function with the characteristic length that depends in a rather complicated way on the width L of the strip or the cross section of a capillary. When $L \rightarrow \infty$, the results in [125–127] are recovered. The form of the stationary density profiles past the TP is more complicated: there is some characteristic length scale λ^* , $\lambda^* \sim VL^2/D$, where V is the TP terminal velocity and D —the diffusion coefficient of the environment particles, such that at distances $\lambda < \lambda^*$ the evolution of the density profile in the wake of the TP is described by a power-law, equation (16), where $d = 2$ for strips and $d = 3$ for the capillaries. Beyond this distance, the approach of the density to its unperturbed value ρ becomes exponential, with the characteristic length dependent on L and diverging when $L \rightarrow \infty$. This overall behaviour has been confirmed numerically for lattice systems using Monte Carlo simulations [210, 211] and also for off-lattice, strip-like geometries with hard-core particles with Langevin dynamics [211]. Returning to the discussion of the entrainment properties, here, as in infinitely large systems, we encounter a *partial* entrainment of the environment particles by the biased TP but the amount of the entrained particles is bigger.

To close this subsection, we briefly mention recent observations made in [212], in which dynamics of several biased TPs in presence of the environment particles in strip-like geometries has been analysed via extensive molecular dynamics simulations. It was observed that for sufficiently large driving forces, a strong clogging of the channel takes place—the biased TPs cluster together such that the particles of the traffic jam region in front of them cannot circumvent the cluster. In consequence, the cluster of the TPs propagates in proportion to $\sqrt{t}$, i.e. the velocity of each individual TP vanishes, such that one is led back to an effectively single-file behaviour (see section 2.2.1).

3.2. Mean velocity of a biased tracer particle

In this subsection we present available analytical and numerical results on the mean velocity V of a biased TP in strip-like and capillary-like geometries. We note that the time-evolution of the TP displacement shows in such a confinement a bit richer behaviour than on infinite lattices with spatial dimension $d \geq 2$. A salient feature here is an emergence of a plateau-like behaviour at intermediate time scales.

3.2.1. Transient mean velocity. Before we proceed to the discussion of the forms of V in narrow channels, we briefly recall one of the results of [121], which concerns the form of the

diffusion coefficient D_L for self-diffusion of an unbiased TP on an infinitely long strip of width L , densely populated by the environment particles. In [121], within the model with the vacancy-assisted dynamics, the following exact, in the first order in the density ρ_0 of vacancies, result has been presented

$$D_L = \frac{\rho_0}{4} \frac{1 - \alpha'}{1 + \alpha'}, \quad (25)$$

where

$$\alpha' = \frac{1}{4} \lim_{z \rightarrow 1^-} [P(0|0; z) - P(2\mathbf{e}_x|0; z)], \quad (26)$$

with $\mathbf{e}_x$ being a unit vector in the positive x -direction and $P(\mathbf{r}|0; z)$ —the generating function associated with the propagator of a symmetric random walk on a 2D strip of width L , starting at the origin and ending at position $\mathbf{r} = (n_1, n_2)$,

$$P(\mathbf{r}|0; z) = \frac{1}{2\pi L} \sum_{k=0}^{L-1} \int_{-\pi}^{\pi} dq \frac{e^{-in_1 q - 2i\pi k n_2 / L}}{1 - z(\cos q + \cos(2\pi k/L)) / 2}. \quad (27)$$

Note that α' here has essentially the same meaning as α in equation (7), except that now it is defined for confined geometries.

This situation has been re-examined in [137] in case of a TP biased by a constant external force, to show that for both 2D strips ($d = 2$) and 3D capillaries ($d = 3$) the velocity of the TP is given, in the same order in the density of vacancies, by

$$\tilde{V} = \rho_0 a'_0, \quad a'_0 = \frac{(1 - \alpha') \sinh(\beta F/2)}{(1 + (2d - 1)\alpha') \cosh(\beta F/2) + 1 - \alpha'}, \quad (28)$$

where α' is defined in equation (26) (with the factor $1/4$ replaced by $1/(2d)$) and with $P(\mathbf{r}|0; z)$ given in equation (27) in case of 2D strips, and by a bit more complicated expression describing the propagator of a random walk in case of 3D capillaries (see [137] for more details). One may directly verify, upon comparing the diffusion coefficient defined in equation (25) and the mobility deduced from equation (28) with $d = 2$, that the Einstein relation holds.

3.2.2. Terminal mean velocity. A surprising observation has been made subsequently in [213], in which it was realised that $\tilde{V}$ in equation (28) does not describe the true velocity of the biased TP in the limit $n \rightarrow \infty$, but only a constant velocity appearing at some transient stage, for times less than the cross over time $t_x \sim 1/\rho_0^2$ (which is however quite large when $\rho_0 \ll 1$). The point is that in the underlying derivation the limit ρ_0 is taken first and then the analysis focuses on the asymptotic behaviour in the limit $n \rightarrow \infty$. It appears, however, that due to some very subtle circumstances these limits do not commute and, in essence, the velocity $\tilde{V}$ in equation (28) corresponds to the temporal regime when the TP interacts with just a single vacancy. Recall that we have already encountered this phenomenon while describing the behaviour of the variance of the TP displacement on an infinite square lattice. The key difference consists in the fact that for a *fixed* small ρ_0 the random walk performed by any of the vacancies between two successive visits to the TP position is a *biased* random walk, in the frame of reference moving with the TP, due to the

interactions of the TP with other vacancies. In consequence, to get the expression for the ultimate, terminal velocity V , one has to examine the general expressions taking the limit $n \rightarrow \infty$ first, and only then to concentrate on the leading behaviour in the limit $\rho_0 \rightarrow 0$. In doing so, it was shown in [213] that for $n \gg 1/\rho_0^2$ the true terminal velocity obeys

$$V = \rho_0 a_0'', \quad \frac{1}{a_0''} = \frac{1}{a_0'} + \frac{2d}{1 - \alpha'} \frac{1}{L^{d-1}}, \quad (29)$$

where a_0' is defined in equation (28) and α' —in equation (26).

The exact expression in equation (29) permits to draw several important conclusions, which all were also confirmed in [213] by extensive Monte Carlo simulations:

- in strips and capillaries with a finite L the true terminal velocity is lower than the velocity $\tilde{V}$ at the transient stage;
- this effect does not exist on infinite lattices when $L \rightarrow \infty$, even for $d = 2$ when the variance shows two distinct behaviours;
- the effect is more pronounced in strip-like geometry than in the capillaries;
- it requires quite long times ($\sim 1/\rho_0^2$) to observe the terminal velocity;
- the jump of the velocities $\tilde{V} - V \sim F^2$ for small forces F , meaning that this anomaly takes place beyond the linear response regime;
- the latter also ensures that the Einstein relation holds on both transient and terminal stages.

Lastly, the analysis in [213] culminated at the derivation of a complete expression for the time evolution of the TP mean displacement, which is valid at an arbitrary discrete time n :

$$\frac{\bar{X}_n}{\rho_0 n} \sim g\left((a_0'')^2 \rho_0^2 n\right), \quad (30)$$

with

$$g(\tau) = a_0' \left[\frac{b}{b^2 - 1} \left(\operatorname{erf}(\sqrt{\tau}) + \frac{e^{-\tau}}{\sqrt{\pi\tau}} \right) + \frac{b}{2} \frac{b^2 + 1}{(b^2 - 1)^2} \frac{\operatorname{erf}(\sqrt{\tau})}{\tau} - \frac{1}{b^2 - 1} + \frac{1}{\tau} \left(\frac{b}{b^2 - 1} \right)^2 \left(e^{(b^2 - 1)\tau} \operatorname{erf}(b\sqrt{\tau}) - 1 \right) \right], \quad (31)$$

where $b = a_0'/a_0'' - 1$. This scaling function reproduces correctly both regimes and has been numerically verified in [213] for both strip-like and capillary-like geometries.

3.2.3. General force-velocity relation. General force-velocity relation for a biased TP moving in crowded strip-like or capillary-like geometries, with an arbitrary density ρ of the environment particles and different characteristic jump times of the TP, τ , and of the environment particles, τ^* , has been derived in [210] using the decoupling approximation. In [210], the terminal velocity has been obtained as an implicit solution of a rather complicated non-linear equation involving matrix determinants, which we do not present here. We merely depict in figure 4 the curves V versus F for a particular case of strip-like geometries with the width $L = 3$, for several densities ρ of the environment particles, fixed characteristic

jump-time of the TP, $\tau = 1$, and two different jump-times of the environment particles, $\tau^* = 1$ and $\tau^* = 10$. We observe that for the former case, when the characteristic jump-times of the TP and of the environment particles are equal to each other, the terminal velocity V is a monotonic function of the force F , which was also noticed in [211]. Conversely, in case of a slowly varying environment, e.g. for $\tau^* = 10$, the velocity V exhibits a strongly non-monotonic behaviour. For small F , a linear dependence $V \sim F$ is observed, as dictated by the linear response, then, V reaches a peak value V^* and then gradually decreases upon a further increase of F . This means that the negative differential mobility phenomenon takes place also in confinement.

3.3. Variance of the biased tracer particle displacement: from super-diffusion to giant diffusion

The first, to the best of our knowledge, analysis, which focused specifically on the time-evolution of the variance of the displacement of a biased TP in a crowded environment in a capillary-like geometry was performed in [144]. In standard settings of active micro-rheology, the authors tracked in molecular dynamics simulations trajectories of a single TP pulled by a constant, sufficiently large force F in a glass-forming Yukawa mixture. To a great surprise, it was realised that the variance of the displacement of this driven TP along the direction of the force grows in proportion to t^z with z very close to 1.5, i.e. the fluctuations in the position exhibit a super-diffusive broadening. In [144], it was suggested that the physical mechanism underlying this remarkable broadening of fluctuations in the nonlinear scaling regime is associated with a hopping motion of the TP from cage to cage: it was indeed observed that the TP in such a system is localised for some time in a cage formed by the surrounding environment particles before it quickly moves to the next cage. Whereas in the cage the motion of the TP exhibits only small anisotropies, it becomes strongly anisotropic with respect to the motion out of the cages. It was also argued that the essential feature for the motion of the TP in the direction of force is a broad waiting time distribution, which was shown to exhibit a fat tail. The authors therefore suggested a scenario which is reminiscent of a certain class of trap models, relating it to the glass-forming properties of the binary Yukawa mixture and the ensuing complex energy landscape.

This analysis has been revisited in the numerical simulations performed in [137], which studied dynamics of a TP pulled by a constant force in mono-disperse dense fluids in a continuum strip-like geometry. More specifically, in [137], molecular dynamics simulations of colloidal fluids and Brownian dynamics of granular fluids (with only one value of the restitution parameter) have been performed, evidencing a super-diffusive broadening of fluctuations, $\sigma_x^2 \sim t^{1.5}$, over a very extended transient regime, which ultimately crossed over at longer times to a diffusive growth of fluctuations in the TP displacement, $\sigma_x^2 \sim t$, with a prefactor—an effective diffusion coefficient—being much bigger than the diffusion coefficient of the environment particles. The mono-disperse

Figure 4. Strip-like geometry with $L = 3$. Analytical predictions (continuous curves) in [210] against the results of numerical simulations (symbols). Left panel: the terminal velocity V versus force F for different densities and $\tau = \tau^* = 1$. Right panel: analogous results for $\tau = 1$ and $\tau^* = 10$ (slowly evolving environment).

colloidal fluids, as well mono-disperse granular fluids are not glass-formers, and there are no complex energy landscape in these systems, but still the super-diffusive regime exists. This naturally questions the scenario proposed in [144] and calls for a different explanation.

In [137] a different physical mechanism has been proposed, which entails—for dense lattice gases in narrow channels (both strip-like and capillary-like)—a long-lived super-diffusive broadening of fluctuations with the dynamical exponent $3/2$ which is ultimately followed by a crossover to the diffusive regime with a giant diffusion coefficient. This mechanism is based on the diffusive motion of rare (in dense systems or at a sufficiently low temperatures) vacancies—defects of packing—which are present at very low density¹² ρ_0 and permit the particles to move by exchanging their positions with the vacancies. In other words, all the environment particles and the TP are completely blocked most of time and make a single jump only when any of the vacancies arrives to their location. It was claimed in [137] that the transient super-diffusive regime exists for times less than $1/\rho_0^2$ and is associated with the interactions of the TP with just a single vacancy, which may be the initially closest one. This vacancy, once it arrives to the TP position, is certain to return to it again many times, and the vacancy indeed keeps on returning to the TP carrying the latter along the direction of the force. It was emphasised in [137] that the statistics of these multiple returns of a given vacancy to the instantaneous position of the TP in quasi-one-dimensional systems is compatible with the observations made in [144]. In consequence, the displacements of the TP are correlated due to multiple interactions with this single vacancy which entails the super-diffusive broadening of fluctuations. At longer times, however, other vacancies will appear at the location of the TP and carry it along the direction of the force. These ‘renewal’ events de-correlate the consecutive displacements of the TP, such that the fluctuations in its position will ultimately grow diffusively.

An exact solution of the model with vacancy-assisted dynamics on strip-like or capillary-like lattices has been provided in [137]. It was shown that in the linear in the density of

vacancies order, the variance σ_x^2 of the TP position along the direction of the force F obeys for sufficiently large (but less than $1/\rho_0^2$) discrete time n :

$$\sigma_x^2 \sim \frac{8(a'_0)^2 \rho_0}{3\sqrt{\pi}L} n^{3/2} + O(n) \quad (32)$$

for strip-like geometries and for the capillary-like ones one has

$$\sigma_x^2 \sim \frac{4\sqrt{2}(a'_0)^2 \rho_0}{\sqrt{3}\pi L^2} n^{3/2} + O(n), \quad (33)$$

where the omitted terms grow linearly with n , parameter a'_0 is defined in equation (28) and is associated with the TP’s transient mean velocity in these systems.

These results demonstrate the super-diffusive growth of fluctuations with an exponent $3/2$, which is in a perfect agreement with the observations made in numerical simulations and also show that this striking behaviour emerges beyond the linear response regime. Indeed, the prefactor in front of $n^{3/2}$ is proportional to the second power of F for small values of F . Interestingly, the validity of the Einstein relation is ensured here by the subdominant in time terms in equations (32) and (33), as verified in [137].

In turn, for times $n \gg 1/\rho_0^2$, the variance σ_x^2 attains a different form, which was also calculated exactly in [137] (see also recent [206]):

$$\sigma_x^2 \sim \frac{2}{L^{d-1}} \left[\frac{1}{a'_0} + \frac{2d}{(1-\alpha')L^{d-1}} \right]^{-1} n, \quad (34)$$

where α' is defined in equation (26), $d = 2$ in case of the strip-like geometries and $d = 3$ for the capillary-like ones.

Remarkably, the effective diffusion coefficient in equation (34) is independent of the density of vacancies ρ_0 . The physical origin of such a behaviour has been explained in [137] and is a very peculiar feature of transport in crowded narrow (quasi-one-dimensional) channels. Note, as well, that the diffusion coefficient of the environment particles is linearly proportional to ρ_0 in the low vacancy density limit. This means that the diffusion coefficient of the biased TP can be, in fact, orders of magnitude bigger than the diffusion coefficient of the environment particles. In consequence, one may claim

¹² Note that there is no direct relation between ρ_0 , defined for lattice systems, and the specific free volume in continuous-space settings.

that the super-diffusion in equations (32) and (33) paves the way to giant diffusion. We also note a recent [214], which studied dynamics of a biased passive tracer in a diffusive environment and revealed a similar super-diffusive transient behaviour.

4. Summary and outlook

In this review we have summarised numerous achievements made within the last several decades in the field of unbiased and biased tracer diffusion in lattice gases with stochastic dynamics on infinite in all directions lattices, and in crowded, infinitely long narrow channels with an embedded lattice structure and periodic boundary conditions in the directions perpendicular to the bias. For dense environments, a very good understanding of different facets of the dynamical behaviour has been achieved via a combined effort of exact theoretical and extensive numerical analysis, and in some instances, i.e. for the single-file systems, also via an experimental analysis. It was realised that in narrow channels surprisingly rich, sometimes quite unexpected and even counter-intuitive behaviours emerge, which are absent in unbounded systems. This is, of course, a well-known anomalous diffusion in single-files, both in the absence and in the presence of an external bias acting on the tracer particle only, in which the initial order in placement of the particles is preserved at all times. Strikingly, in narrow channels which are broader than single-files permitting thus for an effective mixing of the particles, as well, a wealth of anomalous behaviours takes place: these are the velocity anomaly of the biased tracer particle, super-diffusive at transient stages and ultimate giant diffusive broadening of fluctuations of the position of the tracer particle, spectacular multi-tracer effects resulting in self-clogging, and a variety of dynamic laws appearing in ramified channels, as exemplified here by the dynamics of a tracer particle on crowded comb-like structures. Interactions between biased tracer particles and a confined crowded environment also produce peculiar non-equilibrium patterns in the distribution of the environment particles, very different from the ones appearing in unbounded systems. For moderate density systems, similar effects have been evidenced via theories based on an appropriate decoupling of third-order correlation functions and numerical simulations. In particular, a surprising effect of a negative differential mobility has been unveiled showing that the ultimate velocity of a biased tracer particle can be a non-monotonic function of the force. In some parameter ranges, both the velocity and the diffusion coefficient of a biased tracer particle can be non-monotonic functions of the density. Such a behaviour pertains, as well, in unbounded systems but becomes most apparent in confinement. Despite a general good understanding of the behaviour in confinement, there still remain some gaps and some room for a further research. In our view, several stray questions which still await answers are as follows:

- our presentation was focused entirely on the situations in which the walls can be considered as geometrically perfect. In reality, of course, the walls can be geometrically

rough or contain some contaminants acting as temporal traps for the environment and for the tracer particle.

- the cross section of the channel may not be constant but be periodically varying along the channel. It is particularly interesting to analyse the case when this variation is sufficiently strong such that a channel consists of rather broad chambers separated by narrow passage tunnels. In this case, one expects interesting effects to emerge due to the hindered passages of the tracer particle through the bottlenecks (see, e.g. [215] for more details).
- an interesting and experimentally-relevant generalisation of the analysis presented here concerns the situations when the tracer particle is not subject to a constant external force but is active, being either a molecular motor which carries a cargo or is self-propelling itself in a crowded environment (see, e.g. [216] and references therein).
- dynamics in ramified narrow channels still remains rather poorly understood. In particular, it is not very clear what will happen in situations when a biased tracer particle moves in a quiescent environment evolving in branching narrow channels.
- the numerical analysis in [212] of the case when several biased tracer particles move randomly in a narrow crowded channel seems to focus exclusively on the situations with a big driving force. On the other hand, there are all reasons to believe that self-clogging will not emerge at lower forces. The question of specifying appropriate conditions for an effective functioning of such narrow channel devices with respect to the driven component is certainly important for many applications.

Acknowledgments

The authors thank Profs P Krapivsky and T Ooshida for many helpful comments.

ORCID iDs

G Oshanin <https://orcid.org/0000-0001-8467-3226>

References

- [1] Burada P S, Hänggi P, Marchesoni F, Schmid G and Talkner P 2009 Diffusion in confined geometries *Chem. Phys. Chem.* **10** 45
- [2] Nestorovich E M, Danelon C, Winterhalter M and Bezrukov S M 2002 Designed to penetrate: time-resolved interaction of single antibiotic molecules with bacterial pores *Proc. Natl Acad. Sci. USA* **99** 9789
- [3] van den Berg B, Pathyusha Bhamidimarri S, Dahyabhai Prajapati J, Kleinekathöfer U and Winterhalter M 2015 Outer-membrane translocation of bulky small molecules by passive diffusion *Proc. Natl Acad. Sci. USA* **112** E2991
- [4] Bajaj H, Acosta Gutierrez S, Bodrenko I, Mallocci G, Scorciapino M A, Winterhalter M and Ceccarelli M 2017 Bacterial outer membrane porins as electrostatic nanosieves: exploring transport rules of small polar molecules *ACS Nano* **11** 5465

- [5] Rout M P, Magnasco M O, Chait B T and Aitchison J D 2003 Virtual gating and nuclear transport: the hole picture *Trends Cell Biol.* **13** 622
- [6] Wente S R and Rout M P 2010 The nuclear pore complex and nuclear transport *Cold Spring Harb. Perspect. Biol.* **2** a000562
- [7] Kabachinski G and Schwartz T U 2015 The nuclear pore complex structure and function at a glance *J. Cell Sci.* **128** 423
- [8] Welte M A 2004 Bidirectional transport along microtubules *Curr. Biol.* **14** R525
- [9] Nimchinsky E A, Sabatini B L and Svoboda K 2002 Structure and function of dendritic spines *Annu. Rev. Physiol.* **64** 313
- [10] Berg H C and Turner L 1990 Chemotaxis of bacteria in glass-capillary arrays—*Escherichia coli*, motility, microchannel plate, and light-scattering *Biophys. J.* **58** 919
- [11] Hernandez-Ortiz J P, Stoltz C G and Graham M D 2005 Transport and collective dynamics in suspensions of confined swimming particles *Phys. Rev. Lett.* **95** 204501
- [12] Margaretti P and Stark H 2017 Model microswimmers in channels with varying cross section *J. Chem. Phys.* **146** 174901
- [13] Muthukumar M 2011 *Polymer Translocation* (Boca Raton, FL: CRC Press)
- [14] Kafri Y, Lubensky D K and Nelson D R 2004 Dynamics of molecular motors and polymer translocation with sequence heterogeneity *Biophys. J.* **86** 3373
- [15] Sakaue T 2016 Dynamics of polymer translocation: a short review with an introduction of weakly-driven regime *Polymers* **8** 424
- [16] Palyulin V V, Ala-Nissila T and Metzler R 2014 Polymer translocation: the first two decades and the recent diversification *Soft Matter* **45** 9016
- [17] Aksoyoglu M A, Podgornik R, Bezrukov S M, Gurnev P A, Muthukumar M and Parsegian V A 2016 Size-dependent forced PEG partitioning into channels: VDAC, OmpC and α -hemolysin *Proc. Natl Acad. Sci. USA* **113** 9003
- [18] Sarabadani J and Ala-Nissila T 2018 Theory of pore-driven and end-pulled polymer translocation dynamics through a nanopore: an overview *J. Phys.: Condens. Matter* **30** 274002
- [19] Chinappi M and Cecconi F 2018 Protein sequencing via nanopore based devices: a nanofluidics perspective *J. Phys.: Condens. Matter* **30** 204002
- [20] Zorkot M, Golestanian R and Bonthuis D J 2016 The power spectrum of ionic nanopore currents: the role of ion correlations *Nano Lett.* **16** 2205
- [21] Zorkot M and Golestanian R 2018 Current fluctuations across a nano-pore *J. Phys.: Condens. Matter* **30** 134001
- [22] Kondrat S, Wu P, Qiao R and Kornyshev A A 2014 Accelerating charging dynamics in subnanometre pores *Nat. Mater.* **13** 387
- [23] Kondrat S and Kornyshev A A 2011 Superionic state in double-layer capacitors with nanoporous electrodes *J. Phys.: Condens. Matter* **23** 022201
- [24] Mattia D and Gogotsi Y 2008 Review: static and dynamic behavior of liquids inside carbon nanotubes *Microfluid. Nanofluid.* **5** 289
- [25] Lee A, Kondrat S, Oshanin G and Kornyshev A A 2014 Charging dynamics of supercapacitors with narrow cylindrical nanopores *Nanotechnology* **25** 315401
- [26] Dudka M, Kondrat S, Kornyshev A A and Oshanin G 2016 Phase behaviour and structure of a superionic liquid in nonpolarized nanoconfinement *J. Phys.: Condens. Matter* **28** 464007
- [27] Comtet J, Niguès A, Kaiser V, Coasne B, Bocquet L and Siria A 2017 Nanoscale capillary freezing of ionic liquids confined between metallic interfaces and the role of electronic screening *Nat. Mater.* **16** 634
- [28] Futamura R, Iiyama T, Takasaki Y, Gogotsi Y, Biggs M J, Salanne M, Ségolini J, Simon P and Kaneko K 2017 Partial breaking of the coulombic ordering of ionic liquids confined in carbon nanopores *Nat. Mater.* **16** 1225
- [29] Mossa S 2018 Re-entrant phase transitions and dynamics of a nanoconfined ionic liquid *Phys. Rev. X* **8** 031062
- [30] Stone H A, Stroock A D and Ajdari A 2004 Engineering flows in small devices: microfluidics toward a lab-on-a-chip *Annu. Rev. Fluid Mech.* **36** 381
- [31] Mukhopadhyay A and Granick S 2001 Micro- and nanorheology *Curr. Opin. Colloid Interface Sci.* **6** 423
- [32] Liot O, Socol M, Garcia L, Thiéry J, Figarol A, Mingotaud A F and Joseph P 2018 Transport of nano-objects in narrow channels: influence of Brownian diffusion, confinement and particle nature *J. Phys.: Condens. Matter* **30** 234001
- [33] Bet B, Samin S, Georgiev R, Burak Eral H and van Roij R 2018 Steering particles by breaking symmetries *J. Phys.: Condens. Matter* **30** 224002
- [34] Sarracino A, Cecconi F, Puglisi A and Vulpiani A 2016 Nonlinear response of inertial tracers in steady laminar flows: differential and absolute negative mobility *Phys. Rev. Lett.* **117** 174501
- [35] Cecconi F, Puglisi A, Sarracino A and Vulpiani A 2017 Anomalous force-velocity relation of driven inertial tracers in steady laminar flows *Eur. Phys. J. E* **40** 81
- [36] Cecconi F, Puglisi A, Sarracino A and Vulpiani A 2018 Anomalous mobility of a driven active particle in a steady laminar flow *J. Phys.: Condens. Matter* **30** 264002
- [37] Margaretti P, Pagonabarraga I and Rubi M J 2013 Entropic transport in confined media: a challenge for computational studies in biological and soft-matter systems *Front. Phys.* **1** 21
- [38] Katz S, Lebowitz J L and Spohn H 1983 Phase transitions in stationary nonequilibrium states of model lattice systems *Phys. Rev. B* **28** 1655
- [39] Derrida B, Domany E and Mukamel D 1992 An exact solution of a one-dimensional asymmetric exclusion model with open boundaries *J. Stat. Phys.* **69** 667
- [40] Schmittmann B and Zia R K P 1995 Statistical mechanics of driven diffusive systems *Phase Transitions and Critical Phenomena* vol 17, ed C Domb and J L Lebowitz (New York: Academic)
- [41] Evans M R, Kafri Y, Koduvely H M and Mukamel D 1998 Phase separation in one-dimensional driven diffusive systems *Phys. Rev. Lett.* **80** 425
- [42] Bauer W R and Nadler W 2006 Molecular transport through channels and pores: effects of in-channel interactions and blocking *Proc. Natl Acad. Sci. USA* **103** 11446
- [43] Chou T 1998 How fast do fluids squeeze through microscopic single-file pores? *Phys. Rev. Lett.* **80** 85
- [44] Chou T and Lohse D 1999 Entropy-driven pumping in zeolites and biological channels *Phys. Rev. Lett.* **82** 3552
- [45] Berezhkovskii A and Hummer G 2002 Single-file transport of water molecules through a carbon nanotube *Phys. Rev. Lett.* **89** 064503
- [46] Zilman A 2009 Effects of multiple occupancy and interparticle interactions on selective transport through narrow channels: theory versus experiment *Biophys. J.* **96** 1235
- [47] Chou T, Mallick K and Zia R K P 2011 Non-equilibrium statistical mechanics: from a paradigmatic model to biological transport *Rep. Prog. Phys.* **74** 1
- [48] Ai B-Q and Wu J-C 2013 Transport of finite size particles in confined narrow channels: diffusion, coherence, and particle separation *J. Chem. Phys.* **139** 034114

- [49] Curatolo A I, Evans M R, Kafri Y and Tailleur J 2016 Multilane driven diffusive systems *J. Phys. A: Math. Theor.* **49** 095601
- [50] Marconi U M B, Magaretti P and Pagonabarraga I 2015 Tracer diffusion of hard-sphere binary mixtures under nano-confinement *J. Chem. Phys.* **143** 184501
- [51] Baek Y, Kafri Y and Lecomte V 2017 Dynamical symmetry breaking and phase transitions in driven diffusive systems *Phys. Rev. Lett.* **118** 030604
- [52] Baek Y, Kafri Y and Lecomte V 2018 Dynamical phase transitions in the current distribution of driven diffusive channels *J. Phys. A: Math. Theor.* **51** 105001
- [53] Bollinger J A, Carmer J, Jain A and Truskett T M 2016 Impact of solvent granularity and layering on tracer hydrodynamics in confinement *Soft Matter* **12** 9561
- [54] Gardel M L, Valentine M T and Weitz D A 2005 *Microscale Diagnostic Techniques* ed K Breuer (Berlin: Springer)
- [55] Cicuta P and Donald A M 2007 Microrheology: a review of the method and applications *Soft Matter* **3** 1449
- [56] Squires T M and Mason T G 2009 Fluid mechanics of microrheology *Annu. Rev. Fluid Mech.* **42** 413
- [57] Wilson L G, Harrison A W, Schofield A B, Arlt J and Poon W C K 2009 Passive and active microrheology of hard-sphere colloids *J. Phys. Chem. B* **113** 3806
- [58] Wilson L G and Poon W C K 2011 Small-world rheology: an introduction to probe-based active microrheology *Phys. Chem. Chem. Phys.* **13** 1061
- [59] Puertas A M and Voigtman T 2014 Microrheology of colloidal systems *J. Phys.: Condens. Matter* **26** 243101
- [60] Nazockdast E and Morris J F 2016 Microstructural theory and the rheology of concentrated colloidal suspensions *J. Rheol.* **60** 733
- [61] Dullens R P A and Bechinger C 2011 Shear thinning and local melting of colloidal crystals *Phys. Rev. Lett.* **107** 138301
- [62] Cugliandolo L F 2011 The effective temperature *J. Phys. A: Math. Theor.* **44** 483001
- [63] Seifert U 2012 Stochastic thermodynamics, fluctuation theorems and molecular machines *Rep. Prog. Phys.* **75** 126001
- [64] Puglisi A, Sarracino A and Vulpiani A 2017 Temperature in and out of equilibrium: a review of concepts, tools and attempts *Phys. Rep.* **709–10** 1
- [65] Compaan K and Haven Y 1956 Correlation factors for diffusion in solids *Trans. Faraday Soc.* **52** 786
- [66] Howard R E 1966 Random-Walk method for calculating correlation factors: tracer diffusion by divacancy and impurity-vacancy Pairs in cubic crystals *Phys. Rev.* **144** 650
- [67] Manning J R 1968 *Diffusion Kinetics for Atoms in Crystals* (Princeton: van Nostrand)
- [68] Hodgkin A L and Keynes R D 1955 The potassium permeability of a giant nerve fibre *J. Physiol. London* **128** 61
- [69] Harris T E 1965 Diffusion with collisions between particles *J. Appl. Prob.* **2** 323
- [70] Ooshida T and Otsuki M 2018 Two-tag correlations and nonequilibrium fluctuation? Response relation in ageing single-file diffusion *J. Phys.: Condens. Matter* **30** 374001
- [71] Bénichou O, Cazabat A M, Lemarchand A, Moreau M and Oshanin G 1999 Biased diffusion in a one-dimensional adsorbed monolayer *J. Stat. Phys.* **97** 351
- [72] Komorowski T and Olla S 2005 On mobility and Einstein relation for tracers in time-mixing random environments *J. Stat. Phys.* **118** 407
- [73] Rouse P E 1953 A theory of the linear viscoelastic properties of dilute solutions of coiling polymers *J. Chem. Phys.* **21** 1272
- [74] Levitt D G 1973 Dynamics of a single-file pore: non-Fickian behavior *Phys. Rev. A* **8** 3050
- [75] Richards P M 1977 Theory of one-dimensional hopping conductivity and diffusion *Phys. Rev. B* **16** 1393
- [76] Fedders P A 1978 Two-point correlation functions for a distinguishable particle hopping on a uniform one-dimensional chain *Phys. Rev. B* **17** 40
- [77] Alexander S and Pincus P 1978 Diffusion of labeled particles on one-dimensional chains *Phys. Rev. B* **18** 2011
- [78] Arratia R 1983 The motion of a tagged particle in the simple symmetric exclusion system on *Z Ann. Probab.* **2** 362
- [79] van Beijeren H, Kehr K W and Kutner R 1983 Diffusion in concentrated lattice gases. III. Tracer diffusion on a one-dimensional lattice *Phys. Rev. B* **28** 5711
- [80] Majumdar S N and Barma M 1991 Tag diffusion in driven systems, growing interfaces, and anomalous fluctuations *Phys. Rev. B* **44** 5306
- [81] Taloni A and Lomholt M A 2008 Langevin formulation for single-file diffusion *Phys. Rev. E* **78** 051116
- [82] Barkai E and Silbey R 2010 Diffusion of tagged particle in an exclusion process *Phys. Rev. E* **81** 041129
- [83] Ryabov A 2015 *Stochastic Dynamics and Energetics of Biomolecular Systems* (Berlin: Springer)
- [84] Taloni A, Flomenbom O, Castaneda-Priego R and Marchesoni F 2017 Single file dynamics in soft materials *Soft Matter* **13** 1096
- [85] Hahn K, Kärger J and Kukla V 1996 Single-file diffusion observation *Phys. Rev. Lett.* **76** 2762
- [86] Wei Q-H, Bechinger C and Leiderer P 2000 Single-file diffusion of colloids in one-dimensional channels *Science* **287** 625
- [87] Meersmann T, Logan J W, Simonutti R, Caldarelli S, Comotti A, Sozzani P, Kaiser L G and Pines A 2000 Exploring single-file diffusion in one-dimensional nanochannels by laser-polarized ^{129}Xe NMR spectroscopy *J. Phys. Chem. A* **104** 11665
- [88] Lin B, Meron M, Cui B, Rice S A and Diamant H 2005 From random walk to single-file diffusion *Phys. Rev. Lett.* **94** 216001
- [89] Illien P, Bénichou O, Mejía-Monasterio C, Oshanin G and Voituriez R 2013 Active transport in dense diffusive single-file systems *Phys. Rev. Lett.* **113** 038102
- [90] Metzler R, Sanders L, Lomholt M A, Lizana L, Fogelmark K and Ambjörnsson T 2014 Ageing single file motion *Eur. Phys. J. Spec. Top.* **223** 3287
- [91] Hegde C, Sabhapandit S and Dhar A 2014 Universal large deviations for the tagged particle in single-file motion *Phys. Rev. Lett.* **113** 120601
- [92] Krapivsky P L, Mallick K and Sadhu T 2015 Dynamical properties of single-file diffusion *J. Stat. Mech.* **P09007**
- [93] Sadhu T and Derrida B 2015 Large deviation function of a tracer position in single file diffusion *J. Stat. Mech.* **P09008**
- [94] Krapivsky P L, Mallick K and Sadhu T 2014 Large deviations in single-file diffusion *Phys. Rev. Lett.* **113** 078101
- [95] Imamura T, Mallick K and Sasamoto T 2017 Large deviations of a tracer in the symmetric exclusion process *Phys. Rev. Lett.* **118** 160601
- [96] Majumdar S and Barma M 1991 Two-tag correlation functions in one dimensional lattice gases *Physica A* **177** 366
- [97] Rajesh R and Majumdar S N 2001 Exact tagged particle correlations in the random average process *Phys. Rev. E* **64** 036103
- [98] Ooshida T, Goto S, Matsumoto T, Nakahara A and Otsuki M 2013 Analytical calculation of four-point correlations for a simple model of cages involving numerous particles *Phys. Rev. E* **88** 062108

- [99] Sabhapandit S and Dhar A 2015 Exact probability distribution for the two-tag displacement in single-file motion *J. Stat. Mech.* **P07024**
- [100] Poncet A, Bénichou O, Démercy V and Oshanin G 2018 N -tag probability law of the symmetric exclusion process *Phys. Rev. E* **97** 062119
- [101] Leibovich N and Barkai E 2013 Everlasting effect of initial conditions on single-file diffusion *Phys. Rev. E* **88** 032107
- [102] Cividini J and Kundu A 2017 Tagged particle in single-file diffusion with arbitrary initial conditions *J. Stat. Mech.* **083203**
- [103] Lizana L, Lomholt M A and Ambjörnsson T 2014 Single-file diffusion with non-thermal initial conditions *Physica A* **395** 148
- [104] Lizana L, Ambjörnsson T, Taloni A, Barkai E and Lomholt M A 2010 Foundation of fractional Langevin equation: harmonization of a many-body problem *Phys. Rev. E* **81** 051118
- [105] Bénichou O, Illien P, Oshanin G, Sarracino A and Voituriez R 2015 Diffusion and subdiffusion of interacting particles on Comblike structures *Phys. Rev. Lett.* **115** 220601
- [106] Ben Avraham D and Havlin S 2005 *Diffusion and Reactions in Fractals and Disordered Systems* (Cambridge: Cambridge University Press)
- [107] Weiss G H and Havlin S 1986 Some properties of a random walk on a comb structure *Physica A* **134** 474–82
- [108] Galanti M, Fanelli D, Maritan A and Piazza F 2015 Diffusion of tagged particles in a crowded medium *Europhys. Lett.* **107** 20006
- [109] Ghosh S K, Cherstvy A G and Metzler R 2015 Non-universal tracer diffusion in crowded media of non-inert obstacles *Phys. Chem. Chem. Phys.* **17** 1847
- [110] Ghosh S K, Cherstvy A G, Grebenkov D S and Metzler R 2016 Anomalous, non-Gaussian tracer diffusion in crowded two-dimensional environments *New J. Phys.* **18** 013027
- [111] Fedders P A and Sankey O F 1977 The generalized atomic hopping problem occupancy correlation functions *Phys. Rev. B* **15** 3580
- [112] Nakazato K and Kitahara K 1980 Site blocking effect in tracer diffusion on a lattice *Prog. Theor. Phys.* **64** 2261
- [113] Ishioka S and Koiwa M 1980 On the correlation effect in self-diffusion via the vacancy mechanism *Phil. Mag. A* **41** 385
- [114] Kehr K W, Kutner R and Binder K 1981 Diffusion in concentrated lattice gases. Self-diffusion of noninteracting particles in three-dimensional lattices *Phys. Rev. B* **23** 4931
- [115] Kitahara K and Nakazato K 1982 *Point Defects and Defect Interactions in Metals* (Tokyo: University of Tokyo Press)
- [116] Koiwa M and Ishioka S 1983 Random walk properties of lattices and correlation factors for diffusion via the vacancy mechanism in crystals *J. Stat. Phys.* **30** 477
- [117] Tahir-Kheli R A and Elliott R J 1983 Correlated random walk in lattices: tracer diffusion at general concentration *Phys. Rev. B* **27** 844
- [118] van Beijeren H and Kutner R 1985 Mean square displacement of a tracer particle in a hard-core lattice gas *Phys. Rev. Lett.* **55** 238
- [119] Tahir-Kheli R A 1987 Monte Carlo evaluation of the diffusion correlation factor in concentrated lattices and comparison with an improved theory *Phys. Rev. B* **35** 5503
- [120] Brummelhuis M J A M and Hilhorst H J 1988 Single-vacancy induced motion of a tracer particle in a two-dimensional lattice gas *J. Stat. Phys.* **53** 249
- [121] Brummelhuis M J A M and Hilhorst H J 1989 Tracer particle motion in a two-dimensional lattice gas with low vacancy density *Physica A* **156** 575
- [122] Kehr K W and Binder K 1987 *Application of the Monte Carlo Method in Statistical Physics* (Berlin: Springer)
- [123] Pigeon S, Fogelmark K, Söderberg B, Mukhopadhyay G and Ambjörnsson T 2017 Tracer particle diffusion in a system with hardcore interacting particles *J. Stat. Mech.* **043036**
- [124] Zucker I J 2011 70 + years of the Watson integrals *J. Stat. Phys.* **145** 591
- [125] Bénichou O, Cazabat A M, De Coninck J, Moreau M and Oshanin G 2000 Stokes formula and density perturbances for driven tracer diffusion in an adsorbed monolayer *Phys. Rev. Lett.* **84** 511
- [126] Bénichou O, Cazabat A M, De Coninck J, Moreau M and Oshanin G 2001 Force-velocity relation and density profiles for biased diffusion in an adsorbed monolayer *Phys. Rev. B* **63** 235413
- [127] Bénichou O, Klafter J, Moreau M and Oshanin G 2000 Generalized model for dynamic percolation *Phys. Rev. E* **62** 3327
- [128] Bénichou O and Oshanin G 2001 Atomic slide puzzle: self-diffusion of an impure atom *Phys. Rev. E* **64** 020103
- [129] van Gastel R, Somfai E, van Saarloos W and Frenken J W M 2000 A giant atomic slide-puzzle *Nature* **408** 665
- [130] van Gastel R, Somfai E, van Albada S B, van Saarloos W and Frenken J W M 2001 Nothing moves a surface: vacancy mediated surface diffusion *Phys. Rev. Lett.* **86** 1562
- [131] Prostnev A S and Shub B R 2009 Migration of vacancies and impurities on the (1 1 1) surface of a face-centered cubic lattice *Russ. J. Phys. Chem. B* **3** 830
- [132] Prostnev A S and Shub B R 2012 Kinetics of the diffusion of an impurity atom on a fcc (1 1 1) surface *Russ. J. Phys. Chem. B* **6** 65
- [133] Prostnev A S and Shub B R 2013 Percolation mechanism of the diffusion of impurity atoms in dense surface layers *Russ. J. Phys. Chem. B* **7** 568
- [134] Prostnev A S and Shub B R 2014 Diffusion of atoms incorporated in the surface layer of the fcc (1 1 1) face *Russ. J. Phys. Chem. B* **8** 420
- [135] Prostnev A S and Shub B R 2016 Diffusion of atoms in a dense adsorbed layer with a hexagonal structure *Russ. J. Phys. Chem. B* **10** 547
- [136] Prostnev A S and Shub B R 2016 Simulation of the diffusion of atoms in a dense adsorbed layer with a hexagonal structure *Russ. J. Phys. Chem. B* **10** 1022
- [137] Bénichou O, Bodrova A, Chakraborty D, Illien P, Law A, Mejía-Monasterio C, Oshanin G and Voituriez R 2013 Geometry-induced superdiffusion in driven crowded systems *Phys. Rev. Lett.* **111** 260601
- [138] Sriram I, Meyer A and Furst E M 2010 Active microrheology of a colloidal suspension in the direct collision limit *Phys. Fluids* **22** 062003
- [139] Meyer A, Marshall A, Bush B G and Furst E M 2006 Laser tweezer microrheology of a colloidal suspension *J. Rheol.* **50** 77
- [140] Berner J, Muller B, Gomez-Solano J R, Kruger M and Bechinger C 2018 Oscillating modes of driven colloids in overdamped systems *Nat. Commun.* **9** 999
- [141] Harrer Ch J, Winter D, Horbach J, Fuchs M and Voigtmann T 2012 Force-induced diffusion in microrheology *J. Phys.: Condens. Matter* **24** 464105
- [142] Zia R N and Brady J F 2010 Single-particle motion in colloids: force-induced diffusion *J. Fluid Mech.* **658** 188
- [143] Reichhardt C and Olson Reichhardt C J 2004 Local melting and drag for a particle driven through a colloidal crystal *Phys. Rev. Lett.* **92** 108301
- [144] Winter D, Horbach J, Virnau P and Binder K 2012 Active nonlinear microrheology in a glass-forming Yukawa fluid *Phys. Rev. Lett.* **108** 028303

- [145] Winter D and Horbach J 2013 Nonlinear active micro-rheology in a glass-forming soft-sphere mixture *J. Chem. Phys.* **138** 12A512
- [146] Dzubiella J, Löwen H and Likos C N 2003 Depletion forces in nonequilibrium *Phys. Rev. Lett.* **91** 248301
- [147] Sriram I and Furst E M 2012 Out-of-equilibrium forces between colloids *Soft Matter* **8** 3335
- [148] Kliushnychenko O V and Lukyanets S P 2014 Effect of the blockage and concentration switching of the direction of the trace of an inclusion in a lattice gas *J. Exp. Theor. Phys.* **118** 976
- [149] Kliushnychenko O V and Lukyanets S P 2017 Effects of gas interparticle interaction on dissipative wake-mediated forces *Phys. Rev. E* **95** 012150
- [150] Burlatsky S F, Oshanin G, Mogutov A V and Moreau M 1992 Directed walk in a one-dimensional lattice gas *Phys. Lett. A* **166** 230
- [151] Burlatsky S F, Oshanin G, Moreau M and Reinhardt W P 1996 Motion of a driven tracer particle in a one-dimensional symmetric lattice gas *Phys. Rev. E* **54** 3165
- [152] Landim C, Olla S and Volchan S B 1998 Driven tracer particle in one dimensional symmetric simple exclusion *Commun. Math. Phys.* **192** 287
- [153] Antal T, Krapivsky P L and Redner S 2009 Shepherd model for knot-limited polymer ejection from a capsid *J. Theor. Biol.* **261** 488
- [154] Schiessel S, Oshanin G and Blumen A 1995 Polyampholytes in external electric fields: dynamics and conformation properties *J. Chem. Phys.* **103** 5070
- [155] Schiessel S, Oshanin G and Blumen A 1996 Dynamics and conformation properties of polyampholytes in external electrical fields: influence of the charge distribution *Macromol. Theory Simul.* **5** 45
- [156] Lacoste D and Lomholt M A 2015 Stochastic thermodynamics of a tagged particle within a harmonic chain *Phys. Rev. E* **91** 022114
- [157] Huveneers F and Simenhaus F 2015 Random walk driven by the simple exclusion process *Electron. J. Probab.* **20** 1
- [158] Villamaina D, Puglisi A and Vulpiani A 2008 The fluctuation-dissipation relation in sub-diffusive systems: the case of granular single-file diffusion *J. Stat. Mech.* **L10001**
- [159] Villamaina D, Sarracino A, Gradenigo G, Puglisi A and Vulpiani A 2011 On anomalous diffusion and the out of equilibrium response function in one-dimensional models *J. Stat. Mech.* **L01002**
- [160] Oshanin G, Bénichou O, Burlatsky S F and Moreau M 2004 Biased tracer diffusion in hard-core lattice gases: some notes on the validity of the Einstein relation *Instabilities and Nonequilibrium Structures IX* ed O Descalzi, J Martínez and S Rica (Berlin: Springer)
- [161] Bénichou O, Démery V and Poncet A 2018 Unbinding transition of probes in single-file systems *Phys. Rev. Lett.* **120** 070601
- [162] Illien P, Bénichou O, Oshanin G and Voituriez R 2013 Fluctuations and correlations of a driven tracer in a hard-core lattice gas *Phys. Rev. E* **87** 032164
- [163] Illien P, Bénichou O, Oshanin G and Voituriez R 2015 Distribution of the position of a driven tracer in a hardcore lattice gas *J. Stat. Mech.* **P11016**
- [164] Kundu A and Cividini J 2016 Exact correlations in a single-file system with a driven tracer *Europhys. Lett.* **115** 54003
- [165] Cividini J, Kundu A, Majumdar S N and Mukamel D 2016 Correlation and fluctuation in a random average process on an infinite line with a driven tracer *J. Stat. Mech.* **053212**
- [166] Burlatsky S F, Oshanin G, Cazabat A M and Moreau M 1996 Microscopic model of upward creep of an ultrathin wetting film *Phys. Rev. Lett.* **76** 86
- [167] Burlatsky S F, Oshanin G, Cazabat A M, Moreau M and Reinhardt W P 1996 Spreading of a thin wetting film: microscopic approach *Phys. Rev. E* **54** 3832
- [168] Popescu M N and Dietrich S 2004 Model for spreading of liquid monolayers *Phys. Rev. E* **69** 061602
- [169] Abraham D B, Cuerno R and Esteban M 2002 Microscopic model for thin film spreading *Phys. Rev. Lett.* **88** 206101
- [170] Popescu M N, Oshanin G, Dietrich S and Cazabat A M 2012 Precursor films in wetting phenomena *J. Phys.: Condens. Matter* **24** 243102
- [171] Oshanin G, De Coninck J, Cazabat A M and Moreau M 1998 Dewetting, partial wetting, and spreading of a two-dimensional monolayer on solid surface *Phys. Rev. E* **58** 20
- [172] Démery V, Bénichou O and Jacquin H 2014 Generalized Langevin equations for a driven tracer in dense soft colloids: construction and applications *New J. Phys.* **16** 053032
- [173] Candelier R and Dauchot O 2010 Journey of an intruder through the fluidization and jamming transitions of a dense granular media *Phys. Rev. E* **81** 011304
- [174] Ivlev A V, Bartnick J, Heinen M, Du C-R, Nosenko V and Löwen H 2015 Statistical mechanics where Newton's third law is broken *Phys. Rev. X* **5** 011035
- [175] Wulfert R, Seifert U and Speck T 2017 Nonequilibrium depletion interactions in active microrheology *Soft Matter* **13** 9093
- [176] Mejía-Monasterio C and Oshanin G 2011 Bias- and bath-mediated pairing of particles driven through a quiescent medium *Soft Matter* **7** 993
- [177] Sriram I and Furst E M 2015 Two spheres translating in tandem through a colloidal suspension *Phys. Rev. E* **91** 042303
- [178] Dolata B E and Zia R N 2018 Non-equilibrium pair interactions in colloidal dispersions *J. Fluid Mech.* **836** 694
- [179] Furukawa A, Gambassi A, Dietrich S and Tanaka H 2013 *Phys. Rev. Lett.* **111** 055701
- [180] Reichhardt C and Olson Reichhardt C J 2006 Cooperative behavior and pattern formation in mixtures of driven and nondriven colloidal assemblies *Phys. Rev. E* **74** 011403
- [181] Ladadwa I and Heuer A 2013 Nonlinear response and crowding effects in microrheology *Phys. Rev. E* **87** 012302
- [182] Vasilyev O A, Bénichou O, Mejía-Monasterio C, Weeks E R and Oshanin G 2017 Cooperative behavior of biased probes in crowded interacting systems *Soft Matter* **13** 7617
- [183] Rex M and Löwen H 2007 Lane formation in oppositely charged colloids driven by an electric field: chaining and two-dimensional crystallization *Phys. Rev. E* **75** 051402
- [184] Rex M and Löwen H 2008 Influence of hydrodynamic interactions on lane formation in oppositely charged driven colloids *Eur. Phys. J. E* **26** 143
- [185] Glanz T and Löwen H 2012 The nature of the laning transition in two dimensions *J. Phys.: Condens. Matter* **24** 464114
- [186] Poncet A, Bénichou O, Démery V and Oshanin G 2017 Universal long ranged correlations in driven binary mixtures *Phys. Rev. Lett.* **118** 118002
- [187] Reichhardt C and Reichhardt C J O 2018 Velocity force curves, laning, and jamming for oppositely driven disk systems *Soft Matter* **14** 490
- [188] Vissers T, Wysocki A, Rex M, Löwen H, Royall C P, Imhof A and van Blaaderen A 2011 Lane formation in driven mixtures of oppositely charged colloids *Soft Matter* **7** 235
- [189] Sütterlin K R *et al* 2009 Dynamics of lane formation in driven binary complex plasmas *Phys. Rev. Lett.* **102** 085003

- [190] Helbing D 2001 Traffic and related self-driven many-particle systems *Rev. Mod. Phys.* **73** 1067
- [191] Bénichou O and Oshanin G 2002 Ultraslow vacancy-mediated tracer diffusion in two dimensions: the Einstein relation verified *Phys. Rev. E* **66** 031101
- [192] Bénichou O, Illien P, Oshanin G, Sarracino A and Voituriez R 2014 Microscopic theory for negative differential mobility in crowded environments *Phys. Rev. Lett.* **113** 268002
- [193] Zia R K P, Praestgaard E L and Mouritsen O G 2002 Getting more from pushing less: negative specific heat and conductivity in nonequilibrium steady states *Am. J. Phys.* **70** 384
- [194] Eichhorn R, Regtmeier J, Anselmetti D and Reimann P 2010 Negative mobility and sorting of colloidal particles *Soft Matter* **6** 1858
- [195] Jack R L, Kelsey D, Garrahan J P and Chandler D 2008 Negative differential mobility of weakly driven particles in models of glass formers *Phys. Rev. E* **78** 011506
- [196] Sellitto M 2008 Asymmetric exclusion processes with constrained dynamics *Phys. Rev. Lett.* **101** 048301
- [197] Turci F, Pitard E and Sellitto M 2012 Driving kinetically constrained models into nonequilibrium steady states: structural and slow transport properties *Phys. Rev. E* **86** 031112
- [198] Baerts P, Basu U, Maes C and Safaverdi S 2013 Frenetic origin of negative differential response *Phys. Rev. E* **88** 052109
- [199] Basu U and Maes C 2014 Mobility transition in a dynamic environment *J. Phys. A: Math. Theor.* **47** 255003
- [200] Baiesi M, Stella A L and Vanderzande C 2015 Role of trapping and crowding as sources of negative differential mobility *Phys. Rev. E* **92** 042121
- [201] Cividini J, Mukamel D and Posch H A 2018 Driven tracer with absolute negative mobility *J. Phys. A: Math. Theor.* **51** 085001
- [202] Reichhardt C and Reichhardt C J O 2018 Negative differential mobility and trapping in active matter systems *J. Phys.: Condens. Matter* **30** 015404
- [203] Chatterjee A K, Basu U and Mohanty P K 2018 Negative differential mobility in interacting particle systems *Phys. Rev. E* **97** 052137
- [204] Bénichou O, Mejía-Monasterio C and Oshanin G 2013 Anomalous field-induced growth of fluctuations in dynamics of a biased intruder moving in a quiescent medium *Phys. Rev. E* **87** 020103
- [205] Bénichou O, Illien P, Mejía-Monasterio C and Oshanin G 2013 A biased intruder in a dense quiescent medium: looking beyond the force-velocity relation *J. Stat. Mech.* **P05008**
- [206] Illien P, Bénichou O, Oshanin G, Sarracino A and Voituriez R 2018 Nonequilibrium fluctuations and enhanced diffusion of a driven particle in a dense environment *Phys. Rev. Lett.* **120** 200606
- [207] Leitmann S and Franosch T 2013 Nonlinear response in the driven lattice Lorentz gas *Phys. Rev. Lett.* **111** 190603
- [208] Leitmann S and Franosch T 2017 Time-dependent fluctuations and superdiffusivity in the driven lattice Lorentz gas *Phys. Rev. Lett.* **118** 018001
- [209] Leitmann S, Schwab T and Franosch T 2018 Time-dependent perpendicular fluctuations in the driven lattice Lorentz gas *Phys. Rev. E* **97** 022101
- [210] Bénichou O, Illien P, Oshanin G, Sarracino A and Voituriez R 2016 Nonlinear response and emerging nonequilibrium microstructures for biased diffusion in confined crowded environments *Phys. Rev. E* **93** 032128
- [211] Cividini J, Mukamel D and Posch H A 2017 Driven tracers in narrow channels *Phys. Rev. E* **95** 012110
- [212] Kusters R and Storm C 2017 Dynamic phase separation of confined driven particles *Europhys. Lett.* **118** 58004
- [213] Illien P, Bénichou O, Oshanin G and Voituriez R 2014 Velocity anomaly of a driven tracer in a confined crowded environment *Phys. Rev. Lett.* **113** 030603
- [214] Huveneers F 2018 Response to a small external force and fluctuations of a passive particle in a one-dimensional diffusive environment *Phys. Rev. E* **97** 042116
- [215] Reguera D and Rubí J M 2001 Kinetic equations for diffusion in the presence of entropic barriers *Phys. Rev. E* **64** 061106
- [216] Dey S, Ching K and Das M 2018 Active and passive transport of cargo in a corrugated channel: a lattice model study *J. Chem. Phys.* **148** 134907