



HAL
open science

Odd skipped-related 1 (Osr1) identifies muscle-interstitial fibro-adipogenic progenitors (FAPs) activated by acute injury

Jürgen Stumm, Pedro Vallecillo-García, Sophie Vom Hofe-Schneider, David Ollitrault, Heinrich Schrewe, Aris N Economides, Giovanna Marazzi, David A. Sassoon, Sigmar Stricker

► To cite this version:

Jürgen Stumm, Pedro Vallecillo-García, Sophie Vom Hofe-Schneider, David Ollitrault, Heinrich Schrewe, et al.. Odd skipped-related 1 (Osr1) identifies muscle-interstitial fibro-adipogenic progenitors (FAPs) activated by acute injury. *Stem Cell Research*, 2018, 32, pp.8 - 16. 10.1016/j.scr.2018.08.010 . hal-01908913

HAL Id: hal-01908913

<https://hal.sorbonne-universite.fr/hal-01908913v1>

Submitted on 30 Oct 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Odd skipped-related 1 (Osr1) identifies muscle-interstitial fibro-adipogenic progenitors (FAPs) activated by acute injury

Jürgen Stumm^{a,1}, Pedro Vallecillo-García^{a,1}, Sophie Vom Hofe-Schneider^a, David Ollitrault^{b,c}, Heinrich Schrewe^d, Aris N. Economides^e, Giovanna Marazzi^{b,c}, David A. Sassoon^{b,c}, Sigmar Stricker^{a,*}

^a Institute of Chemistry and Biochemistry, Freie Universität Berlin, 14195 Berlin, Germany

^b Stem Cells and Regenerative Medicine, ICAN-UMRS 1166, Université de Pierre et Marie Curie, Sorbonne Universités, 75013 Paris, France

^c ICAN, Institute of Cardiometabolism and Nutrition, 75013 Paris, France

^d Department of Developmental Genetics, Max Planck Institute for Molecular Genetics, 14195 Berlin, Germany

^e Regeneron Pharmaceuticals, Tarrytown, NY 10591, USA

ARTICLE INFO

Summary statement: Expression of Osr1 specifically in muscle interstitial fibro-adipogenic progenitors (FAPs) activated by acute injury provides a tool to isolate and trace this population.

Keywords:

Skeletal muscle
Fibro-adipogenic progenitors
Mesenchymal progenitors, Muscle interstitium
Muscle regeneration

ABSTRACT

Fibro-adipogenic progenitors (FAPs) are resident mesenchymal progenitors in adult skeletal muscle that support muscle repair, but also give rise to fibrous and adipose infiltration in response to disease and chronic injury. FAPs are identified using cell surface markers that do not distinguish between quiescent FAPs and FAPs actively engaged in the regenerative process. We have shown previously that FAPs are derived from cells that express the transcription factor Osr1 during development. Here we show that adult FAPs express Osr1 at low levels and frequency, however upon acute injury FAPs reactivate Osr1 expression in the injured tissue. Osr1⁺ FAPs are enriched in proliferating and apoptotic cells demonstrating that Osr1 identifies activated FAPs. *In vivo* genetic lineage tracing shows that Osr1⁺ activated FAPs return to the resident FAP pool after regeneration as well as contribute to adipocytes after glycerol-induced fatty degeneration. In conclusion, reporter LacZ or eGFP-CreERT2 expression from the endogenous Osr1 locus serves as marker for FACS isolation and tamoxifen-induced manipulation of activated FAPs.

1. Introduction

The remarkable regenerative potential of skeletal muscle relies on myogenic stem cells (satellite cells), however other interstitial populations play a critical supportive role (Bentzinger et al., 2013; Uezumi et al., 2014a; Pannerec et al., 2012). Amongst these, fibro-adipogenic progenitors (FAPs) have attracted immense attention in the past years. FAPs are muscle-interstitial resident mesenchymal progenitor cells that have the capacity to provide a pro-myogenic environment for muscle regeneration (Joe et al., 2010) but also contribute directly to fibrotic degeneration and fatty infiltration in diseased or degenerating muscle (Uezumi et al., 2011; Uezumi et al., 2010; Lemos et al., 2015; Kopinke et al., 2017). As such, FAPs are important cell targets for therapeutic approaches (Lemos et al., 2015; Mozzetta et al., 2013; Contreras et al., 2016; Gonzalez et al., 2017). FAPs are activated upon injury to proliferate (Joe et al., 2010; Uezumi et al., 2010) and are cleared by apoptosis in the course of regeneration (Lemos et al., 2015). The

intrinsic mechanisms of activation and pro-myogenic function as well as the mechanisms that promote fibrotic or adipogenic conversion are not well understood. Murine FAPs were characterized using different cell surface marker combinations. Joe et al. (Joe et al., 2010) used the combination of lin⁻;Sca1⁺;CD34⁺ or equivalently lin⁻;α7-integrin⁺;Sca1⁺ to isolate FAPs, while Uezumi et al. (Uezumi et al., 2010; Uezumi et al., 2014b) used lin⁻;PDGFRα⁺ to isolate FAPs from mouse and human muscle. Both, the Sca1⁺ and PDGFRα⁺ populations appear to largely overlap (Uezumi et al., 2014a). In addition, FAPs show overlap with Tcf4⁺ cells originally defined as muscle connective tissue fibroblasts (Murphy et al., 2011; Vallecillo-García et al., 2017) as well as with the PDGFRα⁺ subpopulation of PICs (PW1⁺ interstitial cells). PICs are Sca1⁺ and CD34⁺ and are marked by expression of the paternally imprinted gene PW1 (Peg3), which is a general stem cell/progenitor cell marker (Besson et al., 2011; Berg et al., 2011). PICs were originally characterized as an interstitial multipotent population distinct from satellite cells (Mitchell et al., 2010). Later it was shown that

* Corresponding author.

E-mail address: sigmar.stricker@fu-berlin.de (S. Stricker).

¹ These authors contributed equally to this work.

PICs can be divided into PDGFR α ⁺ adipogenic PICs that completely overlap with FAPs as well as PDGFR α ⁻ myogenic PICs (Pannerec et al., 2013). The above mentioned markers label tissue-resident quiescent FAPs as well as FAPs activated upon injury or disease. To date, no molecular marker has been found to identify injury-activated FAPs, and no tool exists to specifically purify or manipulate this population, precluding analyses as well as genetic manipulation or *in vivo* lineage tracing of injury-activated FAPs. The identification of an activated FAP-specific molecular marker promises to greatly facilitate our understanding of the intrinsic mechanisms of FAP activation, function, and differentiation under normal or pathological conditions. Here we show that FAPs become positive for the transcription factor Osr1 (Odd skipped-related 1) in response to injury and that Osr1 expression can be used to follow, isolate, and genetically mark activated FAPs during pathological and normal muscle repair.

2. Results and discussion

2.1. Osr1 is expressed in a small number of adult FAPs

During development, Osr1 marks a lateral plate mesoderm-derived population of fibro-adipogenic cells that is also a source for adult Sca1⁺ and PDGFR α ⁺ FAPs (Vallecillo-García et al., 2017). However, Osr1 expression declines during development and early postnatal life in mice, and eGFP expressed from the Osr1 locus [Osr1^{GCE} mouse line; 19] is only detectable *in situ* by antibody staining during early postnatal life but is below detectable levels in adults (Vallecillo-García et al., 2017). To increase detection sensitivity, we inserted a β -galactosidase (β -Gal) reporter into the Osr1 locus (Osr1^{LacZ}, Fig. S1A), which allows for enzymatic signal amplification. The expression of the Osr1^{LacZ} allele recapitulated the developmental expression pattern of Osr1 (Fig. S1B). Using the Osr1^{LacZ} line, we observed the presence of a low number of Osr1⁺ cells in the interstitium of several muscles examined (Fig. 1A). We noted that these cells are also positive for PDGFR α (Fig. 1B). FACS-cytospin of FAPs (lin⁻; α 7-integrin⁻;Sca1⁺) isolated from whole hindlimb muscle of Osr1^{LacZ} mice revealed that approx. 4.5% of adult FAPs were Osr1- β -Gal⁺ (Fig. 1C, Fig. S2). No β -Gal signal was detected in lin⁻; α 7-integrin⁻;Sca1⁻ cells (double negative, DN cells) or lin⁻; α 7-integrin⁺;Sca1⁻ cells (mainly satellite cells, SC) (Fig. 1C). We complemented this approach by isolating FAPs from PW1^{LacZ} animals (lin⁻;PW1⁺;PDGFR α ⁺); this protocol yielded a population that completely overlapped FAPs (Pannerec et al., 2013). Low abundance of Osr1 mRNA in adult FAPs was confirmed by semiquantitative PCR (Fig. 1D). This suggests that Osr1 is expressed in a small proportion of adult FAPs and is consistent with deep RNA sequencing data from adult resident FAPs showing low Osr1 mRNA expression (Ollitrault et al. in preparation).

2.2. Osr1 expression is induced to high levels upon acute injury

Freeze-pierce injury performed on Osr1^{LacZ} tibialis anterior (TA) muscle led to an accumulation of Osr1⁺ cells in the injured region 3 and 5 days post injury (dpi) (Fig. 2A). We used both Osr1^{GCE} and Osr1^{LacZ} alleles to analyze which cells initiated Osr1 expression. First, FAPs were FACS isolated from Osr1^{GCE} mice as lin⁻; α 7-integrin⁻;Sca1⁺ [4; Fig. S3A, B] and analyzed for Osr1-eGFP expression. Using this allele, we found that in uninjured muscle approx. 3.3% of FAPs expressed Osr1-eGFP (Fig. 2B), in agreement with results obtained from the Osr1^{LacZ} line (Fig. 1C). The numbers of Osr1-eGFP⁺ FAPs increased upon injury to 17–19% (approx. 5–6 fold increase) at 3, 5 and 7 dpi (Fig. 2B). At 10 dpi, the fraction of Osr1-eGFP⁺ FAPs decreased (8.5%; Fig. 2B). We note that the freeze-pierce injury we used results in a relatively small lesion leaving the larger part of the muscle uninjured. Cell isolation from the whole TA muscle yields FAPs mainly from non-injured and to a minor fraction only from injured regions. Therefore these numbers reflect the relative increase in Osr1⁺ FAPs amongst all

(injured and uninjured region-derived) FAPs, but does not reflect the absolute number of Osr1⁺ FAPs specifically within the regenerating region.

Osr1-eGFP was exclusively expressed in lin⁻; α 7-integrin⁻;Sca1⁺ FAPs and not detected in lin⁻; α 7-integrin⁺;Sca1⁻ SCs or lin⁻; α 7-integrin⁻;Sca1⁻ DN cells at 5 dpi (Fig. S3C). The induction of Osr1 was confirmed by detection of Osr1 protein in whole TA muscle lysates at 3dpi. We note that Osr1 protein levels are abundant in E13.5 limbs and injured adult TA muscles whereas Osr1 protein levels were below the level of detection in uninjured muscles (Fig. S4). Despite testing 5 commercially available antibodies (see Materials and Methods) we were not able to detect Osr1 protein on tissue sections after injury.

Cytospin analysis of all lin⁻; α 7-integrin⁻; Sca1⁺ FAPs (*i.e.* GFP⁺ and GFP⁻) followed by immunolabeling for PDGFR α confirmed that 18–20% of all PDGFR α ⁺ FAPs (from injured and uninjured parts of the TA muscle) were Osr1-GFP⁺ at 3, 5 and 7 dpi, while this ratio declined to 10.7% at 10 dpi (Fig. 2C). In contrast, more than 85% of Osr1⁺ FAPs were positive for PDGFR α at all time points analyzed (Fig. 2C, Fig. S5A). Both the Osr1⁺ and the Osr1⁻ fraction of Sca1⁺ FAPs overlapped partly with Tcf4 expression (Fig. 2C, Fig. S5A) indicating that there is no correlation between Osr1 and Tcf4 expression. This was confirmed by comparing Tcf4 and PDGFR α expression at 3 and 7 dpi as early and late stages after injury (Fig. S5B). Based on the finding that Tcf4 cells express PDGFR α (Murphy et al., 2011) it was suggested that FAPs and the Tcf4 pool largely overlap (Uezumi et al., 2014a). Of note, Tcf4 is expressed in interstitial cells in uninjured as well as in injured muscle parts (Murphy et al., 2011). We found a variable overlap of PDGFR α and Tcf4, suggesting that (at least in an injured muscle) only a proportion of Sca1⁺ and/or PDGFR α ⁺ FAPs express Tcf4, and that FAPs are a dynamic population following injury. Whether Osr1 and Tcf4 mark FAP subpopulations or represent different stages of FAP activation during the course of regeneration remains to be investigated. Interestingly, the fraction of Tcf4⁺/PDGFR α ⁺ cells decreased between 3 and 7 dpi specifically in the Osr1⁺ FAP pool (Fig. S5B). We note that part of the Tcf4⁺ cells in the Osr1⁺ FAP pool appeared to downregulate PDGFR α over time (Fig. S5B); the identity of Tcf4⁺/PDGFR α ⁻ cells remains to be determined. We further noted expression of Tcf4 in α 7-integrin⁺ myogenic cells (Fig. S5D) consistent with previous studies (Mathew et al., 2011).

To corroborate these findings, FAPs were isolated from Osr1^{LacZ} animals as lin⁻;Sca1⁺;CD34⁺ and analyzed *via* FACS for LacZ and PDGFR α expression (Fig. S6). Since LacZ staining can generate high background, we gated conservatively likely leading to exclusion of positive cells (Fig. S6A, B). Regardless, we could confirm Osr1-LacZ expression in 3 dpi lin⁻;Sca1⁺ FAPs, and that LacZ⁺;Sca1⁺ FAPs strongly overlapped with PDGFR α expression (Fig. S6C). Adult FAPs originate from a developmental Osr1⁺ lineage (Vallecillo-García et al., 2017). Consistent with these findings, long-term lineage tracing using Osr1^{GCE};R26^{LacZ} animals Tamoxifen pulsed at p0 and p1 shows that the progeny of Osr1⁺ developmental cells expand in the injury region upon acute muscle injury in adult animals (Fig. 2D). Taken together, these data show that Osr1 expression is induced upon muscle injury specifically in FAPs within the injured region.

2.3. Osr1 expression identifies injury-activated FAPs that contribute to adipogenic infiltration and post-injury resident FAPs

An initial rapid induction of proliferation is a hallmark of FAP activation in response to injury (Joe et al., 2010; Uezumi et al., 2010; Lemos et al., 2015) which is followed by apoptosis (Lemos et al., 2015). Following injury, we noted Ki67-stained Osr1⁺ cells in tissue sections from Osr1^{LacZ} animals (Fig. 3A). Next, FAPs were isolated (lin⁻; α 7-integrin⁻;Sca1⁺) from Osr1^{GCE} animals before injury and at 3, 5 and 7 dpi. FAPs showed virtually no proliferative activity in uninjured muscle (Fig. 3B). In fact, amongst overall 19,710 non-injury FAPs counted, only 62 Ki67⁺ cells (0.31%) were detected. After injury, Osr1⁺ FAPs




Fig. 1. *Osr1* is sporadically expressed in adult FAPs. (A) X-Gal staining on several *Osr1^{LacZ/+}* adult mouse muscles shows sporadic X-Gal⁺ cells. (B) X-Gal positive cells also express PDGFR α . (C) Isolation of CD45⁻;CD31⁻;Ter119⁻; α 7⁺ integrin⁺;Sca1⁺ FAPs from adult uninjured tibialis anterior of *Osr1^{LacZ}* mice followed by cytospin shows that only FAPs express *Osr1* ($n = 3$ animals). Data are represented as means \pm s.e.m. (D) Semiquantitative PCR on interstitial populations isolated from adult PW1^{LacZ} mice. Faint *Osr1* expression can be detected in PDGFR α ⁺ PICs.

showed a significantly higher fraction of Ki67⁺ cells than *Osr1*⁻ FAPs (Fig. 3B). Nevertheless, *Osr1*⁻ FAPs, which likely represent FAPs from the uninjured part of the TA muscle, showed considerable proliferative activity after injury. This observation may reflect an activation of FAPs to an “alert” state that has been reported in the contralateral muscles of injured mice including increased *in vitro* as well *in vivo* cycling (Rodgers et al., 2014). Notably, using the *Osr1^{GCE}* reporter, *Osr1*⁺ FAPs increased about twofold in numbers in contralateral muscle (6,9% in contralateral as compared to 3,3% in uninjured muscle; see Figs. 2B and S7). *Osr1*⁺ contralateral FAPs were 4,3% positive for Ki67 as compared to 1,6% of *Osr1*⁻ contralateral FAPs, although the difference did not reach statistical significance (Fig. S7). Thus, *Osr1* expression is activated upon induction of the alert state, but not all alert FAPs express *Osr1*. This together with the observation that ~10% of *Osr1*⁻ FAPs in injured muscle express Ki67 suggests that *Osr1* does not mark the entire activated FAPs population at any given time point. Whether this is due to dynamic changes in gene expression concomitant with FAPs differentiation or whether different FAPs subsets exists remains to be investigated. Since apoptosis is a default fate of the majority of injury-activated FAPs (Lemos et al., 2015), we tested *Osr1*⁺ and *Osr1*⁻ FAPs by immunolabeling for cleaved caspase-3 at 3 and 7 dpi representing early and late time points of recovery. At 3 dpi cleaved caspase-3 could only be detected in a small proportion of FAPs (70 out of 68,685 FAPs counted; approx. 0,1%) with no significant difference between *Osr1*⁺

and *Osr1*⁻ FAPs (Fig. 3C). However, at 7 dpi apoptotic cells were significantly increased in the *Osr1*⁺ FAPs population (Fig. 3C) consistent with the proposal that *Osr1*⁺ cells are activated in response to injury to undergo forced cell cycle entry as well as apoptosis.

We next tested the suitability of the *Osr1^{GCE}* allele (expressing an inducible CreErt2 recombinase from the *Osr1* locus) to trace the fate of activated FAPs. We genetically labelled *Osr1*⁺ cells in *Osr1^{GCE};R26^{mTmG}* mice for five consecutive days beginning with the day of injury (Fig. 3D). Since the *Osr1*⁺ FAPs population expands in the injured region during this period, we would anticipate labeling of the activated FAPs pool as compared to FAPs in uninjured muscle. Consistent with this prediction, we observed that pulsing *Osr1*⁺ cells for five days before injury resulted in low levels of labeling as compared to labeling post injury (Fig. S8). This also suggests that *Osr1*⁺ cells in adult uninjured muscle do not represent a specific subpopulation prone to quick expansion upon injury, rather, we propose that sporadic *Osr1* expression in uninjured adult muscle results from activated FAPs engaged in focal repair, however this requires further investigation. Lineage tracing in *Osr1^{GCE};R26^{mTmG}* mice induced after injury was performed at 28 days after injury, where regeneration is almost completed, however the regenerated tissue can be recognized by centrally located myonuclei. The majority of *Osr1*⁺ cell progeny after injury was traced to interstitial PDGFR α ⁺ cells representing resident FAPs (Fig. 3D). *Osr1*⁺ cells also gave rise to interstitial Tcf4⁺ cells (Fig. 3D). This suggests that




Fig. 2. Osr1 expression is reactivated in FAPs upon acute injury. (A) X-Gal staining on *Osr1^{LacZ/+}* adult tibialis anterior muscle 3 and 5 dpi shows accumulation of *Osr1-LacZ* positive cells in the injury region. (B) FACS analysis of *lin⁻;α7-integrin⁻;Sca1⁺* FAPs isolated before injury or at 3, 5, 7 and 10 dpi from *Osr1^{GCE/+}* mice for *Osr1-GFP* expression (n = 3 animals for each time point). Data are represented as means ± s.e.m. (C) Quantification of cytospin analysis of GFP⁺ and GFP⁻ FAPs isolated at indicated time points after injury from *Osr1^{GCE/+}* mice (n = 3 animals). Data are represented as means ± s.e.m. (D) Long-term lineage tracing using *Osr1^{GCE};R26^{LacZ}* animals Tamoxifen pulsed at p0 and p1; progeny of perinatal *Osr1⁺* cells stains positive for X-Gal.

Osr1⁺ activated FAPs return to the resident FAP pool after regeneration as well as to the Tcf4⁺ muscle connective tissue fibroblasts consistent with the proposal that FAPs are a primary source for fibrosis in degenerative disease (Uezumi et al., 2011; Lemos et al., 2015; Contreras et al., 2016; Mueller et al., 2016). It is tempting to speculate that Tcf4 might mark FAPs more prone for fibrogenic differentiation during muscle regeneration, which is supported by the notion that Tcf4 can inhibit adipogenesis *in vitro* (Cawthorn et al., 2007). Whether *Osr1* labels cells more likely to differentiate along the adipogenic lineage will be an interesting avenue to follow.

In addition to being a source of fibrotic tissue in pathologically remodeled skeletal muscle, FAPs are also proposed to be a source of fatty

infiltration (Uezumi et al., 2011; Uezumi et al., 2010; Kopinke et al., 2017; Lemos et al., 2012). Therefore, we tested the adipogenic potential of *Osr1⁺* injury-activated FAPs by glycerol injury in *Osr1^{GCE};R26^{mTmG}* mice, which results in fatty infiltration (Pisani et al., 2010) (Fig. 3E). 14 days after injury, the regenerating region contains infiltrating adipocytes which are mG positive (Fig. 3E). Taken together, these results provide a key confirmation for endogenous FAP fibrotic and adipogenic fates *in situ*.

2.4. *Osr1* marks a transient population of juvenile *Sca1⁺* cells

We noted previously that *Osr1* expression fades in early postnatal




Fig. 3. *Osr1* marks injury-activated fibro-adipogenic progenitors (FAPs). (A) X-Gal⁺ (*Osr1* expressing) cells in the injury region (*Osr1*^{LacZ/+} animal; 3dpi) are positive for the cell cycle marker Ki67. (B, C) Cytospin of lin⁻;α7-integrin⁺;Sca1⁺ FAPs FACS-isolated at indicated days post injury (dpi) and separated into GFP⁺ and GFP⁻ populations were stained for the proliferation marker Ki67 (B) and the apoptosis marker cleaved caspase-3 (C). Quantification shown right (n = 3 independent experiments from 3 animals). (D) Lineage tracing of *Osr1*⁺ cells in *Osr1*^{GCE};R26^{mTmG} animals Tamoxifen-pulsed for five consecutive days after freeze-pierce injury. Contribution of *Osr1*⁺ cells to PDGFRα⁺ and Tcf4⁺ interstitial cells was analyzed 28 days post injury (dpi). Quantification is shown below (n = 3 independent experiments from 3 animals). (E) Lineage tracing of *Osr1*⁺ cells in *Osr1*^{GCE};R26^{mTmG} animals Tamoxifen-pulsed for five consecutive days after glycerol injection injury. Contribution of *Osr1*⁺ cells to ectopic adipose tissue labelled for Perilipin (PLIN) was analyzed at 14 dpi. Data are represented as means +/- s.e.m. *t*-test: * = *P* < 0,05, ** = *p* < 0,01, *** = *p* < 0,005, n.s. = not significant.

life (Vallecillo-García et al., 2017), which is a period of still active myogenesis. Furthermore, early postnatal development represents a dynamic phase in which the transition of developmental progenitors to resident stem cells is accomplished. On this background, we finally re-analyzed *Osr1* expression in juvenile mice. In young mice (p7) β-Gal from the *Osr1*^{LacZ} allele was expressed in muscle interstitial cells (Fig. 4A), most of which were PDGFRα⁺ (86,1%; Fig. 4B). This indicates that *Osr1* is predominantly expressed in juvenile interstitial PDGFRα⁺ cells that likely are FAPs / FAP progenitors and hence should also express Sca1 (Uezumi et al., 2010; Pannerec et al., 2013).

Juvenile Sca1⁺ cells had been analyzed on the background of muscle interstitial PICs (Pannerec et al., 2013). This study showed that juvenile Sca1⁺ cells can be subdivided into two populations: one with low/medium Sca1 expression levels (Sca1^{med}) and a population with high Sca1 expression (Sca1^{hi}) that persists throughout life and completely overlaps with FAPs. We therefore FACS-isolated Sca1^{med};PW1⁺

and Sca1^{hi};PW1⁺ PICs from 7 day old *PW1*^{LacZ} mice as previously described (Pannerec et al., 2013) and analyzed *Osr1* expression by semi-quantitative PCR. Strong *Osr1* expression was seen in Sca1^{med}, whereas lower levels of expression were detected in Sca1^{hi} PICs. No detectable levels of expression were found in Sca1^{hi}PW1⁻ cells or in satellite cells (Fig. 4C).

To further characterize *Osr1*⁺ juvenile cells, we first FACS isolated lin⁻;CD34⁺;Sca1^{med} and lin⁻;CD34⁺;Sca1^{hi} cells from *Osr1*^{LacZ} mice (Fig. 4D). The majority of Sca1^{med} cells were β-Gal positive (79%), of which almost all Sca1^{med};β-Gal⁺ cells were PDGFRα positive (95%; Fig. 4E). In contrast, Sca1^{med};PDGFRα⁺ cells were mostly β-Gal positive (94%; Fig. 4F). The Sca1^{hi} population contained distinct and separable β-Gal⁺ and β-Gal⁻ subpopulations. Interestingly, of the Sca1^{hi};β-Gal⁺ population only approx. 50% expressed PDGFRα (Fig. 4G). In contrast, most of the Sca1^{hi};PDGFRα⁺ cells were β-Gal⁺ (78%; Fig. 4H), showing a prevalence for *Osr1* expression in the




Fig. 4. In juvenile mice *Osr1* is expressed in a transient *Sca1*^{med} population. (A) X-Gal staining on 7 day old *Osr1*^{LacZ/+} tibialis anterior muscle shows *Osr1*-LacZ⁺ interstitial cells. (B) *Osr1*-LacZ⁺ cells are mostly positive for PDGFR α *in situ* (n = 3 animals). (C) Semiquantitative PCR on interstitial populations isolated from 7 day old *PW1*^{LacZ} mice. Strong *Osr1* expression can be detected in *Sca1*^{med} PICs, while faint *Osr1* expression can be detected in *Sca1*^{hi} PICs. (D) Isolation of CD34⁺/*Sca1*⁺ cells from 7 day old tibialis anterior muscles; the *Sca1*^{med} and *Sca1*^{hi} populations are discernible. (E, F) Analysis of the *Sca1*^{med} population. (E) *Sca1*^{med} cells are mostly PDGFR α ⁺; *Sca1*^{med} PDGFR α ⁺ cells mostly express β -Gal. (F) *Sca1*^{med}; β -Gal⁺ cells express PDGFR α . (G, H) Analysis of *Sca1*^{hi} cells. (G) *Sca1*^{hi} cells are only in part positive for PDGFR α ; *Sca1*^{hi}; PDGFR α ⁺ cells mostly express β -Gal. (H) Only approx. Half of *Sca1*^{hi} cells express β -Gal. (*Sca1*^{hi}; β -Gal⁺ cells are also only in part PDGFR α ⁺. (I-L) Lineage analysis of perinatal *Osr1*⁺ cells. (I) Lineage tracing strategy. Tamoxifen was administered at p0/p1, analysis was performed at 3 weeks of age; adipogenic differentiation was analyzed at 7 weeks of age. (J) *Osr1*⁺ progeny majorly express PDGFR α , but in part also Tcf4. Quantification shown right (n = 2 animals). (K) *Osr1*⁺ cells do not give rise to CD31⁺ endothelial cells or α SMA⁺ smooth muscle cells. (L) *Osr1*⁺ cells give rise to muscle interstitial adipocytes labelled for Perilipin (PLIN). Data are represented as means \pm s.e.m.

PDGFR α fraction. Taken together, these results show that *Osr1* is expressed primarily in interstitial *Sca1*^{med};PDGFR α ⁺ cells and to lower extent in *Sca1*^{hi};PDGFR α ⁺ cells in juvenile muscle.

We next re-analyzed the developmental fate of perinatal *Osr1*⁺ cells by genetically labeling the *Osr1* lineage in *Osr1*^{GCE};R26^{mTmG} mice. Administration of Tamoxifen (TAM) at p0 and p1 was used to developmentally trace *Osr1*⁺ cells (Fig. 4I). At day p21, *Osr1*⁺ progenitors contributed to the muscle interstitial PDGFR α cell pool (approx. 70%; Fig. 4J) in line with our previous demonstration that adult FAPs derive from these developmental *Osr1*⁺ cells (Vallecillo-García et al., 2017). *Osr1* descendants were also present in cells expressing the muscle connective tissue fibroblast marker, Tcf4, although at a markedly lower number (Fig. 4J). No contribution of *Osr1*⁺ progenitors was seen in α SMA⁺ vascular smooth muscle or CD31⁺ endothelium (Fig. 4K). In addition, *Osr1*⁺ progenitors gave rise to muscle interstitial adipocytes (Fig. 4L). This shows that juvenile *Osr1*⁺ cells maintain plasticity and that interstitial fibroblasts as well as interstitial adipocytes derive from this pool in addition to FAPs.

During embryonic development, *Osr1* is expressed in *Sca1*⁺ cells whereas *Sca1*-expressing muscle interstitial cells arise during late fetal development, concomitant with the appearance of *Sca1*⁺/*Osr1*⁺ cells (Vallecillo-García et al., 2017). While it remains unresolved whether *Osr1*⁺ cells acquire *Sca1* expression during fetal development or whether the *Sca1*⁺ cells arise *de novo*, our observation that *Osr1* is predominantly expressed in a transient population of *Sca1*^{med} FAPs during postnatal development has interesting implications. Specifically, we note that *Osr1* is expressed in muscle interstitial mesenchymal cells throughout embryonic and postnatal myogenesis, first in *Sca1*⁺, later in *Sca1*^{med} and to a lesser degree in *Sca1*^{hi} cells. This pattern of expression suggests a lineage continuum in which *Sca1*^{hi} cells are derived from *Sca1*^{med} cells, and that downregulation of *Osr1* expression is required for this differentiation step. Interestingly, both the disappearance of *Sca1*^{med} cells and the downregulation of *Osr1* expression correlate with the termination of postnatal muscle growth driven by incorporation of new nuclei into myofibers (Pannerec et al., 2013; White et al., 2010). These data together with the observation of re-activation of *Osr1* expression in injury-activated FAPs leads to the proposal that *Osr1* expression marks FAPs or their progenitors during periods of active myogenesis. During embryonic myogenesis, *Osr1*⁺ cells provide a pro-myogenic niche for myogenic progenitors that promotes myogenic cell proliferation and survival. (Vallecillo-García et al., 2017). Similarly, adult FAPs promote myogenesis *in vitro* (Joe et al., 2010) and Tcf4⁺ cells, which in part overlap with FAPs, are required for efficient muscle regeneration (Murphy et al., 2011). The re-activation of *Osr1* expression further suggests that adult FAPs reactivate a developmental program to support tissue regeneration.

In summary, we show that *Osr1* is the first specific marker identified for injury activated FAPs. Given the key role for FAPs in promoting proper muscle regeneration, the ability to lineage track these cells will be valuable for designing approaches to optimize muscle repair and target the muscle stem cell niche.

3. Materials and methods

3.1. Mice

Mice were maintained in an enclosed, pathogen-free facility. The targeting construct for the *Osr1* multifunctional allele was electroporated into G4 mouse ES cells (George et al., 2007). The transgenic locus was confirmed by Southern blotting and Sanger sequencing. Successfully recombined clones were subjected to tetraploid aggregation. Mice were crossed back to wild type C57BL/6j mice for at least 6 generations before establishing the line. The following mouse lines were described before: *Osr1*^{GCE} (Mugford et al., 2008); R26^{LacZ} (Soriano, 1999); R26^{mTmG} (Muzumdar et al., 2007). Mouse experiments were performed in accordance with European Union regulations

(Directive 2010/63/EU) and under permission from the Landesamt für Gesundheit und Soziales (LaGeSo) Berlin, Germany (Permission numbers ZH120, G0240/11, G0114/14, G0209/15, G0268/16).

3.2. Cell lineage tracing

Lineage tracing of neonatal *Osr1*⁺ cells was performed in *Osr1*^{GCE/+};R26^{LacZ/+} neonates by subcutaneous injection of Tamoxifen (Sigma Aldrich; solved in 90% (v/v) sunflower oil/ 10% (v/v) ethanol) into the neck fold (75 μ g/ g body weight). Lineage tracing of injury-activated *Osr1*⁺ FAPs was performed in *Osr1*^{GCE/+};R26^{mTmG/+} or *Osr1*^{GCE/+};R26^{LacZ/+} mice by intraperitoneal injection of Tamoxifen at the day of injury and the next 4 following days (3 mg per injection time point). For lineage tracing of *Osr1*⁺ cells before injury, *Osr1*^{GCE/+};R26^{LacZ/+} and *Osr1*^{GCE/+};R26^{mTmG/+} animals were injected with 3 mg Tamoxifen for 5 consecutive days one week before injury.

3.3. Muscle injury

Injury was applied to the tibialis anterior muscle of 3–5 months old *Osr1*^{GCE}, *Osr1*^{LacZ}, *Osr1*^{GCE};R26^{mTmG} or *Osr1*^{GCE};R26^{LacZ} mice using the freeze-pierce technique. Mice were anesthetized by intraperitoneal injection of 10% (v/v) ketamine / 2% (v/v) xylazine (Rompun® 2%) in sterile PBS (5 μ l / g body weight). Mice were kept on a heating plate warmed to 37 °C throughout the procedure. The skin above the Tibialis anterior muscle was opened and the muscle was longitudinally pierced 5 times using a 0.7 mm liquid nitrogen cooled syringe needle.

For glycerol injury mice were anesthetized as described above. Skin above the Tibialis anterior muscle was opened and 25 μ l of 50% v/v glycerol/sterile PBS were injected into the muscle.

3.4. Histology, antibody labeling, western blot

Muscle was dissected, immediately embedded in 6% (w/v) gum tragacanth (Sigma Aldrich) dissolved in H₂O and snap frozen in liquid nitrogen cooled isopentane (–160 °C). Frozen tissue was sectioned at 7 μ m and fixed using 4% (w/v) PFA in PBS for 5 min at RT.

Permeabilization of sections was performed in 0.3% (v/v) Triton X-100 (Sigma Aldrich) in phosphate buffer (PBS) for 10 min. Sections from adult tissues were blocked with 5% (w/v) bovine serum albumin (Sigma Aldrich) in PBS. Sections then preincubated with 5% horse serum, 5 mg mL⁻¹ blocking reagent (Perkin Elmer) and 0.1% Triton X-100 in PBS for 1 h at RT. Primary antibodies were applied in the same solution and incubated at 4 °C overnight, followed by secondary antibody staining for 1 h at room temperature. Counterstaining was performed with 5 μ g μ L⁻¹ 4',6-diamidino-2-phenylindole (DAPI; Invitrogen), slides were mounted with FluoromountG (SouthernBiotech). Antibodies are listed in Supplementary tables 1 and 2.

For beta galactosidase detection, sections or cells were fixed as above and incubated at 37 °C overnight in X-Gal (0.16% (w/v) X-Gal, 5 mM K₃Fe(CN)₆, 5 mM K₄Fe(CN)₆, 2 mM MgCl₂ in PBS). Beta galactosidase development was stopped by washing in PBS. For whole-mount X-Gal stainings embryos were fixed in X-Gal fixing solution (1% (w/v) Formaldehyde, 0.2% (w/v) Glutaraldehyde, 0.02% (v/v) NP-40, 1% (v/v) PBS in bidest H₂O) for 1 h at 4 °C, then stained for 24 h at 37 °C in X-Gal staining solution.

Protein extraction and western blot were performed according to standard procedures. In brief, muscle tissue was homogenized, RIPA buffer (150 mM sodium chloride; 1.0% NP-40 or Triton X-100; 0.5% sodium deoxycholate; 0.1% SDS (sodium dodecyl sulfate); 50 mM Tris, pH 8.0) was added (0.4 ml per 0.1 g of tissue), and incubated on ice for 15 min. After brief centrifugation supernatant was mixed 5:1 with loading buffer (2.4 g SDS; 12 mg bromophenol blue; 9.4 ml glycerol; 2.4 ml Tris 6.8 (0.5 M); 1.8 ml beta-mercaptoethanol; 4.2 ml dd H₂O) and heated at 98 °C for 10 min. An aliquot of the supernatant was used

for determination of protein concentration (Pierce / Thermo Scientific BSA Protein Assay Kit). 50 µg of Protein was loaded per lane on an 8% SDS polyacrylamid gel. Gels were blotted to Amersham Hybond P 0.45 PVDF blotting membrane (activated in methanol for 1 min before use). Membranes were blocked with 5% milk powder (Roth) in TBST, antibody incubation (antibodies see supplementary table 1) was performed in blocking solution at 4 °C over night. Membranes were washed three times with TBST, antibody detection was performed with the ECL Kit (Pierce / Thermo Scientific) according to the manufacturer's instructions. Imaging and quantification were performed on a Fusion FX7 (Vilber Lourmat).

Antibodies used for attempting *Osr1* immunolabeling on fresh frozen sections of injured muscle: Sigma-Aldrich #HPA015525, #WH0130497M4; Abnova # H00130497-M04; Santa Cruz #sc-376,545; Abcam #ab74003.

3.5. Cell isolation and flow cytometry

Cell isolation and labeling was essentially performed as described in (Vallecillo-García et al., 2017). Isolation of cells from *Osr1*^{GCE/+} mice was performed as follows: Briefly, whole hind limb or *tibialis anterior* muscles were carefully isolated, roughly minced and digested in high-glucose DMEM medium containing 10% fetal calf serum (FCS, Biochrom), 1% Penicillin Streptomycin solution (P/S; 10,000 U/ml) and 2,5 mg/ml Collagenase A (Roche) for 75 min at 37 °C with vigorous shaking. 2 IU/ml of Dispase II (Sigma Aldrich) were added to the digestion solution and muscle lysates were digested for further 30 min. Muscle slurries were passed 10 times through a 20G syringe (BD Bioscience) and a 70-µm cell strainer. Cells were collected by centrifugation at 300 g for 5 min and resuspended in staining buffer consisting of 500 µl Hank's balanced salt solution (HBSS, Thermo Fisher scientific), 0.4% bovine serum albumin (Sigma Aldrich) and 20 µg/ml Gentamycin (Serva Electrophoresis). Cells were stained on ice for 30 min and washed twice with staining buffer previous to FACS sorting. Propidium iodide was used as a viability dye. Isolation of cells from *Osr1*^{LacZ/+} animals was performed as follows: freshly dissected muscle tissue was minced and digested in HBSS (Gibco) supplemented with 2.4 U/ml Dispase II (Roche), 2 µg/ml Collagenase A (Roche), 0.4 mM CaCl₂, 5 mM MgCl₂, 10 ng/ml DNase I (Roche) for 2 h at 37 °C under agitation. Single cell suspension was obtained after 3 successive cycles of straining and washing in Washing Buffer consisting of HBSS containing 0.2% (w/v) BSA (Sigma Aldrich), 1% (v/v) penicillin-streptomycin, 10 ng/ml DNase I and 10% (v/v) mouse serum. Cells were incubated with primary antibodies for 1 h on ice. The suspension was subjected to 2 washing steps, resuspended in Washing Buffer and to LacZ reporter staining using C₁₂FDG (Life Technologies). For lacZ staining C₁₂FDG was added to the cell suspension to a final concentration of 600 µM and incubated for 30 min at 37 °C. 2 washing steps followed before proceeding to FACS analysis. Cells purified by sorting were cytospun to coverslips. Coverslips were coated with poly-L-lysine by incubation with a 10-fold solution of poly-lysine in bidest H₂O for 1 h at RT, rinsed twice in bidest H₂O and air dried. Purified cells were added to prepared slides and allowed to adhere for 1 h at 4 °C. Supernatant was removed by centrifugation at 50 rcf for 5 min at 4 °C, cells were fixed using 4% (w/v) PFA in PBS for 5 min at RT and permeabilized with 0.3% (v/v) Triton X-100 in PBS for 5 min at RT. Antibodies are listed in Additional file 1, Supplementary tables 1 and 2. Sorts and analyses were performed on a FACS Aria II (BD Biosciences). Sorting gates were defined based on unstained controls. Data were collected using FACSDIVA software. Biexponential analyses were performed using FlowJo 10 (FlowJo LLC) software. Analysis was performed on three independent biological replicates.

3.6. Cell quantification

Quantification of FACS-isolated *Osr1*⁺ cells after immunolabeling

(cytospin) was performed on at least three independent biological replicates (*i.e.* cells FACS isolated from three different animals). Quantifications of cytospun cells for X-Gal staining and PDGFRα, Tcf4 were made from two areas of 0.81 mm² per replicate. Quantification of Ki67⁺ and cleaved caspase 3⁺ cells was performed on a tile scan image comprising the whole cover slip. The image analysis module of the Leica Las X software was used for cell counting. Student's *t*-test was performed using Prism 5 (GraphPad) software. Error bars in all figures, including supplementary information, represent the mean ± standard error of the mean (s.e.m.).

3.7. Microscopy

Images were acquired using a Zeiss LSM700 confocal microscope, a Leica DMR or Leica DMI8 microscope. Bright field images of whole-mount embryos were taken with a Leica Leica MZ12 stereo microscope. Images were captured using Axio Vision Rel. 4.8 and Zen 2010 (Zeiss) or LAS X (Leica).

Acknowledgements

We thank the animal facility of the Max Planck Institute for Molecular Genetics, Berlin for expert support, especially Karol Macura, Judith Fiedler, Lars Wittler, Andrea König, Katja Zill and Ludger Hartmann.

Competing interests

No competing interests declared.

Funding

This work was funded by the German Research Foundation (DFG; grant GK1631), French-German University (UFA-DFH; grant CDFa-06-11), the Association Française contre les Myopathies (AFM 16826), and the Fondation pour la Recherche Médicale (FRM DEQ20140329500) as part of the MyoGrad International Research Training Group for Myology. This work was funded by the Focus Area DynAge of the Freie Universität Berlin. We acknowledge support by the German Research Foundation and the OpenAccess Publication Fund of the Freie Universität Berlin.

Author's contributions

DAS and SS conceived and designed the study. JS, PVG, SvHS and DO performed experiments and collected data. JS, PVG, SvHS, DO, GM, DAS and SS performed data analysis and interpretation. ANE generated the *Osr1*-LacZ knock-in construct and HS generated the *Osr1*-LacZ knock-in ES cells. DAS and SS wrote the manuscript with help from JS, PVG and GM.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scr.2018.08.010>.

References

- Bentzinger, C.F., Wang, Y.X., Dumont, N.A., Rudnicki, M.A., 2013. Cellular dynamics in the muscle satellite cell niche. *EMBO Rep.* 14, 1062–1072.
- Berg, J.S., Lin, K.K., Sonnet, C., Boles, N.C., Weksberg, D.C., Nguyen, H., Holt, L.J., Rickwood, D., Daly, R.J., Goodell, M.A., 2011. Imprinted genes that regulate early mammalian growth are coexpressed in somatic stem cells. *PLoS One* 6, e26410.
- Besson, V., Smeriglio, P., Wegener, A., Relaix, F., Nait Oumesmar, B., Sassoon, D.A., Marazzi, G., 2011. PW1 gene/paternally expressed gene 3 (PW1/Peg3) identifies multiple adult stem and progenitor cell populations. *Proc. Natl. Acad. Sci. U. S. A.* 108, 11470–11475.
- Cawthorn, W.P., Heydt, F., Hegyi, K., Sethi, J.K., 2007. Tumour necrosis factor-alpha

- inhibits adipogenesis via a beta-catenin/TCF4(TCF7L2)-dependent pathway. *Cell Death Differ.* 14, 1361–1373.
- Contreras, O., Rebolledo, D.L., Oyarzun, J.E., Olguin, H.C., Brandan, E., 2016. Connective tissue cells expressing fibro/adipogenic progenitor markers increase under chronic damage: relevance in fibroblast-myofibroblast differentiation and skeletal muscle fibrosis. *Cell Tissue Res.* 364, 647–660.
- George, S.H., Gertsenstein, M., Vintersten, K., Korets-Smith, E., Murphy, J., Stevens, M.E., Haigh, J.J., Nagy, A., 2007. Developmental and adult phenotyping directly from mutant embryonic stem cells. *Proc. Natl. Acad. Sci. U. S. A.* 104, 4455–4460.
- Gonzalez, D., Contreras, O., Rebolledo, D.L., Espinoza, J.P., van Zundert, B., Brandan, E., 2017. ALS skeletal muscle shows enhanced TGF-beta signaling, fibrosis and induction of fibro/adipogenic progenitor markers. *PLoS One* 12, e0177649.
- Joe, A.W., Yi, L., Natarajan, A., Le Grand, F., So, L., Wang, J., Rudnicki, M.A., Rossi, F.M., 2010. Muscle injury activates resident fibro/adipogenic progenitors that facilitate myogenesis. *Nat. Cell Biol.* 12, 153–163.
- Kopinke, D., Roberson, E.C., Reiter, J.F., 2017. Ciliary hedgehog signaling restricts injury-induced Adipogenesis. *Cell* 170, 340–351 (e312).
- Lemos, D.R., Paylor, B., Chang, C., Sampaio, A., Underhill, T.M., Rossi, F.M., 2012. Functionally convergent white adipogenic progenitors of different lineages participate in a diffused system supporting tissue regeneration. *Stem Cells* 30, 1152–1162.
- Lemos, D.R., Babaeijandaghi, F., Low, M., Chang, C.K., Lee, S.T., Fiore, D., Zhang, R.H., Natarajan, A., Nedospasov, S.A., Rossi, F.M., 2015. Nilotinib reduces muscle fibrosis in chronic muscle injury by promoting TNF-mediated apoptosis of fibro/adipogenic progenitors. *Nat. Med.* 21, 786–794.
- Mathew, S.J., Hansen, J.M., Merrell, A.J., Murphy, M.M., Lawson, J.A., Hutcheson, D.A., Hansen, M.S., Angus-Hill, M., Kardon, G., 2011. Connective tissue fibroblasts and Tcf4 regulate myogenesis. *Development* 138, 371–384.
- Mitchell, K.J., Pannerec, A., Cadot, B., Parlakian, A., Besson, V., Gomes, E.R., Marazzi, G., Sassoon, D.A., 2010. Identification and characterization of a non-satellite cell muscle resident progenitor during postnatal development. *Nat. Cell Biol.* 12, 257–266.
- Mozzetta, C., Consalvi, S., Saccone, V., Tierney, M., Diamantini, A., Mitchell, K.J., Marazzi, G., Borsellino, G., Battistini, L., Sassoon, D., Sacco, A., Puri, P.L., 2013. Fibroadipogenic progenitors mediate the ability of HDAC inhibitors to promote regeneration in dystrophic muscles of young, but not old mdx mice. *EMBO molecular medicine* 5, 626–639.
- Mueller, A.A., van Velthoven, C.T., Fukumoto, K.D., Cheung, T.H., Rando, T.A., 2016. Intronic polyadenylation of PDGFRalpha in resident stem cells attenuates muscle fibrosis. *Nature* 540, 276–279.
- Mugford, J.W., Sipila, P., McMahon, J.A., McMahon, A.P., 2008. Osr1 expression demarcates a multi-potent population of intermediate mesoderm that undergoes progressive restriction to an Osr1-dependent nephron progenitor compartment within the mammalian kidney. *Dev. Biol.* 324, 88–98.
- Murphy, M.M., Lawson, J.A., Mathew, S.J., Hutcheson, D.A., Kardon, G., 2011. Satellite cells, connective tissue fibroblasts and their interactions are crucial for muscle regeneration. *Development* 138, 3625–3637.
- Muzumdar, M.D., Tasic, B., Miyamichi, K., Li, L., Luo, L., 2007. A global double-fluorescent Cre reporter mouse. *Genesis* 45, 593–605.
- Pannerec, A., Marazzi, G., Sassoon, D., 2012. Stem cells in the hood: the skeletal muscle niche. *Trends Mol. Med.* 18, 599–606.
- Pannerec, A., Formicola, L., Besson, V., Marazzi, G., Sassoon, D.A., 2013. Defining skeletal muscle resident progenitors and their cell fate potentials. *Development* 140, 2879–2891.
- Pisani, D.F., Bottema, C.D., Butori, C., Dani, C., Dechesne, C.A., 2010. Mouse model of skeletal muscle adiposity: a glycerol treatment approach. *Biochem. Biophys. Res. Commun.* 396, 767–773.
- Rodgers, J.T., King, K.Y., Brett, J.O., Cromie, M.J., Charville, G.W., Maguire, K.K., Brunson, C., Mastey, N., Liu, L., Tsai, C.R., Goodell, M.A., Rando, T.A., 2014. mTORC1 controls the adaptive transition of quiescent stem cells from G0 to G(Alert). *Nature* 510, 393–396.
- Soriano, P., 1999. Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat. Genet.* 21, 70–71.
- Uezumi, A., Fukada, S., Yamamoto, N., Takeda, S., Tsuchida, K., 2010. Mesenchymal progenitors distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle. *Nat. Cell Biol.* 12, 143–152.
- Uezumi, A., Ito, T., Morikawa, D., Shimizu, N., Yoneda, T., Segawa, M., Yamaguchi, M., Ogawa, R., Matev, M.M., Miyagoe-Suzuki, Y., Takeda, S., Tsujikawa, K., Tsuchida, K., Yamamoto, H., Fukada, S., 2011. Fibrosis and adipogenesis originate from a common mesenchymal progenitor in skeletal muscle. *J. Cell Sci.* 124, 3654–3664.
- Uezumi, A., Ikemoto-Uezumi, M., Tsuchida, K., 2014a. Roles of nonmyogenic mesenchymal progenitors in pathogenesis and regeneration of skeletal muscle. *Front. Physiol.* 5, 68.
- Uezumi, A., Fukada, S., Yamamoto, N., Ikemoto-Uezumi, M., Nakatani, M., Morita, M., Yamaguchi, A., Yamada, H., Nishino, I., Hamada, Y., Tsuchida, K., 2014b. Identification and characterization of PDGFRalpha+ mesenchymal progenitors in human skeletal muscle. *Cell Death Dis.* 5, e1186.
- Vallecillo-García, P., Orgeur, M., Vom Hofe-Schneider, S., Stumm, J., Kappert, V., Ibrahim, D.M., Börno, S.T., Hayashi, S., Relaix, F., Hildebrandt, K., Sengle, G., Koch, M., Timmermann, B., Marazzi, G., Sassoon, D.A., Duprez, D., Stricker, S., 2017. Odd skipped-related 1 identifies a population of embryonic fibro-adipogenic progenitors regulating myogenesis during limb development. *Nature Communications* 8, 1218.
- White, R.B., Bierinx, A.S., Gnocchi, V.F., Zammit, P.S., 2010. Dynamics of muscle fibre growth during postnatal mouse development. *BMC Dev. Biol.* 10, 21.