



**HAL**  
open science

## Draft genomes of “*Pectobacterium peruviense*” strains isolated from fresh water in France

Pierre Faye, Claire Bertrand, Jacques Pédrón, Marie-Anne Barny

### ► To cite this version:

Pierre Faye, Claire Bertrand, Jacques Pédrón, Marie-Anne Barny. Draft genomes of “*Pectobacterium peruviense*” strains isolated from fresh water in France. *Standards in Genomic Sciences*, 2018, 13, pp.27. 10.1186/s40793-018-0332-0 . hal-01922694

**HAL Id: hal-01922694**

<https://hal.sorbonne-universite.fr/hal-01922694v1>

Submitted on 14 Nov 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

SHORT GENOME REPORT

Open Access



# Draft genomes of “*Pectobacterium peruvienne*” strains isolated from fresh water in France

Pierre Faye, Claire Bertrand, Jacques Pédrón and Marie-Anne Barny\* 

## Abstract

Bacteria belonging to the genus *Pectobacterium* are responsible for soft rot disease on a wide range of cultivated crops. The “*Pectobacterium peruvienne*” specie, recently proposed inside the *Pectobacterium* genus, gathers strains isolated from potato tubers cultivated in Peru at high altitude. Here we report the draft genome sequence of two strains belonging to “*P. peruvienne*” isolated from river water in France indicating that the geographic distribution of this specie is likely to be larger than previously anticipated. We compared these genomes with the one published from the “*P. peruvienne*” specie type strain isolated in Peru.

**Keywords:** *Pectobacterium peruvienne*, Soft rot, Plant pathogen, Water, France

## Introduction

The *Pectobacterium* genus [1] gathers important plant pathogens that cause soft rot disease on a large variety of plant species [2]. Given their ability to cause disease on major crops, such as potato, *Pectobacterium* sp. have mainly been isolated from diseased plant during initial outbreak or sustained epidemic and their descriptions outside of agricultural context is scarce [3].

The classification of the *Pectobacterium* genus has been subject to extensive revision over the last decade. It is currently subdivided in 7 species; *P. carotovorum* [1], *P. atrosepticum* [4], *P. betavascularum* [4], *P. wasabiae* [4], *P. aroidearum* [5] *P. polaris* [6], *P. parmentieri* [7] and the recently proposed “*P. peruvienne*” [8]. The *P. carotovorum* specie is heterogeneous and is currently subdivided several recognized subspecies, *P. carotovorum* subsp. *carotovorum* [9, 10], *P. carotovorum* subsp. *odoriferum* [9, 10] and proposed subspecies “*P. carotovorum* subsp. *actinidiae*” [11] and “*P. carotovorum* subsp. *brasiliense*” [12]. This heterogeneity led to assignation of many *Pectobacterium* isolates to *P. carotovorum*. One example is the strain UG32 (also named IFB5232, SCRI179, LMG30269 and PCM2893) that was initially described as *P. carotovorum* subsp. *carotovorum*

and is now the proposed type strain of the “*P. peruvienne*” specie [8, 13]. All the strains described so far in the “*P. peruvienne*” specie have been isolated in Peru in the seventies during the twentieth century from potato plants cultivated at high altitude (2400–3800 m). Here we described the draft genome sequence of two strains A97-S13-F16 and A350-S18-N16 isolated in February and November 2016 at different altitudes in the Durance river stream in France.

## Organism information

### Classification and features

Strain A97-S13-F16 was isolated in february 2016 from fresh water sampled in the river Durance while strain A350-S18-N16 was isolated in november 2016 from fresh water sampled in river Bléone, close to the confluent with river Durance. The fresh water parameters measured at the sampling times respectively were the following respectively for A97-S13-F16 and A350-S18-N16 sampled water: temperature 6.4 °C and 10.4 °C; turbidity 2.69 NTU and 145 NTU, conductivity 629 μS and 629 μS. Following sampling, 500 ml of fresh water was filtered through 0.2 μm pore filters (Sartorius cellulose acetate filters), the bacteria present on the filters were suspended in 1 ml sterile distilled water and 100 μl of the suspension were poured onto semi selective modified single-layers CVP<sub>AG366</sub> plates (same medium as

\* Correspondence: [marie-anne.barny@sorbonne-universite.fr](mailto:marie-anne.barny@sorbonne-universite.fr)  
Institute of Ecology and Environmental sciences-Paris, Sorbonne Université,  
INRA, 4 place Jussieu, F-75 252 Paris, France



described in [14] except that tryptone was not added to the medium, hereafter described as CVP). After 2 days of growth at 28 °C, two strains forming pits on CVP medium were further isolated, named A97-S13-F16 and A350-S18-N16 and stored in 40% /60% glycerol/ LB liquid medium (10 g tryptone, 5 g yeast extract, 10 g NaCl per one liter of medium) at – 80 °C.

Cells of both strains are rod shaped with length of approximately 2 µm in the exponential growth phase on LB medium (Fig. 1) and both strains are macerating potato tubers (Additional file 1: Figure S1). They are forming isolated colonies after 24 h at 28 °C on LB-15 g agar medium and after 48 h at 28 °C on TSA 10% medium (1,5 g tryptone, 0,5 g soy peptone, 0,5 g NaCl, 15 g agar per one liter of medium) and are inducing pits in CVP medium after 48 h at 28 °C.

Amplification and sequencing of the *gapA* house keeping gene was recently described to rapidly characterize the different *Pectobacterium* species [15]. The *gapA* sequences of strains A97-S13-F16 and A350-S18-N16 clustered with the one of proposed “*P. peruvienne*” type strain (Fig. 2A) and the clusterization of both strains with “*P. peruvienne*” was confirmed through MLSA analysis of full genomes (Fig. 2B).

General feature of A97-S13-F16 and A350-S18-N16 are indicated in Table 1.

## Genome sequencing information

### Genome project history

The aim of the project was to described *Pectobacterium* sp. isolated from environmental samples outside agricultural context. Fresh water sampling was performed in the river Durance and its tributaries in 2016. Amongst the isolated strains, the two strains A97-S13-F16 and


A350-S18-N16, isolated in different locations and at different months in the river stream, were selected for sequencing following amplification and sequencing of their *gapA* house keeping gene because phylogenetic analysis of their *gapA* sequences positioned both *gapA* sequences close to the *gapA* sequence of the recently proposed “*P. peruvienne*” type strain UGC32 [8, 13, 15].

### Growth conditions and DNA isolation


After isolation from fresh water in 2016, strains A97-S13-F16 and A350-S18-N16 have been stored in 40%/60% glycerol /LB medium at – 80 °C. For preparation of genomic DNA, the strains were first grown overnight at 28 °C on solid LB medium. A single colony was then pick up and grown overnight in 2 ml of liquid LB medium at 28 °C with 120 rpm shaking. Bacterial cells were harvested by centrifugation (5 min at 12,000 rpm) and DNA was extracted with the wizard® genomic DNA extraction kit (Promega) following the supplier specification. DNA was suspended in 100 µl of sterile distilled water and the quantity and quality of DNA was assessed by nano-drop measurement, spectrophotometry analysis and gel analysis.

### Genome sequencing and assembly

Genome sequencing was performed at the next generation sequencing core facilities of the Institute for Integrative Biology of the Cell, Bât. 21, Avenue de la Terrasse 91,190 Gif-sur-Yvette Cedex France. Nextera DNA libraries were prepared from 50 ng of high quality genomic DNA. Paired end 2×75 bp sequencing was performed on an Illumina NextSeq500 instrument, with a High Output 150 cycle kit.



**Fig. 1** Photomicrographs of Gram stained exponentially growing “*P. peruvienne*” cells. (a) strain A97-S13-F16, (b) A350-S18-N16. A light microscope with 100X magnification was used. These photomicrographs show the rod shaped forms of both strains. The bar scale represent 5 µm



CLC Genomics Workbench (Version 9.5.2, Qiagen Bioinformatics) was used to assemble 30,066,500 (mean length 53 bp) and 8,174,334 reads (mean length 52 bp) for strains A97-S13-F16 and A350-S18-N16 respectively. Final sequencing coverages were 331× and 86× with 61 and 73 scaffolds for strains A97-S13-F16 and A350-S18-N16 respectively (Table 2).

#### Genome annotation

Coding sequences were predicted using the RAST server [16] with the Glimmer 3 prediction tool [17]. COG assignments and Pfam domain predictions were done using the Web CD-Search Tool [18]. CRISPRFinder [19] was used to detect CRISPRs. Signal peptide and transmembrane domain were detected with the SignalP 4.1

Server [20] and transmembrane helices were predicted with TMHMM [21].

#### Genomes properties

The "*P. peruvienne*" A97-S13-F16 draft genome contains 4,775,191 bp with a GC content of 51%. Total predicted genes are 4503 while predicted protein coding genes are 4459 and RNA genes 44. The final assembly comprised 61 scaffolds. Among the predicted genes, 72.21% have a predicted function, 79.91% were assigned to COG and 85.40% have a predicted Pfam domain. Among the predicted proteins, 392 have a predicted signal peptide while 1090 contain a predicted transmembrane helix. Three CRISPR repeats array were detected in this genome.

**Table 1** Classification and general features of strains A97-S13-F16 and A350-S18-N16

MIGS ID	Property	Term	Evidence code <sup>a</sup>	
	Classification	Domain <i>Bacteria</i>	TAS [14]	
		Phylum <i>Proteobacteria</i>	TAS [15]	
		Class <i>Gammaproteobacteria</i>	TAS [16]	
		Order <i>Enterobacterale</i>	TAS [17]	
		Family <i>Pectobacteriaceae</i>	TAS [17]	
		Genus <i>Pectobacterium</i>	TAS [18]	
		Species <i>Pectobacterium peruvienne</i> strains: A97-S13-F16 and A350-S18-N16	NAS	
		Gram stain	<i>Negative</i>	NAS
		Cell shape	<i>Rod</i>	IDA
		Motility	<i>Motile</i>	IDA
	Sporulation	<i>Non sporulating</i>	NAS	
	Temperature range	<i>Able to grow at 25 °C–28 °C</i>	IDA	
	Optimum temperature	<i>Able to grow at 20 °C–30 °C</i>	NAS	
	pH range; Optimum	<i>Unknown</i>	NAS	
	Carbon source	<i>Ability to degrade polypectate</i>	IDA	
MIGS-6	Habitat	<i>Isolated from water,</i>	IDA	
MIGS-6.3	Salinity	<i>Able to grow in 1% NaCL</i>	IDA	
MIGS-22	Oxygen requirement	<i>Facultative anaerobic</i>	NAS [17]	
MIGS-15	Biotic relationship	<i>Free-living</i>	IDA	
MIGS-14	Pathogenicity	<i>Ability to rot potato tubers</i>	IDA	
MIGS-4	Geographic location	<i>A97-S13-F16, river Durance France</i>	IDA	
		<i>A359-S18-N18, river Bléone France</i>		
MIGS-5	Sample collection	<i>A97-S13-F16, 1st February 2016</i>	IDA	
		<i>A359-S18-N18, 19th November 2016</i>		
MIGS-4.1	Latitude	<i>A97-S13-F16, 44.701667</i>	IDA	
		<i>A359-S18-N18, 44.038631</i>		
MIGS-4.2	Longitude	<i>A97-S13-F16, 6.599693</i>	IDA	
		<i>A359-S18-N18, 5.967728</i>		
MIGS-4.4	Altitude	<i>A97-S13-F16, 907 m</i>	IDA	
		<i>A359-S18-N18, 438 m</i>		

<sup>a</sup> Evidence codes - IDA Inferred from Direct Assay, TAS Traceable Author Statement (i.e., a direct report exists in the literature), NAS Non-traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [20]

The “*P. peruvienne*” A350-S18-N16 draft genome contains 4,871,019 bp with a GC content of 51,1%. Total predicted genes are 4635 while predicted protein coding genes are 4487 and RNA genes 48. The final assembly comprised 73 scaffolds. Among the predicted genes, 72.01% have a predicted function, 78.77% were assigned to GOG and 85.09% have a predicted Pfam domain. Among the predicted proteins, 395 have a predicted signal peptide while 1095 contain a predicted transmembrane helix. Two CRIPS repeats array were detected in this genome.

The properties and the statistics of the two draft genomes are summarized in Tables 3 and 4.

### Insight from genome sequences

#### Genome comparison between A97-S13-F16 and A350-S18-N16 and the genome of representative species of the *Pectobacterium* genus

A phylogenetic tree, constructed from concatenated sequences of 1266 homologs proteins, clustered the A97-S13-F16 and A350-S18-N16 strains together, close to UGC32 the proposed “*P. peruvienne*” type strain

**Table 2** Genome sequencing project information

MIGS ID	Property	<i>P. Peruvienne</i>	
		A97-S13-F16	A350-S18-N16
	Finishing quality	61 scaffolds	73 scaffolds
MIGS-28	Libraries used	Nextera DNA Library	Nextera DNA Library
MIGS 29	Sequencing platforms	Illumina NS500	Illumina NS500
MIGS 31.2	Fold coverage	331X	86X
MIGS 30	Assemblers	CLC Genomics Workbench V 9.5.2	CLC Genomics Workbench V 9.5.2
MIGS 32	Gene calling method	Glimmer 3	Glimmer 3
	Locus Tag	A97-S13-F16	A350-S18-N16
	Genbank ID	PYUO01000000	PYUO01000000
	GenBank Date of Release	10th July 2018	10th July 2018
	GOLD ID		
	BIOPROJECT	PRJNA445781	PRJNA445781
MIGS 13	Source Material Identifier	CFBP8625 <sup>a</sup>	CFBP8626 <sup>a</sup>
	Project relevance	Environment	Environment

<sup>a</sup> Strains A97-S13-F16 and A350-S18-N16 are available at the CIRN-CFBP Collection under the indicated numbers

(Fig. 1B). ANIb were further calculated between genomes of strains A97-S13-F16 and A350-S18-N16 and the genomes of described *Pectobacterium* species and subspecies (Additional file 2: Table S1). Pairwise ANIb values between the three “*P. peruvienne*” genomes, A97-S13-F16 and A350-S18-N16 and UGC32, were above 97,5%. Pairwise ANIb values of these three “*P. peruvienne*” genomes with genomes of other *Pectobacterium* species and subspecies were below 94%. dDDH is an in silico method to approach the wet-lab DDH method as closely as possible [22]. dDDH were

calculated between the genomes of A97-S13-F16 and A350-S18-N16 and *Pectobacterium* genomes representative of known species and subspecies (Additional file 2: Table S1). dDDH values between A350-S18-N16, A97-S13-F16 genomes and the proposed “*P. peruvienne*” UGC32 genomes were above 79%, well above the 70% species boundary. When pairwise calculations were performed between these three genomes with those of known *Pectobacterium* species and subspecies the estimated dDDH values dropped below 54%, well below the species boundary. This confirmed that

**Table 3** Genome statistics

Attribute	<i>P. peruvienne</i> A97-S13-F16		<i>P. peruvienne</i> A350-S18-N16	
	Value	% of Total	Value	% of Total
Genome size (pb)	4,755,191	100.00	4,871,019	100.00
DNA coding (bp)	4,108,775	86.41	4,211,847	86.47
DNA G + C (pb)	2,425,147	51.00	2,489,091	51.10
DNA scaffolds	61		73	
Total genes	4503	100.00	4635	100.00
Protein coding genes	4459	99.02	4587	98.96
RNA genes	44	0.97	48	1.03
Pseudo genes	NA		NA	
Genes in internal clusters	NA		NA	
Genes with function prediction	3252	72.21	3338	72.01
Genes assigned to COGs	3563	79.91	3613	78.77
Genes with Pfam domains	3808	85.40	3903	85.09
Genes with signal peptides	392	8.79	395	8.52
Genes with transmembrane helices	1090	24.44	1095	23.62
CRISPR repeats	3		2	

**Table 4** Number of genes associated with the 25 COG functional categories

Code	<i>P. peruvienne</i>		<i>P. peruvienne</i>		Description
	A97-S13-F16		A350-S18-N16		
	Value	%age	Value	%age	
E	366	8.21	367	8.00	Amino acid transport and metabolism
G	362	8.12	356	7.76	Carbohydrate transport and metabolism
D	41	0.92	43	0.94	Cell cycle control, cell division, chromosome partitioning
N	110	2.47	107	2.33	Cell motility
M	239	5.36	244	5.32	Cell wall/membrane/envelope biogenesis
H	167	3.75	167	3.64	Coenzyme transport and metabolism
Z	1	0.02	1	0.02	Cytoskeleton
V	85	1.91	90	1.96	Defense mechanisms
C	226	5.07	225	4.91	Energy production and conversion
W	4	0.09	4	0.09	Extracellular structures
S	202	4.53	209	4.56	Function unknown
G	204	4.58	206	4.49	General function prediction only
P	242	5.43	240	5.23	Inorganic ion transport and metabolism
U	82	1.84	91	1.98	Intracellular trafficking, secretion, and vesicular transport
I	103	2.31	102	2.20	Lipid transport and metabolism
X	23	0.52	57	1.24	Mobilome: prophages, transposons
F	90	2.02	91	1.98	Nucleotide transport and metabolism
O	152	3.41	152	3.31	Posttranslational modification, protein turnover, chaperones
L	127	2.85	127	2.77	Replication, recombination and repair
A	1	0.02	1	0.02	RNA processing and modification
Q	59	1.32	59	1.29	Secondary metabolites biosynthesis, transport and catabolism
T	146	3.27	148	3.23	Signal transduction mechanisms
K	291	6.53	287	6.26	Transcription
J	238	5.34	239	5.21	Translation, ribosomal structure and biogenesis
–	898	20.14	974	21.23	Not in COGs


The total %age is based on the total number of protein coding genes in the genome

A97-S13-F16 and A350-S18-N16 belong to the “*P. peruvienne*” specie.

#### Genomes comparison between the “*P. peruvienne*” strains

The phylogenetic trees (Fig. 2) indicate that strains A97-S13-F16 and A350-S18-N16 are more closely related to each other than they are from the “*P. peruvienne*” type strain UGC32. To further gain insight into the distance between the three “*P. peruvienne*” strains, we looked for shared and unique genes between genomes of strains A97-S13-F16, A350-S18-N16 and UGC32 type strain (Fig. 3). A97-S13-F16, A350-S18-N16 and UGC32 strains contain respectively a pool of specific

genes of 292, 414 and 346. The slightly higher pool of specific genes observed in strain A350-S18-N16 could be partly related to its higher content of mobile genetic elements inserted in its genome as described in Table 4. Indeed, we observed 3 clusters of phage-related genes in strain A350-S18-N16, only one being also detected in strain A97-S13-F16. The Venn diagram indicated that 4129 genes are shared between strains A97-S13-F16 and A350-S18-N16 while only 3757 and 3765 genes are respectively shared between the type strain UGC32 and A97-S13-F16 / A350-S18-N16. This confirmed that A97-S13-F16 and A350-S18-N16 genomes are more closely related to each other than they are with the genome of the proposed type strain UGC32.



**Fig. 3** Venn diagram. Shared and unique genes between the genomes of “*P. peruvienne*” A97-S13-F16 and A350-S18-N16 and the proposed “*P. peruvienne*” type strain UGC32. Orthology was assumed using a threshold of 80% identity on at least 80% of the protein length

## Conclusions

In this study we presented the draft genome sequences of two strains of “*P. peruvienne*” isolated from fresh water in river stream in France. The “*P. peruvienne*” specie has recently been proposed and, until our study, the described strains belonging to the “*P. peruvienne*” specie have all been isolated on potato tubers in the altiplano in Peru [8]. The presence of strains belonging to the “*P. peruvienne*” specie in two independent environmental samples in France indicates that the geographic distribution of this specie is likely to be larger than previously anticipated. Both French strains are able to rot potato tubers like the proposed type strain UG32. The two French isolates are more closely related to each other than they are with the type strain UGC32. Whether this reflects the geographic provenance (France vs Peru) or the niche provenance (water vs diseased plants) is unknown.

## Additional files

**Additional file 1: FigureS1.** Symptoms observed on potato tubers. Overnight cultures of bacterial strains A97-S13-F16 and A350-S18-N16 were suspended in 50 mM phosphate buffer pH 6.8 and adjusted to 1.0 at OD<sub>580nm</sub>. Tubers of *S. tuberosum* var. charlotte were inoculated with 10 µl of the cell suspension and placed at room temperature on wet paper towel in a plastic box. Six days post-infection, tubers were cut in half and representative symptoms are shown: A: A97-S13-F16, B: A350-S18-N16, C: 50 mM phosphate buffer pH 6.8. (DOCX 9779 kb)

**Additional file 2: Table S1.** ANIb and dDDH pairwise values. dDDH and ANIb are respectively presented in the upper and lower part of the matrix triangle. Strains belonging to the same species are highlighted in red. Specific threshold value is 96% for ANIb and 70% for DDH. ANIb values were computed using the Blast algorithm of the Jspecies package [23]. dDDH were calculated according to [22]. (DOCX 79 kb)

## Abbreviations

ANI: Average Nucleotide Identity; COG: Clusters of Orthologous Groups; CRISPR: Clustered Regularly Interspaced Short Palindromic Repeats; dDDH: digital DNA-DNA hybridization; gapA: glyceraldehyde-3-phosphate dehydrogenase A; MLSA: Multi Locus Sequence Analysis; NTU: Normalized Turbidity Unit; sp.: specie; subsp.: subspecie

## Acknowledgements

This work has benefited from the expertise of the High-Throughput Sequencing Platform of I2BC, Gif sur Yvette, France. We thank our British colleague Emma Rochelle Newall for english editing of the manuscript. We thank Odile Berge and Frédérique Van Gijsegem for their help during river samplings and Ariane Toussaint and Antoine Pourbaix for hosting us during the field samplings.

## Funding

This work is supported by Agence Nationale de la Recherche (COMBICONTROL, grant ANR-15-CE21-0003) and CNRS program (EC2CO- Biohefect/Ecodyn//Dril/MicrobiEenCARTOBACTER).

## Availability of data and materials

This Whole Genome Shotgun project has been deposited at DDBJ/ENA/GenBank under the accessions PYUO00000000 and PYUP00000000. The versions described in this paper are versions PYUO01000000 and PYUP01000000. The strains are available at the CIRM CFBP.



**Authors' contributions**

BMA initiated the study and provided background information. BMA and CB isolated the strains. CB performed the *gapA* amplification and phylogenetic analysis, isolated the DNA for sequencing and performed the microscopy analysis. JP and FP assembled, analyzed the genomes and conducted the MLSA analysis. BMA and JP wrote the manuscript. All authors read and approved the final manuscript.

**Ethics approval and consent to participate**

not applicable.

**Consent for publication**

not applicable.

**Competing interests**

The authors declare that they have no competing interests.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 20 May 2018 Accepted: 28 September 2018

Published online: 12 October 2018

**References**

- Skerman VB, McGowan V, Sneath PH. Approved lists of bacterial names. *Int J Syst Bacteriol.* 1980;30:225–420.
- Ma B, Hibbing ME, Kim H-S, Reedy RM, Yedidia I, Breuer J, et al. Host range and molecular phylogenies of the soft rot Enterobacterial genera *Pectobacterium* and *Dickeya*. *Phytopathology.* 2007;97:1150–63.
- Pérombelon M, Kelman A. Ecology of the soft rot *Erwinias*. *Annu Rev Phytopathol.* 1980;18:361–87.
- Gardan L, Gouy C, Christen R, Samson R. Elevation of three subspecies of *Pectobacterium carotovorum* to species level: *Pectobacterium atrosepticum* sp. nov., *Pectobacterium betavasculorum* sp. nov. and *Pectobacterium wasabiae* sp. nov. *Int J Syst Evol Microbiol.* 2003;53:381–91.
- Nabhan S, De Boer SH, Maiss E, Wydra K. *Pectobacterium aroidearum* sp. nov., a soft rot pathogen with preference for monocotyledonous plants. *Int J Syst Evol Microbiol.* 2013;63:2520–5.
- Dees MW, Lysøe E, Rossmann S, Perminow J, Brurberg MB. *Pectobacterium polaris* sp. nov., isolated from potato (*Solanum tuberosum*). *Int J Syst Evol Microbiol.* 2017;67:5222–9.
- Khayri S, Cigna J, Chong T, Quêtu-Laurent A, Chan K, Helias V, et al. Transfer of the potato plant isolates of *Pectobacterium wasabiae* to *Pectobacterium parmentieri* sp. nov. *Int J Syst Evol Microbiol.* 2016;66:5379–83.
- Waleron M, Misztak A, Waleron M, Franczuk M, Wielgomas B, Waleron K. Transfer of *Pectobacterium carotovorum* subsp. *carotovorum* strains isolated from potatoes grown at high altitudes to *Pectobacterium peruviane* sp. nov. *Syst Appl Microbiol.* 2018;41:85–93.
- Hauben L, Moore ERB, Vauterin L, Steenackers M, Mergaert J, Verdonck L, et al. Phylogenetic position of Phytopathogens within the *Enterobacteriaceae*. *Syst Appl Microbiol.* 1998;21:384–97.
- List Editor: Validation List no. 68. Validation of publication of new names and new combinations previously effectively published outside the IJSB. *Int J Syst Bacteriol.* 1999;49:1–3.
- Koh Y, Kim G, Lee Y, Sohn S, Koh H, Kwon S, et al. *Pectobacterium carotovorum* subsp. *actinidiae* subsp. nov., a new bacterial pathogen causing canker-like symptoms in yellow kiwifruit, *Actinidia chinensis*. *N Z J Crop Horticult Sci.* 2012;40:269–79.
- Nabhan S, De Boer SH, Maiss E, Wydra K. Taxonomic relatedness between *Pectobacterium carotovorum* subsp. *carotovorum*, *Pectobacterium carotovorum* subsp. *odoriferum* and *Pectobacterium carotovorum* subsp. *brasilense* subsp. nov. *J Appl Microbiol.* 2012;113:904–13.
- Panda P, Fiers MWEJ, Lu A, Armstrong KF, Pitman AR. Draft genome sequences of three *Pectobacterium* strains causing blackleg of potato: *P. carotovorum* subsp. *brasilensis* ICMP 19477, *P. atrosepticum* ICMP 1526, and *P. carotovorum* subsp. *carotovorum* UGC32. *Genome Announc.* 2015;3:e00874–15.
- Hélias V, Hamon P, Huchet E, Wolf JVD, Andrivon D. Two new effective semiselective crystal violet pectate media for isolation of *Pectobacterium* and *Dickeya*: isolating pectolytic bacteria on CVP. *Plant Pathol.* 2012;61:339–45.
- Cigna J, Dewaegeneire P, Beury A, Gobert V, Faure D. A *gapA* PCR-sequencing assay for identifying the *Dickeya* and *Pectobacterium* potato pathogens. *Plant Dis.* 2017;101:1278–82.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST server: rapid annotations using subsystems technology. *BMC Genomics.* 2008;9:75.
- Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res.* 1999;27:4636–41.
- Marchler-Bauer A, Bryant SH. CD-search: protein domain annotations on the fly. *Nucleic Acids Res.* 2004;32:W327–31.
- Grissa I, Vergnaud G, Pourcel C. CRISPRFinder: a web tool to identify clustered regularly interspaced short palindromic repeats. *Nucleic Acid Res.* 35: W52–W57.
- Petersen TN, Brunak S, von HG, Nielsen H. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods.* 2011;8:785–6.
- Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden markov model: application to complete genomes. *J Mol Biol.* 2001;305:567–80.
- Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics.* 2013;14:60.
- Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinforma Oxf Engl.* 2016;32:929–31.
- Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics.* 2004;5:113.
- Castresana J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol.* 2000;17:540–52.
- Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol.* 2003;52:696–704.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

