



HAL
open science

RanBP2 regulates the anti-retroviral activity of TRIM5 α by SUMOylation at a predicted phosphorylated SUMOylation motif

Ghizlane Maarifi, Juliette Fernandez, Débora Portilho, Aude Boulay, Jacques Dutrieux, Stéphane Oddos, Gillian Butler-Browne, Sébastien Nisole, Nathalie Arhel

► **To cite this version:**

Ghizlane Maarifi, Juliette Fernandez, Débora Portilho, Aude Boulay, Jacques Dutrieux, et al.. RanBP2 regulates the anti-retroviral activity of TRIM5 α by SUMOylation at a predicted phosphorylated SUMOylation motif. *Communications Biology*, 2018, 1 (1), pp.193. 10.1038/s42003-018-0198-0 . hal-01930335

HAL Id: hal-01930335

<https://hal.sorbonne-universite.fr/hal-01930335v1>

Submitted on 21 Nov 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ARTICLE

DOI: 10.1038/s42003-018-0198-0

OPEN

RanBP2 regulates the anti-retroviral activity of TRIM5 α by SUMOylation at a predicted phosphorylated SUMOylation motif

Ghizlane Maarifi¹, Juliette Fernandez¹, Débora M. Portilho^{2,3}, Aude Boulay², Jacques Dutrieux², Stéphane Oddos³, Gillian Butler-Browne⁴, Sébastien Nisole¹ & Nathalie J. Arhel^{1,2}

TRIM5 α is a cytoplasmic restriction factor that blocks post-entry retroviral infection. Evidence suggests that its antiviral activity can be regulated by SUMO, but how this is achieved remains unknown. Here, we show that TRIM5 α forms a complex with RanGAP1, Ubc9, and RanBP2 at the nuclear pore, and that RanBP2 E3 SUMO ligase promotes the SUMOylation of endogenous TRIM5 α in the cytoplasm. Loss of RanBP2 blocked SUMOylation of TRIM5 α , altered its localization in primary cells, and suppressed the antiviral activity of both rhesus and human orthologs. In cells, human TRIM5 α is modified on K84 within a predicted phosphorylated SUMOylation motif (pSUM) and not on K10 as found *in vitro*. Non-modified TRIM5 α lacked antiviral activity, indicating that only SUMOylated TRIM5 α acts as a restriction factor. This work illustrates the importance of the nuclear pore in intrinsic antiviral immunity, acting as a hub where virus, SUMO machinery, and restriction factors can meet.

¹Institut de Recherche en Infectiologie de Montpellier (IRIM), Université de Montpellier, CNRS, 34090 Montpellier, France. ²Saint-Louis Hospital, INSERM, Paris 7 University, 75010 Paris, France. ³Bioaxial SAS, 75014 Paris, France. ⁴Center for Research in Myology, INSERM, CNRS, Paris 6 University, 75013 Paris, France. Correspondence and requests for materials should be addressed to N.J.A. (email: nathalie.arhel@irim.cnrs.fr)

Post-translational modifications by small ubiquitin-related modifiers (SUMO) can modulate protein stability, localization, and activity, to regulate numerous eukaryotic cellular functions such as gene expression, DNA repair, and chromatin organization^{1,2}. In vertebrates, three SUMO paralogues can be covalently added onto substrate proteins: SUMO1, -2, and -3. SUMO2 and SUMO3 are almost identical in amino acid sequence and are therefore referred to as SUMO2/3. A SUMO4 variant exists but does not appear to be conjugated to cellular proteins³, and a SUMO5 paralogue has recently been described⁴. SUMO conjugation, known as SUMOylation, is brought about by the sequential action of three enzymes: an E1 SUMO-activating enzyme (SAE1/SAE2 heterodimer, Aos1/Uba2 in yeast), an E2 SUMO conjugating enzyme (Ubc9), and an E3 SUMO ligase such as protein inhibitor of activated STAT proteins (PIAS proteins)⁵, polycomb group protein Pc2⁶, and RanBP2⁷. The host SUMOylation system can play an important role in response to viral infection, both by activating cellular defenses that control viral replication and, conversely, serving as a target for viral evasion strategies that boost infection. Examples from the literature indicate that viral proteins can undergo SUMOylation or influence the SUMOylation of cellular proteins, and that the infection of some viruses is dependent on pathways that are regulated by SUMOylation⁸.

Restriction factors are antiviral proteins produced by the host to antagonise viral infection. They are considered the first line of defence against viruses, and can intervene literally within minutes of cell entry, to either degrade the virus or block its replication. A fine example is provided by the alpha isoform of the tripartite motif-containing protein 5 (TRIM5 α), a cytoplasmic protein that prevents retroviruses from jumping the species barrier by intercepting viral capsids before they reach the nucleus. Although TRIM5 α is constitutively expressed, experiments that modify the global cellular SUMO machinery profoundly affect TRIM5 α -mediated restriction^{9–11}, suggesting that the SUMO system provides a way for a host to modulate the antiviral activity of restriction factors. TRIM5 α itself can be directly modified both by SUMO1 and SUMO2¹¹, however no effect on its antiviral activity could so far be demonstrated.

Enzymes of the SUMO machinery are enriched in the nucleus and most SUMO substrates are nuclear proteins involved in nuclear processes such as transcription and DNA repair. Although TRIM5 α forms cytoplasmic bodies, it can also shuttle to the nucleus^{12,13}, suggesting that it could interact with SUMO machinery there. Some have shown co-localization of over-expressed TRIM5 α with Promyelocytic Leukaemia Nuclear Bodies (PML-NB)^{13,14}, which are a preferred site of SUMOylation and harbor many proteins of the SUMO machinery, including SUMO and PIAS1^{15,16}, however we could not reproduce these findings with endogenous TRIM5 α ¹³. Therefore, although PIAS1 can SUMO modify TRIM5 α in vitro¹¹, their distinct endogenous localizations in cells suggests that PIAS1 is unlikely to act as the main E3 SUMO ligase of TRIM5 α in vivo. In addition, treatment with ginkgolic acid, an inhibitor of the E1 SUMO-activating enzyme, reduces nuclear residency of TRIM5 α ¹³, suggesting that TRIM5 α is SUMO modified in the cytoplasm and shuttles to the nucleus as a SUMOylated protein. Cytoplasmic SUMOylation has already been described for some proteins with restricted cytoplasmic localization, such as RanGAP1, indicating that enzymes of the SUMO machinery are also present in the cytoplasm¹⁷. We therefore set out to identify elements of the SUMO machinery that account for SUMO conjugation of TRIM5 α in the cytoplasm.

Here, we show that TRIM5 α associates with RanGAP1, RanBP2, and Ubc9 at the nuclear pore, and is SUMO conjugated by the nucleoporin and E3 SUMO ligase RanBP2 within this complex. In SUMO overexpressing cells, RanBP2 promoted both

SUMO1 and SUMO2/3 conjugation on at least two sites. A 33 kDa C-terminal domain of RanBP2 previously shown to contain E3 ligase activity was sufficient to modify TRIM5 α in an in vitro SUMO assay. RanBP2 dramatically impacted the localization of TRIM5 α in two primary cell models and was required for efficient restriction of HIV-1 by rhesus TRIM5 α in HeLa cells. This is the first report of RanBP2 participating in cellular intrinsic immunity mechanisms by regulating the antiviral activity of a restriction factor via its E3 SUMO ligase domain.

Results

TRIM5 α binds to RanGAP1 and RanBP2 at the nuclear pore.

Previous work showed that TRIM5 α can undergo SUMO modification in vitro in the presence of recombinant SUMO, E1, E2, and E3 ligase PIAS1¹¹. A TRIM5 α mutant at the SUMOylation consensus site (K10R) is a poor substrate in this in vitro reaction, but maintains wild-type antiviral activity in cells, casting doubt on the relevance of PIAS1-mediated SUMOylation on TRIM5 α function. To address this apparent discrepancy, we assessed the SUMOylation status of both wild-type and K10R mutant TRIM5 α in cells. Flag-tagged TRIM5 α was overexpressed and purified by immunoprecipitation from HEK 293T cell extracts. Blotting with an α -SUMO1 antibody revealed a ~70 kDa band, i.e., ~15 kDa above unmodified TRIM5 α , in both wild-type and K10R transfected cells, indicating that the K10R mutant is SUMOylated in cells (Fig. 1a). These results, together with the fact that PIAS1 and TRIM5 α are present in distinct subcellular compartments, indicates that PIAS1 cannot SUMOylate TRIM5 α in vivo.

We reasoned that the E3 SUMO ligase responsible for TRIM5 α SUMOylation in vivo is likely to be cytoplasmic since our previous work suggested that TRIM5 α undergoes SUMOylation in the cytoplasm¹³. To identify elements of the SUMO machinery involved in the regulation of TRIM5 α function, we searched for cell types in which endogenous TRIM5 α did not form typical cytoplasmic aggregates, and investigated whether this altered localization correlated with a proximity to elements of the SUMO pathway. We identified an unusual pattern for TRIM5 α in human myotubes, where the majority of the signal accumulated around the nuclear envelope (Fig. 1b). Since the nuclear pore is a privileged site for SUMOylating and deSUMOylating enzymes¹⁸, we asked whether TRIM5 α co-localized with elements of the nuclear pore in these cells. Super-resolution imaging revealed a co-localization of TRIM5 α with the Ran-specific GTPase activating protein RanGAP1, which is present on cytoplasmic filaments of the nuclear pore, but not with Nup153, which is found in the nuclear basket (Fig. 1c), indicating that TRIM5 α is present at the cytoplasmic face of the nuclear pore in these cells.

An accumulation of TRIM5 α around the nuclear membrane was not observed in other tested cell types, suggesting that this phenomenon might be transient in most systems. To determine whether TRIM5 α can co-localize with RanGAP1 in other cells, we performed proximity ligation assay (PLA) in HeLa and HEK 293T cells. TRIM5 α ×RanGAP1 PLA signals were detected at a frequency of 4 and 1.5 spots/cell, respectively, and localized to the nuclear membrane in 55% of cases in both cell types, indicating that TRIM5 α can co-localize with RanGAP1 at the nuclear pore in these cells (Fig. 1d).

To assess whether TRIM5 α can interact with RanGAP1 in HeLa cells, we purified endogenous RanGAP1 by direct immunoprecipitation and blotted for possible interacting partners. RanGAP1 immunoprecipitated RanBP2, as expected since these have been shown to form a stable complex in cells¹⁹ (Fig. 1e). Endogenous TRIM5 α was also found in RanGAP1 immunoprecipitates, both as the unmodified 55 kDa species, and a ~70 kDa protein that also appeared in SUMO1 blots identifying




Fig. 1 TRIM5 α and SUMO-modified TRIM5 α form a complex with RanGAP1 and RanBP2 at the nuclear pore. **a** The consensus site for in vitro SUMOylation by PIAS1 is not a target for SUMOylation in human cells. HEK 293T cells were transfected with Flag-tagged human TRIM5 α or K10R mutant, previously shown to abrogate PIAS1-mediated SUMOylation in vitro. After 3 days, TRIM5 α was immunoprecipitated using α -Flag antibodies and probed with α -SUMO1 or α -Flag antibodies. SUMO-modified TRIM5 α is detected at ca. 70 kDa in the SUMO1 blots for both WT and K10R lanes (TRIM5 α *S1). Western blots are representative of 3 independent experiments (original blots in Supplementary Figure 2a). **b** TRIM5 α localizes at the nuclear envelope in myotubes. To identify the E3 SUMO ligase responsible for TRIM5 α SUMOylation in cells, we searched for cell types in which endogenous TRIM5 α did not form typical cytoplasmic aggregates, and probed for proximity with SUMO machinery. In human myotubes, the majority of TRIM5 α signal accumulated around the nuclear envelope. Scale bar = 20 μ m. **c** TRIM5 α co-localizes with RanGAP1 in myotubes. Super-resolution imaging of human myotubes shows TRIM5 α co-localization with Ran-specific GTPase activating protein RanGAP1, present on cytoplasmic filaments of the nuclear pore, but not with Nup153, which is found in the nuclear basket. Donor 1: neonatal myoblasts (CHQ), donor 2: adult myoblasts (160 M). Scale bar = 2 μ m. **d** TRIM5 α co-localizes with RanGAP1 in HeLa and HEK 293T cells. Proximity ligation assay was performed in HeLa and 293T cells, only HeLa are shown as an example. TRIM5 α × RanGAP1 PLA signals were detected at a frequency of 4 and 1.5 spots/cell, respectively, and localized to the nuclear membrane in 55% of cases in both cells types, confirming that TRIM5 α co-localizes with RanGAP1 at the nuclear envelope in these cells. Scale = 10 μ m. **e** TRIM5 α and SUMO-modified TRIM5 α form a complex with RanGAP1 and RanBP2 in HeLa cells. RanGAP1 was immunoprecipitated from HeLa cytoplasmic extracts and precipitates were probed for TRIM5 α and SUMO1. RanGAP1 immunoprecipitates contained RanBP2, unmodified TRIM5 α (55 kDa), and a ~70 kDa TRIM5 α species that also appeared in SUMO1 blots identifying it as SUMO1-modified TRIM5 α (TRIM5 α *S1). SUMOylated RanGAP1 (~90 kDa) is present at very low levels in cytoplasmic extracts and is therefore not detected here. Western blots are representative of 2 independent experiments (original blots in Supplementary Figure 2b)

it as SUMO1-modified TRIM5 α (TRIM5 α *S1) (Fig. 1e). Results indicated that both unmodified and SUMO-modified TRIM5 α are present in a complex with two major components of the nucleocytoplasmic transport machinery and key elements of the SUMO system, RanGAP1 and RanBP2.

RanBP2 E3 SUMO ligase promotes SUMO1 conjugation of TRIM5 α in cells. To address whether RanBP2 can promote TRIM5 α SUMO modification within the RanGAP1/RanBP2 complex, we isolated endogenous TRIM5 α by RanGAP1 immunoprecipitation from cells expressing or not RanBP2. Transduction of HeLa cells with a specific RanBP2 shRNA²⁰ resulted in efficient knockdown of RanBP2 (Fig. 2a, input). RanGAP1 immunoprecipitates from non-transduced and control vector transduced cells contained RanBP2, Ubc9, TRIM5 α , and TRIM5 α *S1, indicating that these are present together in a complex in cells. As previously, TRIM5 α *S1 was identified as a ~70 kDa

band that was visible in both TRIM5 α and SUMO1 blots. Both Ubc9 and RanBP2 were absent from RanGAP1 immunoprecipitates in RanBP2 depleted cells, indicating that Ubc9's association with the RanGAP1/RanBP2 complex is dependent upon its interaction with RanBP2. This was not found to be the case for TRIM5 α , which was still detected in RanGAP1 immunoprecipitates from RanBP2 knockdown cells, suggesting that its association with RanGAP1 can be maintained in the absence of RanBP2. Critically, TRIM5 α *S1 levels were reduced and unmodified TRIM5 α levels were increased in the absence of RanBP2, indicating that RanBP2 promotes SUMO1 modification of TRIM5 α in cells (Fig. 2a).

To confirm that TRIM5 α is a target for SUMO1 modification by RanBP2 E3 SUMO ligase, we performed the reverse experiment and immunoprecipitated endogenous TRIM5 α directly from HeLa cells treated with shRNAs against RanBP2 or TRIM5 α (Fig. 2b). RanBP2 was present in TRIM5 α immunoprecipitates, and disappeared as expected in the absence




Fig. 3 RanBP2 promotes multiple conjugation of both SUMO1 and SUMO2/3 on TRIM5 α . **a** Schematic representation of the experimental procedure. **b** HeLa-SUMO1, and **c** HeLa-SUMO2/3 cells were transduced with shBP2, shE, or not transduced (NT). After 3 days, RanBP2 depletion was assessed by qPCR. TRIM5 α was immunoprecipitated from **d** HeLa-SUMO1 and **e** HeLa-SUMO2/3 cells. Input samples were analyzed by immunoblotting with a His antibody to reveal SUMO-modified proteins, while TRIM5 α immunoprecipitates were probed with α -SUMO1 or α -SUMO2/3 antibodies. Higher molecular weight bands indicate SUMO-modified TRIM5 α (TRIM5 α *S1 and TRIM5 α *S2/3). IgG heavy (HC) and light chains (LC) are indicated. Western blots are representative of 3 independent experiments (original blots in Supplementary Figure 4). **f** TRIM5 α *S1 bands were quantified from 10 independent SUMO blots of TRIM5 α immunoprecipitates. The graph shows the mean relative intensity of SUMO bands normalized for NT controls \pm SEM. Statistical analysis was performed using a paired t test

perform reporter assays since RanBP2 knockdown blocks HIV-1 nuclear entry²⁰. Therefore the effect of RanBP2 on rhTRIM5 anti-HIV activity may only be reliably measured before the nuclear import step. In our assay, expression of rhTRIM5 α led to a decrease in Pol copy numbers in control samples, indicating TRIM5 α -mediated restriction (Fig. 4a). Transduction with a shRNA targeting RanBP2 did not affect reverse transcription in control cells, as shown previously²⁰. Strikingly, however, depletion of RanBP2 cancelled the ability of TRIM5 α to block HIV-1 reverse transcription. Similar results were obtained if rhTRIM5 α was transfected prior to RanBP2 knockdown, and at all tested multiplicities of infection (MOI) (Fig. 4b).

To obtain a greater restriction phenotype, we generated HeLa clones overexpressing rhTRIM5 α . In these cells, HIV-1 reverse transcription reached background levels similarly to those obtained with Nevirapine. Remarkably, the simple lipofection of an siRNA against RanBP2 in these stable clones, compared to control siRNA, blocked the ability of rhTRIM5 α to restrict HIV-1 indicating that RanBP2 is key in mediating efficient rhTRIM5 α restriction (Fig. 4c).

RanBP2 modulates the antiviral activity of endogenous TRIM5 α . We previously showed that SUMOylation of TRIM5 α can regulate its subcellular localization in specific cell types¹³. To determine whether SUMOylation of TRIM5 α by RanBP2 can impact its subcellular localization, we returned to our muscle cell model in which TRIM5 α localizes at the nuclear envelope (Fig. 1b). Previous work showed that skeletal muscle differentiation is associated with changes in nuclear pore architecture and composition²², including a dramatic up-regulation of RanBP2 protein expression and residence in nuclear pore complexes (NPC)²³. We obtained primary myoblasts in which RanBP2 was barely detectable, and differentiated them into

myotubes to induce strong expression of RanBP2 at the nuclear envelope. Interestingly, the absence of RanBP2 in the myogenic precursor cells was associated with the accumulation of TRIM5 α in the cytoplasm away from the nuclear envelope (Fig. 5a). Although a direct correlation cannot be proven given that the differentiation of myoblasts to myotubes changes the expression of many genes²⁴, these results suggest that induction of RanBP2 expression alters TRIM5 α localization.

As a parallel example, we tested primary monocyte-derived DCs, in which we described a SUMO-dependent nuclear sequestration of TRIM5 α ¹³. Primary DCs were transduced with either empty or shRanBP2 lentiviral vectors and monitored for TRIM5 α localization by confocal microscopy (Fig. 5b). In control cells, TRIM5 α was almost exclusively nuclear, as previously reported¹³. In contrast, loss of RanBP2 led to a dramatic relocation of some TRIM5 α signal to the cytoplasm as bright aggregates, suggesting that SUMOylation of TRIM5 α by RanBP2 contributes to its nuclear targeting in DCs.

Having shown that RanBP2 mediates the antiviral activity of overexpressed rhTRIM5 α , we examined whether it also affects the ability of N-MLV infection to be restricted by endogenous huTRIM5 α . We lipofected human cells with an siRNA against RanBP2 or control, and infected with increasing doses of N-MLV-YFP. As expected, human TRIM5 α restricts N-MLV at low doses and is saturated with increasing MOI. However, the depletion of RanBP2 led to a measurable increase in N-MLV infection efficiency at all tested viral inoculums, with a 5-fold increase in fluorescence intensity at MOI 1, indicating that RanBP2 regulates the antiviral activity of endogenous human TRIM5 α (Fig. 5c).

TRIM5 α is SUMOylated on a pSUM consensus motif at position K84. Having established that the human TRIM5 α K10R




Fig. 4 RanBP2 regulates the anti-HIV-1 activity of rhesus TRIM5α. **a** HeLa cells were transduced with shBP2 or shE at d0, transduced with rhesus TRIM5α (T5) or empty vector (EV) at d1, and infected with VSV-G pseudotyped HIV-1 at MOI 1 at d5 in the absence or presence of the reverse transcription inhibitor Nevirapine (Nev). RanBP2 and TRIM5α transcripts were quantified by qRT-PCR and normalized for β2-microglobulin. To determine the efficiency of reverse transcription, Pol DNA transcripts were quantified at 6 hpi by qPCR and normalized for actin. Results show the mean of 3 experiments performed with independent lentiviral vector stocks ± SD. Unpaired t test was performed on HIV-1 data sets using Prism 6. Expression of rhTRIM5α led to a decrease in Pol copy numbers in control samples following HIV-1 infection, indicating efficient TRIM5α-mediated restriction. Depletion of RanBP2 cancelled the ability of TRIM5α to block HIV-1 reverse transcription, indicating that RanBP2 is key in mediating efficient TRIM5α restriction. **b** The same experiment was repeated but by first overexpressing rhTRIM5α then depleting RanBP2. TRIM5α expression was monitored by flow cytometry using HA labelling. Graphs are representative of two independent experiments. **c** To obtain a greater restriction phenotype, the experiment was repeated using rhTRIM5α cell clones. Graphs are representative of two independent experiments

mutant is SUMOylated to wild-type levels in cells, we sought to identify the site of SUMO modification on TRIM5α. We generated 4 lysine mutants in strong direct and inverted SUMO consensus motifs as predicted by the JASSA bioinformatic tool²⁵ (Fig. 6a) and tested them for their ability to be SUMOylated. In two different cell types, the K10R, K111R, and K167R mutants were all SUMOylated as wild-type TRIM5α. In contrast, the K84R mutant showed significantly reduced SUMOylation, indicating that this residue is the preferred site for SUMO modification in cells (Fig. 6b, c). This residue is located in the linker 1 region that connects the RING and B-box2 domains (Fig. 6a). Remarkably, K84 resides in a pSUM phosphorylated SUMOylation motif (yKXpSP)^{25,26} in addition to an antisense SUMOylation consensus motif²⁷, similarly to the corresponding K85 residue in rhesus TRIM5α. Experimental evidence²⁸ and prediction algorithms (PhosphoSite Plus) both confirm that S86 on human TRIM5α and S87 on rhesus TRIM5α are phosphorylated,

suggesting that TRIM5α is SUMOylated on a consensus pSUM motif next to a phosphorylated serine.

We next tested the K84R mutant for its antiviral activity. Non-restrictive MDTF cells were transfected with the different human TRIM5α constructs. Transfection efficiency was found to be around 30% for all samples (Fig. 6d). The K10R, K111R, and K167R mutants all restricted N-MLV efficiently. In contrast however, the K84R mutant cancelled the antiviral activity of TRIM5α, indicating that SUMOylation at K84 is essential for TRIM5α function (Fig. 6d).

Discussion

Restriction factors are considered the first line of defense against viruses, however little is known about how their antiviral function is regulated in cells. Evidence suggests that the antiviral activity of TRIM5α can be regulated by the SUMO machinery^{9–11,13} but




Fig. 5 RanBP2 regulates the localization and antiviral restriction activity of human endogenous TRIM5 α . **a** RanBP2 regulates the subcellular localization of TRIM5 α in human myotubes. The localization of RanBP2 (top), and TRIM5 α (bottom) were assessed in myoblasts and myotubes using confocal imaging. Scale bar = 20 μ m. **b** RanBP2 regulates the subcellular localization of TRIM5 α in human dendritic cells. Human monocyte-derived DCs were transduced with a RanBP2 shRNA (shBP2) or empty vector (shE). Knock-down was assessed at d7 by qRT-PCR and DCs were labelled with an α -TRIM5 α antibody. Three representative fields are shown for each condition. Scale bar = 5 μ m. **c** RanBP2 regulates the antiviral activity of endogenous TRIM5 α . HeLa cells, which naturally restrict N-MLV, were lipofected with an siRNA against RanBP2 or control (siBP2 and siCtrl, respectively) and after 2 days were infected with increasing doses of N-MLV-YFP (MOI 1, 5, 10). Images were acquired at 2 dpt by widefield microscopy. The graph shows the mean fluorescence intensity (MFI) in representative fields at MOI 1. Scale bar = 100 μ m

how this is achieved was until now unknown. To our knowledge, we show for the first time that a nuclear pore component, Nup358 also called RanBP2, SUMOylates the retroviral restriction factor TRIM5 α and promotes its antiviral activity.

RanBP2 was previously shown to promote SUMO conjugation of Sp100⁷, HDAC4²⁹, and Mdm2³⁰. This study identifies TRIM5 α as a new substrate for RanBP2 E3 SUMO ligase. RanBP2 promoted single conjugation of SUMO1 moieties onto TRIM5 α in HeLa cells, as well as multiple SUMO1 and SUMO2/3 conjugation in SUMO-overexpressing HeLa cells. Although the lysine at position 10 of TRIM5 α is in a strong SUMOylation consensus motif, it is unlikely that RanBP2 mediates SUMOylation at this

position, since K10R mutants are still SUMOylated in HeLa cells, and are unaffected in their restriction activity¹¹. Indeed, we show that the lysine at position 84, which is in a predicted pSUM consensus motif and an antisense SUMO conjugation pattern, is the preferred site for SUMO1 modification in cells. Interestingly, the adjacent serine within the pSUM domain, S86 in human TRIM5 α (PhosphoSite Plus algorithm) and S87 in rhesus TRIM5 α ²⁸, is a major site for phosphorylation, suggesting that SUMO modification of TRIM5 α at K84 is phosphorylation-dependent.

The primary effect of SUMO modification of TRIM5 α by RanBP2 is likely to be to turn on TRIM5 α 's activity as an




Fig. 6 TRIM5 α is SUMOylated on residue K84 within a predicted pSUM consensus motif. **a** Schematic representation of the human TRIM5 α protein domains (to scale). The 4 lysines within putative consensus SUMO modification sequences are indicated in red. The corresponding sequences in rhesus TRIM5 α are indicated on the right for comparison purposes. **b** HEK 293T (original blots in Supplementary Figure 5a) and **c** MDTF cells (original blots in Supplementary Figure 5b) were transfected with the indicated pcDNA human TRIM5 α expression constructs. Inputs were monitored by Western blotting using anti-Flag and -GAPDH antibodies. TRIM5 α was immunoprecipitated using anti-Flag beads as in Fig. 1a. TRIM5 α SUMOylation was assessed by Western blotting on immunoprecipitated TRIM5 α using anti-Flag and -SUMO1 antibodies. The TRIM5 α *S1 band was readily detectable for all TRIM5 α mutants but was noticeably reduced for the K84R mutant, indicating that SUMO is conjugated at this residue. Western blots are representative of 2 independent experiments. **d** SUMOylation on K84 is required for the antiviral function of TRIM5 α . MDTF cells were transfected with the wild-type and mutant human TRIM5 α constructs then infected with N-MLV at MOI 2. Top panels show representative fields of wide-field microscopy. Equal transfection efficiency was monitored by flow cytometry (plots below). Mean fluorescence intensity (MFI) was quantified by Fiji software. The graph shows the values from 3 independent fields \pm SD and is representative of 2 independent experiments. Human TRIM5 α and the K10R, K111R, and K167R mutants all restricted N-MLV, whereas the K84R mutant lost the ability to do so. Scale bar = 50 μ m

anti-retroviral factor. Depletion of RanBP2 abolished restriction of HIV-1 in rhesus TRIM5 α overexpressing cells and blocked restriction of N-MLV by human endogenous TRIM5 α . Moreover, the K84R mutant lost its ability to restrict incoming retroviruses, indicating that TRIM5 α must be SUMOylated to carry antiviral activity. The presence of K84 within a phosphorylation-dependent SUMOylation pSUM domain implies that the activity of TRIM5 α is turned on sequentially first by phosphorylation then SUMOylation. Further work is needed to determine whether mutating the phosphoserine within the pSUM blocks

SUMOylation, and how TRIM5 α phosphorylation is regulated. Intriguingly, only SUMOylated TRIM5 α can antagonise incoming retroviruses, suggesting that a constitutively active restriction might be disadvantageous to the cell. Our results point to an elaborate intrinsic immunity mechanism that is mounted by the cell to regulate the antiviral function of a key restriction factor, suggesting that a constitutively active restriction might be disadvantageous to the cell.

In addition, RanBP2 modulates the subcellular localization of TRIM5 α . In dendritic cells, where efficient sensing of invading

pathogens is essential, SUMO modification of TRIM5 α leads to its relocation to the nucleus where it is no longer able to neutralize incoming cytoplasmic viruses¹³. The absence of RanBP2 in these cells leads to a partial relocation of TRIM5 α to the cytoplasm where it forms aggregates. Given this, it is likely that SUMOylation of TRIM5 α by RanBP2 can both promote and inhibit (via nuclear sequestration) its activity as a retroviral restriction factor, and that this is fine-tuned according to the cell type (lymphocytes versus DC¹³) and whether the cell is undergoing infection. Similarly, other restriction factors such as PML/TRIM19 and Mx proteins have been shown to be regulated by SUMO, either by direct conjugation or via SIM domains³¹, underlining the importance of the SUMO machinery in intrinsic antiviral defense mechanisms.

SUMOylation is a highly transient post-translational modification, which likely explains why TRIM5 α SUMOylated bands were more apparent when TRIM5 α was enriched, either by RanGAP1 co-immunoprecipitation or direct TRIM5 α immunoprecipitation, and in SUMO-overexpressing HeLa cells. Similarly, TRIM5 α localization at the nuclear pore was rarely observed except by TRIM5 α ×RanGAP1 PLA, suggesting that its residency at the nuclear pore is a transient phenomenon. A striking exception was provided by myotubes, in which TRIM5 α forms a perinuclear ring, which may be accounted for by the fact that RanBP2 is strongly induced during myogenic differentiation. The transient nature of SUMO modification of TRIM5 α in all other systems is likely to be important to fine-tune its function and localization.

Besides direct conjugation of TRIM5 α by SUMO variants¹¹, which we confirm in this study, the SUMO machinery is likely to contribute in additional ways to TRIM5 α function. Several SUMO-interacting motifs (SIM) identified in TRIM5 α ^{9,10,14} may mediate interaction with SUMOylated proteins. For instance, the physical association between TRIM5 α and RanGAP1 that we describe in this study likely involves an interaction between SUMOylated RanGAP1 and SIM domains on TRIM5 α , and may be key to bring TRIM5 α in proximity of its E3 SUMO ligase, RanBP2.

Viruses that replicate in the nucleus, such as lentiviruses, exploit NPCs and their associated proteins for nuclear entry. However, the NPC is also a privileged site for SUMOylating and deSUMOylating enzymes¹⁸, indicating that the nuclear pore is likely to both facilitate and impair viral nuclear import, acting both as gate and gatekeeper of viral passage into the nucleus³². In the case of HIV, the nucleoporin RanBP2 promotes capsid docking at the nuclear envelope and infection^{20,33,34}, but can also potentiate the antiviral functions of the restriction factor TRIM5 α by mediating its SUMOylation.

Interestingly, some HIV proteins have been shown to interact with the SUMO machinery and undergo SUMOylation^{35–37}, and further work will determine whether RanBP2 or more generally the SUMO machinery present at the NPC are involved in SUMOylating viral proteins during their transit into the nucleus.

Equally interesting is the presence of TRIM5 α with RanGAP1/RanBP2/Ubc9 at the nuclear pore. Although TRIM5 α is traditionally thought to intercept viral capsids during their rapid cytoplasmic trafficking towards the nucleus, the association of TRIM5 α with cytoplasmic bodies is not required for its anti-retroviral activity³⁸. Given our findings, it is possible that TRIM5 α can intercept capsids at the nuclear pore, where they accumulate for several hours prior to nuclear entry^{39,40}, a bottleneck that constitutes a point of vulnerability in HIV infections. The possible presence of TRIM5 α at the NPCs is reminiscent of MxB, another restriction factor that is also reported to operate at the cytoplasmic face of NPCs^{41,42}.

Results obtained here and elsewhere point to an essential role for the nuclear pore and more specifically RanBP2 in lentiviral infections, acting as a hub where virus, SUMO machinery, and restriction factors can meet.

Methods

Cells. HeLa-SUMO1 and -3 cell lines were obtained by transducing HeLa cells (ATCC) with Puro-His-SUMO1, or -3 lentiviral vector. HEK 293T cells were obtained from the ATCC. *Mus dunni* tail fibroblast (MDTF) cells devoid of TRIM5 α , APOBEC3G, and tetherin restriction factors, were used to titer the N-MLV vector⁴³. Myoblasts were isolated from the quadriceps of donor 1 CHQ and donor 2 160M (aged 5 days and 53 years, respectively⁴⁴), and differentiated into myotubes in low-serum DMEM 2% FBS for 5 days. Myotubes appear as polynucleated desmin-positive cells. Blood was obtained from healthy donors who signed an informed consent that their blood could be used for research purposes (Etablissement de Sang Français (EFS)). Dendritic cells were differentiated from blood monocytes as previously described¹³. The rhTRIM5 α cell clones A7 and B4 (for EV and T5, respectively) were obtained by transducing HeLa cells with the empty LPCX or LPCX-HA-rhTRIM5 α retroviral expression vectors²¹, selecting for puromycin resistance and performing limiting dilutions of the parent population. The clones were screened by flow cytometry (Novocyte, Ozyme) for HA expression.

Antibodies. Primary antibodies used were mouse anti-6×His (Clontech), -desmin (Dako), -Nup153 (gift from B. Burke), -3xFlag (Sigma), -actin (Sigma), -GAPDH (Sigma), -Ubc9 (Abgent) and α -tubulin (Sigma); rabbit anti-RanBP2 (Pierce), -TRIM5 α (gift from P. Bieniasz)^{13,45}, -SUMO1 and -SUMO3 (Cell Signaling); goat anti-RanGAP1 (Santa Cruz Biotechnology); rat anti-HA (3F10, Roche). Secondary antibodies used were anti-mouse, -rabbit, -goat, -rat HRP (GE Healthcare), protein G-HRP used to avoid IgG detection when revealing immunoprecipitated TRIM5 α , and Alexa-488, -555, and -647 (Invitrogen).

RNA interference and quantitative PCR. Lentiviral vectors shE, shBP2, and shTRIM5 α have been described previously^{13,20}. The following siRNAs were used: siCtrl GAGAAGGUAAGCUGGACAUU and siBP2 CACAGACAAAGCCUGUAAUU. HeLa, HeLa-SUMO1, and HeLa-SUMO3 cells (0.5×10^6) seeded at d0 were transduced at d1 with MOI 50 lentiviral vector. Alternatively, cells were lipofected with 50 nM siRNA using RNAiMAX transfection reagent (Invitrogen). Realtime quantitative PCR (qPCR) reactions were performed on a QuantStudio 5 (Thermo Fisher Scientific) using the following primers and probes.

RBP2-F: ACAATGGAATTAAGCCCTTAAATGT,
RBP2-R: GAAACAATCAGCTACTTCTTTAGTTTTTA,
RBP2-P: FAM-TTGGACTGCCTCAGATTATGCTGATGGAGAAGCAA-3'
BHQ1,
 β 2M-F: GTGCTCGCGCTACTCTCTCT,
 β 2M-R: CTCTGCTGGATGACGTGAGT,
 β 2M-P: HEX-CGCTGGATAGCCTCCAGGCC-BHQ1,
Pol-F: TTTAGATGGAATAGATAAGGCCAA,
Pol-R: CAGCTGGCTAACTATTTCTTTTGCTA,
Pol-P: FAM-AATCACTAGCCATTGCTCTCCAATTAC-TAMRA,
Actin-F: AACACCCAGCCATGTACGT,
Actin-R: CGGTGAGGATCTTCATAGGTAGT,
Actin-P: FAM-CCAGCCAGGTCCAGACGCAGGA-BHQ,
TRIM5-F: TTGGATCCTGGGGGTATGTGCTGG,
TRIM5-R: TGATATTGAAGAATGAGACAGTGCAG,
RPL13A-F: CCTGGAGGAGAAGAGGAAAGAGA,
RPL13A-R: TTGAGGACCTCTGTATTGTCAA.

Viruses. HIV-1 was produced by transient co-transfection of 293T cells by calcium phosphate precipitation with the envelope defective proviral plasmid (pLAI- Δ env) and the VSV-G envelope expression plasmid. Virus supernatants were titered by p24 ELISA assay (Clontech). The N-MLV vector was produced by co-transfection of pCFG2-YFP, pCIG3N, and pVSV-G. Vector supernatants were collected at 2 dpt, cleaned up through 0.45 μ m filter, and concentrated by ultracentrifugation for 1 h at 22,000 rpm using a SW32Ti Beckman rotor. Vector stocks were titered in MDTF cells by flow cytometry.

Fluorescence imaging. Widefield microscopy was performed using an Evos microscope (Life Technologies). Confocal microscopy and PLA were performed on a LSM880 microscope as previously described^{11,13}. Quantification of PLA spots per cells was performed using Imaris v 9.0 (ImarisXT, Bitplane) on 4 different z-stacks (~200 nuclei) for each condition \pm SD. Proximity of spots to the nuclear envelope was assessed using a nuclear cache within a 1 μ m distance of the nuclear envelope. Super-resolution (80 nm) imaging was performed on a CoDiM 100 system (Bioaxial France) equipped with 488, 561, and 640 lasers (Toptica, Germany). The CoDiM system was mounted on a Nikon Eclipse upright confocal microscope (Nikon, Japan) equipped with 1.2NA water immersion and 1.49NA oil immersion objectives (Nikon, Japan). CoDim super-

resolution imaging uses a biaxial crystal to project a variety of high-contrast Airy-disc sized excitation patterns with high spatial-frequency content on the sample. Resulting fluorescence images are then recorded on a low-noise Hamamatsu Orca flash 4 sCMOS camera (Hamamatsu, Japan) and final images are obtained using a proprietary maximum likelihood a posteriori image reconstruction algorithm as described in ref. 46.

Site-directed mutagenesis. The K84R, K111R, and K167R human TRIM5 α mutants were generated by site-directed mutagenesis using the QuikChange II site-directed mutagenesis kit (Agilent).

Co-immunoprecipitation experiments. RanGAP1 and TRIM5 α co-immunoprecipitation were carried out at 4 days post-transduction with lentiviral vectors shE or shBP2 using cytosolic extracts (prepared using NE-PER kit (Pierce)) and whole cell lysates, respectively, in the presence of 20 mM NEM (N-Ethylmaleimide, deSUMOylase inhibitor) and EDTA-free protease inhibitor cocktail (Roche). Supernatants were pre-cleared by incubation with 40 μ l Novex recombinant protein G sepharose and 3 μ l normal goat serum (Invitrogen) for 90 min at 4 °C followed by centrifugation at 350g for 5 min. Supernatants were then incubated with 100 μ l goat α -RanGAP1 or 10 μ l rabbit α -TRIM5 α antibody and 40 μ l protein G sepharose overnight at 4 °C. Beads were washed 5 times in lysis buffer before elution.

To detect SUMOylation of WT and mutant human TRIM5 α constructs, HEK 293T or MDTF cells ($2\text{--}3 \times 10^6$) seeded at d0 were transfected with 10 μ g of pcDNA3.1 empty vector, or Flag-tagged TRIM5 α constructs at d1 using calcium phosphate precipitation and JetPEI (PolyPlus), respectively. Cells were lysed at d4 in 500 μ l of IP lysis buffer and immunoprecipitation was performed as described above with anti-Flag antibody. The bound proteins Flag, TRIM5 α , and SUMO1 were detected by Western blotting. All blots were acquired using the Chemidoc XRS+ imaging system and band quantifications were performed using the Image LabTM software.

In vitro SUMO assay. Human TRIM5 α recombinant protein was produced in *Escherichia coli* as a hexa-His fusion protein and purified by Ni-affinity chromatography (NIH AIDS Reagent Program). In vitro SUMO assay was performed using a SUMOylation kit by Enzo according to manufacturer's protocol. Reactions containing SUMO1, E1, E2 (Ubc9) 200 nM of His-tagged HR-TRIM5 α , and increasing amounts of RanBP2 Δ FG-GST (50, 100, or 200 nM) were incubated 60 min at 37 °C. Controls without ATP or E2 were included. Proteins were analyzed by western blotting using rabbit α -SUMO1 antibody provided by the Kit or mouse α -6xHis antibody.

Data availability

Authors confirm that all relevant data are available if requested.

Received: 2 March 2018 Accepted: 19 October 2018

Published online: 15 November 2018

References

- Hay, R. T. SUMO: a history of modification. *Mol. Cell* **18**, 1–12 (2005).
- Seeler, J. S. & Dejean, A. Nuclear and unclear functions of SUMO. *Nat. Rev. Mol. Cell Biol.* **4**, 690–699 (2003).
- Gareau, J. R. & Lima, C. D. The SUMO pathway: emerging mechanisms that shape specificity, conjugation and recognition. *Nat. Rev. Mol. Cell Biol.* **11**, 861–871 (2010).
- Liang, Y. C. et al. SUMO5, a novel poly-SUMO isoform, regulates PML nuclear bodies. *Sci. Rep.* **6**, 26509 (2016).
- Johnson, E. S. & Gupta, A. A. An E3-like factor that promotes SUMO conjugation to the yeast septins. *Cell* **106**, 735–744 (2001).
- Kagey, M. H., Melhuish, T. A. & Wotton, D. The polycomb protein Pc2 is a SUMO E3. *Cell* **113**, 127–137 (2003).
- Pichler, A., Gast, A., Seeler, J. S., Dejean, A. & Melchior, F. The nucleoporin RanBP2 has SUMO1 E3 ligase activity. *Cell* **108**, 109–120 (2002).
- Everett, R. D., Boutell, C. & Hale, B. G. Interplay between viruses and host SUMOylation pathways. *Nat. Rev. Microbiol.* **11**, 400–411 (2013).
- Arriagada, G., Muntean, L. N. & Goff, S. P. SUMO-interacting motifs of human TRIM5 α are important for antiviral activity. *PLoS Pathog.* **7**, e1002019 (2011).
- Lukic, Z., Goff, S. P., Campbell, E. M. & Arriagada, G. Role of SUMO-1 and SUMO interacting motifs in rhesus TRIM5 α -mediated restriction. *Retrovirology* **10**, 10 (2013).
- Dutrieux, J., Portillo, D. M., Arhel, N. J., Hazan, U. & Nisole, S. TRIM5 α is a SUMO substrate. *Retrovirology* **12**, 28 (2015).
- Diaz-Griffero, F., Gallo, D. E., Hope, T. J. & Sodroski, J. Trafficking of some old world primate TRIM5 α proteins through the nucleus. *Retrovirology* **8**, 38 (2011).
- Portillo, D. M. et al. Endogenous TRIM5 α function is regulated by SUMOylation and nuclear sequestration for efficient innate sensing in dendritic cells. *Cell Rep.* **14**, 355–369 (2016).
- Nepveu-Traversy, M. E. et al. A putative SUMO interacting motif in the B30.2/SPRY domain of rhesus macaque TRIM5 α important for NF- κ B/AP-1 signaling and HIV-1 restriction. *Heliyon* **2**, e00056 (2016).
- Hofmann, T. G. & Will, H. Body language: the function of PML nuclear bodies in apoptosis regulation. *Cell Death Differ.* **10**, 1290–1299 (2003).
- Sachdev, S. et al. PIASy, a nuclear matrix-associated SUMO E3 ligase, represses LEF1 activity by sequestration into nuclear bodies. *Genes Dev.* **15**, 3088–3103 (2001).
- Melchior, F., Schergaut, M. & Pichler, A. SUMO: ligases, isopeptidases and nuclear pores. *Trends Biochem. Sci.* **28**, 612–618 (2003).
- Palancade, B. & Doye, V. Sumoylating and desumoylating enzymes at nuclear pores: underpinning their unexpected duties? *Trends Cell Biol.* **18**, 174–183 (2008).
- Saitoh, H., Pu, R., Cavenagh, M. & Dasso, M. RanBP2 associates with Ubc9p and a modified form of RanGAP1. *Proc. Natl Acad. Sci. USA* **94**, 3736–3741 (1997).
- Di Nunzio, F. et al. Human nucleoporins promote HIV-1 docking at the nuclear pore, nuclear import and integration. *PLoS ONE* **7**, e46037 (2012).
- Stremlau, M. et al. The cytoplasmic body component TRIM5 α restricts HIV-1 infection in Old World monkeys. *Nature* **427**, 848–853 (2004).
- Markiewicz, E., Ledran, M. & Hutchison, C. J. Remodelling of the nuclear lamina and nucleoskeleton is required for skeletal muscle differentiation in vitro. *J. Cell Sci.* **118**, 409–420 (2005).
- Asally, M. et al. Nup358, a nucleoporin, functions as a key determinant of the nuclear pore complex structure remodeling during skeletal myogenesis. *FEBS J.* **278**, 610–621 (2011).
- Moran, J. L., Li, Y., Hill, A. A., Mounts, W. M. & Miller, C. P. Gene expression changes during mouse skeletal myoblast differentiation revealed by transcriptional profiling. *Physiol. Genomics* **10**, 103–111 (2002).
- Beauchair, G., Bridier-Nahmias, A., Zagury, J. F., Saib, A. & Zamborlini, A. JASSA: a comprehensive tool for prediction of SUMOylation sites and SIMs. *Bioinformatics* **31**, 3483–3491 (2015).
- Picard, N. et al. Identification of estrogen receptor beta as a SUMO-1 target reveals a novel phosphorylated sumoylation motif and regulation by glycogen synthase kinase 3beta. *Mol. Cell Biol.* **32**, 2709–2721 (2012).
- Melchior, F. SUMO—nonclassical ubiquitin. *Annu. Rev. Cell. Dev. Biol.* **16**, 591–626 (2000).
- Langelier, C. R. et al. Biochemical characterization of a recombinant TRIM5 α protein that restricts human immunodeficiency virus type 1 replication. *J. Virol.* **82**, 11682–11694 (2008).
- Kirsh, O. et al. The SUMO E3 ligase RanBP2 promotes modification of the HDAC4 deacetylase. *EMBO J.* **21**, 2682–2691 (2002).
- Miyachi, Y., Yogosawa, S., Honda, R., Nishida, T. & Yasuda, H. Sumoylation of Mdm2 by protein inhibitor of activated STAT (PIAS) and RanBP2 enzymes. *J. Biol. Chem.* **277**, 50131–50136 (2002).
- Hannoun, Z., Maarifi, G. & Chelbi-Alix, M. K. The implication of SUMO in intrinsic and innate immunity. *Cytokine Growth Factor Rev.* **29**, 3–16 (2016).
- Flatt, J. W. & Greber, U. F. Misdelivery at the nuclear pore complex—stopping a virus dead in its tracks. *Cells* **4**, 277–296 (2015).
- Schaller, T. et al. HIV-1 capsid-cyclophilin interactions determine nuclear import pathway, integration targeting and replication efficiency. *PLoS Pathog.* **7**, e1002439 (2011).
- Brass, A. L. et al. Identification of host proteins required for HIV infection through a functional genomic screen. *Science* **319**, 921–926 (2008).
- Gurer, C., Berthou, L. & Luban, J. Covalent modification of human immunodeficiency virus type 1 p6 by SUMO-1. *J. Virol.* **79**, 910–917 (2005).
- Jaber, T. et al. Human Ubc9 contributes to production of fully infectious human immunodeficiency virus type 1 virions. *J. Virol.* **83**, 10448–10459 (2009).
- Zamborlini, A. et al. Impairment of human immunodeficiency virus type-1 integrase SUMOylation correlates with an early replication defect. *J. Biol. Chem.* **286**, 21013–21022 (2011).
- Song, B. et al. TRIM5 α association with cytoplasmic bodies is not required for antiretroviral activity. *Virology* **343**, 201–211 (2005).
- Arhel, N. et al. Quantitative four-dimensional tracking of cytoplasmic and nuclear HIV-1 complexes. *Nat. Methods* **3**, 817–824 (2006).
- Arhel, N. J. et al. HIV-1 DNA Flap formation promotes uncoating of the pre-integration complex at the nuclear pore. *EMBO J.* **26**, 3025–3037 (2007).
- Kane, M. et al. MX2 is an interferon-induced inhibitor of HIV-1 infection. *Nature* **502**, 563–566 (2013).
- Goujon, C. et al. Human MX2 is an interferon-induced post-entry inhibitor of HIV-1 infection. *Nature* **502**, 559–562 (2013).

43. Carthagen, L. et al. Implication of TRIM alpha and TRIMCyp in interferon-induced anti-retroviral restriction activities. *Retrovirology* **5**, 59 (2008).
44. Edom, F., Mouly, V., Barbet, J. P., Fiszman, M. Y. & Butler-Browne, G. S. Clones of human satellite cells can express in vitro both fast and slow myosin heavy chains. *Dev. Biol.* **164**, 219–229 (1994).
45. Zhang, F., Perez-Caballero, D., Hatzioannou, T. & Bieniasz, P. D. No effect of endogenous TRIM5alpha on HIV-1 production. *Nat. Med.* **14**, 235–236 (2008).
46. Caron, J. et al. Conical diffraction illumination opens the way for low phototoxicity super-resolution imaging. *Cell Adh. Migr.* **8**, 430–439 (2014).

Acknowledgements

This work was supported by grants from the Agence Nationale de Recherches sur le SIDA et les hépatites virales (ANRS), the ATIP-Avenir program, Bioaxial, and the EpiGenMed Labex (“Investissements d’avenir” ANR-10-LABX-12-01). The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: Human TRIM5 Alpha Recombinant Protein. Flag-tagged wild-type and K10R TRIM5 α were provided by Stephen Goff (Columbia University, USA). The Nup153 antibody was provided by Brian Burke (Institute of Medical Biology, Singapore). The following online sequence prediction tools were used: JASSA (<http://www.jassa.fr/>) and PhosphoSite Plus (www.phosphosite.org). We thank Nathalie Eisenhardt, Andrea Pichler (Max Planck Institute of Immunobiology and Epigenetics, Freiburg, Germany), and Guillaume Bossis (Institut de Génétique Moléculaire de Montpellier) for expert advice with the SUMO assays. We acknowledge the imaging facility MRI, member of the national infrastructure France-BioImaging supported by the French National Research Agency (ANR-10-INBS-04, “Investments for the future”).

Author contributions

G.M., J.F., D.M.P., A.B., J.D., S.O., S.N., and N.J.A. performed experiments. J.D. and G.B.-B. provided reagents and expertise. S.N. and N.J.A. designed the experiments. G.M., J.F., S.N., and N.J.A. wrote the manuscript.

Additional information

Supplementary Information accompanies this paper at <https://doi.org/10.1038/s42003-018-0198-0>.

Competing interests: At the time of the study, D.M.P. and S.O. were employees of BioAxial (Paris) that commercialises the CoDiM system. The other authors declare no competing interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

Publisher’s note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018