



HAL
open science

Effect of temperature on an alga-grazer trophic transfer: A dual stable isotope (^{13}C , ^{15}N) labeling experiment

Erwann Legrand, Sophie Martin, Cédric Leroux, Pascal Riera

► To cite this version:

Erwann Legrand, Sophie Martin, Cédric Leroux, Pascal Riera. Effect of temperature on an alga-grazer trophic transfer: A dual stable isotope (^{13}C , ^{15}N) labeling experiment. *Marine Ecology*, 2018, 39 (2), pp.e12495. 10.1111/maec.12495 . hal-01949447

HAL Id: hal-01949447

<https://hal.sorbonne-universite.fr/hal-01949447>

Submitted on 10 Dec 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of temperature on an alga-grazer trophic transfer: A dual stable isotope (^{13}C , ^{15}N) labeling experiment

Erwann Legrand^{1,2}  | Sophie Martin^{1,2} | Cédric Leroux^{1,3} | Pascal Riera^{1,2}

¹Sorbonne Universités, UPMC Univ Paris 06, Station Biologique, Roscoff, France

²CNRS, UMR7144, Station Biologique, Roscoff, France

³CNRS, FR2424, Station Biologique, Roscoff, France

Correspondence

Erwann Legrand, Sorbonne Universités, UPMC Univ Paris 06, Station Biologique, Place Georges Teissier, Roscoff Cedex, France.

Email: erwann.legrand@sb-roscoff.fr

Funding information

Brittany Regional Council; French National Research Agency, Grant/Award Number: ANR-10-BTBR-04; Écosphère Continentale et Côtière

Abstract

In this study, we examined the impact of temperature on the carbon and nitrogen trophic transfers from a macroalga to a macro-grazer by the use of dual ^{13}C - and ^{15}N -labeling. Using an experimental approach in mesocosms, individuals of the urchin *Psammechinus miliaris* were maintained for 1 month at 17°C (mean summer temperature in the Bay of Brest) and at 20°C (maximum summer temperature) and fed with ^{13}C - and ^{15}N -labeled *Solieria chordalis*. The results showed that the urchins' ^{13}C uptake was $0.30 \mu\text{g}^{13}\text{C g dry weight (DW)}^{-1}$ at 17°C and $0.14 \mu\text{g}^{13}\text{C g DW}^{-1}$ at 20°C at the end of the experiment. The lower uptake at the higher temperature may be attributed to a decrease in metabolic activity at 20°C, involving lower feeding and/or respiration rates. Conversely, no significant effect of temperature was detected on ^{15}N uptake. At the end of the experiment, the urchins' ^{15}N uptake was $0.04 \mu\text{g}^{15}\text{N g DW}^{-1}$ at 17°C and $0.03 \mu\text{g}^{15}\text{N g DW}^{-1}$ at 20°C. This suggests that temperature may affect carbon and nitrogen trophic fluxes differently. The use of dual isotope labeling offers interesting prospects and needs to be further extended in order to better understand trophic interactions in marine communities and the consequences of current environmental changes, such as global warming.

KEYWORDS

carbon, macroalga, nitrogen, stable isotope labeling, temperature, trophic interaction, uptake, urchin

1 | INTRODUCTION

Temperature is one of the most important environmental factors affecting biological processes in marine ectothermic species (Nguyen et al., 2011; Uthicke, Liddy, Nguyen, & Byrne, 2014; Zuo, Moses, West, Hou, & Brown, 2012). Several studies demonstrated the effect of temperature on physiological processes, including feeding rates (Frey & Gagnon, 2015; Gooding, Harley, & Tang, 2009; Thomas, Crear, & Hart, 2000). However, the response to changes in temperature in terms of assimilation rates remains poorly documented, despite the importance of this process in species ecophysiology. In benthic ecosystems, urchins are considered as strong drivers of community structure, mainly as a result of their ability to control macroalgal biomass (Steneck, 2013). Within maerl beds of the Bay of Brest (Britany, France), the purple-tipped sea urchin, *Psammechinus miliaris* (Müller,

1771), can reach high densities and is known to be a major factor regulating macroalgal biomass (Guillou, Grall, & Connan, 2002).

Several studies have already evidenced the usefulness of stable isotope labeling for quantifying the importance of microalgal or bacterial production as food source for grazers (Herman, Middelburg, Widdows, Lucas, & Heip, 2000; Leroy et al., 2012; Middelburg et al., 2000). However, studies using dual stable isotope labeling (^{13}C , ^{15}N) to examine trophic transfers from macroalgae to macro-grazers under the influence of environmental factors, such as temperature, are still lacking.

The present study aimed to investigate whether temperature can influence macroalgal-derived carbon and nitrogen assimilation into grazer tissues. Carbon and nitrogen trophic transfers from the red seaweed *Solieria chordalis* (J. Agardh, 1842) to the urchin *P. miliaris* were experimentally examined using dual ^{13}C - and ^{15}N -labeling.

In the Bay of Brest, the alga *S. chordalis* is abundant (Hily, Potin, & Floch, 1992) and reaches the highest biomass in the summer (Guillou et al., 2002). Therefore, trophic transfers were investigated at 17°C (mean summer temperature, recorded by Service d'Observation en Milieu Littoral, SOMLIT) and at 20°C (maximum summer temperature, Martin, Clavier, Chauvaud, & Thouzeau, 2007).

2 | MATERIAL AND METHODS

2.1 | Species collection

Individuals of *Psammechinus miliaris* were collected from a maerl bed in the bay of Brest, France (48°17'18"N, 04°23'46"W) in April 2016 using a naturalist's dredge (width: 1 m, height: 0.2 m, net: 1.5 m long). Organisms were then transported in seawater tanks to the Roscoff Marine Station. To mitigate the stress experienced by the species during sampling and transport, organisms were maintained in open-flow aquaria pending experiments. Thalli of the seaweed *Solieria chordalis* were collected from maerl beds in the Bay of Brest (48°19'56"N, 04°19'56"W) in early June 2016 and transported to the Roscoff Marine Station.

2.2 | Experimental design

In May 2016, 36 *Psammechinus miliaris* individuals of medium size (26 ± 3 mm) were randomly assigned to 10 15-l aquaria (Supporting Information Fig. S1). In all aquaria, the temperature was gradually increased by 0.5°C per day over 4 days to reach 17°C, corresponding to the mean summer temperature in the Bay of Brest (SOMLIT). Then, in five aquaria, the temperature was increased again for 6 days to reach 20°C, the mean maximum summer temperature in the Bay of Brest (Martin et al., 2007). The temperature was controlled in two 100-l tanks, continuously supplied with filtered (5 µm) open natural seawater, with a high water flow rate of 150 L/hr per tank. The temperature was maintained by an offline feedback system (IKS Aquastar, Karlsbad, Germany) that activated or stopped heaters, controlling the temperature in the tanks. Each 100-L tank provided seawater to five aquaria. During the experiment, the temperature was at 17.2 ± 0.3 and at 19.9 ± 0.2°C, according to the treatments (17 and 20°C). For both temperature treatments, urchins were starved 10 days before the feeding experiment to empty their gut. During the experiment, the irradiance was 90–100 µmol photons m⁻² s⁻¹, corresponding to the mean in situ summer irradiance at 5 m depth on maerl beds in the Bay of Brest (Martin, Castets, & Clavier, 2006).

2.3 | Seaweed labeling

Thalli of *Solieria chordalis* were cleaned of epiphytes, rinsed with filtered seawater (on 0.7 µm GF/F glass microfiber filters) and placed in a 9-L closed-circuit aquarium. Five liters of filtered (0.7 µm) seawater was added in the aquarium. A solution of ¹³C-HCO₃⁻ at 1 g/L was prepared from NaH¹³CO₃ (99 atom% ¹³C) and 1 L of filtered 0.7 µm GF/F seawater. Similarly, a solution of ¹⁵N-NH₄ at 0.25 g/L

was prepared by diluting ¹⁵NH₄Cl (99 atom% ¹⁵N) in filtered seawater. The seaweeds were fertilized with the solutions of NaH¹³CO₃ and ¹⁵NH₄Cl for a period of 7 days. Each day, 10 ml of a solution of phosphate (PO₄³⁻) at 1 g/L was also added to ensure that P was not limiting. A circulation pump ensured nutrient homogeneity in the aquarium. At the end of the labeling period, some of the thalli were rinsed with filtered seawater and immediately frozen at -20°C until isotope measurements. In order to provide sufficient food for urchins for the duration of the experiment, a second stock of *S. chordalis* was labeled after 8 days, following the same protocol.

2.4 | Feeding experiment

For each temperature treatment, three urchins were collected at day 0 (T0) to measure the initial isotopic signature. Labeled thalli of *Solieria chordalis* from the same labeled stock were provided every 2 days to urchins of eight aquaria (four per condition) at an amount of 1 g of fresh weight per urchin. In one control aquarium per temperature treatment, urchins were fed with non-labeled thalli at the same amount (1 g per urchin). After 6 days, one urchin was collected per aquarium. A second urchin was collected after 15 days and the last urchin after 30 days (T30). Just before collection, urchins were starved for 48 hr to allow the assimilation of ingested algae and to allow evacuation of their digestive content. All samples were immediately frozen at -20°C pending isotope measurement.

2.5 | Isotope analysis

For each individual, the digestive system was removed and samples were acidified under a magnifying glass to remove calcareous structures (Kamp & Witte, 2005). Urchin soft tissues and algae were dried in an oven for 48 hr at 60°C, then ground to a fine powder. Subsamples were weighed into tin cups for isotope analyses. Stable isotope ratios for carbon (R = ¹³C/¹²C) and nitrogen (R = ¹⁵N/¹⁴N) were determined using a carbon, hydrogen and nitrogen (CHN) analyser (ThermoFinnigan 1112 Series) interfaced with a mass spectrometer (ThermoFinnigan MAT Deltaplus) via a Conflow III open split interface.

2.6 | Uptake of ¹³C and ¹⁵N by urchins

Carbon and nitrogen ratios were used to calculate δ-values (in ‰):

$$\delta X = [(R_{\text{sample}}/R_{\text{standard}}) - 1] \times 1,000$$

where X is ¹³C or ¹⁵N. Sample ratios (R_{sample}) were reported relative to international standards (R_{standard}) Vienna Pee Dee Belemnite (R_{V-PDB} = 0.0112372) for carbon and atmospheric N₂ (R_{air} N₂ = 0.0036765) for nitrogen.

Incorporation of ¹³C or ¹⁵N labels in urchins was calculated as excess ¹³C or ¹⁵N, expressed in terms of specific uptake (i.e., Δδ¹³C = δ¹³C_{T30} - δ¹³C_{T0} and Δδ¹⁵N = δ¹⁵N_{T30} - δ¹⁵N_{T0}) and total uptake (I), in accordance with previous studies (Herman et al., 2000;

Leroy et al., 2012). I corresponds to the product of excess ^{13}C or ^{15}N (E) and carbon and nitrogen biomass (B), with I in $\mu\text{g}^{13}\text{C}$ or ^{15}N , B in μg C or N. E is calculated as the difference between the fraction ^{13}C or ^{15}N of the initial value (F_{T_0}) and the final value ($F_{T_{30}}$):

$$E = F_{T_{30}} - F_{T_0}$$

with $F = ^{13}\text{C}/(^{13}\text{C} + ^{12}\text{C}) = R/(R + 1)$ for ^{13}C and $F = ^{15}\text{N}/(^{15}\text{N} + ^{14}\text{N}) = R/(R + 1)$ for ^{15}N . The carbon and nitrogen isotopic ratios (R) were obtained from the measured $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, respectively, as:

$$R = (\delta^{13}\text{C}/1,000 + 1) \times R_{\text{VPDB}}$$

with R_{VPDB} the Vienna Pee Dee Belemnite-limestone standard, and,

$$R = (\delta^{15}\text{N}/1,000 + 1) \times R_{\text{airN}_2}$$

with R_{airN_2} the atmospheric dinitrogen standard.

Total carbon and nitrogen transfer (^{12}C and ^{13}C ; ^{14}N and ^{15}N) from algae to grazers was obtained from ^{13}C and ^{15}N incorporation, dividing $I^{13}\text{C}$ or $I^{15}\text{N}$ of urchins by $F^{13}\text{C}$ or $F^{15}\text{N}$ of the source.

2.7 | Data analysis

All data were expressed as the mean \pm SE. Statistical analysis was completed using the statistical software R version 3.2.2 (R Core Team, 2014). $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of labeled *Solieria chordalis* were compared between the beginning (0 day) and the end (8 days) using the non-parametric Wilcoxon-Mann-Whitney test, as assumptions of normality (Shapiro test) and homogeneity of variances (Bartlett test) were not met. The differences in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ between labeled and non-labeled urchins were also examined using the Wilcoxon-Mann-Whitney test. The effects of temperature and time on $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of urchins supplied with labeled *S. chordalis* were examined using the non-parametric Scheirer-Ray-Hare test, as assumptions of normality and homogeneity of variances (Bartlett test) were not met. Similarly, the Scheirer-Ray-Hare test

was performed to examine the effect of increased temperature and time on urchin ^{13}C and ^{15}N uptake.

3 | RESULTS

After 8 days, stocks of *Solieria chordalis* thalli were labeled at a level of $\delta^{13}\text{C} = 1152 \pm 106\%$, ($n = 3$) and $2131 \pm 617\%$, ($n = 3$) for stocks 1 and 2, respectively. These levels were significantly higher than non-labeled thalli ($\delta^{13}\text{C} = -17.6 \pm 0.4\%$; Wilcoxon-Mann-Whitney test, $p = .005$). Similarly, labeled *S. chordalis* were significantly ^{15}N -enriched ($\delta^{15}\text{N} = 5191 \pm 701\%$, $n = 3$ and $4604 \pm 319\%$, $n = 3$, for stocks 1 and 2, respectively) compared with non-labeled seaweeds ($\delta^{15}\text{N} = 16.2 \pm 0.3\%$; Wilcoxon-Mann-Whitney test, $p = .005$).

At the beginning of the experiment, *Psammechinus miliaris* exhibited an initial $\delta^{13}\text{C}$ level of $-17.2 \pm 0.7\%$ ($n = 3$) and $-18.4 \pm 0.3\%$ ($n = 3$) at 17 and 20°C, respectively. In urchins supplied with labeled *S. chordalis*, $\delta^{13}\text{C}$ reached $941 \pm 203\%$ ($n = 4$) at 17°C and $623 \pm 118\%$ ($n = 4$) at 20°C after 30 days (Figure 1a). This enrichment was significant compared with the initial isotopic signatures (Wilcoxon-Mann-Whitney, $p < .001$). Throughout the experiment, higher $\delta^{13}\text{C}$ values were observed at 17 than at 20°C, whereas no significant effect of time was observed (Table 1, Figure 1a).

Psammechinus miliaris initial $\delta^{15}\text{N}$ values were $5.3 \pm 0.8\%$ ($n = 3$) and $4.4 \pm 0.3\%$ ($n = 3$) at 17 and 20°C, respectively. After 30 days, $\delta^{15}\text{N}$ in urchins supplied with labeled *S. chordalis* reached $1405 \pm 109\%$ ($n = 4$) at 17°C and $1514 \pm 335\%$ ($n = 4$) at 20°C (Figure 1b). The ^{15}N enrichment after 30 days was significant in comparison with initial isotopic values (Wilcoxon-Mann-Whitney, $p < .001$; Figure 1b). Temperature and time had no significant effect on urchins' $\delta^{15}\text{N}$ (Table 1).

At 17°C, mean ^{13}C specific uptake in urchins supplied with labeled *S. chordalis* ranged from $0.22 \mu\text{g}^{13}\text{C}$ (day 15) to $0.36 \mu\text{g}^{13}\text{C}$ (day 6) and reached $0.30 \mu\text{g}^{13}\text{C}$ after 30 days (Figure 2a). At 20°C, ^{13}C uptake ranged from $0.14 \mu\text{g}^{13}\text{C}$ (day 30) to $0.19 \mu\text{g}^{13}\text{C}$ (day 6). ^{13}C uptake was significantly higher at 17 than at 20°C (Table 2), whereas no significant effect of time was evidenced (Table 2). *Psammechinus*




FIGURE 1 Variations in $\delta^{13}\text{C}$ (a) and $\delta^{15}\text{N}$ (b; mean \pm SE) in *Psammechinus miliaris* maintained for 6, 15 and 30 days at 17°C (black dots) and at 20°C (white dots) and supplied with labeled (full lines) and non-labeled (dotted lines) *Solieria chordalis* ($n = 4$)

TABLE 1 Results of Scheirer–Ray–Hare tests to assess the effect of temperature (17 and 20°C) and time (days 6, 15 and 30) and their combined effect on the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in *Psammechinus miliaris* supplied with labeled *Solieria chordalis* ($n = 4$)

Factor	df	$\delta^{13}\text{C}$ (‰)		$\delta^{15}\text{N}$ (‰)	
		H	<i>p</i>	H	<i>p</i>
Temperature	1	7.3	.007**	1.2	.27
Time	2	1.3	.53	4.6	.10
Temperature × time	2	1.0	.60	2.2	.32

Significant *p*-values are in bold.
p* < .05; *p* < .01; ****p* < .001.

miliaris supplied with labeled *S. chordalis* exhibited ^{15}N specific uptake values ranging from 0.04 $\mu\text{g}^{15}\text{N}$ (day 30) to 0.11 $\mu\text{g}^{15}\text{N}$ (day 6) at 17°C (Figure 2b). At 20°C, ^{15}N uptake varied between 0.03 $\mu\text{g}^{15}\text{N}$ (day 30) and 0.05 $\mu\text{g}^{15}\text{N}$ (day 6). Temperature had no significant effect on ^{15}N specific uptake in *P. miliaris*, whereas ^{15}N uptake decreased significantly over time (Table 2).

4 | DISCUSSION

In this experiment, the urchin *Psammechinus miliaris* consumed substantial amounts of the alga *Solieria chordalis* at 17 and 20°C. The gut transit time of *P. miliaris* varies according to the food source ingested (Bedford & Moore, 1985) and may reach 42 hr in small individuals (Otero-Villanueva, 2004). The present feeding experiment was performed over 30 days, which is sufficient to measure carbon and nitrogen assimilation rates in *P. miliaris*.

At 17°C, an increase in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ occurred during the first six days, corresponding to a significant ^{13}C and ^{15}N uptake. This may be a result of a high level of consumption of the labeled *S. chordalis* following the starving period that occurred before the experiments, as recently observed by Rubilar et al. (2016). In addition, several studies have reported a link between feeding and other metabolic processes in echinoids, such as respiration or excretion (Brockington & Peck, 2001; Carr & Bruno, 2013). These physiological processes can influence the $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$ ratios. Among the physiological processes that can influence the $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$ ratios, the

TABLE 2 Results of Scheirer–Ray–Hare tests on the effect of temperature (17 and 20°C) and time (days 6, 15 and 30) and their combined effect on ^{13}C and ^{15}N uptake ($I^{13}\text{C}$ and $I^{15}\text{N}$, respectively) in *Psammechinus miliaris* supplied with labeled *Solieria chordalis* ($n = 4$)

Factor	df	$I^{13}\text{C}$ $\mu\text{g}^{13}\text{C}$ g DW ⁻¹		$I^{15}\text{N}$ $\mu\text{g}^{15}\text{N}$ g DW ⁻¹	
		H	<i>p</i>	H	<i>p</i>
Temperature	1	5.8	.016*	0.74	.39
Time	2	2.3	.32	11.9	.003**
Temperature × time	2	1.0	.62	1.5	.47

Significant *p*-values are in bold.
p* < .05; *p* < .01; ****p* < .001.
DW, dry weight.

increase in respiration rates induced by higher levels of feeding may contribute to ^{13}C enrichment due to a preferential loss of ^{12}C (Rau et al., 1983), and higher ^{15}N enrichment may be due to an increase in ^{15}N depleted nitrogen excretion (Minagawa & Wada, 1984) in the form of urea (Basuyaux & Mathieu, 1999). At 20°C, although the food source was identical, the increases in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of *P. miliaris* during the first 6 days were less pronounced as compared to those at 17°C. Hence, from the present study, temperature appears to play an important role in the trophic fluxes from algae to grazers, especially regarding carbon uptake.

^{13}C uptake was significantly higher at 17 than at 20°C throughout the experiment. At the end of the experiment, urchins supplied with labeled ^{13}C showed an assimilation of 10.1 and 4.9 $\text{ng}^{13}\text{C day}^{-1}$ at 17 and 20°C, respectively. The effect of temperature increase on the ^{13}C uptake of *P. miliaris* appeared to significantly alter the total carbon uptake, as estimated total carbon fluxes after 30 days were 0.29 $\mu\text{g C g dry weight (DW)}^{-1} \text{day}^{-1}$ at 17°C and 0.14 $\mu\text{g C g DW}^{-1} \text{day}^{-1}$ at 20°C. These differences in carbon uptake observed between the two temperatures may be attributed to the higher metabolic activity at 17 than at 20°C, involving higher feeding and respiration rates. The lower metabolism at 20°C contrasts with the findings of several studies on diverse marine taxa (e.g., bivalves, gastropods, echinoderms), which showed higher feeding and respiration rates as the temperature increases (Cáceres-Puig,




FIGURE 2 Variations in ^{13}C (a) and ^{15}N (b) uptake ($I^{13}\text{C}$ and $I^{15}\text{N}$, respectively; mean \pm SE) in *Psammechinus miliaris* maintained for 6, 15 and 30 days at 17°C (black dots) and at 20°C (white dots) and supplied with labeled *Solieria chordalis* ($n = 4$)

Abasolo-Pacheco, Mazón-Suastegui, Maeda-Martínez, & Saucedo, 2007; Carr & Bruno, 2013; Gooding et al., 2009; Noisette et al., 2014). However, a previous experiment on the high-latitude urchin *Strongylocentrotus droebachiensis* evidenced a decrease in feeding rates above 15°C, which is close to the maximum summer temperature (16°C, Frey & Gagnon, 2015). Similarly in the lobster *Jasus edwardsii*, lower feeding and respiratory rates were observed above the upper thermal limit at 24°C (Thomas et al., 2000).

Interestingly, ¹³C and ¹⁵N uptake responded differently during the experiment. Indeed, conversely to ¹³C uptake, no effect of temperature increase was observed on ¹⁵N uptake in *P. miliaris*. At the end of this feeding experiment, ¹⁵N incorporation was 1.2 ng ¹⁵N day⁻¹ at 17°C and 0.8 ng ¹⁵N day⁻¹ at 20°C, which corresponded to estimated total nitrogen fluxes of 17.8 and 13.2 ng N g DW⁻¹ day⁻¹, respectively. This suggests an effect of temperature increase on carbon and nitrogen stoichiometry in *P. miliaris*. Although heterotrophs are considered as more homeostatic than autotrophs (Persson et al., 2010), several biological and environmental factors have been evidenced to influence stoichiometric homeostasis. Some studies have shown that changes in elemental composition of food sources may induce imbalance in consumer elemental homeostasis (Malzahn & Boersma, 2012; Prado, Heck, & Cebrian, 2014). Moreover, in the bacteria *Pseudomonas fluorescens*, Chrzanowski and Grover (2008) demonstrated that temperature can influence their elemental content regardless of their resource nutrient content, which appears consistent with the present results. This imbalance in carbon and nitrogen fluxes in *P. miliaris* between 17 and 20°C may have major implications for physiological processes, such as growth and reproduction (Frost, Xenopoulos, & Larson, 2004; Heflin et al., 2012).

Some studies have shown that temperature increase may alter macroalgal palatability to grazers (Poore et al.,), through changes in nutritional qualities or levels of deterrent secondary metabolites (Staehr & Wernberg, 2009; Sudatti, Fujii, Rodrigues, Turra, & Pereira, 2011). In this context, further labeling experiments should be conducted to examine the trophic transfer from macroalgae to grazers at different temperatures. In addition, these results highlight that quantitative estimation of trophic transfers between the two first trophic levels should not be based solely on carbon fluxes as is often the case.

In conclusion, the use of dual stable isotope labeling shows promise for testing the influence of various environmental factors on major macroalga-grazer trophic interactions within marine habitats, including their response to climate change. Further similar experimental studies should also couple stable isotope data related to assimilated nutrients with other physiological processes associated with feeding, such as ingestion, excretion or respiration.

ACKNOWLEDGEMENTS

The authors thank the Center of Marine Biologic Resources at the Station Biologique de Roscoff for its kind permission to use their premises for the duration of the experiments. We are grateful to Olivier Bohner and Jérôme Coudret for their field and laboratory

assistance and for the help in system maintenance. We acknowledge the crew of the research vessel *Albert Lucas* for their help with species collection. We also thank the SOMLIT (Institut National des Sciences de l'Univers - Centre National de la Recherche Scientifique) program for the temperature data sets provided. This work was supported by the Brittany Regional Council, the French National Research Agency via the 'Investment for the Future' program IDEALG (no. ANR-10-BTBR-04), and by the French national EC2CO program ('Écosphère Continentale et Côtière', project MAERLCHANGE).

ORCID

Erwann Legrand  <http://orcid.org/0000-0001-5224-5227>

REFERENCES

- Basuyaux, O., & Mathieu, M. (1999). Inorganic nitrogen and its effect on growth of the abalone *Haliotis tuberculata* Linnaeus and the sea urchin *Paracentrotus lividus* Lamarck. *Aquaculture*, 174, 95–107. [https://doi.org/10.1016/S0044-8486\(98\)00510-9](https://doi.org/10.1016/S0044-8486(98)00510-9)
- Bedford, A. P., & Moore, P. G. (1985). Macrofaunal involvement in the sublittoral decay of kelp debris: The sea urchin *Psammechinus miliaris* (Gmelin) (Echinodermata: Echinoidea). *Estuarine, Coastal and Shelf Science*, 20, 19–40. [https://doi.org/10.1016/0272-7714\(85\)90116-7](https://doi.org/10.1016/0272-7714(85)90116-7)
- Brockington, S., & Peck, L. S. (2001). Seasonality of respiration and ammonium excretion in the Antarctic echinoid *Sterechinus neumayeri*. *Marine Ecology Progress Series*, 219, 159–168. <https://doi.org/10.3354/meps219159>
- Cáceres-Puig, J. I., Abasolo-Pacheco, F., Mazón-Suastegui, J. M., Maeda-Martínez, A. N., & Saucedo, P. E. (2007). Effect of temperature on growth and survival of *Crassostrea corteziensis* spat during late-nursery culturing at the hatchery. *Aquaculture*, 272, 417–422. <https://doi.org/10.1016/j.aquaculture.2007.06.030>
- Carr, L. A., & Bruno, J. F. (2013). Warming increases the top-down effects and metabolism of a subtidal herbivore. *PeerJ*, 1, e109. <https://doi.org/10.7717/peerj.109>
- Chrzanowski, T. H., & Grover, J. P. (2008). Element content of *Pseudomonas fluorescens* varies with growth rate and temperature: A replicated chemostat study addressing ecological stoichiometry. *Limnology and Oceanography*, 53, 1242–1251. <https://doi.org/10.4319/lo.2008.53.4.1242>
- Frey, D. L., & Gagnon, P. (2015). Thermal and hydrodynamic environments mediate individual and aggregative feeding of a functionally important omnivore in reef communities. *PLoS ONE*, 10, e0118583. <https://doi.org/10.1371/journal.pone.0118583>
- Frost, P. C., Xenopoulos, M. A., & Larson, J. H. (2004). The stoichiometry of dissolved organic carbon, nitrogen, and phosphorus release by a planktonic grazer, *Daphnia*. *Limnology and Oceanography*, 49, 1802–1808. <https://doi.org/10.4319/lo.2004.49.5.1802>
- Gooding, R. A., Harley, C. D. G., & Tang, E. (2009). Elevated water temperature and carbon dioxide concentration increase the growth of a keystone echinoderm. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 9316–9321. <https://doi.org/10.1073/pnas.0811143106>
- Guillou, M., Grall, J., & Connan, S. (2002). Can low sea urchin densities control macro-epiphytic biomass in a north-east Atlantic maerl bed ecosystem (Bay of Brest, Brittany, France)? *Journal of the Marine Biological Association of the United Kingdom*, 82, 867–876. <https://doi.org/10.1017/S0025315402006276>
- Heflin, L. E., Gibbs, V. K., Powell, M. L., Makowsky, R., Lawrence, A. L., & Lawrence, J. M. (2012). Effect of diet quality on nutrient allocation to the test and Aristotle's lantern in the sea urchin *Lytechinus*

- variegatus* (Lamarck, 1816). *Journal of Shellfish Research*, 31, 867–874. <https://doi.org/10.2983/035.031.0335>
- Herman, P. M. J., Middelburg, J. J., Widdows, J., Lucas, C. H., & Heip, C. H. R. (2000). Stable isotopes' as trophic tracers: combining field sampling and manipulative labelling of food resources for macrobenthos. *Marine Ecology Progress Series*, 204, 79–92. <https://doi.org/10.3354/meps204079>
- Hily, C., Potin, P., & Floch, J. Y. (1992). Structure of subtidal algal assemblages on soft-bottom sediments: fauna/flora interactions and role of disturbances in the Bay of Brest, France. *Marine Ecology Progress Series*, 85, 115–130. <https://doi.org/10.3354/meps085115>
- Kamp, A., & Witte, U. (2005). Processing of ^{13}C -labelled phytoplankton in a fine-grained sandy-shelf sediment (North Sea): relative importance of different macrofauna species. *Marine Ecology Progress Series*, 297, 61–70. <https://doi.org/10.3354/meps297061>
- Leroy, F., Riera, P., Jeanthon, C., Edmond, F., Leroux, C., & Comtet, T. (2012). Importance of bacterivory and preferential selection toward diatoms in larvae of *Crepidula fornicata* (L.) assessed by a dual stable isotope (^{13}C , ^{15}N) labeling approach. *Journal of Sea Research*, 70, 23–31. <https://doi.org/10.1016/j.seares.2012.02.006>
- Malzahn, A. M., & Boersma, M. (2012). Effects of poor food quality on copepod growth are dose dependent and non-reversible. *Oikos*, 121, 1408–1416. <https://doi.org/10.1111/j.1600-0706.2011.20186.x>
- Martin, S., Castets, M. D., & Clavier, J. (2006). Primary production, respiration and calcification of the temperate free-living coralline alga *Lithothamnion corallioides*. *Aquatic Botany*, 85, 121–128. <https://doi.org/10.1016/j.aquabot.2006.02.005>
- Martin, S., Clavier, J., Chauvaud, L., & Thouzeau, G. (2007). Community metabolism in temperate maerl beds. I. Carbon and carbonate fluxes. *Marine Ecology Progress Series*, 335, 19–29. <https://doi.org/10.3354/meps335019>
- Middelburg, J. J., Barranguet, C., Boschker, H. T. S., Herman, P. M. J., Moens, T., & Heip, C. H. R. (2000). The fate of intertidal microphytobenthos carbon: An in situ ^{13}C -labeling study. *Limnology and Oceanography*, 45, 1224–1234. <https://doi.org/10.4319/lo.2000.45.6.1224>
- Minagawa, M., & Wada, E. (1984). Stepwise enrichment of ^{15}N along food chains: Further evidence and the relation between $\delta^{15}\text{N}$ and animal age. *Geochimica et Cosmochimica Acta*, 48, 1135–1140. [https://doi.org/10.1016/0016-7037\(84\)90204-7](https://doi.org/10.1016/0016-7037(84)90204-7)
- Nguyen, K. D. T., Morley, S. A., Lai, C.-H., Clark, M. S., Tan, K. S., Bates, A. E., & Peck, L. S. (2011). Upper temperature limits of tropical marine ectotherms: Global warming implications. *PLoS ONE*, 6, e29340. <https://doi.org/10.1371/journal.pone.0029340>
- Noisette, F., Richard, J., Le Fur, I., Peck, L. S., Davoult, D., & Martin, S. (2014). Metabolic responses to temperature stress under elevated pCO_2 in *Crepidula fornicata*. *Journal of Molluscan Studies*, 81, 238–246. <https://doi.org/10.1093/mollus/eyu084>
- Otero-Villanueva, M. d. M., Kelly, M. S., Burnell, G. (2004). How diet influences energy partitioning in the regular echinoid *Psammechinus miliaris*: constructing an energy budget. *Journal of Experimental Marine Biology and Ecology*, 304, 159–181. <https://doi.org/10.1016/j.jembe.2003.12.002>
- Persson, J., Fink, P., Goto, A., Hood, J. M., Jonas, J., & Kato, S. (2010). To be or not to be what you eat: regulation of stoichiometric homeostasis among autotrophs and heterotrophs. *Oikos*, 119, 741–751. <https://doi.org/10.1111/j.1600-0706.2009.18545.x>
- Poore, A. G. B., Graba-Landry, A., Favret, M., Brennand, H. S., Byrne, M., & Dworjanyn, S. A. (2013). Direct and indirect effects of ocean acidification and warming on a marine plant-herbivore interaction. *Oecologia*, 173, 1113–1124. <https://doi.org/10.1007/s00442-013-2683-y>
- Poore, A. G. B., Graham, S. E., Byrne, M., & Dworjanyn, S. A. (2016). Effects of ocean warming and lowered pH on algal growth and palatability to a grazing gastropod. *Marine Biology*, 163, 1–11. <https://doi.org/10.1007/s00227-016-2878-y>
- Prado, P., Heck, K. L., & Cebrian, J. (2014). Moderate stoichiometric homeostasis in the sea urchin *Lytechinus variegatus*: effects of diet and growth on C:N: P ratios. *Marine Biology*, 161, 2869–2883. <https://doi.org/10.1007/s00227-014-2552-1>
- R Core Team (2014). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <https://www.R-project.org/>.
- Rau, G. H., Mearns, A. J., Young, D. R., Olson, R. J., Schafer, H. A., & Kaplan, I. R. (1983). Animal $^{13}\text{C}/^{12}\text{C}$ correlates with trophic level in pelagic food webs. *Ecology*, 64, 1314–1318. <https://doi.org/10.2307/1937843>
- Rubilar, T., Epherra, L., Deias-Spreng, J., Díaz De Vivar, M. E., Avaro, M., Lawrence, A. L., & Lawrence, J. M. (2016). Ingestion, absorption and assimilation efficiencies, and production in the sea urchin *Arbacia dufresnii* fed a formulated feed. *Journal of Shellfish Research*, 35, 1083–1093. <https://doi.org/10.2983/035.035.0431>
- Staehr, P. A., & Wernberg, T. (2009). Physiological responses of *Ecklonia radiata* (Laminariales) to a latitudinal gradient in ocean temperature. *Journal of Phycology*, 45, 91–99. <https://doi.org/10.1111/j.1529-8817.2008.00635.x>
- Steneck, R. S. (2013). Sea urchins as drivers of shallow benthic marine community structure. In J. M. Lawrence (Ed.), *Developments in aquaculture and fisheries science. Sea urchins: Biology and ecology*, 3rd ed. (pp. 195–212). San Diego, CA: Academic Press.
- Sudatti, D. B., Fujii, M. T., Rodrigues, S. V., Turra, A., & Pereira, R. C. (2011). Effects of abiotic factors on growth and chemical defenses in cultivated clones of *Laurencia dendroidea* J. Agardh (Ceramiales, Rhodophyta). *Marine Biology*, 158, 1439–1446. <https://doi.org/10.1007/s00227-011-1660-4>
- Thomas, C. W., Crear, B. J., & Hart, P. R. (2000). The effect of temperature on survival, growth, feeding and metabolic activity of the southern rock lobster, *Jasus edwardsii*. *Aquaculture*, 185, 73–84. [https://doi.org/10.1016/S0044-8486\(99\)00341-5](https://doi.org/10.1016/S0044-8486(99)00341-5)
- Uthicke, S., Liddy, M., Nguyen, H. D., & Byrne, M. (2014). Interactive effects of near-future temperature increase and ocean acidification on physiology and gonad development in adult Pacific sea urchin, *Echinometra* sp. A. *Coral Reefs*, 33, 831–845. <https://doi.org/10.1007/s00338-014-1165-y>
- Zuo, W., Moses, M. E., West, G. B., Hou, C., & Brown, J. H. (2012). A general model for effects of temperature on ectotherm ontogenetic growth and development. *Proceedings of the Royal Society B: Biological Sciences*, 279, 1840–1846. <https://doi.org/10.1098/rspb.2011.2000>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.