



HAL
open science

Adult-onset Still's disease biological treatment strategy may depend on the phenotypic dichotomy

François Vercausse, Thomas Barnetche, Estibaliz Lazaro, Emilie Shipley,
François Lifermann, Alexandre Balageas, Xavier Delbrel, Bruno Fautrel,
Christophe Richez, Thierry Schaeffer, et al.

► **To cite this version:**

François Vercausse, Thomas Barnetche, Estibaliz Lazaro, Emilie Shipley, François Lifermann, et al..
Adult-onset Still's disease biological treatment strategy may depend on the phenotypic dichotomy.
Arthritis Research and Therapy, 2019, 21, pp.53. 10.1186/s13075-019-1838-6 . hal-02050492

HAL Id: hal-02050492

<https://hal.sorbonne-universite.fr/hal-02050492>

Submitted on 27 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Open Access



Adult-onset Still's disease biological treatment strategy may depend on the phenotypic dichotomy

François Vercruyse^{1*} , Thomas Barnetche¹, Estibaliz Lazaro², Emilie Shipley³, François Lifermann⁴, Alexandre Balageas⁵, Xavier Delbrel⁶, Bruno Fautrel⁷, Christophe Richez¹, Thierry Schaeffer¹ and Marie-Elise Truchetet¹

Abstract

Objectives: Adult-onset Still's disease (AOSD) phenotype appears to be dichotomized in systemic or chronic articular forms. As biologicals and particularly interleukin (IL)-1 and IL-6 blockers play a more and more prominent role in the treatment, their place requires clarification. This study aimed to identify factors predictive of treatment response to anakinra or tocilizumab and investigate whether the choice of biotherapy and delays in the initiation of biotherapy influenced the likelihood of steroid discontinuation.

Methods: A multicenter exploratory retrospective study included all patients diagnosed with AOSD and receiving biological treatments in three regional hospitals until 2018. Clinical and biological characteristics at diagnosis and treatment-related data were collected. The nonparametric Mann-Whitney test was used to perform univariate analysis for quantitative variables, and Fisher's exact test was used for qualitative variables.

Results: Twenty-seven patients were included. All but one patient achieved remission with either anakinra or tocilizumab. Treatment responses depended on disease phenotype: the presence of arthritis and a chronic articular phenotype were associated with a substantial response to tocilizumab with $p = 0.0009$ (OR 36 [2.6–1703]) and $p = 0.017$ (OR 10 [1.22–92.6]), respectively, whereas the systemic form and the absence of arthritis were associated with a substantial response to anakinra with $p = 0.0009$ (OR 36 [2.6–1703]) and $p = 0.017$ (OR 10 [1.22–92.6]), respectively. Tocilizumab increased the likelihood of corticosteroid withdrawal ($p = 0.029$) regardless of delays in initiation or when it was initiated relative to other treatment in the overall therapeutic strategy.

Conclusion: This study highlights the therapeutic implications of the phenotypic dichotomy of AOSD and should help us better codify AOSD treatment.

Keywords: Adult-onset Still's disease, Tocilizumab, Anakinra, Treatment strategy, Biologics

Key messages

- What is already known about this subject?
 - The phenotype of adult-onset Still's disease appears to be dichotomized into systemic or chronic articular forms.
- Biological treatments are dominated by IL-1 or IL-6 receptor inhibitors.
- What are the new lines of research suggested by this study?
 - The presence of arthritis and a chronic articular phenotype may be associated with a preferential response to tocilizumab, whereas the systemic form may be associated with a preferential response to anakinra.
 - Tocilizumab increases the likelihood of corticosteroid withdrawal regardless of delays in its initiation or when it is initiated relative

* Correspondence: francois.vercruyse@chu-bordeaux.fr

¹Rheumatology Department, Centre Hospitalier Universitaire de Bordeaux, FHU ACRONIM, Service de Rhumatologie, Place Amélie Raba Léon, 33076 Bordeaux, France

Full list of author information is available at the end of the article



to other treatment in the overall therapeutic strategy

- How might this impact clinical practice?
 - These results should help us better codify the treatment strategy for adult-onset Still's disease.

Introduction

Adult-onset Still's disease (AOSD) is a rare systemic inflammatory disorder. Its annual incidence is estimated to be between 0.16 and 0.62 per 100,000 persons globally [1], and its prevalence approximately 1–24 per million [2].

AOSD is an autoinflammatory disease resulting from multifactorial causes such as genetic factors and immunological imbalances involving macrophage, neutrophil, and inflammasome hyperactivation or dysfunction [3].

Recent studies suggest that there is a dichotomy between two major phenotypes of AOSD: a systemic pattern that presents with an acute onset associating high fever, skin rash, hematopoietic signs, and sometimes visceral damage; and a chronic articular profile in which polyarthritis prevails [4, 5]. This phenotypic dichotomy is highlighted by recent studies showing that there are different cytokine imbalances in these two phenotypes. Although results may vary, this imbalance would sometimes favor interleukin (IL)-1 β , IL-18, and interferon (IFN)- γ production in the systemic form, whereas in the chronic articular form, IL-6, IL-8, tumor necrosis factor (TNF)- α , and IL-17 production would predominate [6–12].

The prognosis of AOSD is unpredictable as it can manifest with polyarticular erosive disease or life-threatening manifestations such as reactive hemophagocytic lymphohistiocytosis (rHLH). The disease course is also very polymorphic, ranging from a monophasic flare to polycyclic or chronic forms. This clinical and prognostic polymorphism presents a major challenge and makes it difficult to predict the outcome at the time of diagnosis and to determine appropriate management.

Therapy is therefore usually empirical, and beside methotrexate, biologics have played an increasing role in the chronic or refractory forms. A recent meta-analysis suggested that IL-1 and IL-6 receptor inhibitors are the most effective biotherapies for AOSD [13] and have a strong steroid-sparing effect. The systemic phenotype has been suggested to respond preferentially to IL-1 receptor inhibitors, such as anakinra, while the chronic articular form shows better responses to IL-6 receptor inhibitors, such as tocilizumab [6, 7, 14–17].

This exploratory study therefore aimed to identify initial clinical and/or biological factors that were predictive of a preferential response to anakinra or tocilizumab. Secondary objectives were to investigate whether the choice of biotherapy (anakinra or tocilizumab) and their delay in initiation influenced the likelihood of steroid

discontinuation and to describe the clinical and biological phenotypes of patients with AOSD.

Methods

Study design and data collection

A multicenter exploratory retrospective study of all patients diagnosed with AOSD at Bordeaux teaching hospital and Pau and Dax peripheral hospitals until 2018 was conducted. Patients included had to be 16 years old or older, to be diagnosed with AOSD, to meet the Yamaguchi and/or Fautrel criteria, to have a chronic or refractory disease course, and to receive or have received a biological disease-modifying antirheumatic drugs (bDMARD). Exclusion criteria were age younger than 16 years old, a diagnosis of systemic-onset juvenile idiopathic arthritis (SoJIA), an exclusion of AOSD diagnosis during disease course, Yamaguchi and Fautrel criteria not met, lack of data, and absence of bDMARD during disease course.

Patients with AOSD were identified via ICD-10 code, via a diagnostic grid that automatically references autoimmune and inflammatory diseases at Bordeaux University Hospital, via the bioinformatics services of each hospital, and via direct contact with the specialists treating the patients. Data regarding clinical and biological characteristics at diagnosis and treatment details were collected using a standard data extraction form. All the information was double-checked for any discrepancies.

The following demographic and clinical AOSD-related characteristics were collected: sex, date of birth, age at diagnosis, year of diagnosis, duration of disease, duration of disease at first bDMARD initiation, smoking habits, personal or familial history of rheumatic autoimmune inflammatory or neoplastic disease, type of Still's disease-related rash if present, arthralgia, arthritis, hands affected or not, early morning stiffness, myalgia, fever characteristics, general deterioration and weight loss > 4 kg, lymphadenopathy, pharyngitis, angina, and visceral involvement (pleuritis, pericarditis, pulmonary hypertension).

The following biological characteristics were recorded: leukocytosis > 10,000/mm³, neutrophilia > 8000/mm³, eosinophilia > 500/mm³, anemia with hemoglobin < 12 g/dl, thrombocytosis > 400,000/mm³, increased C-reactive protein (CRP) levels, maximum CRP at diagnosis, decreased albumin, increased liver enzymes, increased creatine kinase, increased lactate dehydrogenase (LDH), increased ferritin, decreased glycosylated ferritin, autoimmunity markers (antinuclear antibodies (ANA), antideoxyribonucleic acid, anti-extractable nuclear antigen, rheumatoid factor, anti β 2GP1, lupus anticoagulant, anticardiolipin), disseminated intravascular coagulation markers, and rHLH markers (cytopenia, hypertriglyceridemia, hypofibrinogenemia, hemophagocytosis in the bone marrow).

Patients were classified as having two distinct disease patterns: the systemic form or the chronic articular form. In the absence of consensual criteria in this regard, this classification was based on expert opinions that was often spelled out in the medical reports and was confronted to disease course outcome, which was available due to the retrospective nature of this study.

Outcome measurements

The effectiveness of treatment was defined as follows: effective treatment was considered when all initial symptoms and biological anomalies had resolved, partially effective treatment was considered when all but one initial symptom or biological anomaly had resolved, and ineffective treatment was considered when two or more initial symptoms or biological anomaly persisted. The attending doctors assessed disease flares and relapses.

Treatment lines were assessed as follows: nonsteroidal anti-inflammatories (NSAIDs) were not considered a treatment modality for chronic and refractory AOSD, but corticosteroids, conventional synthetic disease-modifying antirheumatic drugs (csDMARD) and especially methotrexate (both galenic forms taken together) and cyclosporine, and bDMARD (anakinra, tocilizumab, canakinumab, infliximab, and etanercept) were considered treatment lines.

The outcome of interest for the univariate analysis of predictive factors was the full effectiveness of the treatment as defined above, including both systemic and articular manifestations.

Statistical analysis

Quantitative variables are presented as mean with standard deviation or as median with interquartile range, and qualitative variables are presented as percentages. Analysis was performed on the following variables: all the clinical and biological manifestations included in the Yamaguchi or Fautrel criteria, common demographic characteristics, and all the symptoms or biological abnormalities usually observed during AOSD according to the literature and our personal experience. The nonparametric Mann-Whitney test was used to perform a univariate analysis of quantitative variables, and the Fisher exact test was used for qualitative variables. Calculations were performed with STATA 13.1 SE software. *P* values less than 0.05 were considered significant.

Results

Patients' characteristics

Twenty-seven patients were included in this study: 21 from Bordeaux teaching hospital, 4 from Dax hospital, and 2 from Pau hospital. Patient characteristics at diagnosis are described in Table 1.

All patients fulfilled both the Yamaguchi and Fautrel criteria. No patients had any personal or familial history

of rheumatic, autoimmune, inflammatory, or neoplastic disease. All patients presented with a fever above 38 °C, mainly intermittent and vesperal fevers. The next most prevalent symptoms were arthralgia (96%), arthritis (52%), skin rash (81%), general deterioration (72%), and ear-nose-throat (ENT) symptoms (69%).

Eleven (40.7%) patients were classified as systemic form of disease, and 16 (59.3%) as chronic articular form.

All patients presented with elevated CRP levels. The next most commonly observed anomalies were neutrophilia (96%), low albumin (75%), and increased ferritin (73%). No patients presented with increased creatine kinase (CK), positive extractable nuclear antigen (ENA), positive antineutrophil cytoplasmic antibody (ANCA), positive anticitrullinated protein antibody (ACPA), or positive anticardiolipin. Nine patients were either ANA-positive (five, 1/160), anti-DNA positive (one), or rheumatoid factor (RF)-positive (three) at diagnosis, but these results were judged clinically insignificant because lower than two times the upper normal range and never associated with symptoms concordant with SLE, APLS, or RA and were all controlled to be negative. Furthermore, all patients included in the study met both Yamaguchi and Fautrel criteria.

The median duration of disease was 5 years (IQR 2–7) and 6.5 years (IQR 5–9) for the systemic and chronic articular forms, respectively.

All patients presented a chronic refractory form of the disease, which required the off-label use of bDMARD because of failures of corticosteroids and csDMARD or necessity to reduce or stop corticosteroids.

Biological treatment efficacy

Table 2 presents the detailed responses to anakinra and tocilizumab among the patients included in the study. Twenty-six of 27 patients (96.3%) achieved remission with either anakinra or tocilizumab. The median delay in the introduction of the first biotherapy was 5 months (interquartile range (IQR), 1–21 months): this delay was median 1 month (IQR, 0–6) in the systemic form and median 17 months (IQR, 4–38) in the chronic articular form.

A total of 15 patients received anakinra; 5 as a first-line treatment, 4 as a second-line treatment, 5 as a third-line treatment, and 1 as a fourth-line treatment. The median delay between diagnosis and the introduction of anakinra was 1.5 months (IQR, 0–14). Thirteen (86.7%) patients were responders. The 2 anakinra nonresponders received anakinra as a third-line treatment: one had previously failed to respond to tocilizumab, and the other later responded to tocilizumab and even managed to stop it.

Table 1 Characteristics of the patients included in the study

n/N* (%)	Systemic form 11 (40.7)	Chronic articular form 16 (59.3)	Total 27
Demographic and clinical			
Sex : female	8 (73)	10 (63)	18 (67)
Age at first symptoms	33 (IQR 23-48)	29 (IQR 21-58.5)	30 (IQR 21-48)
Duration of disease (years)	5 (IQR 2-7)	6.5 (IQR 5-9)	6 (IQR 4-9)
Ever smoked	4 (36)	2/9 (22)	6/20 (30)
Still's disease-related rash	10 (91)	12 (75)	22 (81)
Evanescient erythema	4 (36)	8 (50)	12 (44)
Maculopapular rash	7 (64)	7 (44)	14 (52)
Urticaria rash	3 (27)	2 (13)	5 (18.5)
Arthralgia	10 (91)	16 (100)	26 (96)
Arthritis	3 (27)	11 (69)	14 (52)
Hands affected (arthralgia or arthritis)	9 (82)	14 (88)	23 (85)
EMS	7/8 (88)	13/14 (93)	20/22 (90)
Myalgia	9 (82)	8 (50)	17 (63)
Fever>38°C	11 (100)	16 (100)	27 (100)
Fever>39°C for longer than 7 days	4 (36)	4/15 (27)	8/26 (31)
Intermittent fever	9 (82)	10/15 (67)	19/26 (73)
Vesperal fever	8 (73)	9/15 (60)	17/23 (74)
Deterioration of the general state or weight loss>4kg	8 (73)	10/15 (67)	18/25 (72)
Lymphadenopathy	8 (73)	6/15 (40)	14/26 (54)
Pharyngitis	8 (73)	10/15 (67)	18/26 (69)
Angina	4 (36)	1/15 (7)	5/26 (19)
Visceral involvement	2 (18)	0 (0)	2 (8)
Pleuritis	1 (9)	1 (0)	1 (4)
Pericarditis	1 (9)	2 (0)	1 (4)
Pulmonary hypertension	1 (9)	3 (0)	1 (4)
Biological			
Neutrophilia>8000/mm3	11 (100)	14/15 (93)	25/26 (96)
Eosinophilia>500/mm3	1/10 (10)	0 (0)	1/25 (4)
Anemia<12 g/dl	6/10 (60)	9 (56)	15/26 (58)
Thrombocytosis>400 000/mm3	3/10 (30)	10 (63)	13/26 (50)
Increased C-reactive protein (CRP) levels	11 (100)	16 (100)	27 (100)
Maximum CRP at diagnosis	200 (IQR 160-300)	200 (IQR 67.5-254.5)	200 (IQR 104.5-267.5)
Low albumin	9/10 (90)	9/14 (64)	18/24 (75)
Increased AST or ALT	8 (73)	5/15 (33)	13/26 (50)
Increased ferritin	10 (91)	9/15 (60)	19/26 (73)
Ferritin 1000-5000	4 (36)	6/15 (40)	10/26 (38.5)
Ferritin 5000-10000	2 (18)	1/15 (7)	3/26 (11.5)
Ferritin>10 000	4 (36)	2/15 (13)	6/26 (23)
Glycosylated ferritin<20%	8 (73)	10 (63)	18 (66.7)
Glycosylated ferritin 20-40%	2 (18)	6 (38)	8 (29.6)
Glycosylated ferritin>40%	1 (9)	0 (0)	1 (3.7)
ANA-positivity	2 (18)	3 (19)	5 (18.5)
anti-DNA positivity	0 (0)	1 (6)	1 (3.7)
RF positivity	0 (0)	3 (19)	3 (11)
β2GP1 positivity	1/5 (20)	0 (0)	1/5 (20)
Lupus anticoagulant positivity	1/5 (20)	0 (0)	1/5 (20)
Disseminated intravascular coagulation	0 (0)	0 (0)	0 (0)
rHLH**	3 (27)	0 (0)	3 (11)
Increased triglycerids	5/7 (71)	1/10 (10)	6/17 (35)
Hemophagocytosis	3 (27)	0/5 (0)	3/8 (37.5)

*N is indicated when data was not available for all 27 patients

** reactive haemophagocytic lymphohistiocytosis

A total of 17 patients received tocilizumab, 2 as a first-line treatment, 6 as a second-line treatment, 5 as a third-line treatment, and 4 as a fourth-line treatment. The median delay between diagnosis and the introduction of tocilizumab was 17 months (IQR, 4–39 months). Fourteen patients (82.4%) were responders. Among the 3 tocilizumab nonresponders, one presented with a systemic form and received tocilizumab as a first-line treatment but later responded to anakinra, one presented with a systemic form and received tocilizumab as a fourth-line treatment after responding to anakinra but was switched because of a suspicion of anakinra-induced toxidermia, and one presented with a chronic articular

form and received tocilizumab as a fourth-line treatment after failure of anakinra.

The difference of median duration of disease at the introduction of the first bDMARD in the anakinra-treated patients group and in the tocilizumab-treated patients group was statistically significant ($p = 0.4457$): median 2 months (IQR, 0.25–12 months) in the anakinra-treated patients and median 14.5 (IQR, 3.25–38.25 months) in the tocilizumab-treated patients.

Eight patients (29.6%) managed to stop their biological treatment without relapse at the last known visit which ranged from 6 to more than 24 months after biological treatment discontinuation: 3 of 13 (23%) anakinra responders

Table 2 Delay in corticosteroids withdrawal and detailed responses to anakinra and tocilizumab among the patients included in the study

STILL	Phenotype	Corticosteroids			Anakinra-treated patients				Tocilizumab-treated patients			
		Stopped	Delay between diagnosis and steroid withdrawal (months)	Delay between diagnosis and initiation of first bDMARD (months)	Delay between diagnosis and initiation of anakinra (months)	Managed to		Line of treatment	Delay between diagnosis and initiation of tocilizumab (months)	Tocilizumab responders	Managed to	
						stop anakinra	stop anakinra				stop tocilizumab	Line of treatment
STILL 1	Chronic articular	Yes	12	39	-	-	-	-	39	Yes	No	Third
STILL 2	Systemic	No	-	0	0	Yes	Yes	*	9	No	-	Fourth
STILL 3	Systemic	No	-	1	1	Yes	No	First	-	-	-	-
STILL 4	Chronic articular	Yes	60	38	-	-	-	-	38	Yes	No	Second
STILL 5	Chronic articular	Yes	17	7	-	-	-	-	7	Yes	No	Second
STILL 6	Chronic articular	Yes	71	69	-	-	-	-	69	Yes	No	Second
STILL 7	Systemic	Yes	12	2	2	Yes	No	Second	20	Yes	Yes	Third
STILL 8	Systemic	Yes	20	12	-	-	-	-	12	Yes	No	Third
STILL 9	Chronic articular	No	-	21	-	-	-	-	21	Yes	No	Third
STILL 10	Chronic articular	No	-	4	4	No	-	Third and fifth	17	No	-	Fourth
STILL 11	Systemic	Yes	12	0	0	Yes	Yes	First and third	-	-	-	-
STILL 12	Chronic articular	No	-	**	-	-	-	-	**	Yes	No	Third
STILL 13	Chronic articular	Yes	45	24	24	No	-	Third	36	Yes	Yes	Fourth
STILL 14	Chronic articular	Yes	18	4	-	-	-	-	4	Yes	Yes	Second
STILL 15	Chronic articular	Yes	8	1	-	-	-	-	1	Yes	Yes	Second
STILL 16	Chronic articular	No	-	14	14	Yes	No	Fourth	-	-	-	-
STILL 17	Chronic articular	Yes	29	17	-	-	-	-	17	Yes	Yes	Fourth
STILL 18	Systemic	No	-	6	6	Yes	No	Third	-	-	-	-
STILL 19	Chronic articular	Yes	15	0	-	-	-	-	0	Yes	No	First
STILL 20	Systemic	No	-	0	0	Yes	Yes	First	-	-	-	-
STILL 21	Chronic articular	No	-	252	-	-	-	-	252	Yes	No	Second
STILL 22	Systemic	No	-	0	0	Yes	No	First	-	-	-	-
STILL 23	Systemic	No	-	0	8	Yes	No	Second	0	No	-	First
STILL 24	Chronic articular	No	-	1	1	Yes	No	Second	-	-	-	-
STILL 25	Chronic articular	No	-	24	24	Yes	No	Third	-	-	-	-
STILL 26	Systemic	Yes	6	14	14	Yes	No	Third	-	-	-	-
STILL 27	Systemic	Yes	53	2	2	Yes	No	Second	-	-	-	-

* First, third, fifth and seventh
** Unknown

and 5 of 14 (35.7%) tocilizumab responders. The difference in delays in initiation of anakinra or tocilizumab was not statistically significant between those who managed to stop their biotherapy without relapse versus those who could not stop their biotherapy or those who relapsed, with a median delay of 3 months (IQR, 1.25–13.75) versus 8.5 months (IQR, 1–23.25), respectively ($p = 0.61$).

Five patients received both biotherapies. Two of these patients presented with a systemic form and were tocilizumab nonresponders but anakinra responders, one presented with a chronic articular form and was anakinra nonresponder but tocilizumab responder, one presented with a chronic articular form and did not respond to either biotherapy, and one patient presented with a systemic form and responded to both biotherapies; this patient first responded to anakinra but corticosteroids could not be tapered so he was switched to tocilizumab in order to successfully stop corticosteroids.

One patient received canakinumab and failed to respond but was anakinra responder. Three patients received antitumor necrosis factor (TNF) agents but these treatments were not efficacious (2 etanercept, 1 infliximab): two of them presented a chronic articular form and one a systemic form.

One of 27 (3.7%) patients did not respond to either biotherapy (anakinra then tocilizumab) and died of a probably infectious-induced AOSD flare.

Predictive factors for preferential responses to anakinra or tocilizumab

Patients who responded to either biotherapies or no biotherapy were excluded from this analysis ($n = 2$). Table 3 describes the univariate analysis for potential predictive factors of a good therapeutic response to anakinra or tocilizumab. The outcome of interest was the full effectiveness of the treatment, defined as the resolution of all initial systemic and articular symptoms and biological anomalies.

A chronic articular phenotype and the presence of arthritis were associated with a substantial response to tocilizumab: 12 of the 14 patients presenting a chronic articular phenotype responded to tocilizumab whereas 3 of the 13 patients presenting a systemic phenotype responded to tocilizumab ($p = 0.0009$, OR 36 [2.6–1703]) and 10 of the 14 patients presenting arthritis responded to tocilizumab whereas 3 of the 13 patients without arthritis responded to tocilizumab ($p = 0.017$, OR 10 [1.22–92.6]).

A systemic form and the absence of arthritis were associated with a substantial response to anakinra: 1 of the 14 patients presenting a chronic articular phenotype responded to anakinra whereas 9 of the 13 patients presenting a systemic phenotype responded to anakinra ($p = 0.0009$, OR 36 [2.6–1703]) and 3 of the 14 patients presenting arthritis responded to anakinra whereas 9 of the 13 patients without arthritis responded to anakinra ($p = 0.017$, OR 10 [1.22–92.6]).

Table 3 Univariate analysis of predictive factors for a good therapeutic response to anakinra or tocilizumab ($n = 25$)

	Anakinra responders (n=12)	Tocilizumab responders (n=13)	p value
Clinical variables			
Sex, F	8	10	0.673
Age at first symptoms	42.5 (IQR 22.5-54)	25 (IQR 21-36)	0.276
Ever smoked	4	2	1.000
Never smoked	4	5	0.170
Still's disease-related rash			
Evanescient erythema	6	6	1.000
Maculopapular rash	6	7	1.000
Urticaria	4	1	0.160
Arthralgia	11	13	0.480
Arthritis	3	10	0.017*
Absence of arthritis	9	3	0.017*
Hands affected (arthralgia or arthritis)	10	11	1.000
Early morning stiffness	8	11	1.000
Myalgia	9	7	0.411
Fever>39°C for longer than 7 days	5	3	0.667
Intermittent fever	9	9	1.000
Vesperal fever	7	8	0.635
Deterioration of the general state and/or weight loss>4 kg	7	9	0.371
Adenopathy	6	6	1.000
Pharyngitis	9	8	1.000
Angina	4	1	0.317
Phenotypic variables			
Systemic form	9	1	0.0009**
Chronic articular form	3	12	0.0009**
Biological variables			
Anemia<12g/dl	6	8	1.000
Thrombocytosis>400 000/mm ³	3	8	0.123
Thrombopenia<150 000/mm ³	1	0	0.458
Low albumin	8	8	0.646
Ferritin			
5000-10 000 (µg/L)	1	2	1.000
>10 000	3	2	0.645
Glycosylated ferritin<20%	6	10	0.226
ANA positivity	1	4	0.593
Hemophagocytosis	3	0	1.000

* OR 10 [1.22-92.6]

** OR 36 [2.6-1703]

No difference in biological variables was statistically predictive of treatment response, although the presence of thrombocytosis seemed to be preferentially associated with a better response to tocilizumab than to anakinra ($n = 8/14$ vs. $n = 3/13$, $p = 0.123$).

Likelihood of steroid withdrawal depending on the specific biotherapy and delays in initiation of biotherapy

All patients received corticosteroids at diagnosis. Among the 26 patients in remission, 14 (53.8%) achieved corticosteroid withdrawal within a mean delay of 27 months, 10 (71.4%) were on tocilizumab, and 4 (28.6%) were on anakinra (Table 2).

Table 4 presents the likelihood of corticosteroid withdrawal depending on treatment with anakinra and tocilizumab: tocilizumab was associated with an increased likelihood of corticosteroid withdrawal ($p = 0.029$), whereas anakinra did not seem to increase the likelihood of corticosteroid withdrawal ($p = 1.000$).

A shorter delay before the initiation of anakinra or tocilizumab was not associated with a greater likelihood of corticosteroid withdrawal among biotherapy responders. Among anakinra responders, the difference in the

median delay from diagnosis to anakinra initiation between those who achieved steroid withdrawal ($n = 4$) and those who did not ($n = 10$) was not statistically significant ($p = 0.5767$); the median delay was 2.5 months (IQR, 0–8 months) among those who did not stop corticosteroids versus 2 months (IQR, 2–14 months) among those who stopped corticosteroids. Among tocilizumab responders, the difference in the median delay from diagnosis to tocilizumab initiation between those who achieved steroid withdrawal ($n = 11$) and those who did not ($n = 5$) was not statistically significant ($p = 0.9548$); the median delay was 17 months (IQR, 9–21 months) among the tocilizumab responders who did not stop corticosteroids versus 17 months (IQR, 4–38 months) among those who stopped corticosteroids. Initiating tocilizumab as the first-line treatment rather than as a second-line or later treatment did not improve the likelihood of corticosteroid withdrawal ($n = 1/2$ vs. $n = 11/15$, $p = 1.000$). In addition, initiating tocilizumab as a first- or second-line treatment rather than as a third-line or later treatment also did not improve the likelihood of steroid withdrawal ($n = 6/8$ vs. $n = 5/9$, $p = 0.619$); similarly, initiating tocilizumab as a first-, second- or third-line treatment

Table 4 Likelihood of corticosteroid withdrawal stratified by biotherapy

Anakinra responders	Corticosteroids stopped			p value
	No	Yes	Total	
No	1	1	2	1.000
Yes	9	4	13	
<i>Total</i>	10	5	15	
<hr/>				
Tocilizumab responders	Corticosteroids stopped			p value
	No	Yes	Total	
No	3	0	3	0.029
Yes	3	11	14	
<i>Total</i>	6	11	17	

did not improve the likelihood of steroid withdrawal compared to that from tocilizumab used as a fourth-line treatment ($n = 9/13$ vs. $n = 2/4$, $p = 0.584$).

Discussion

This retrospective study was conducted to evaluate bDMARD strategies in the management of AOSD, with a special attention to predictive factors of their efficacy and their effect on corticosteroid sparing. All but one patient achieved remission with either anakinra or tocilizumab. Our results support the dichotomy of AOSD in two phenotypes: treatment responses to bDMARD seemed to depend on disease phenotype, the presence of arthritis, and a chronic articular phenotype were associated with a substantial response to tocilizumab, whereas the systemic form was associated with a substantial response to anakinra. In our cohort, anakinra did not increase the likelihood of corticosteroid withdrawal but tocilizumab did, regardless of delays in initiation, or when it was initiated relative to other treatment in the overall therapeutic strategy.

The clinical and biological characteristics of the patients included in this study seem similar to those of AOSD patients reported in the literature [18], with the exception of life-threatening conditions, which have been estimated to affect as many as 15–20% of AOSD patients, while only 8% were affected in this study.

Multivariate analysis was not performed because of the sample size, a limitation inherent to rare diseases such as AOSD. As a consequence, we lacked patients who received both biotherapies in each phenotype. But rather than aiming to look for significant differences and to draw definite conclusions, the goal of this study was to establish a reference for future large-scale studies and compile data for registries to confirm the results.

The distinction between systemic and chronic articular forms is not always straightforward, and no consensual

criteria exist to distinguish these two phenotypes. For example, a systemic form progressing for several years, who received corticosteroids for a long time and who benefited from bDMARD late in disease course, might be confused for a chronic articular form.

These results highlight the excellent efficacy of anakinra and tocilizumab in AOSD, regardless of the phenotype, but show some differences between them. However, the retrospective nature of the study is a major limitation to compare these two drugs. The choice of biotherapy may have been different depending on their availability on the year of diagnosis: anakinra was launched on the market several years before tocilizumab and the effects of anakinra and tocilizumab on AOSD were not simultaneously described.

One could argue that clinicians may have influenced the choice of biotherapy by preferentially prescribing tocilizumab to patients presenting with arthritis or a chronic articular form, analogous to rheumatoid arthritis, and therefore distorted the results. Comparing these two bDMARDs is impossible outside a randomized controlled trial (RCT), and none ever demonstrated that tocilizumab should be preferentially prescribed to AOSD patients with arthritis [5, 19, 20].

Yet the following results reinforce our findings. All but three of the patients included in this study were diagnosed in 2009 or later, and these three received anakinra or tocilizumab after both their efficacies were described in AOSD.

A major result of our study is that anakinra responders more commonly did not have arthritis, as the absence of arthritis was statistically associated with a better response to anakinra ($p = 0.017$). Tocilizumab also showed less efficiency on the systemic form. Although our results cannot imply definitive conclusions on the fact that one biotherapy is more effective than the other in each phenotype because of the lack of patients who received

both biotherapy in each phenotype, this result supports our clinical feeling and the notion that even if clinicians' choices may have been influenced by analogy to rheumatoid arthritis, their insights seem to be founded.

While there is no doubt about anakinra's dramatic efficacy on AOSD in general and particularly on the systemic phenotype [2, 18, 19, 21–27], whether anakinra is effective for joint involvement is a subject of debate as data seem to be inconsistent [4, 21–24, 28, 29]. Some studies reported a substantial efficacy [5, 23, 24], when others suggest otherwise [21, 28, 30]. The largest retrospective study investigating anakinra in AOSD showed a significant improvement on articular manifestations (DAS28 falling from 4.7 ± 1.2 to 1.7 ± 0.9 after 12 months) but 14.4% of the patients still presented arthritis after 12 months of treatment [24]. Notably, they found no difference in delay of response to therapy between the two phenotypes and did not observe any difference in the type of response in the two patterns of the disease. In our study, 5 patients presenting with a chronic articular form received anakinra: 3 responded to anakinra and never received tocilizumab, one was an anakinra nonresponder but responded to tocilizumab, and one responded to anakinra but had to be switched to tocilizumab in order to stop the corticosteroid. Definitive conclusions about anakinra's efficacy on AOSD's joint involvement therefore seem still premature.

The absence of demonstrated steroid-sparing with anakinra in this study is questionable as it demonstrated a steroid-sparing effect in several other studies [22, 23, 25]. First, the evaluation in our study was based on the rate of steroid withdrawal, not on a steroid-sparing effect as prednisone dosage was not available. Secondly, a channeling bias exists: the acute onset of the systemic phenotype and its potential life-threatening complications might cause the clinicians to be more cautious while tapering corticosteroids, and the tocilizumab-induced fall in CRP might reassure them. Furthermore, anakinra was made available before tocilizumab: the progressively known dramatic efficacy of anakinra may have reassured the clinicians and enticed them into a faster corticosteroids tapering when tocilizumab was made available. In addition, three patients presenting a systemic form in our study were able to taper corticosteroids down to 5 mg daily but the clinicians opted to maintain this dosage as a long-term therapy. Both the management and the place of corticosteroids in AOSD treatment strategy should maybe differ based on the differences between the two phenotypes.

No patients in this study responded to anti-TNF blockers, whatever the phenotype. The efficacy of these medications in AOSD is controversial, but some studies suggest that they have a place in the treatment of the chronic articular form [31–33], while others have showed rather disappointing results [19, 20, 34]. The

excellent efficacy and clear steroid-sparing effect of tocilizumab determined in our study, concurring with the findings of a recent meta-analysis showing that AOSD patients treated with tocilizumab had a pooled remission rate of approximately 85% regardless of the phenotype [35], and of the first RCT investigating tocilizumab in AOSD [36], should make clinicians prioritize tocilizumab to anti-TNF blockers for patients with a chronic articular form of AOSD. Canakinumab showed some inconsistent results in AOSD [37–39] but is currently being investigated for its efficacy on AOSD's joint involvement (NCT02204293). No patient in this study received abatacept or rituximab; the data are very limited but some authors reported efficacy in patients who fail to respond to TNF, IL-1, or IL-6 inhibition [40–45].

Physiopathological data may highlight the efficacy of IL-17 or IL-18 blockade in AOSD. A recent study showed that IL-18 inhibition using the recombinant human IL-18-binding protein tadekinig alpha is a therapeutic option in patients with AOSD in a phase 2 open-label trial [46]. IL-17 [47] and INF- γ [48] blockade have not been studied yet but should be in the near future. Janus kinase (JAK) inhibitors may also be of interest, particularly in the chronic articular form of AOSD.

The two main goals when initiating a bDMARD in AOSD should be a complete remission and corticosteroids withdrawal or significant tapering.

This study does not reveal definitive answers regarding whether an early introduction of biologics improves the general prognosis of AOSD. We believe that the early use of biologics in AOSD patients is worth considering. It is our impression that the earlier biologics are used, the shorter and smoother the disease course is. Similarly, other authors have suggested that biotherapies should be initiated earlier in the course of AOSD [13, 18], arguing that earlier initiation may improve the prognosis mainly via a steroid-sparing effect. Notably, the results of this study suggest that the steroid-sparing effect of biologics is not influenced by delays in initiation of biological therapy.

One-quarter to one-third (29.6%) of our patients managed to stop biologics despite chronic or refractory forms of AOSD, raising the question of how and when to stop biologics in these patients. A channeling bias exists, as no predefined corticosteroids tapering strategy could be set due to the retrospective nature of this study but patients who managed to stop anakinra had all achieved remission for at least 6 to 12 months, had initially stopped corticosteroids, and then all underwent progressive reduction in their biological treatments before definitive discontinuation. Among patients who managed to stop tocilizumab, corticosteroids had always previously been stopped, and tocilizumab dosage was progressively decreased when it had been administered

by perfusion or tapered when it had been administered by injections. The high percentage of patients who successfully discontinued biological treatments in our study should encourage clinicians to consider tapering biologics among patients without visceral involvement who are in clinical and biological remission for 6 to 12 months.

The first evidence-based clinical practice guidelines for the management of AOSD [49] emphasized the use of IL-6 inhibitors in the treatment of AOSD, but IL-1 inhibitors were judged to be “weakly recommended as a therapeutic agent against refractory AOSD” although one RCT and several case reports suggest otherwise. A proposal may be to treat a first flare of AOSD with corticosteroids and to introduce biologics in cases of steroid-refractory flares or as early as the second flare of AOSD. In that case, anakinra should be preferred when faced with the systemic form of AOSD and tocilizumab in patients with the articular form. The well-known idea of a “window of opportunity” in rheumatoid arthritis may also be valid in AOSD, with the objective being to prevent the development of a self-perpetuating “cytokine storm.”

We need large-scale studies, RCTs, and international collaborative registries regarding this multifaceted disease. As emphasized in a recent editorial, these studies will have to be more precise about evaluating the patients’ disease course and particularly their phenotypes in order to concentrate research on the subgroup of patients with a chronic or refractory form of the disease [50]. A national French cohort, the French Adult and Childhood Onset Still Disease Cohort (RaDiCo-ACOS-TILL), was recently opened for inclusion and will surely help to shed light on these issues in the near future.

Conclusion

This study supports the dichotomy of AOSD in two phenotypes and introduces the idea that treatment response to bDMARD may depend on disease phenotype. The presence of arthritis and a chronic articular phenotype seem to be associated with a substantial response to tocilizumab, whereas the systemic form seems to be associated with a substantial response to anakinra. Although further research is needed, this exploratory study highlights the therapeutic implications of the phenotypic dichotomy of AOSD. These results should help us better codify treatment strategies for AOSD.

Abbreviations

ANA: Antinuclear antibodies; ANCA: Antineutrophil cytoplasmic antibody; ACPA: Positive anticitrullinated protein antibody; Anti-DNA: Antideoxyribonucleic acid; Anti-ENA: Anti-extractable nuclear antigen; AOSD: Adult-onset Still’s disease; APLS: Anti-phospholipid syndrome; bDMARD: Biological disease-modifying antirheumatic drugs; CK: Creatine kinase; CRP: C-reactive protein; csDMARD: Conventional synthetic disease-modifying antirheumatic drugs; ENT: Ear-nose-throat; IL: Interleukin;

INF: Interferon; IQR: Interquartile range; JAK: Janus kinase; LDH: Lactate dehydrogenase; NSAIDs: Nonsteroidal anti-inflammatories; RCT: Randomized controlled trial; RF: Rheumatoid factor; rHLH: Reactive hemophagocytic lymphohistiocytosis; SLE: Systemic lupus erythematosus; SoJIA: Systemic-onset juvenile idiopathic arthritis; TNF: Tumor necrosis factor

Acknowledgements

The authors thank the patients who were involved in this study and the individuals from the rheumatology and internal medicine departments who took part in the care of these patients.

Funding

This study was not funded.

Availability of data and materials

The authors state that they had full access to all the data in the study and they take responsibility for the integrity of the data and the accuracy of the data analysis as well as the decision to submit them for publication.

Authors’ contributions

FV and MT contributed to the conception and design of the study. FV contributed to the acquisition of the data. FV, TB, and MT contributed to the analysis and interpretation of the data. FV, MT, TB, FL, ES, EL, BF, CR, and TS drafted the manuscript. FV, MT, AB, XD, FL, ES, EL, BF, CR, and TS contributed to the critical revision of the manuscript. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

The authors state that they had full access to all the data in the study and they take responsibility for the integrity of the data and the accuracy of the data analysis as well as the decision to submit them for publication.

Competing interests

François Vercauysse has no relevant relationship conditions or circumstances that present a potential conflict of interest regarding this study. Thomas Barnetche has no relevant relationships conditions or circumstances that present a potential conflict of interest regarding this study. Estibaliz Lazaro has received consultancy fees from Roche, Novartis, and UCB. Emilie Shipley has no relevant relationships conditions or circumstances that present a potential conflict of interest regarding this study. François Lifermann has no relevant relationships conditions or circumstances that present a potential conflict of interest regarding this study. Alexandre Balageas has no relevant relationships conditions or circumstances that present a potential conflict of interest regarding this study. Xavier Delbrel has no relevant relationships conditions or circumstances that present a potential conflict of interest regarding this study. Bruno Fautrel, Marie-Elise Truchetet, Christophe Richez, and Thierry Schaeverbeke have received consultancy fees and/or research funding from Abbvie, Bristol-Myers Squibb, Pfizer, Roche, Lilly, Novartis, and UCB.

Publisher’s Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Rheumatology Department, Centre Hospitalier Universitaire de Bordeaux, FHU ACRONIM, Service de Rhumatologie, Place Amélie Raba Léon, 33076 Bordeaux, France. ²Centre Hospitalier Universitaire de Bordeaux, FHU ACRONIM, Service de Médecine Interne, Avenue Magellan, 33600 Pessac, France. ³Centre Hospitalier de Dax, Service de Rhumatologie, Boulevard Yves du Manoir, 40100 Dax, France. ⁴Centre Hospitalier de Dax, Service de Médecine Interne, Boulevard Yves du Manoir, 40100 Dax, France. ⁵Centre Hospitalier de Pau, Service de Rhumatologie, 4 Boulevard Hauterive, 64000 Pau, France. ⁶Centre Hospitalier de Pau, Service de Médecine Interne, 4 Boulevard Hauterive, 64000 Pau, France. ⁷Centre Hospitalier Universitaire de Paris, Hôpital Pitié Salpêtrière, Service de Rhumatologie, Boulevard de l’Hôpital, 75013 Paris, France.

Received: 27 August 2018 Accepted: 30 January 2019

Published online: 12 February 2019

References

- Magadur-Joly G, Billaud E, Barrier JH, Pennec YL, Masson C, Renou P, et al. Epidemiology of adult Still's disease: estimate of the incidence by a retrospective study in West France. *Ann Rheum Dis*. 1995;54(7):587–90.
- Castañeda S, Blanco R, González-Gay MA. Adult-onset Still's disease: advances in the treatment. *Best Pract Res Clin Rheumatol*. 2016;30(2): 222–38.
- Jamilloux Y, Gerfaud-Valentin M, Martinon F, Belot A, Henry T, Sève P. Pathogenesis of adult-onset Still's disease: new insights from the juvenile counterpart. *Immunol Res*. 2015;61(1–2):53–62.
- Maria ATJ, Le Quellec A, Jorgensen C, Touitou I, Rivière S, Guilpain P. Adult onset Still's disease (AOSD) in the era of biologic therapies: dichotomous view for cytokine and clinical expressions. *Autoimmun Rev*. 2014;13(11): 1149–59.
- Gerfaud-Valentin M, Jamilloux Y, Iwaz J, Sève P. Adult-onset Still's disease. *Autoimmun Rev*. 2014;13(7):708–22.
- Fujii T, Nojima T, Yasuoka H, Satoh S, Nakamura K, Kuwana M, et al. Cytokine and immunogenetic profiles in Japanese patients with adult Still's disease. Association with chronic articular disease. *Rheumatology*. 2001;40(12):1398–404.
- Shimizu M, Nakagishi Y, Yachie A. Distinct subsets of patients with systemic juvenile idiopathic arthritis based on their cytokine profiles. *Cytokine*. 2013; 61(2):345–8.
- Shimizu M, Yokoyama T, Yamada K, Kaneda H, Wada H, Wada T, et al. Distinct cytokine profiles of systemic-onset juvenile idiopathic arthritis-associated macrophage activation syndrome with particular emphasis on the role of interleukin-18 in its pathogenesis. *Rheumatology*. 2010;49(9): 1645–53.
- Ichida H, Kawaguchi Y, Sugiura T, Takagi K, Katsumata Y, Gono T, et al. Clinical manifestations of adult-onset Still's disease presenting with erosive arthritis: association with low levels of ferritin and interleukin-18. *Arthritis Care Res*. 2014;66(4):642–6.
- Mavragani CP, Spyridakis EG, Koutsilieris M. Adult-onset Still's disease: from pathophysiology to targeted therapies. *Int J Inflamm [Internet]*. 2012;2012. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3390042/>
- Saiki O, Uda H, Nishimoto N, Miwa T, Mima T, Ogawara T, et al. Adult Still's disease reflects a Th2 rather than a Th1 cytokine profile. *Clin Immunol Orlando Fla*. 2004;112(1):120–5.
- Efthimiou P, Moorthy LN, Mavragani CP, Skokos D, Fautrel B. Adult onset Still's disease and autoinflammation. *Int J Inflamm [Internet]*. 2012;2012. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3437651/>.
- Zhou S, Qiao J, Bai J, Wu Y, Fang H. Biological therapy of traditional therapy-resistant adult-onset Still's disease: an evidence-based review. *Ther Clin Risk Manag*. 2018;14:167–71.
- de Boysson H, Février J, Nicolle A, Auzary C, Geffray L. Tocilizumab in the treatment of the adult-onset Still's disease: current clinical evidence. *Clin Rheumatol*. 2013;32(1):141–7.
- Yoshida Y, Sakamoto M, Yokota K, Sato K, Mimura T. Tocilizumab improved both clinical and laboratory manifestations except for Interleukin-18 in a case of multiple drug-resistant adult-onset Still's disease. *Intern Med*. 2011; 50(16):1757–60.
- Naniwa T, Ito R, Watanabe M, Hayami Y, Maeda S, Sasaki K, et al. Case report: successful use of short-term add-on tocilizumab for multirefractory systemic flare of adult-onset Still's disease. *Clin Rheumatol*. 2013;32(Suppl 1): S103–6.
- Ortiz-Sanjuán F, Blanco R, Calvo-Río V, Narvaez J, Rubio Romero E, Olivé A, et al. Efficacy of tocilizumab in conventional treatment-refractory adult-onset Still's disease: multicenter retrospective open-label study of thirty-four patients: tocilizumab in AOSD refractory to standard treatment. *Arthritis Rheumatol*. 2014;66(6):1659–65.
- Govoni M, Bortoluzzi A, Rossi D, Modena V. How I treat patients with adult onset Still's disease in clinical practice. *Autoimmun Rev*. 2017; 16(10):1016–23.
- Gerfaud-Valentin M, Maucourt-Boulch D, Hot A, Iwaz J, Ninet J, Durieu I, et al. Adult-onset Still disease. *Medicine (Baltimore) [Internet]*. 2014 Mar 4 [cited 2015 Dec 18];93(2). Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4616309/>
- Al-Homood IA. Biologic treatments for adult-onset Still's disease. *Rheumatology*. 2014;53(1):32–8.
- Ortiz-Sanjuán F, Blanco R, Riancho-Zarrabeitia L, Castañeda S, Olivé A, Riveros A, et al. Efficacy of anakinra in refractory adult-onset Still's disease. *Medicine (Baltimore) [Internet]*. 2015 Oct 2 [cited 2015 Dec 17];94(39). Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4616841/>
- Giampietro C, Ridene M, Lequerre T, Chalumeau NC, Amoura Z, Sellam J, et al. Anakinra in adult-onset Still's disease: long-term treatment in patients resistant to conventional therapy. *Arthritis Care Res*. 2013;65(5):822–6.
- Laskari K, Tzioufas AG, Moutsopoulos HM. Efficacy and long-term follow-up of IL-1R inhibitor anakinra in adults with Still's disease: a case-series study. *Arthritis Res Ther*. 2011;13(3):R91.
- Colafrancesco S, Priori R, Valesini G, Argolini L, Baldissera E, Bartoloni E, et al. Response to interleukin-1 inhibitors in 140 Italian patients with adult-onset still's disease: a multicentre retrospective observational study. *Front Pharmacol [Internet]*. 2017 Jun 13:8. Available from: <http://journal.frontiersin.org/article/10.3389/fphar.2017.00369/full>
- Lequerré T, Quartier P, Rosellini D, Alaoui F, Bandt MD, Mejjad O, et al. Interleukin-1 receptor antagonist (anakinra) treatment in patients with systemic-onset juvenile idiopathic arthritis or adult onset Still disease: preliminary experience in France. *Ann Rheum Dis*. 2008;67(3): 302–8.
- Castañeda S, Atienza-Mateo B, Martín-Varillas JL, Serra López-Matencio JM, González-Gay MA. Anakinra for the treatment of adult-onset Still's disease. *Expert Rev Clin Immunol*. 2018;14(12):979–92.
- Ruscitti P, Ursini F, Cipriani P, De Sarro G, Giacomelli R. Biologic drugs in adult onset Still's disease: a systematic review and meta-analysis of observational studies. *Expert Rev Clin Immunol*. 2017;13(11):1089–97.
- Cavalli G, Franchini S, Aiello P, Guglielmi B, Berti A, Campochiaro C, et al. Efficacy and safety of biological agents in adult-onset Still's disease. *Scand J Rheumatol*. 2015;44(4):309–14.
- Priori R, Ceccarelli F, Barone F, Iagnocco A, Valesini G. Clinical, biological and sonographic response to IL-1 blockade in adult onset Still's disease. *Clin Exp Rheumatol*. 2008;26:933–7.
- Gerfaud-Valentin M, Maucourt-Boulch D, Hot A, Iwaz J, Ninet J, Durieu I, et al. Adult-onset still disease: manifestations, treatment, outcome, and prognostic factors in 57 patients. *Medicine (Baltimore)*. 2014;93(2):91–9.
- Kim H-A, Sung J-M, Suh C-H. Therapeutic responses and prognosis in adult-onset Still's disease. *Rheumatol Int*. 2012;32(5):1291–8.
- Pay S, Türkçapar N, Kalyoncu M, Simşek I, Beyan E, Ertenli I, et al. A multicenter study of patients with adult-onset Still's disease compared with systemic juvenile idiopathic arthritis. *Clin Rheumatol*. 2006;25(5):639–44.
- Franchini S, Dagna L, Salvo F, Aiello P, Baldissera E, Sabbadini MG. Efficacy of traditional and biologic agents in different clinical phenotypes of adult-onset Still's disease. *Arthritis Rheum*. 2010;62(8):2530–5.
- Fautrel B, Sibilia J, Mariette X, Combe B. Club Rhumatismes et Inflammation. Tumour necrosis factor alpha blocking agents in refractory adult Still's disease: an observational study of 20 cases. *Ann Rheum Dis*. 2005;64(2):262–6.
- Ma Y, Wu M, Zhang X, Xia Q, Yang J, Xu S, et al. Efficacy and safety of tocilizumab with inhibition of interleukin-6 in adult-onset Still's disease: a meta-analysis. *Mod Rheumatol*. 2017;18:1–25.
- Kaneko Y, Kameda H, Ikeda K, Ishii T, Murakami K, Takamatsu H, et al. Tocilizumab in patients with adult-onset Still's disease refractory to glucocorticoid treatment: a randomised, double-blind, placebo-controlled phase III trial. *Ann Rheum Dis*. 2018;77(12):1720–9.
- Rossi-Semerano L, Fautrel B, Wendling D, Hachulla E, Galeotti C, Semerano L, et al. Tolerance and efficacy of off-label anti-interleukin-1 treatments in France: a nationwide survey. *Orphanet J Rare Dis*. 2015;10(1):19.
- Banse C, Vittecoq O, Benhamou Y, Gauthier-Prieur M, Lequerré T, Lévesque H. Reactive macrophage activation syndrome possibly triggered by canakinumab in a patient with adult-onset Still's disease. *Joint Bone Spine*. 2013;80(6):653–5.
- Barsotti S, Neri R, Iacopetti V, d'Ascanio A, Talarico R, Tripoli A, et al. Successful treatment of refractory adult-onset still disease with canakinumab: a case report. *JCR J Clin Rheumatol*. 2014;20(2):121.
- Ahmadi-Simab K, Lamprecht P, Jankowiak C, Gross WL. Successful treatment of refractory adult onset Still's disease with rituximab. *Ann Rheum Dis*. 2006 Aug 1;65(8):1117–8.
- Ostrowski RA, Tehrani R, Kadanoff R. Refractory adult-onset Still disease successfully treated with abatacept. *J Clin Rheumatol*. 2011;17(6):315–7.

42. Belfeki N, Smiti Khanfir M, Said F, Hamzaoui A, Ben Salem T, Ben Ghorbel I, et al. Successful treatment of refractory adult onset Still's disease with rituximab. *Reumatismo*. 2016;16;68(3):159.
43. Quartuccio L, Maset M, De Vita S. Efficacy of abatacept in a refractory case of adult-onset Still's disease. *Clin Exp Rheumatol*. 2010;28(2):265–7.
44. Padilla-Ibarra J, Sanchez-Ortiz A, Sandoval-Castro C, Ramos-Remus C. Rituximab treatment for pulmonary arterial hypertension in adult-onset Still's disease. *Clin Exp Rheumatol*. 2013;31(4):657–8.
45. Bartoloni E, Alunno A, Luccioli F, Santoboni G, Gerli R. Successful treatment of refractory adult-onset Still's disease with anti-CD20 monoclonal antibody. *Clin Exp Rheumatol*. 2009;27(5):888–9.
46. Gabay C, Fautrel B, Rech J, Spertini F, Feist E, Kötter I, et al. Open-label, multicentre, dose-escalating phase II clinical trial on the safety and efficacy of tadekinig alfa (IL-18BP) in adult-onset Still's disease. *Ann Rheum Dis*. 2018;annrheumdis – 2017–212608.
47. Chen D-Y, Chen Y-M, Lan J-L, Lin C-C, Chen H-H, Hsieh C-W. Potential role of Th17 cells in the pathogenesis of adult-onset Still's disease. *Rheumatol Oxf Engl*. 2010;49(12):2305–12.
48. Choi J-H, Suh C-H, Lee Y-M, Suh Y-J, Lee S-K, Kim S-S, et al. Serum cytokine profiles in patients with adult onset Still's disease. *J Rheumatol*. 2003 Nov; 30(11):2422–7.
49. Mimura T, Kondo Y, Ohta A, Iwamoto M, Ota A, Okamoto N, et al. Evidence-based clinical practice guideline for adult Still's disease. *Mod Rheumatol*. 2018;13:1–53.
50. Guilpain P, Le Quellec A, Maria ATJ. Therapeutic innovation in adult-onset Still's disease (and other rare inflammatory disorders): how to secure evidence-based medicine? *Ann Rheum Dis*. 2018;77:1699–701.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

