[Click here to view linked References](#)

Biocontrol, new questions for Ecotoxicology?

Marcel Amichot¹, Pierre Joly², Fabrice Martin-Laurent³, David Siaussat⁴, Anne-Violette

Lavoir¹

1: ISA, Université Côte d'Azur, INRA, CNRS, ISA, Sophia-Antipolis, France. To whom correspondence should be sent.

marcel.amichot@inra.fr; anne-violette.lavoir@inra.fr

2: Biovitis, 15400 Saint Etienne de Chomeil, France

3: Agroécologie, AgroSup Dijon, INRA, Univ. Bourgogne Franche-Comté, 21000 Dijon, France

4: Sorbonne Université- Institute of Ecology and Environmental Sciences of Paris- UMR: INRA, CNRS, IRD, Paris Diderot, Paris-Est Créteil- Campus Pierre et Marie Curie- 4 Place Jussieu, 75005 Paris

Abstract :

Biocontrol strategies are viewed as an environment friendly alternative to the use of conventional synthetic pesticides to control pests or weeds in agrosystems and are expected to supersede the use of pesticides. Nevertheless, biocontrol solutions are not devoid of drawbacks. As encountered with conventional pesticides, one can expect side-effects of biopesticides on non-targeted organisms and/or ecosystem processes. The “bio-” prefix in the word “biopesticides” does it necessarily guarantee their environment safe profile?

In this context, we call to mind the researchers of the network ECOTOX (French network of ecotoxicology; <https://www6.inra.fr/ecotox>) in the framework of a roundtable entitled “Biocontrol & Ecotoxicology” on the following questions: who is experienced in ecotoxicology in relation with biocontrol in its own research? To what extend would it be useful to introduce ecotoxicology in biocontrol? What is the fate of biopesticides in the environment? What role could we cast to ecotoxicology in biocontrol regulations? We report here a synthesis of the discussions engaged during the roundtable.

Key words:

Biopesticide; biocontrol agent; ERA ecotoxicological risk assessment; agroecology; biopesticide regulation.

A.-V.L. was supported by: ERA-NET ARIMNet2, project STomP, IDEX BOOST Projet structurant Université Côte d'Azur.

A.M. was supported by: IDEX BOOST Projet structurant Université Côte d'Azur

Introduction

Biocontrol is viewed as an environment friendly alternative to the use of conventional (synthetic) pesticides to control pests or weeds in agrosystems. It relies on the use of biopesticides (Table 1) gathering either macro/micro-organisms (also called biocontrol agents), natural substances purified from living organisms (biopesticides *sensus stricto*) or mineral preparations as defined in the Article L253-6 from the "Code rural et de la pêche maritime" (Code rural et de la pêche maritime 2014). The organisms can be insects, nematodes, bacteria, fungi or viruses which could act as predators, parasitoids or pathogens of the pests or weeds (Flint and Dreistadt 1998). The natural substances can be plant extracts at various purification degrees or toxins from animals, plants or bacteria. Depending on their nature and composition, they can induce toxic effects or manipulate the behaviour of targeted organisms due to an attractive (as lures in traps) or repulsive effects. As for synthetic pesticides, the natural substances can be sprayed or associated to inert material or artificial food in order to optimize their efficiency in the field.

For many reasons, the biocontrol solutions which were until a recent time rarely used make their coming out. The politics, responding to the public pressure asking for safer agricultural practices, launched several new policies to reduce the use of pesticides such as the European Union (EU) regulation on the sustainable use of pesticides (Directive 2009/128/EC 2009) or the French law for the Agroecology transition (Loi n° 2014-1170 2014) and the Ecophyto II program (Ecophyto II 2015). This new paradigm paves the way to an important development of biocontrol solutions including biopesticides in the coming decades. However, their regulation is still following the same path than that of synthetic pesticides ((EC) N° 1107/2009/EC 2009) which seriously slow down their release on the market, except for the macro-organisms. The International Biocontrol Manufacturers' Association (IBMA) asked EU for considering an adaptation of the regulation to facilitate the homologation of biocontrol agents, notably by considering the concepts of 'substantial equivalence' and of 'low risk substances'. This trend is pushed in front by the French ministry of agriculture to release on the market biocontrol solutions and facilitate the adoption of the Agroecology transition (Chomienne 2017).

The "bio-" prefix in the word "biopesticides" implies the organic origin of such compounds/organisms (the updated list of the authorized biopesticides in France is edited by the "Direction Générale de l'Alimentation (DGAL 2018) but does it necessarily guarantee their environment safe profile? The advantages of their biodegradation and thus

1 their expected low persistence in the environment are often evoked to elude the key
2 questions on the potential side effects of biopesticides. Indeed, they comprised one or
3 several active ingredients which can be harmful for non-target organisms and/or
4 ecosystem processes. Do we have to assess the ecotoxicological risk of biopesticides in
5 the same way as it is currently done for synthetic pesticides? Unfortunately, there are few
6 available pertinent data on the assessment of the ecotoxicological impact on non-target
7 organisms of biopesticides to feed this debate. Many reasons can be evoked to explain
8 this lack of data among which: (i) biocontrol solutions are not enough present on the
9 market despite their evident interest because of different reasons listed above, (ii) the
10 industrial sector of the biocontrol is principally made of small to medium size companies
11 with low R&D budget and facilities, (iii) very few academic laboratories are active in this
12 domain.

13
14 Within this context, we identified four questions, listed thereafter, that were discussed
15 with scientists in a roundtable during the conference of the French ECOTOX network (Dec.
16 2017, Valence, France), gathering researchers working on ecotoxicology:

17 I. Which biopesticides are in need to be studied today?

18 II. To what extent their side effects should be studied and on which organisms?

19 III. What role for ecotoxicology in regulation and homologation processes for the release
20 of biopesticides on the market?

21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 **I. Which biopesticides are in need to be studied today?**

36 We did not identify a biopesticide to address more urgently than another in the
37 framework of the Ecotoxicological Risk Assessment (ERA; Fig.1). Actually, such a
38 selection should occur as what exists for conventional pesticides: the most used, the most
39 toxic or the most present in the environment have to be the most studied. Furthermore, a
40 compound authorized as biopesticide in one country could be forbidden in another one,
41 depending on national legislation, like for synthetic pesticides. There is a clear need for a
42 global homogenization, at the widest level, of the regulations for bio- and synthetic
43 pesticides. Academic researchers also study biopesticides that are not yet on this market,
44 such as Essential Oils (See Pavela and Benelli 2016 for a review) to test their toxicity on
45 pests and their ecotoxicity towards non-target organisms. Actually, they precede the ERA
46 requirements.

47 The concepts, approaches, methodologies and skills applied for ERA of pesticides are
48 readily available (Fig. 1) and can be immediately remobilized and adapted for developing
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

appropriate ERA of biocontrol solutions. Yet, a problem often encountered when performing ERA of biopesticides lies in the difficulty to identify and have available the active ingredient. In line with this, the development of standardized biopesticides formulations is as crucial as for synthetic pesticides. Indeed, for both bio- and synthetic-pesticides one can have questions relative to the role, the effect and the origin of a potential co-formulant? Attention should be paid to develop suitable formulations to ensure the efficacy of both bio- and synthetic-pesticides and limit as far as possible ecotoxicological risk for the environment.

During the roundtable discussion, it appears that only a few numbers of laboratories of the French Network of Ecotoxicology are dealing with ecotoxicology in biocontrol. The same remark can be done at the EU level and worldwide.

II. To what extent, side effects should be studied and on which organisms

In France the agro-ecology transition is part of the 'law for the future of agriculture (Loi n° 2014-1170 2014). This law was adopted in year 2014 with the objectives to diminish the dependency of agriculture to agrochemicals and to decrease the negative effects of agriculture on environmental and human health. Within this context, the use of biocontrol solutions is promoted by the French Ministry of Agriculture. Despite this political pressure, the healthiness of biopesticides remains a matter of debate as recently underlined in a report from the French General Council for Food, Agriculture (Chomienne 2017). Indeed, there is still the need for a careful toxicological, ecotoxicological and ecological evaluation of the biocontrol products and solutions. Biopesticides as synthetic pesticides both contain one or several active ingredients designed to control a given pest which can harm non-targeted organisms.

Several ecotoxicological studies of various biopesticides on diverse non-targeted organisms have already been published such as arthropods (Biondi et al. 2012a; Biondi et al. 2012b; Nakasu et al. 2014; Renzi et al. 2016; Amichot et al. 2016), soil organisms (Ipsilantis et al. 2012; Romdhane et al. 2016; Chelinho et al. 2017), aquatic organisms (Duchet et al. 2010; Manachini et al. 2013) or mammals (Rahoui et al. 2014). The choice of such non-target organisms reflects on one hand the ecosystem service they provide: the beneficial arthropods are involved in biocontrol and pollination services when the beneficial soil organisms are involved in decomposition or nutrient transfer (e.g. mycorrhizas) services. Looking for the side effect of biopesticides on such organisms implies the potential side effect of biopesticides on ecosystem services. On the other hand, this choice

reflects the different components of the ecosystem (e.g. hydrosphere representing by aquatic organisms, lithosphere by soil organisms and biosphere by all of them) in order to evaluate the healthiness of biopesticides on all the environmental components.

In addition, the simultaneous use of several biopesticides as part of the biocontrol strategy applied on a given crop, such as the combination of parasitoids and micro-organisms or even the combination of biopesticide with conventional pesticides, asks the question of mixtures. Finally, studies performed with laboratory model organisms such as *Drosophila* or Mouse might also be extremely helpful to identify the mechanisms responsible for the putative side effects of biopesticides.

As mentioned above, insects, plants, bacteria, plant or microbial extracts can be used as biopesticides (Table 1). Being diverse by nature, they are diverging by their mode of action with a wide diversity of mechanisms involved in their toxicity, and thus they may have various side effects not only on target-organisms but also on ecosystem functions. ERA for pesticides cannot be readily transferred to other biocontrol solutions involving macro- and microscopic organisms. Regarding biocontrol agents, the risk assessment focus more on invasive species or ecosystem functions malfunctioning. It is noteworthy that ecosystem functions have been proposed by the European Food Safety Agency as specific protection goals for ERA of pesticides. Indeed, release of biocontrol agents in the field can be compared to invasive species thus questioning which is relevant to the ecology field, and more specifically to the community ecology (See De Castro et al 2010 and Sundh & Goettel 2013 for a comparison between macro- and micro-organisms). Furthermore, it was underlined that for micro-organisms releasing toxins, it will be necessary to trace the fate of micro-organisms and of their toxins as they may have different persistence trajectories in the environment (See Sundh & Goettel 2013 for a review).

From this point of view, the discussion points out the need for the academics to clarify the contribution of both ecology and ecotoxicology concepts to environmental risk assessment of biopesticides. As a conclusion we agreed that ecotoxicologists and ecologists should overcome disciplinary boundaries and share their concepts, approaches and tools to remobilize them for a more global environmental risk assessment of biopesticides.

III. What role for ecotoxicology in regulation and homologation processes for the release of biopesticides on the market?

1 The complexity of the biopesticide regulation has been underlined in the previous
2 sections. Indeed, two difficulties were highlighted: the first one is the nature of the
3 biocontrol solution which can either be a substance or a living organism. On one hand, the
4 biopesticides *stricto sensu* and the microorganisms are until now submitted to the same
5 EU-Regulation (i.e. (EC) N° 1107/2009) like the synthetic pesticides, but with different data
6 requirements for microorganisms ((EC) N° 283/2013). On the other hand, macro-organism
7 use is submitted to a peculiar regulation (Loi “Grenelle II” 2010, Décret n° 2012-140 2012),
8 based on their fundamental biology and their ecological interactions.
9

10
11
12
13 The second difficulty comes from the frequent bypass of current regulations by
14 proposing biostimulants doped with biopesticides to improve their intrinsic quality and
15 functions (See (Pavela and Benelli 2016) for the essential oils case and (Sundh and
16 Goettel 2012) for the micro-organism case). In fact, the regulation for biostimulants is
17 much less binding than the one for pesticides and biopesticides ((EC) N° 1069/2009 2009,
18 (EC) N° 1107/2009 2009). Consequently, we concluded that there is an urgent need for
19 the overall reconsideration of the regulations for the use of organisms, biopesticides and
20 biostimulants doped with biopesticides under the general umbrella of biocontrol solutions.
21

22
23
24
25
26
27
28 The process leading to the approval of biopesticides should also question the
29 ecotoxicity of their co-formulants entering in the recipe of their formulation which currently
30 is under the REACH regulation ((EC) N° 1907/2006 2006): the potential side effect of the
31 mixture of co-formulants entering in the composition of the formulation have to be
32 addressed.
33

34
35
36
37 We recognized the importance of ecotoxicology for (i) *a priori* ecotoxicological risk
38 assessment of biopesticides as a part of the homologation process and for (ii) *a posteriori*
39 ecotoxicological risk assessment of formulated biopesticides which may lead to their
40 restriction of use when danger was obvious. Thus, ecotoxicology can play a pivotal role to
41 secure the development of biocontrol solutions.
42
43
44
45

46 47 48 **Conclusion**

49
50 To conclude, we recognized that among the French ECOTOX network and likely more
51 widely, there are today only few academic laboratories working in the field of ecotoxicology
52 involved in the ERA for biocontrol solutions. Thus, it appeared that the *a priori* ERA of
53 biopesticides should be done in a similar manner as carried out for conventional pesticides.
54 The richness of the discussions among the participants to the roundtable confirmed the
55 relevance of the questions asked. We agreed that further discussions have to be planned
56
57
58
59
60
61
62
63
64
65

to define a common ground between ecotoxicological and ecological concepts that Ecotoxicology should definitely help to reach the crucial challenges for the biocontrol.

Given the fact that biopesticides *sensus stricto* follow the same regulation than synthetic pesticides, their ecotoxicological risk assessment is not really a matter of discussion except that the overall process is time consuming and costly like it is for synthetic pesticides. This may result in slowing down the release of biocontrol solutions on the market in particular for small- or medium-size companies who have less budget than pesticide industries to carry out all the requirements needed to claim an authorization to put on the market a new active ingredient. On the contrary, for living organisms used as biocontrol agents, ecotoxicological risk assessment passes beyond the concepts of ecotoxicology and have recourse to ecology concepts and researches on invasive species. Therefore, their mode of action and their putative side effects on non-target organisms are expected to be extremely different. We believe that it is an interesting path to follow. Furthermore, the use of living organisms which produce toxins (as *Bacillus thuringiensis* for instance) should be the object of specific attention since the toxins component can be submitted to ERA whereas the bacteria component can be considered as a putative invasive organism (See Sundh & Goettel 2013 for a review).

We feared a major difficulty in the ecotoxicological risk assessment scheme of biocontrol agents because of the diversity of their nature which can either be living organisms or substances (plant extracts, toxins...). Nevertheless, we believe that well mastered concepts, approaches and methodologies of ecotoxicology and also of ecology can be adapted and transferred for a better ecotoxicological risk assessment of biocontrol agents to truly ensure their safe development and to consolidate them as a credible alternative to conventional plant protection products.

Acknowledgements:

We would like to warmly thank the 50 participants (44 academics (including one Canadian), 2 from the industry, 2 from the stakeholder and 2 from a scientific foundation) to ECOTOX conference who took part to this roundtable. Please note that this text is the transcription of our discussions held during the roundtable and should not be considered as an exhaustive picture of ecotoxicology in biocontrol.

References

Amichot, M., Curty, C., Benguettat-Magliano, O., et al (2016) Side effects of *Bacillus thuringiensis* var. *kurstaki* on the hymenopterous parasitic wasp *Trichogramma chilonis*. *Environ Sci Pollut Res Int* **23**, 3097–3103. doi: 10.1007/s11356-015-5830-7

Arrêté AGRG1502673A (2015):
<https://www.legifrance.gouv.fr/eli/arrete/2015/2/26/AGRG1502673A/jo/texte>

Biondi A, Desneux N, Siscaro G, Zappalà L (2012a) Using organic-certified rather than synthetic pesticides may not be safer for biological control agents: selectivity and side effects of 14 pesticides on the predator *Orius laevigatus*. *Chemosphere* 87:803–812. doi: 10.1016/j.chemosphere.2011.12.082

Biondi A, Mommaerts V, Smaghe G, et al (2012b) The non-target impact of spinosyns on beneficial arthropods. *Pest Manag Sci* 68:1523–1536. doi: 10.1002/ps.3396

Chelinho S, Maleita CMN, Francisco R, et al (2017) Toxicity of the bionematicide 1,4-naphthoquinone on non-target soil organisms. *Chemosphere* 181:579–588. doi: 10.1016/j.chemosphere.2017.04.092.

Chomienne, J.-P., Dutartre, S., Larguier, M., Pinçonnet, D. & Tessier, R. *Les produits de biocontrôle pour la protection des cultures*. 1–87 (2017).

Code rural et de la pêche maritime (2014)
<https://www.legifrance.gouv.fr/affichCodeArticle.do?cidTexte=LEGITEXT000006071367&idArticle=LEGIARTI000006583210&dateTexte=&categorieLien=cid> in French

de Castro VLSS, Jonsson CM, Silva CMM, de Holanda Nunes Maia A (2010) Assessing the safety of *Pseudomonas putida* introduction in the environment: An overview of ecotoxicological tests. *Regulatory Toxicology and Pharmacology* 56:300–305. doi: 10.1016/j.yrtph.2009.09.019.

Décret n° 2012-140 (2012)
<https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000025241913&categorieLien=id>.

DGAL (2018) <https://info.agriculture.gouv.fr/gedei/site/bo-agri/instruction-2018-528> in French. Note that this list is edited bimonthly.

Directive 2009/128/EC (2009) : <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32009L0128>

Duchet C, Coutellec M-A, Franquet E, et al (2010) Population-level effects of spinosad and *Bacillus thuringiensis israelensis* in *Daphnia pulex* and *Daphnia magna*: comparison of laboratory and field microcosm exposure conditions. *Ecotoxicology* 19:1224–1237. doi:

10.1007/s10646-010-0507-y

(EC) N° 1907/2006 (2006) <https://eur-lex.europa.eu/legal-content/en/ALL/?uri=CELEX%3A32006R1907>

(EC) N° 1069/2009 (2009): <https://eur-lex.europa.eu/legal-content/EN/TXT/?qid=1535720961378&uri=CELEX:32009R1069>

(EC) N° 1107/2009 (2009): <http://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32009R1107>

(EC) N° 283/2013 (2013): <https://eur-lex.europa.eu/legal-content/AUTO/?uri=CELEX:32013R0283&qid=1535720740043&rid=1>

Ecophyto II (2015): agriculture.gouv.fr/sites/minagri/files/151022_ecophyto.pdf in French.

Flint ML, Dreistadt SH (1998) Natural Enemies Handbook. Univ of California Press

Ipsilantis I, Samourelis C, Karpouzas DG (2012) The impact of biological pesticides on arbuscular mycorrhizal fungi. Soil Biology and Biochemistry 45:147–155. doi: 10.1016/j.soilbio.2011.08.007.

Loi “Grenelle II” (2010) <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000022470434> in French.

Loi n° 2014-1170 (2014) : <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000029573022&dateTexte=&categorieLien=id> in French.

Manachini B, Arizza V, Rinaldi A, et al (2013) Eco-physiological response of two marine bivalves to acute exposition to commercial Bt-based pesticide. Marine Environmental Research 83:29–37. doi: 10.1016/j.marenvres.2012.10.006

Nakasu EYT, Williamson SM, Edwards MG, et al (2014) Novel biopesticide based on a spider venom peptide shows no adverse effects on honeybees. Proceedings of the Royal Society B: Biological Sciences 281:20140619–20140619. doi: 10.1098/rspb.2014.0619

Pavela R, Benelli G (2016) Essential Oils as Ecofriendly Biopesticides? Challenges and Constraints. Trends in Plant Science 21:1000–1007. doi: 10.1016/j.tplants.2016.10.005

Rahioui I, Eyraud V, Karaki L, et al (2014) Host range of the potential biopesticide Pea Albumin 1b (PA1b) is limited to insects. Toxicon 89:67–76. doi: 10.1016/j.toxicon.2014.07.004

Renzi MT, Amichot M, Pauron D, et al (2016) Chronic toxicity and physiological changes induced in the honey bee by the exposure to fipronil and *Bacillus thuringiensis*

spores alone or combined. *Ecotoxicol Environ Saf* 127:205–213. doi: 10.1016/j.ecoenv.2016.01.028.

Romdhane S, Devers-Lamrani M, Barthelmebs L, et al (2016) Ecotoxicological Impact of the Bioherbicide Leptospermone on the Microbial Community of Two Arable Soils. *Front Microbiol* 7:2762–12. doi: 10.3389/fmicb.2016.00775.

Sundh I, Goettel MS (2012) Regulating biocontrol agents: a historical perspective and a critical examination comparing microbial and macrobial agents. *BioControl* 58:575–593. doi: 10.1007/s10526-012-9498-3

Legends

Figure 1

Flowchart of the Ecotoxicological Risk Assessment for pesticides.

This flowchart is also conceptually suitable to address the putative risks associated with the use of biopesticides or biostimulants. After the problem formulation, laboratory experiments are conducted to address the putative toxicity of the compound on pertinent organisms, for instance organisms present in the ecological niche in which the compound is planned to be used. If toxicity is detected, then field experiments will determine the environmental distribution of the compound and the modes of exposition of the organisms. If a new use is intended for a known compound, it is possible to directly plan exposure measures as shown by the dotted arrow because the toxicity of the compound is already known. We boxed the lower area to emphasize the close relationships that exist between Ecotoxicological Risk Assessment and the Environmental Risks Assessment (the boxed area), although separated in the regulations. As for synthetic pesticides, the choice of the non-targeted organisms to be tested with biopesticides in laboratory experiments is crucial for the relevancy of the tests. Furthermore, there is a gap to fill for the exposition measures relatively to biopesticides as the data are very scarce in the literature.

Table 1: Biopesticides *sensus lato* authorised for use in France

Synthesis of biocontrol solutions authorized in France. Many products are natural extracts from plant or mineral origins. Most of them have a broad range of targets but some are very specific such as the pheromones. The reader may note that not all products are used to kill pests, a few impact the behaviour of the pests (mainly semiochemicals category) or stimulate the ability of plants to defend against pests (included in the plant extract) *.

Based on the list of authorized biopesticides in France for biopesticides *stricto sensu* (DGAL 2018), the mineral solution and the microorganisms and for the macro-organisms (Arrêté AGRG1502673A 2015).

Table 1

	<i>Living Organisms = Biocontrol agents</i>				<i>Natural substances = Biopesticides sensus stricto</i>					<i>Mineral substances</i>	
	Macro-organisms			Microorganisms	Natural extract				Semiochemicals		
	Pathogens	Predators	Parasitoids	Pathogens	Plant origin*	Animal origin	Microorganism origin	Complex	Pheromones		Allelochemicals
	Nematodes	Insects		Bacteria, Fungi, Virus							
Mode of Action	Insecticide			Insecticide, Fungicide, Bactericide, Herbicide					Behavioural effect on insects		Insecticide, Fungicide, Bactericide, Herbicide
Regulation	Loi “Grenelle II” Décret n° 2012-140			Commission Regulation (EU) No 283/2013	Regulation (EC) No 1107/2009 For biostimulants*: (CE) No 1069/2009 (CE) No 1107/2009						
List of official biopesticides in France	Décret 26 Feb 2015 https://www.legifrance.gouv.fr/eli/arrete/2015/2/26/AGRG1502673A/jo/texte			Bimonthly edited list at: https://info.agriculture.gouv.fr/gedei/site/bo-agri/instruction-2018-528							