Antibacterial, anti-adherent and cytotoxic activities of surfactin(s) from a lipolytic strain *Bacillus safensis* F4

Faten Abdelli, Marwa Jardak, Jihene Elloumi, Didier Stien, Slim Cherif, Sami Mnif, Sami Aifa

► To cite this version:

Faten Abdelli, Marwa Jardak, Jihene Elloumi, Didier Stien, Slim Cherif, et al.. Antibacterial, anti-adherent and cytotoxic activities of surfactin(s) from a lipolytic strain *Bacillus safensis* F4. Biodegradation, In press, 10.1007/s10532-018-09865-4 . hal-02093931

HAL Id: hal-02093931

<https://hal.sorbonne-universite.fr/hal-02093931>

Submitted on 9 Apr 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Antibacterial, anti-adherent and cytotoxic activities of surfactin(s)
from a lipolytic strain *Bacillus safensis* F4

¹Faten ABDELLI, ¹Marwa JARDAK, ¹Jihene ELLOUMI, ²Didier STIEN, ³Slim CHERIF,
¹Sami MNIF and ^{1*}Sami AIFA

¹Laboratory of Molecular and Cellular Screening Processes, Centre of Biotechnology of Sfax,
P.O. Box 1177, 3018

²UPMC UNIV PARIS 06, CNRS, Laboratoire de Biodiversité et Biotechnologies
Microbiennes (LBBM), Sorbonne Universities, Observatoire Oceanologique, 66650 Banyuls-
Sur-Mer, FRANCE.

³Faculty of Sciences, Gafsa. University Campus Sidi Ahmed Zarrouk 2112 Gafsa. University
of Gafsa.

*Corresponding author:

Prof. Sami Aifa

Centre of Biotechnology of Sfax (University of Sfax).

Sidi Mansour Road Km 6, BP 1177, 3018 Sfax, Tunisia

Email : sami.aifa@cbs.rnrt.tn

Phone: +216-74871816

Fax: +216-74875818

Abstract

The bacterial strain F4, isolated from olive oil-contaminated soil, has been found to produce biosurfactants as confirmed by oil displacement test and the emulsification index results. The identification of the strain F4, by 16S ribosomal RNA gene, showed a close similarity to *Bacillus safensis*, therefore the strain has been termed *Bacillus safensis* F4. The Thin Layer Chromatography (TLC) and the High Pressure LiquidChromatography-Mass Spectrometry (HPLC-MS/MS) demonstrated that the biosurfactant had a lipopeptide structure and was classified as surfactin. The present study showed also that the produced biosurfactant has an important antibacterial activity against several pathogen strains as monitored with minimum inhibitory concentration (MIC) micro-assays. In particular, it presented an interesting anti-planktonic activity with a MIC of 6.25 mg mL⁻¹ and anti-adhesive activity which exceeded 80% against the biofilm-forming *Staphylococcus epidermidis* S61 strain. Moreover, the produced lipopeptide showed an antitumor activity against T47D breast cancer cells and B16F10 mouse melanoma cells with IC₅₀ of 0.66 mg mL⁻¹ and 1.17 mg mL⁻¹, respectively. Thus, our results demonstrated that *Bacillus safensis* F4 biosurfactant exhibited a polyvalent activity *via* a considerableantibiofilm and antitumoralpotencies.

Keywords:

Anti-adherent; anti-cancer; Biosurfactant; *Bacillussafensis* F4; Surfactin(s).

Introduction

Biosurfactants or bioemulsifiers are amphipathic surface-active molecules, which are produced by micro-organisms, composed of hydrophobic (nonpolar) and hydrophilic (polar) moieties. As a consequence, they have the ability to aggregate at interfaces between fluids with different polarities such as oil/water or air/water, reduce the surface and interfacial tensions and form emulsions (Sen et al. 2017). These compounds are characterized as glycolipids, lipopeptides, lipopolysaccharides, fatty acids, phospholipids and neutral lipids (Bezza and Chirwa 2016; Colla et al. 2010). Biosurfactants are produced by a wide variety of bacteria, actinobacteria and fungi with different chemical structures. Some bacterial genera like *Bacillus* and *Arthrobacter* are known with their production of lipopeptide biosurfactant (Sriram et al. 2011). Some studies have described the biological activities of the biosurfactants including antimicrobial, anti-adhesive and anti-biofilm properties (Silva et al. 2014, Gudiña et al. 2010a). In fact, the bacterial infections and their biofilm formation abilities causing resistance increase against drugs is getting a serious problem for human health. An urgent need for solving this problem is based on the screening of novel drugs eradicating or inhibiting biofilm formation. The adherence is the first step of the infectious process that requires efficient antagonising molecules. Previous studies reported that based on their amphiphilic structures, the biosurfactants reduce the surface tension and therefore affecting the bacterial adherence (Janek et al. 2013). In this context, the lipopeptide biosurfactant produced by *Bacillus subtilis* presented antibacterial, anti-adhesive and anti-biofilm activities on uropathogenic bacteria (Moryl et al. 2015). Moreover, a glycolipid biosurfactant, presented cytotoxic activities on cancer cell lines, was produced by a *Nocardia farcinica* strain (Christova et al. 2015). The biosurfactants, which are selective in nature, act on the surface of liquids and facilitate the action of certain enzymes such as lipases and/or esterases by reducing the surface

tension of liquids and/or improving the solubility of water immiscible substrates (Sekhon et al. 2011, 2012).

Lipases are characterized by their ability to synthesize ester bonds in a non-aqueous media (Ülker and Karaoglu 2012) and their production can be associated with several factors including pH, temperature, carbon source and the presence of inducers such as oils and some biosurfactants (Cherif et al. 2011; Colla et al. 2010).

Nowadays, biosurfactants take an important scientific interest with their interesting proprieties such as the high biodegradability, lower toxicity, better environmental compatibility, and important specific activity at extreme conditions of temperature, pH and salinity (Sriram et al. 2011).

In this context, searching for novel biosurfactant producing strains with potential biosurfactant production is required. For that, lipolytic strains could be a possible original source of biosurfactant production (Sekhon et al. 2012). The present study describes the biosurfactant production by a lipolytic strain *B. safensis* F4 and investigates its antibacterial, anti-adhesive and antitumor activities.

Materials and methods

Bacterial strains

B. safensis F4, *B. subtilis*, *Staphylococcus aureus*, *Enterococcus faecium*, *Micrococcus luteus*, *Agrobacterium tumefaciens*, *Salmonella enterica*, *Escherichia coli* and *Pseudomonas savastanoi* were grown in LB (Luria-Bertani) medium. *S. epidermidis* S61, a biofilm-forming bacterium isolated in our lab from the roof of an old house in Sfax, Tunisia (Jardak et al. 2017), was grown in Tryptic Soy Broth (TSB) medium.

Cell lines and cultures

Breast cancer T47D and mouse melanoma B16F10 cell lines, obtained from the American Type Culture Collection (ATCC), were grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% foetal bovine serum, 50 IU/mL penicillin, 50 mg mL⁻¹ streptomycin at 37°C in a humidified 5% CO₂ atmosphere.

Bacterial biosurfactant activity

The oil displacement assay was performed according to Morikawa et al. (1993) using the Petri plate (90 mm diameter) filled with 25 mL of distilled water then 10 µL of a crude oil was added. 10 µL of a cell free culture supernatant was slowly placed on the center of the oil surface. The diameter of the clear halo zone was measured after 30 seconds of incubation.

The determination of the emulsification index (E24) is carried out according to the following equation (Cooper and Goldenberg 1987). E24 was measured using the cell free culture. Two millilitres of a vegetable oil were added to an equal volume of cell free supernatant and homogenized for 2 minutes at high speed. The height of emulsion layer was measured after 24h. All the experiments were done in triplicate.

$$\text{E24 (\%)} = (\text{Total height of the emulsified layer} / \text{Total height of the liquid layer}) \times 100$$

Surface tension determination

Surface tension of the 24 h culture broth supernatant was measured according to the De Nouy methodology using a tensiometer TD1 (Lauda-Königshofen, Germany). The measurement was performed in triplicate.

Identification of bacterial strain

Strain F4 was identified using the API 20E test *Enterobacteriaceae* (BioMérieux, France) and by sequencing of the 16S rRNA gene. The genomic DNA of the strain F4 was extracted

following the protocol detailed by Wilson et al. (1987). The 16S ribosomal DNA of the strain F4 was amplified by PCR (Polymerase chain reaction) using the universal bacterial primers Fd1 and Rd1 (Fd1, 5'-AGAGTTTGATCCTGGCTCAG-3'; Rd1, 5'-AAGGAGG-TGATCCAGCC-3'), and the following program: denaturation at 94°C for 30 sec, annealing at 55°C for 45 sec and extension at 72°C for 1 min 45 sec for a total of 30 cycles.

The PCR products were purified with a Favor Prep GEL/ PCR Purification Kit (FAVORGEN) and sequenced using the ABI PRISM, 3100. The obtained sequences were compared with other bacterial sequences in the NCBI database using BLAST program. The phylogenetic tree was constructed using the neighbour-joining method (Naruya and Nei 1987) by MEGA 4.0.

Bacterial biosurfactant production

B.safensis F4 strain was retained as the best local strain producing biosurfactant. The strain was incubated overnight at 30°C and 160 rpm in 250 mL shaking flasks with 100 mL of LB medium. Two millilitres of culture were used as inoculum and were cultivated in 500 mL shaking flasks containing 200 mL of the medium with 1% olive oil. The culture was incubated for 24 h at 180 rpm and 30°C to allow maximum biosurfactant production. Cell-free supernatant was obtained by centrifugation at 4°C during 20 min at 4000xg (ROTANTA 460 RF, Hettich). The obtained supernatant was treated by acidification to pH 2.0 using 3M HCl solution and incubated overnight at 4°C. Then, the acidified supernatant was extracted with ethyl acetate and concentrated with a rotary evaporator (Gargouri et al. 2016).

Thin layer chromatography (TLC)

The extracted biosurfactant in ethyl acetate was analysed by TLC. The sample dissolved in methanol was spotted on silica gel TLC plate (TLC Silica gel 60 F₂₅₄, Merck Darmstadt, Germany). The plate was developed with a mobile phase of chloroform/methanol/water

respectively in the ratio of 65:25:4 (v/v/v). The dried plate was sprayed with a solution of 0.25% ninhydrin in acetone and then, incubated at 105°C for 5 min (Janek et al. 2010).

Biosurfactant purification and identification

The extracted biosurfactant in ethyl acetate was fractioned using solid-phase extraction (SPE) (Alajlani et al. 2016). C₁₈ Phenomenex strata-X column (silica gel, 10 g) was conditioned by the elution of 3 volumes of acetonitrile. The sample was deposited on the surface of the silica and drawn through the solvent. For the mobile phase, the HPLC (High Pressure Liquid Chromatography) grade acetonitrile (100% - 3 volume column) was used in first step, then a binary mixture of HPLC grade dichloromethane/ methanol (v/v - 3 volume column) was used. The obtained eluates were collected and dried under vacuum. Finally, the acetonitrile fraction was retained.

Two microliters of acetonitrile fraction diluted at 5 mgmL⁻¹ in methanol, were injected in a Dionex Ultimate 3000 UHPLC-HESI HRMS Q-Exactive focus system (Thermo Scientific) connected to Xcalibur software. The chromatographic separation was conducted followed the protocol of Girard et al. (2017) with slight modifications. The Hypersil GOLD C₁₈ column (150 mm × 2.1 mm) with 1.9 µm particle size (Thermo Scientific) and constant flow rate of 0.5 mL min⁻¹. The column oven was set to 50°C. The water (eluent A) and acetonitrile (eluent B) containing both 0.1% formic acid, were used as mobile phases. A gradient profile was applied, starting with 5% of B and kept constant for 1 min. The percentage of B was linearly increased to 100% in 15 min, and was kept at 100% for 9 min and returned to initial conditions over 1 min. Four minutes of equilibration were followed, giving a total operating time of 30 min. The instrument has been run in the full scan mode with a range of 100 to 1500 m/z equipped with an electrospray interface (ESI). The polarity of the electrospray interface was continuously switched between positive and negative polarity. The LB medium

was used as a control subjected to extraction with ethyl acetate. The common peaks between the chromatographs of the samples and the medium were not retained.

Determination of minimum inhibitory concentration (MIC)

The minimum inhibitory concentration (MIC) for the lipopeptide, produced by *B.safensis* F4, was defined as the lowest concentration that inhibited the growth of microorganisms after 24 h. The test was performed against several human and plant pathogenic strains (*S.aureus*, *E.faecium*, *M. luteus*, *A. tumefaciens*, *S. enterica*, *E. coli* and *P. savastanoi*) and *B. subtilis*. The choice of these strains is justified since we tried to maximize our chance for finding interesting molecules that could be applied to fight against human or plant bacterial infections. The biosurfactant anti-planktonic activity against *S. epidermidis* S61 was performed with the same test. Each bacterium was grown in LB medium overnight at 30°C. Bacterial cultures were then adjusted to an optical density of 0.6 at a wavelength of 600 nm. The crude biosurfactant was dissolved in Dimethylsulfoxide (DMSO) and then filtered. Serial dilutions were made to yield volumes of 100 µL per well with final concentrations ranging from 0.0125 to 25 mg mL⁻¹ in LB medium. Twenty microliters of bacterium overnight culture, with appropriate OD, were added to each well and a final volume of 200 µL per well was adjusted with medium. Wells containing just LB medium with inoculum and these containing medium, inoculum and Ampicillin served as controls. The plate was then incubated at 37°C for 24 h. Twenty microliters of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) solution at 1 mg mL⁻¹ were added to each well. The determination of the biosurfactant MIC was based on the MTT color change. In fact, the viable bacteria were detected by the change of yellow MTT color to purple. For that, the well devoid of bacterial growth (yellow color) was retained as MIC, which was expressed in mg mL⁻¹. The same test was carried out against Gram-positive and Gram-negative strains.

Anti-adhesive activity

The 96-well flat bottom plates were used for biofilm cultures (Mathur et al. 2006).

S. epidermidis S61, a biofilm-forming strain (Jardak et al. 2017), was grown overnight in TSB medium at 30°C and diluted with fresh medium supplemented with 2.25% (w/v) glucose. One hundred microliters of the bacterial culture dilution was added into each well to obtain a final OD_{600 nm} of 0.1. Then, 100 µL of *B. safensis* F4 biosurfactant dissolved in TSB, containing 20% (v/v) of DMSO, at various concentrations, were added into wells to reach final concentrations of 0.039, 0.078, 0.156, 0.312, 0.625, 1.25, 2.5, 5 and 10 mg mL⁻¹. Wells containing only TSB medium supplemented with, 2.25% glucose and 20% (v/v) of DMSO, and bacterial suspension were served as controls.

Plates were incubated for 24 h at 30° C under static conditions. After incubation, the wells were emptied into a container by inverting the plates. Each well was gently washed twice with 250 µL of sterile phosphate buffered saline (PBS: 137 mM NaCl, 2.7 mM KCl; 10 mM Na₂HPO₄ ; 1.76 mM KH₂PO₄; pH 7.2) in order to remove the planktonic cells (Beenken et al. 2003) . After washing, plates were dried at 60°C for 60 min. Then, wells were stained with 150 µL of crystal violet (0.2%) prepared in 20% ethanol for 15 min at room temperature (Vasudevan et al. 2003). After staining incubation, crystal violet was removed and excess dye was washed three times with sterile water. Finally, 200 µL of glacial acetic acid 33% was added to each well and plates were incubated for 1 hour at room temperature. The optical density (OD) was measured at 570 nm using a Varioskan microplate reader (ThermoFisher).

The percentage of the adhesion inhibition was calculated by the following formula:

$$[(\text{OD (control*)} - \text{OD (treated strain)}) / \text{OD (control*)}] \times 100$$

*Control: untreated strain with the extract

The anti-adhesive activities of the crude biosurfactant and the acetonitrile fraction, against *S. epidermidis* S61 were confirmed by microscopic observations using the OLYMPUS fluorescent microscope BX50 equipped with a digital camera OLYMPUS DP70. The biofilms were grown on glass pieces (Ø 10mm) placed in 24-well polystyrene plates treated with the biosurfactant. Non-treated wells, containing TSB supplemented with 10% (v/v) of DMSO, served as controls (Padmavathi and Pandian 2014). The biosurfactant was added at a final concentration of 10 mg mL⁻¹ in TSB with 10% (v/v) of DMSO. The bacterial inoculation was adjusted to an OD_{600nm} of 0.1. Plates were incubated at 30°C for 24 h. The wells were then carefully emptied with pipetting and glass slides were washed with sterile PBS (1X) before the treatment with 500 µL of acridine orange (0.1%, w/v, dissolved in PBS 1X). Visualization was performed through a 40x objective using U-MWB2 filter with excitation at 460-490 nm and emission at 520 nm.

Cytotoxicity assays

T47D breast cancer and B16F10 mouse melanoma cells were grown in 96-well plates (Orange Scientific) until 40% confluence. The biosurfactant, was added at different concentrations (0.1, 1 and 10 mg mL⁻¹) and incubated for 48 hours at 37°C in a humidified atmosphere containing 5% CO₂.

The cell viability was assessed using the MTT assay as previously described by Mosmann (1983). After treatment, the medium was exchanged by a fresh one and 10 µL of MTT solution (5 mg mL⁻¹ in PBS) were added. After incubation for 4 hours, 100 µL of 10% SDS (Sodium dodecyl sulfate) solution were added to each well to dissolve the formazan. The optical density was measured at 570 nm using a Varioskan microplate reader (ThermoFisher). The growth inhibition was expressed according to the following formula:

$$(\%) \text{ cell survival} = (AT/A0) \times 100$$

A0: control absorbance; AT: treated cells absorbance.

Statistical Analysis

All experiments were done in triplicate. The obtained results are expressed as mean values with the standard error. The statistical analyses were performed using Student's t-test to compare the controls and treated samples at a significance level of 5%.

Results and discussion

Selection of biosurfactant producing strain

Morphological and biochemical tests showed that the rod-shaped strain F4 was motile, Gram-positive, catalase-positive and oxidase-positive. Based on the phylogenetic analysis of the 16S rRNA gene sequences, the strain F4 was affiliated to the genus *Bacillus* with 99% of similarity to *Bacillus safensis* FO-36b^T (AF234854) (Fig. 1) and was termed as *B. safensis* F4. The 16S rRNA gene sequence, including 1378 nucleotides, was deposited in the GenBank nucleotide database under the accession number MF927780.

In fact, *B. safensis* F4 was retained after laboratory screening of lipolytic strains for their ability to produce biosurfactant during growth on olive oil. The oil displacement assay showed that the selected strain presented the highest clear halo zone (about $21.08 \pm 1.46 \text{ cm}^2$). The emulsification activity of the selected strain against sunflower oil was 74.99%. Previous results showed that *Bacillus cereus* NK1 biosurfactant presented a clear halo zone of 2.95 cm^2 and 62% in the oil displacement test and emulsification activity against *n*-hexadecane, respectively (Sriram et al. 2011). Ibrahim (2018) claimed that biosurfactant produced by *Ochrobactrum anthropi* HM-1 culture showed a clear halo zone 38.5 cm^2 , while 33.17 cm^2 was presented by *Citrobacter freundii* HM-2 biosurfactant. The cell-free culture broths of

HM-1 and HM-2 strains successfully emulsified sunflower oil with approximately 70% and 60%, respectively.

Surface tension determination

Surface tension is a key parameter for the evaluation of biosurfactant production. In fact, a microorganism is considered as a promising biosurfactant producer, if it could reduce the surface tension to less than 40 mN m^{-1} (Shete et al., 2006). The obtained results showed that our biosurfactant is able to reduce surface tension until $30.73 \text{ mN m}^{-1} \pm 0.48$ which is lower than results obtained by Ghazala et al. (2017) during the characterization of an anionic lipopeptide produced by *Bacillus mojavensis* I4 where the surface tension of the culture supernatant was $31.5 \pm 0.8 \text{ mN m}^{-1}$. Moreover, our results are very close to those obtained by Jemil et al., (2016) which showed that the best result in decreasing surface tension was observed with *Bacillus methylotrophicus* DCS1 strain (31 mN m^{-1}). Likewise, other study showed that biosurfactants produced by *O. anthropi* HM-1 and *C. freundii* HM-2 were able to reduce surface tension until 30.8 ± 0.6 and $32.5 \pm 1.3 \text{ mN m}^{-1}$, respectively (Ibrahim 2018). While, compared to surface tensions of some chemical surfactants studied by Ghazala et al. (2017), *B. safensis* F4 cell free broth showed lower surface tension than SDS ($34.8 \pm 1.3 \text{ mN m}^{-1}$) and Triton X-100 ($32 \pm 0.9 \text{ mN m}^{-1}$).

Characterization of *B. safensis* F4 biosurfactant

TLC analysis showed that *B. safensis* F4 biosurfactant is a lipopeptide. The relative front (R_f) value was 0.56 (Fig. 2) which confirmed that the biosurfactant extract is a lipopeptide as reported by similar previous studies (Fernandes et al. 2007).

In order to identify our biosurfactant, the acetonitrile fraction was collected, and then analysed by LC-MS (Liquid chromatography–mass spectrometry) (Fig. 3). The details of the

obtained masses have been identified according to previous reported studies. Results showed the presence of two surfactinderivates ($M+H^+=1022.6668$ and $1008.6513/ M-H+=1020.6579$ and 1006.6436 , respectively) at retention time of 15.98 min with the presence of adducts ($M+Na$) (Table 2). The presence of surfactin was confirmed by the positive and negative ionizationmode (Jasim et al. 2016). At the same retention time of 15.98 min, the two compounds were identified as Leu/Ile-7, C_{14} surfactin and Leu/Ile-7, C_{13} surfactin with different masses of 1021.66 m/z and 1007.65 m/z (Price et al. 2007). Another peak at 9.20 min has been depicted ($M-H^+ = 329.2328$), which could correspond to pinellic acid. According to literature, pinellic acid is mainly known with its anti-allergic (Arulselvan et al. 2016) and anti-inflammatory (Nagai et al. 2004) activities.

Determination of minimum inhibitory concentration (MIC)

Das et al. (2008) reported that some types of biosurfactants produced by many *Bacillus* species present antimicrobial activity against many bacteria including pathogenic strains.

Our lipopeptide showed limited activity against Gram-negative bacteria compared to that obtained against the Gram-positive tested strains. The tested biosurfactant has a MIC of 0.78 mg mL^{-1} against *B. subtilis* and 1.56 mg mL^{-1} against *S. aureus*, *E. faecium* and *M. luteus*. However, it presented a MIC value of 3.125 mg mL^{-1} against *A. tumefaciens*, *S. enterica*, *E. coli* and 1.56 mg mL^{-1} against *P. savastanoi* (Table 1). Moreover, Singh and Cameotra (2004) reported that the lipopeptide produced by *B. subtilis* C1 was found to be active against several Gram-positive bacteria.

In another study, biosurfactant produced by the *Lactobacillus paracasei* ssp. *paracasei* A20 showed significant antimicrobial activities against pathogenic *E. coli*, *S. aureus* with MIC higher values ranging between 25 and 50 mg mL^{-1} comparing with our results (Gudiña et al. 2010 b). Likewise, a high level of growth inhibition was observed against different

pathogens with a biosurfactant produced by *Lactobacillus helveticus* at a concentration of 25 mg mL⁻¹ (Sharma and Saharan 2016). Furthermore, many lipopeptides produced by *Bacillus licheniformis*, (Yakimov et al. 2007; Fiechter 1992) and *B. subtilis* (Vollenbroich et al. 1997) were known by their important antimicrobial activities. In other studies, the crude biosurfactant produced by *Lactobacillus jensenii* presented approximately 100% activity against *E. coli*, and *S. aureus* with a MIC of 50 mg mL⁻¹ which are higher than our MIC values (Sambanthamoorthy et al. 2014).

Concerning the anti-planktonic activity, the crude biosurfactant and the acetonitrile fraction were tested against *S. epidermidis* S61. Results showed that the crude biosurfactant and acetonitrile fraction effectively inhibited its growth with MIC of 12.5 mg mL⁻¹ and 6.25 mg mL⁻¹, respectively. However, biosurfactant produced by *L. helveticus* showed a high percentage of growth inhibition (98.4%) against *S. epidermidis* with a concentration of 25 mg mL⁻¹ (Sharma and Saharan 2016).

Anti-adhesive activity

The ability of the crude biosurfactant and the acetonitrile fraction to inhibit the early biofilm formation at various concentrations was carried out against *S. epidermidis* S61. According to the Figure 4, the crude biosurfactant and the tested fractions significantly ($P < 0.001$) inhibited the biofilm formation with approximately the same percentages of 90% and 80% at the concentrations of 10 and 5 mg mL⁻¹, respectively. However, at the concentration of 2.5 mg mL⁻¹, the acetonitrile fraction, containing the surfactin, showed higher anti-adherence activity with a percentage of inhibition of 64% against 53% of the crude biosurfactant. Comparing with our results, the purified biosurfactant produced by *B. cereus* NK1 presented lower percentages of biofilm inhibition of *S. epidermidis* at the ratios of 33.55% and 26.46% at concentrations of 10 and 5 mg mL⁻¹, respectively (Sriram et al. 2011). In similar studies, the

anti-adhesive activity of *B.methylophilus* DCS1 crude lipopeptide was evaluated against different strains using biosurfactant pre-treated polystyrene surfaces. Results showed that the highest anti-adhesive effect was observed against *C. albicans* with an inhibition percentage of about 89.3% when biosurfactant was applied at a concentration of 1 mg mL⁻¹ (Jemil et al., 2017). In another study, the crude biosurfactant isolated from *L. paracasei ssp. paracasei* A20 inhibited the adherence of *S. epidermidis* at the concentration of 50 mg mL⁻¹ with a percentage of 72.9% (Gudiña et al. 2010 b) compared to 90 and 80% at 10 and 5 mg mL⁻¹ respectively of our present biosurfactant.

Moreover, biosurfactant produced by *L. helveticus* showed a potential anti adhesive activity against *S.epidermidis* with a percentage of 85% which is similar to our results but at higher concentration of 25 mg mL⁻¹ (Sharma and Saharan 2016). Furthermore, biosurfactants produced by *L. jensenii* and *L. rhamnosus* presented anti-adhesive and anti-biofilm activities against the pathogen strains *A. baumannii*, *E. coli*, and *S. aureus* at concentrations ranging between 25 and 50 mg mL⁻¹ (Sambanthamoorthy et al. 2014).

The anti-adherence activity of the two extracts was confirmed by fluorescence microscopy. The images of the acridine orange staining treated slides with extracts showed the reduction in the biofilm covered surface compared to the control (Fig. 5). Lipopeptides are able to decrease biofilm surface and interfacial tension (Zhao et al. 2017). Previous studies demonstrated that biosurfactants had the ability to alter the surface characteristics of bacterial cells and reduce their adhesive properties. In fact, the application of biosurfactant to a substratum surface can decrease its hydrophobicity, interfere with the microbial adhesion and microorganisms adsorption process (Rodrigues et al. 2006a).

Cytotoxicity assays

363 The cytotoxicity assay of the crude biosurfactant and the acetonitrile fraction was performed
 364 against T47D breast cancer cells and B16F10 mouse melanoma cells.

365 Figures 6 showed that the crude biosurfactant and the acetonitrile fraction showed high
 366 inhibition against T47D and B16F10 cells at 10 mg mL⁻¹ ($P < 0.001$). Furthermore, at the
 367 concentration of 1 mg mL⁻¹, the acetonitrile fraction was more toxic against B16F10 cells
 368 with a survival of 59.75% than T47D cells, whereas, at concentration of the 0.1 mg mL⁻¹, both
 369 tested samples did not show any toxicity against both cell lines.

370 The acetonitrile fraction inhibited significantly cancer cell growth at almost all the tested
 371 concentrations ($P < 0.01$). It presented an IC₅₀ of 1.17 mg mL⁻¹ and 0.66 mg mL⁻¹ against
 372 B16F10 cells and T47D cells, respectively (Fig. 7).

373 These results can be correlated with the composition of the acetonitrile fraction, which mainly
 374 consists of surfactin, belonging to lipopeptides. According to literature, lipopeptides can act
 375 as antitumor agents (Rodrigues et al. 2006b).

376 Previous studies reported that a biosurfactant extracted from *Lactobacillus casei* showed anti-
 377 proliferative potencies against an epithelial cell line with an IC₅₀ (The half-maximal inhibitory
 378 concentration) ranging from 109.1 ± 0.84 mg mL⁻¹ to 129.7 ± 0.52 mg mL⁻¹ (Merghni et al.
 379 2017) which are higher than the IC₅₀ values obtained by our biosurfactant.

380 Moreover, it was previously demonstrated that surfactin could disrupt the membrane structure
 381 via two main mechanisms which are insertion into lipid bilayers, modification of membrane
 382 permeabilization via channel formation or diffusion of ions across the membrane barrier and
 383 membrane solubilization by a detergent-like mechanism (Deleu et al. 2013, Wu et al.
 384 2017). Interestingly, Gudiña et al. (2016) reported that the surfactin anticancer activity is in
 385 relation with its hydrophobic nature. In fact, the fatty acid moiety of surfactin strongly
 386 interacts with the acyl chain of the phospholipids in order to penetrate the outer sheet of lipid

bilayer, while the peptide moiety interacts with the polar head group of the lipids in cancer cells.

Conclusion

In the present study, the best producing biosurfactant strain has been screened and selected. Termed *B. safensis* F4, it is a lipolytic bacterial strain that has the propriety to produce surfactin with important surface-active properties. Crude and purified biosurfactant showed important antibacterial activity under planktonic conditions, preventing also bacterial adherence through inhibiting early stage biofilm formation. Interestingly, surfactin from *Bacillus* sp. F4 has potent cytotoxic activity against cancer cell lines, T47D breast cancer cells and B16F10 mouse melanoma cells. These findings make this studied surfactin a good candidate for potential applications in preventing infectious diseases and treating cancer.

Acknowledgement

The authors are grateful to Dr. Dorra Driss for her valuable critical reading of the manuscript. This work was funded by the Tunisian Ministry of Higher Education and Scientific Research.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Alajlani M, Shiekh A , Hasnain S, Brantner A (2016) Purification of Bioactive Lipopeptides Produced by *Bacillus subtilis* Strain BIA. *Chromatographia* 79: 1527-1532. doi: 10.1007/s10337-016-3164-3
- Arulselvan P, Fard MT, Tan WS, Gothai S, Fakurazi S, Norhaizan ME, Kumar SS (2016) Role of Antioxidants and Natural Products in Inflammation. *Oxid Med Cell Longev*. doi:10.1155/2016/5276130

410 Beenken KE, Blevins JS, Smeltzer MS (2003) Mutation of *sarA* in *Staphylococcus aureus*
 411 limits biofilm formation. *Infect Immun* 71: 4206-4211. doi: 10.1128/IAI.71.7.4206–
 412 4211.2003

413 Bezza FA, Chirwa EMN (2016) Pyrene Biodegradation Enhancement Potential of
 414 Lipopeptide Biosurfactant Produced by *Paenibacillus dendritiformis* CN5 Strain. *J Hazard*
 415 *Mater.* doi:10.1016/j.jhazmat.2016.08.035

416 Cherif S, Mnif S, Hadrich F, Abdelkafi S, Sayadi S (2011) Strategy for improving
 417 extracellular lipolytic activities by a novel thermotolerant *Staphylococcus* sp. strain. *Lipids*
 418 *Health Dis* 10: 209- 216. doi:10.1186/1476-511X-10-209

419 Christova N, Lang S, Wray V, Kaloyanov K, Konstantinov S, Stoineva I (2015) Production,
 420 Structural Elucidation, and In Vitro Antitumor Activity of Trehalose Lipid Biosurfactant from
 421 *Nocardia farcinica* Strain. *J Microbiol Biotechnol* 25:439-447. doi: 10.4014/jmb.1406.06025

422 Colla LM, Rizzardi J, Pinto MH, Reinehr CO, Bertolin TE, Costa JA (2010) Simultaneous
 423 production of lipases and biosurfactants by submerged and solid-state bioprocesses.
 424 *Bioresour Technol* 101:8308–8314. doi:10.1016/j.biortech.2010.05.086

425 Cooper DG, Goldenberg BG (1987) Surface-Active Agents from Two *Bacillus* Species. *Appl*
 426 *Environ Microb* 53: 224-229

427 Das P, Mukherjee S, Sen R (2008) Antimicrobial potential of a lipopeptide biosurfactant
 428 derived from a marine *B. circulans*. *J Appl Microbiol* 104:1675-1684. doi: 10.1111/j.1365-
 429 2672.2007.03701.x

430 Deleu M, Lorent J, Lins L, Brasseur R, Braun N, El Kirat K, Nylander T, Dufrêne YF,
 431 Mingeot-Leclercq M-P (2013) Effects of surfactin on membrane models displaying lipid
 432 phase separation. *BBA - Biomembranes* 1828: 801–815. doi:10.1016/j.bbamem.2012.11.007

433 Fernandes PA, Arruda IR, Santos AFA, Araújo AA, Maior AMS, Ximenes EA (2007)
 434 Antimicrobial activity of surfactants produced by *Bacillus subtilis* R14 against multidrug-
 435 resistant bacteria. *Braz J Microbiol* 38:704-709. doi: 10.1590/S1517-83822007000400022
 436 Fiechter A (1992) Biosurfactants: moving towards industrial application. *Trends Biotechnol*
 437 10: 208- 217. doi: 10.1016/0167-7799(92)90215-H
 438 Gargouri B, Contreras MM, Ammar S, Segura-Carretero A, Bouaziz M (2016) Biosurfactant
 439 production by the crude oil degrading *Stenotrophomonas* sp. B-2: chemical characterization,
 440 biological activities and environmental applications. *Environ Sci Pollut Res* 24: 3769–3779.
 441 doi: 10.1007/s11356-016-8064-4
 442 Ghazala I, Bouassida M, Krichen F, Benito JM, Ellouz-Chaabouni S, Haddar A (2017)
 443 Anionic lipopeptides from *Bacillus mojavensis* I4 as effective antihypertensive agents:
 444 Production, characterization, and identification. *Eng Life Sci* 0: 1–10. doi:
 445 10.1002/elsc.201700020
 446 Girard L, Blanchet É , Intertaglia L , Baudart J , Stien D , Suzuki M , Lebaron P and Lami R
 447 (2017) Characterization of N-Acyl Homoserine Lactones in *Vibrio tasmaniensis* LGP32 by a
 448 Biosensor-Based UHPLC-HRMS/MS Method. *Sensors* 17: 906-918. doi: 10.3390/s17040906
 449 Gudiña EJ, Teixeira JA, Rodrigue LR (2010a) Isolation and functional characterization of a
 450 biosurfactant produced by *Lactobacillus paracasei*. *Colloid Surface B* 76: 298–304. doi:
 451 10.1016/j.colsurfb.2009.11.008
 452 Gudiña EJ, Rocha V, Teixeira JA, Rodrigues LR (2010b) Antimicrobial and antiadhesive
 453 properties of a biosurfactant isolated from *Lactobacillus paracasei*ssp. *paracasei* A20. *Lett*
 454 *Appl Microbiol* 50 : 419-424. doi : 10.1111/j.1472-765X.2010.02818.x

455 Gudiña EJ, Teixeira JA, Rodrigues LR (2016) Biosurfactants produced by marine
 456 microorganisms with therapeutic applications. *Mar Drugs* 14: 38-52. doi:
 457 10.3390/md14020038
 458 Ibrahim HMM (2018) Characterization of biosurfactants produced by novel strains of
 459 *Ochrobactrum anthropi* HM-1 and *Citrobacter freundii* HM-2 from used engine oil-
 460 contaminated soil. *Egyptian Journal of Petroleum* 27: 21-29. doi: 10.1016/j.ejpe.2016.12.005
 461 Janek T, Łukaszewicz M, Rezanka T, Krasowska A (2010) Isolation and characterization of
 462 two new lipopeptide biosurfactants produced by *Pseudomonas fluorescens* BD5 isolated from
 463 water from the Arctic Archipelago of Svalbard. *Bioresource Technol* 101: 6118–6123. doi:
 464 10.1016/j.biortech.2010.02.109
 465 Janek T, Łukaszewicz M, Krasowska A (2013) Identification and characterization of
 466 biosurfactants produced by the Arctic bacterium *Pseudomonas putida* BD2. *Colloid Surface B*
 467 110: 379-386. doi: 10.1016/j.colsurfb.2013.05.008
 468 Jardak M, Abdelli F, Laadhar R, Lami R , Stien D , Aifa S, Mnif S (2017) Evaluation of
 469 biofilm-forming ability of bacterial strains isolated from the roof of an old house. *J Gen Appl*
 470 *Microbiol* 63: 186-194. doi: 10.2323/jgam.2016.10.005
 471 Jasim B, Sreelakshmi KS, Mathew J, Radhakrishnan EK (2016) Surfactin, Iturin, and
 472 Fengycin Biosynthesis by Endophytic *Bacillus* sp. from *Bacopamonneri*. *MicrobEcol* 72:
 473 106–119. doi: 10.1007/s00248-016-0753-5
 474 Jemil N, Ben Ayed H, Hmidet N, Nasri M (2016) Characterization and properties of
 475 biosurfactants produced by a newly isolated strain *Bacillus methylotrophicus* DCS1 and their
 476 applications in enhancing solubility of hydrocarbon. *World J MicrobiolBiotechnol* 32:175-
 477 187. doi 10.1007/s11274-016-2132-2

478 Jemil N, Ben AyedH , Manresa A, Nasri M, Hmidet N (2017) Antioxidant properties,
 479 antimicrobial and anti-adhesive activities of DCS1 lipopeptides from *Bacillus*
 480 *methylophilus* DCS1. BMC Microbiol 17:144-154.doi :10.1186/s12866-017-1050-2

481 Mathur T, Singhal S, Khan S, Upadhyay DJ, Fatma T, Rattan A (2006) Detection of biofilm
 482 formation among the clinical isolates of staphylococci: an evaluation of three different
 483 screening methods. Indian J MedMicrobi. 24: 25-29. doi: 10.4103/0255-0857.19890

484 Merghni A, Dallel I , Noumi E , Kadmi Y, Hentati H , Tobji S, Ben Amor A , Mastouri M
 485 (2017) Antioxidant and antiproliferative potential of biosurfactants isolated from
 486 *Lactobacillus casei* and their anti-biofilm effect in oral *Staphylococcus aureus* strains.
 487 MicrobPathog 104: 84-89. doi: 10.1016/j.micpath.2017.01.017

488 Morikawa M, Daido H, Takao T, Murata S, Shimonishi Y, Imanaka T (1993) A new
 489 lipopeptide biosurfactant produced by *Arthrobacter* sp. strain MIS38. J Bacteriol 175: 6459-
 490 6466. doi: 10.1128/jb.175.20.6459-6466.1993

491 Moryl M, Spętana M, Dziubek K, Paraszkiwicz K, Różalska S, Płaza GA, Różalski A (2015)
 492 Antimicrobial, antiadhesive and antibiofilm potential of lipopeptidessynthesised by *Bacillus*
 493 *subtilis*, on uropathogenic bacteria. ActaBiochimPol62:725-732.doi:
 494 10.18388/abp.2015_1120

495 MosmannT (1983) Rapid colorimetric assay for cellular growth and survival: application to
 496 proliferation and cytotoxicity assays. J Immunol Methods 65: 55-63. doi: 10.1016/0022-
 497 1759(83)90303-4

498 Nagai T, Arai Y, Emori M , Nunome S, Yabe T, Takeda T, Yamada H (2004) Anti-allergic
 499 activity of a Kampo (Japanese herbal) medicine “Sho-seiryu-to (Xiao-Qing-Long-Tang)” on
 500 airway inflammation in a mouse model. IntImmunopharmacol 4:1353 -1365. doi:
 501 10.1016/j.intimp.2004.05.021

502 Naruya S, Nei M (1987) The neighbor-joining method: a new method for reconstructing
 503 phylogenetic trees. *MolBiolEvol* 4: 406-425. doi: 10.1093/oxfordjournals.molbev.a040454
 504 Padmavathi AR, Pandian SK (2014) Antibiofilm activity of biosurfactant producing coral
 505 associated bacteria isolated from Gulf of Mannar, *IndianJ Microbiol* 54: 376–382. Doi:
 506 10.1007/s12088-014-0474-8
 507 Price NPJ, Rooney AP, Swezey JL, Perry E, Cohan FM (2007) Mass spectrometric analysis
 508 of lipopeptides from *Bacillus* strains isolated from diverse geographical locations. *FEMS*
 509 *Microbiol Lett* 271: 83-89. doi: 10.1111/j.1574-6968.2007.00702.x
 510 Rodrigues L, Banat IM, Vandermis HC, Teixeira JA, Oliveira R (2006a) Interference in
 511 adhesion of bacteria and yeasts isolated from explanted voice prostheses to silicone rubber by
 512 rhamnolipid biosurfactants. *J Appl Microbiol* 100: 470–480. doi:10.1111/j.1365-
 513 2672.2005.02826.x
 514 Rodrigues L, Banat IM, Teixeira J, Oliveira R (2006b) Biosurfactants: potential applications
 515 in medicine. *J Antimicrob Chemother* 57: 609-618. doi: 10.1093/jac/dkl024
 516 Sambanthamoorthy K, Feng X, Patel R, Patel S, Parnavitana C (2014) Antimicrobial and
 517 antibiofilm potential of biosurfactants isolated from lactobacilli against multi drug resistant
 518 pathogens. *BMC Microbiol* 14: 197- 205. doi: 10.1186/1471-2180-14-197
 519 SekhonKK, Khanna S, Cameotra SS (2011) Enhanced biosurfactant production through
 520 cloning of three genes and role of esterase in biosurfactant release. *Microb Cell Fact* 10: 49-
 521 58. doi: 10.1186/1475-2859-10-49
 522 Sekhon KK, Khanna S, Cameotra SS (2012) Biosurfactant Production and Potential
 523 Correlation with Esterase Activity. *J Pet EnvironBiotechnol* 3: 7- 16. doi: 10.4172/2157-
 524 7463.1000133

525 Sharma D, Saharan BS (2016) Functional characterization of biomedical potential of
 526 biosurfactant produced by *Lactobacillus helveticus*. Biotechnology Reports 11: 27-35. doi:
 527 10.1016/j.btre.2016.05.001

528 Sen S, Borah SN, Bora A, Deka S (2017) Production, characterization, and antifungal
 529 activity of a biosurfactant produced by *Rhodotorulababjevae* YS3. Microb Cell Fact 16:95-
 530 108. doi: 10.1186/s12934-017-0711-z

531 Shete AM, Wadhawa G, Banat IM, Chopade BA (2006) Mapping of patents on bioemulsifier
 532 and biosurfactant: A review. J Sci Ind Res 65: 91–115.

533 Silva NRA, Luna MAC, Santiago ALCMA, Franco LO, Silva GKB, Souza PM, Okada K,
 534 Albuquerque CDC, Silva CAA, Takaki GMC (2014) Biosurfactant-and-bioemulsifier
 535 produced by a promising *Cunninghamella echinulata* isolated from Caatinga soil in the
 536 northeast of Brazil. Int J Mol Sci 15:15377-15395. doi: 10.3390/ijms150915377

537 Singh P, Cameotra SS (2004) Potential applications of microbial surfactants in biomedical
 538 sciences. Trends Biotechnol 22:142-146. Doi:10.1016/j.tibtech.2004.01.010

539 Sriram MI, Kalishwaralal K, Deepak V, Gracerosep R, Srisakthi K, Gurunathan S (2011)
 540 Biofilm inhibition and antimicrobial action of lipopeptide biosurfactant produced by heavy
 541 metal tolerant strain *Bacillus cereus* NK1. Coll Surf B 85:174–181. doi:
 542 10.1016/j.colsurfb.2011.02.026

543 Ülker S, Karaoglu S A (2012) Purification and characterization of an extracellular lipase from
 544 *Mucorhiemalis f. corticola* isolated from soil. J BiosciBioeng 114:385-390. Doi:
 545 10.1016/j.jbiosc.2012.04.023

546 Vasudevan P, Nair M K, Annamalai T, Venkitanarayanan K S (2003) Phenotypic and
 547 genotypic characterization of bovine mastitis isolates of *Staphylococcus aureus* for biofilm
 548 formation. Vet Microbiol 92:179-185. doi: 10.1016/S0378-1135(02)00360-7

549 Vollenbroich D, Pauli G, Ozel M, Vater J (1997) Antimycoplasma Properties and Application
 550 in Cell Culture of Surfactin, a Lipopeptide Antibiotic from *Bacillus subtilis*. Appl Environ
 551 Microb 63: 44-49
 552 Wilson K (1987) Preparation of genomic DNA from bacteria, in Ausubel F M, Brent R,
 553 Kingston R E, Moore D D, Seidman J G, Smith J A, Struhl K (ed.), Current protocols in
 554 molecular biology, John Wiley and Sons, New York, 2003, p2.4.1.-2.4.2
 555 Wu YS, Ngai SC, Goh BH, Chan KG, Lee LH, Chuah LH (2017) Anticancer Activities of
 556 Surfactin and Potential Application of Nanotechnology Assisted Surfactin Delivery. Front.
 557 Pharmacol 8:761. doi: 10.3389/fphar.2017.00761
 558 Yakimov MM, Timmis KN, Golyshin PN (2007) Obligate oil-degrading marine bacteria.
 559 Curr Opin Biotech 18: 257-266. doi: 10.1016/j.copbio.2007.04.006
 560 Zhao H, Shao D, Jiang C, Shi J, Li Q, Huang Q, Rajoka MSR, Yang H, Jin M (2017)
 561 Biological activity of lipopeptides from *Bacillus*. Appl Microbiol Biotechnol 101: 5951–5960.
 562 doi: 10.1007/s00253-017-8396-0
 563
 564
 565