



**HAL**  
open science

## **Long-term changes in greenhouse gas emissions from French agriculture and livestock (1852–2014): From traditional agriculture to conventional intensive systems**

Josette Garnier, Julia Le Noë, Audrey Marescaux, Alberto Sanz-Cobena, Luis Lassaletta, Marie Silvestre, Vincent Thieu, Gilles Billen

### ► To cite this version:

Josette Garnier, Julia Le Noë, Audrey Marescaux, Alberto Sanz-Cobena, Luis Lassaletta, et al.. Long-term changes in greenhouse gas emissions from French agriculture and livestock (1852–2014): From traditional agriculture to conventional intensive systems. *Science of the Total Environment*, 2019, 660, pp.1486-1501. 10.1016/j.scitotenv.2019.01.048 . hal-02139279

**HAL Id: hal-02139279**

**<https://hal.sorbonne-universite.fr/hal-02139279>**

Submitted on 24 May 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Long-term changes in greenhouse gas emissions from French agriculture and livestock (1852–2014): From traditional agriculture to conventional intensive systems

Josette Garnier <sup>a,\*</sup>, Julia Le Noë <sup>a</sup>, Audrey Marescaux <sup>a</sup>, Alberto Sanz-Cobena <sup>b</sup>, Luis Lassaletta <sup>b</sup>, Marie Silvestre <sup>c</sup>, Vincent Thieu <sup>a</sup>, Gilles Billen <sup>a</sup>

<sup>a</sup> SU CNRS EPHE UMR 7619 Metis, BP 123, Tour 56-55, Etage 4, 4 Place Jussieu, 75005 Paris, France


<sup>b</sup> CEIGRAM-Agricultural Production, Universidad Politécnica de Madrid, Ciudad Universitaria Madrid, E-28040 Madrid, Spain

<sup>c</sup> SU CNRS FR3020 FIRE, BP 123, Tour 56-55, Etage 4, 4 Place Jussieu, 75005 Paris, France

## HIGHLIGHTS

- French GHG emissions from agricultural and forest sectors were estimated.
- A long-term trajectory (1852–2014) was reconstructed for N<sub>2</sub>O, CH<sub>4</sub>, CO<sub>2</sub>.
- GHG emissions have grown four-fold since 1852, to 120,000 CO<sub>2</sub> Eq yr<sup>-1</sup> in the 2000s.
- GHG emissions have only stabilised, in spite of agro-environmental measures.
- Deep changes in the agro-food system would reduce agricultural GHG emissions.

## GRAPHICAL ABSTRACT



## ABSTRACT

France was a traditionally agricultural country until the first half of the 20th century. Today, it is the first European cereal producer, with cereal crops accounting for 40% of the agricultural surface area used, and is also a major country for livestock breeding with 25% of the European cattle livestock. This major socioecological transition, with rapid intensification and specialisation in an open global market, has been accompanied by deep environmental changes. To explore the changes in agricultural GHG emissions over the long term (1852–2014), we analysed the emission factors of N<sub>2</sub>O from field experiments covering major land uses, in a gradient of fertilisation and within a range of temperature and rainfall, and used CH<sub>4</sub> emission coefficients for livestock categories, in terms of enteric and manure management, considering the historical changes in animal excretion rates. We also estimated indirect CO<sub>2</sub> emissions, rarely accounted for in agricultural emissions, using coefficients found in the literature for the dominant energy consumption items (fertiliser production, field work and machinery, and feed import). From GHG emissions of ~30,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup> in 1852, reaching 54,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup> in 1955, emissions more than doubled during the ‘Glorious thirties’ (1950–1980), and peaked around 120,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup> in the early 2000s. For the 2010–2014 period, French agriculture GHG emissions stabilised at ~114,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup>, distributed into 49% methane (CH<sub>4</sub>), 22% carbon dioxide (CO<sub>2</sub>) and 29% nitrous oxide (N<sub>2</sub>O). A regional approach through 33 regions in France shows a diversity of agriculture reflecting the hydro-ecoregion distribution and the agricultural specialisation of local areas. Exploring

### Keywords:

French agriculture  
GHG emissions  
Historical reconstruction  
Spatial distribution  
Scenario analysis

\* Corresponding author.

E-mail address: josette.garnier@upmc.fr (J. Garnier).

## 1. Introduction

Atmospheric greenhouse gas (GHG) concentrations such as carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) have rapidly increased, especially since the mid-20th century, as shown by ice core and modern data (IPCC, 2008). At the global scale, agricultural activities and land use changes accounted for 24% of the total emissions (IPCC, 2008) (not including indirect emissions such as those linked to fertiliser manufacture). For the European Union (EU-28), agricultural emissions are considered as the second largest GHG contributors after fossil fuel combustion (respectively, 10% vs. 78% of total EU emissions, amounting 4300 million tons CO<sub>2</sub> equivalent, in 2016) (EEA, 2018). For France (out of the 520 million tons CO<sub>2</sub> equivalent yr<sup>-1</sup>), GHG from agriculture and fossil fuel combustion represented 19% vs. 71%, respectively in 2010, with a decreasing trend since the Kyoto protocol (CITEPA, 2012). Fugitive emissions from fuels represented only 1% of the total GHG emissions, industrial processes accounted for about 3.7% and wastes for 4.8% (CITEPA, 2012). Surprisingly, according to CITEPA, whereas 66% and 87% of CH<sub>4</sub> and N<sub>2</sub>O emissions are issued from French agriculture, agriculture is not considered as a source of CO<sub>2</sub>, its emissions being included in others sectors (machinery, chemicals, ...). Therefore, the role of agriculture in overall GHG emissions, including direct and indirect CO<sub>2</sub> emissions linked to fossil fuel-driven farming practices still present a number of uncertainties (Lemke et al., 2007).

The control factors of agricultural N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> emissions differ greatly from each other. N<sub>2</sub>O fluxes are often reported to be associated with mineral fertiliser applications (Bouwman, 1996; Skiba et al., 1996; Smith et al., 1997; ENA, 2011), particularly on wet non-saturated soils (Clayton et al., 1994; Aguilera et al., 2013), but manure and other organic fertilisers also contribute to N<sub>2</sub>O emissions (Aguilera et al., 2013). CH<sub>4</sub> fluxes come essentially from livestock (Moss et al., 2000; Vermorel et al., 2008; Springmann et al., 2018), manure management and enteric fermentation (particularly ruminants), although some soils can be a CH<sub>4</sub> sink through CH<sub>4</sub> oxidation (Boeckx and Van Cleemput, 2001), a capacity that managed agricultural soils have partly lost due to nitrogen fertiliser application, which is unfavourable to methane oxidizing micro-organisms (Ojima et al., 1993; Dobbie and Smith, 1996). In contrast, paddy soils could be net emitters of this gas. While net soil CO<sub>2</sub> emissions result from the balance between humified organic matter input and mineralisation or leaching, CO<sub>2</sub> emissions by agriculture stem more from CO<sub>2</sub> emitted from the fossil fuel use related to fertiliser manufacture, use of machinery for farm work and feed imports (Gingrich et al., 2007; Dyer et al., 2010; Aguilera et al., 2015). How these different control factors combine with each other to determine the variations in time and space of GHG emissions by agricultural systems remains a question that is difficult to answer.

France is currently the world's fourth largest agricultural exporter ([www.fao.org/faostat](http://www.fao.org/faostat)), with a rather diversified mosaic of regional agricultural systems. It therefore constitutes a good case study for analysing the relationships between the structural characteristics of agriculture and its GHG emissions.

Based on the concepts of socio-ecological trajectories (Fischer-Kowalski and Haberl, 2007) and territorial ecology (Barles, 2010; Barles, 2017), the long-term trends of the French agro-food system have been described by Le Noé et al. (2018) in terms of N, P and C fluxes over the period from 1852 to 2014. A gradual intensification and specialisation of regional systems was shown, all characterised by integrated crop and livestock farming until the beginning of the 20th century, toward either specialised cropping systems fueled by synthetic


fertilisation or intensive livestock farming highly dependent on external feed imports (Le Noé et al., 2016). Within the country, the Seine watershed is one emblematic example of the former specialised cropping systems, while Bretagne is a region of typical intensive livestock farming systems highly disconnected from croplands. Marescaux et al. (2018) established the GHG budget for the Seine Basin, including the hydrosystem network, the agricultural and non-agricultural sectors. Of the approximately 61,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup> emitted from the whole Seine Basin, non-agriculture GHGs were shown to dominate the total emissions (73%), while the agricultural sector amounted to 23% and emissions by rivers reached 4%. The agricultural emissions found for the Seine Basin were 30% higher than those provided by official French GHG emission inventories.

The first aim of this paper is to establish a spatially distributed long-term budget of GHG emissions by the French agricultural sector. A major issue behind this effort is to identify the effect of the structural changes from mixed crop and livestock farming systems to specialised systems on the GHG emissions account. Indeed, the shift of agricultural and livestock management practices which occurred with mechanisation and intensification are expected to cause increased GHG emissions. Moreover, specific emission rates of CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O are expected to depend either on crop or livestock typologies. Another objective is to determine the levers for future mitigation of N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> emissions by agricultural practices. In this line, we explored two contrasting scenarios recently developed by Billen et al. (2018): (i) continuing the current trends of specialisation into either cropping systems based on chemically synthesised inputs or intensive livestock farming highly dependent on feed import; and (ii) shifting to organic farming and reconnection of crop and livestock farming, while reducing the animal proteins in human diets by half. These two prospective scenarios of French agriculture are tested in order to evaluate to which extent less intensive agriculture and livestock breeding can allow a reduction of GHG emissions.

## 2. Material and methods

### 2.1. Major physiographic and agricultural characteristics of France

France is a heterogeneous country with a mountainous region in the South-East and low relief in the West (Bretagne) as well as large sedimentary plains in the lower part of the Seine, Loire and Garonne rivers flowing to the Atlantic façade. From north to south, there is a climate gradient: temperate, oceanic, temperate warm, and Mediterranean climatic zones (Fig. 1a). These regions are dominated by field crops in the South-West and North-West parts of the country, with Bretagne dominated by intensive livestock farming (Fig. 1b). The rest of France (in the South-East) is either characterised by large vineyard domains in the areas bordering the Mediterranean Sea or in the Rhone alluvial corridor, and by varied mixed crop and livestock in the mountainous regions (the Alps, the Jura and the Massif Central). With 33 regions defined by aggregation of the 94 metropolitan départements (Le Noé et al., 2017), we defined three supra-regions, with a similar surface area, with homogenous agricultural patterns and climate for each: (i) the Seine Basin, due to its intensive cereal cropping agriculture and temperate climate, (ii) the oceanic-temperate Great West, including the Bretagne region and the lower Loire basin with intensive livestock, and the Great South-West covering the Garonne basin, with a temperate warm climate, particularly conducive to growing maize for feed purposes (Fig. 1c).



**Fig. 1.** General characteristics for France of a. climate from hydroecoregion; b. agricultural patterns; c. homogenous agricultural regions (33) and perimeter of three supra-regions chosen for their identical surface area and their trend for agricultural specialisation.

## 2.2. Reconstruction of past land use and agricultural system

### 2.2.1. The GRAFS approach for characterising the agricultural structure over the long term

The GRAFS approach (Generalized Representation of Agro-Food System), firstly developed and applied on a global scale (Billen et al., 2013, 2014; Lassaletta et al., 2014a) to local scales (Garnier et al., 2016) for nitrogen (N) circulation, was enlarged to phosphorus (P) (Garnier et al., 2015) and now to N, P and carbon (C), (Le Noë et al., 2017). Briefly, the GRAFS approach describes the agro-food system of a given geographical area by considering four main compartments exchanging nutrient flows: cropland, grassland, livestock system and local population. The potential losses to the environment associated with these exchanges have been estimated for hydrosystems (Garnier et al., 2015, 2016, 2018), but not yet for losses to the atmosphere. This functional representation links arable land productivity, semi-natural or managed grassland that contributes to livestock feeding, and finally, human food requirements. The agro-food system is driven by (i) nutrient inputs to the soil (synthetic and/or organic fertilisation, atmospheric deposition [hereafter referred to as exogenous fertilisation] as well as symbiotic fixation), (ii), the size of the livestock and its feed requirement and (iii) the size of the human population and its dietary preferences, including feed and food imports/exports. The GRAFS approach does not take into account forested areas.

Based on the long-term agricultural statistics available at the scale of the 94 French administrative “département” units for metropolitan France (equivalent to the European Union NUTS3 statistical division) and additional data gathering (e.g. the forested areas below), the GRAFS approach was documented for 22 dates from 1852 to 2014, and for the 33 French regions (Le Noë et al., 2017, 2018), (Fig. 1).

### 2.2.2. Past evolution of forested areas

The surface forest and wooded areas were documented by “département” for the years 1929 (Ministère Agriculture – Enquête Agricole), 1946, 1950, 1955, 1960 and 1965 (Ministère Agriculture – SAA). All years were available from 1970 to 1988 based on data provided by Agreste-SSP. The recent years from 1989 to 2014 were retrieved from the AGRESTE database on agricultural statistics (<https://stats.agriculture.gouv.fr/disar-web/>). For France, whereas the total surface areas amounted to  $10.67 \cdot 10^6$  ha in 1929, examining the chronicle by Cinotti (1996) for the 19th century showed values around  $9.3 \cdot 10^6$  ha in 1850 (compared to the  $10.67 \cdot 10^6$  ha in 1929) and a linear trend for the intermediate dates, 1885 and 1906. The same changes were applied to the 33 regions, for the dates before 1929.

### 2.3. Reconstruction of greenhouse gas emissions

As a whole, the GHG budget of the agriculture sector can be estimated as the result of  $N_2O$  emissions from cropped soils, grassland and forest,  $CH_4$  released from livestock production (enteric fermentation and manure management) and  $CO_2$  emitted from fossil fuel directly used by farming practices as well as indirectly for manufacture and transport of agricultural inputs. Due to their specific origin, each of these GHG emissions required an adapted methodology. Agricultural soil C sequestration has been estimated elsewhere and will be compared to GHG emissions (Le Noë et al., 2019).

#### 2.3.1. Reconstruction of $N_2O$ emissions

To infer  $N_2O$  emissions back to 1852 at the scale of the 33 regions of France, on the basis of the recent knowledge gained from field measurements, we first established an empirical relationship linking yearly  $N_2O$  emissions to mineral and organic fertilisation, temperature and rainfall, and then assumed that this relationship could be extrapolated to past

situations within the timeframe of this study (e.g. Bouwman et al., 2013).

For N<sub>2</sub>O emissions, a literature review from Garnier et al. (2009) and Cayuela et al. (2017) was completed for a total of 208 yearly cropland N<sub>2</sub>O emissions values and their associated explicative variables, namely N fertiliser (organic and mineral) inputs, annual mean temperature and rainfall. A similar set of data was gathered for 138 cases of grasslands. For forests, we found 48 cases for N<sub>2</sub>O emissions that were only associated with temperature and rainfall data, since they had not been fertilised. Additional early field measurements from the Seine watersheds were also included in this analysis (Benoit et al., 2015a, 2015b). The complete data set is presented in the supplementary material (SM1).

We searched the best fit parameters for the following relationship:

$$N_2O_{em} = (a + b N_{inp}^d) * (Rain/Rain_{ref})^c * Q_{10}^{T/10}$$

where N<sub>2</sub>O emissions (N<sub>2</sub>O<sub>em</sub>) and N inputs (N<sub>inp</sub>) are in kg N<sub>2</sub>O-N ha<sup>-1</sup> yr<sup>-1</sup>, rainfall (Rain) in mm yr<sup>-1</sup> and temperature (T) in °C (Table 1), and a, b, c, Rain<sub>ref</sub>, d and Q<sub>10</sub> are parameters to be calibrated within a range of realistic values.

This relationship assumes a power function for the relationship with N inputs and rainfall, and a classical Q10 exponential relationship with temperature.

Since most grassland was fertilised, no significant difference was found between cropland and grassland so that the two data series were merged. Regarding forests, N<sub>inputs</sub> are restricted to atmospheric deposition and a specific relationship was established (Table 1).

The six parameter values (a, b, c, d, Rain<sub>ref</sub> and Q10) were determined by a systematic optimisation procedure searching the combination of parameter values providing the best fit of the calculated

**Table 1**

a. Summary of the data gathered for establishing relationships between N<sub>2</sub>O emissions and its controlling factors, n = 394, number of data. Relationships, and associated parameter values: b. for cropland and grassland and c. for forest. NRMSE and bias are calculated for evaluation of the fitted relationships.

a.					
N = 394	Nb of values	N inputs, kgN ha <sup>-1</sup> yr <sup>-1</sup>	Rainfall, mm yr <sup>-1</sup>	Temperature, °C	N <sub>2</sub> O emission, kgN-N <sub>2</sub> O ha <sup>-1</sup> yr <sup>-1</sup>
Cropland	208	0-450	327-1250	2.75-18.5	0.01-11.0
Grassland	138	0-753	400-1837	1.00-16.0	-0.50-18.9
Forest	48	0	607-1239	3.60-10.1	0.17-4.9

b.			
N <sub>2</sub> O <sub>em</sub> = (a + b N <sub>inp</sub> <sup>d</sup> ) * (Rain / Rain <sub>ref</sub> ) <sup>c</sup> * Q <sub>10</sub> <sup>T/10</sup>			
	Units	Value	± step
a	kgN/ha/yr	0.15	0.05
b	dimless	0.016	0.001
c	dimless	1.2	0.05
d	dimless	1.0	0.1
Rain <sub>ref</sub>	mm yr <sup>-1</sup>	1000	100
Q <sub>10</sub>	dimless	1.2	0.2
nRMSE = 14%			
bias = 5%			

c.			
N <sub>2</sub> O <sub>em</sub> = a * (Rain / Rain <sub>ref</sub> ) <sup>c</sup> * Q <sub>10</sub> <sup>T/10</sup>			
	Units	Value	± step
a	kgN/ha/yr	1.9	0.05
c	dimless	1.2	0.1
Rain <sub>ref</sub>	mm yr <sup>-1</sup>	1400	100
Q <sub>10</sub>	dimless	1.2	0.2
nRMSE = 23%			
bias = 10%			

emissions to the observed N<sub>2</sub>O emission values. The resulting relationship fits the data with an acceptable % bias (ratio of mean calculated values to mean observed values) and Normalised RMSE (root mean square error normalised against the range of observed values) (Table 1). Further details on the procedure are provided in Supplementary material (SM2).

These relationships were applied to cropland and grassland on one hand and to forest on the other hand for each region and 22 dates from 1852 to 2014. Temperature and rainfall were reconstructed from EOPS data for the 1950-2017 period (Version 17, 0.25 degrees resolution (<https://www.ecad.eu/download/ensembles/download.php>) and spatially averaged by French "département" (NUTS-3 equivalent). For the period prior to 1950, temperature and rainfall data were downloaded for 28 towns spread over the country, back to the dates available in the past, between 1850 and 2017 (<http://meteo-climat-bzh.dyndns.org/meteo100-1783-2018-3-tn-1-0-0.php>). To avoid any discrepancy between the two series, the anomaly for each town was calculated compared to the mean over the long-term period and then added to the mean calculated for the 1950-2017 period. The values prior to 1950 for the towns were then assigned to their respective regions. The same procedure was applied for both temperature and rainfall. The average differences between the two data series for the 1950-2017 period for all 33 regions is close to zero (i.e., no systematic bias) with a standard deviation of 15%, for temperature and rainfall.

### 2.3.2. Reconstruction of CH<sub>4</sub> emissions

CH<sub>4</sub> emissions were estimated based on livestock numbers and specific emission factors for each animal and age class category, corrected for past variations in excretion rates.

Current CH<sub>4</sub> emission factors (kg-CH<sub>4</sub> head<sup>-1</sup> year<sup>-1</sup>) from enteric fermentation and manure storage and management were taken from Garnier et al. (2013, see Table 1SM) compiled principally from Vermorel et al. (2008), IPCC (1997) and Zhou et al. (2007). An emission factor for humans was also taken into account, following Crutzen et al. (1986).

These CH<sub>4</sub> emissions concerned five animal sub-categories for cattle, three for sheep, three for pig, five for poultry, while goat, horse and rabbit represented one category each. Knowing the number of heads per category and the associated manure produced, the total amount of CH<sub>4</sub> emitted by livestock was calculated using the corresponding specific emission factors.

Livestock numbers per category were taken directly from agricultural statistics (Agreste, or Gallica (<https://gallica.bnf.fr>) when these numbers were not available from Agreste). However, the evolution of animal size and physiology changed over the period studied (Chatzimpiros, 2011) and this must be taken into account. Historical variations of excretion rates of the major livestock categories were established by Le Noë et al. (2018, SM1 & SM2) (Table 2). The correction factors found during the period for the animal categories reported in Table 2 were also used for their corresponding sub-categories. For poultry and rabbit, no change was considered over time.

**Table 2**

Empirical relationship for calculating excretion rates (y, in kgN head<sup>-1</sup> yr<sup>-1</sup>) of cattle, sheep, pig and horses as a function of time (t, in year) over the period 1850-2014, and corresponding values of the parameters b, a, a', t<sub>max</sub> and dt, calibrated against historical and current animal excretion data.

(From Le Noë et al., 2018.)

	b	a	a'	t <sub>max</sub>	dt
	kgN head <sup>-1</sup> yr <sup>-1</sup>	kgN head <sup>-1</sup> yr <sup>-1</sup>	yr	yr	yr
Cattle	45	0.05	65	2010	40
Sheep	4	0.02	9	2020	45
Pig	56	0.034	-	-	-
Horse	480	0.3	-	-	-

General formula:  $y = b + a(t - 1850) + a' \exp[-(t - t_{max})^2 / dt^2]$ .

### 2.3.3. Reconstruction of CO<sub>2</sub> emissions

Current CO<sub>2</sub> emissions by direct or indirect fossil fuel combustion by the agricultural sector were calculated using the official French CLIMAGRI approach (Doublet, 2011), based on CO<sub>2</sub> emission factors calculated for mechanised field work and livestock activities, as well as for synthetic fertiliser manufacture. The coefficients used are gathered in Table 3.

The coefficients related to mechanisation (fossil fuel combustion for machinery, field work in cropland and grasslands, and livestock management and feed to livestock) were applied pro-rata to the usable agricultural area or the total livestock units for each region, and extrapolated to the past taking into account the degree of mechanisation of each region. The proxy for establishing this degree of mechanisation was based on the observed evolution of the numbers of horses between 1906 (zero mechanisation) and 1980 (100% mechanisation) (Fig. 2).

### 2.4. Exploring scenarios


The two contrasting scenarios recently developed by Billen et al. (2018) are explored herein in terms of GHG emissions for their divergent assumptions: one continuing the trends of **O**pening to distant markets and **S**pecialisation into either cropping systems based on synthesised external inputs or intensive livestock farming (O/S), the other with an agricultural system shifting to **A**utonomy through organic farming, crop and livestock **R**econnection and a **D**emitarian diet (A/R/D). The O/S scenario reflects a main stream vision mainly driven by the desire for economic growth, very present in the official discourse, which is accompanied by the strengthening of territorial specialisation in a globalised economy, the continuation of agricultural intensification and the concentration of the population in large cities. On the contrary, the A/R/D scenario assumes a radical rupture toward agro-ecology (Billen et al., 2018) and organic farming, searching for the autonomy of farmers with respect to agricultural inputs such as N fertilisers and animal feed, and reduction of the share of animal protein in the human diet as recommended by, for example, WHO for health reasons and Springmann et al. (2018) for environmental reasons, all trends already detectable although far from being fully underway. We here assumed reducing the animal proteins in human diets by half following the so-called demitarian diet (see the Barsac declaration, in 2009; <https://en.wikipedia.org/wiki/Demitarian>). Both scenarios were tested for the 2040 horizon and we therefore considered an increase of 1.5 °C for both scenarios in addition to the 1.5 °C already observed since the end of the 19th century. We kept the rainfall as it is at present because no general trend has been observed, in spite of yearly oscillations over the studied period. These projections are well in the ranges of those reported in Jouzel et al. (2014) for France for 2050, with four different models. Changes in rainfall might be more visible at the seasonal scale, not investigated here, with wetter winters and drier summers, and increased occurrence of extreme weather events in all seasons. Regarding

**Table 3**

Coefficient applied for the calculations of CO<sub>2</sub> emissions according to major emitter sectors. N fertilisers and P fertilisers concern the production of fertilisers actually used, feed to livestock is the imported feed, machinery corresponds to the manufacture of agricultural equipment, energy for cropland, grassland and livestock, the fuel or electricity necessary for fieldwork and for livestock breeding. (From Doublet, 2011.)

Major sectors emitting CO <sub>2</sub>	Units	Coefficients
Fertilisers N	tonC-CO <sub>2</sub> /tonN	1.12
Fertilisers P	tonC-CO <sub>2</sub> /tonP	0.46
Feed to livestock	tonC-CO <sub>2</sub> /tonN imported	1.339
Machinery	tonC-CO <sub>2</sub> /ha/yr	0.026
Energy for cropland	tonC-CO <sub>2</sub> /ha UAA/yr	0.077
Energy for grassland	tonC-CO <sub>2</sub> /ha UAA/yr	0.055
Energy for livestock	tonC-CO <sub>2</sub> /LU/yr	0.056

UAA: utilised agricultural area; LU: livestock unit.



**Fig. 2.** Evolution of the coefficient of mechanisation from 1906 (zero mechanisation) to 1980 (100% mechanisation) as calculated from the decreasing number of horses for each of the 33 regions. The coefficient is given here for France and three contrasted regions. Agriculture modernisation appeared earlier in Ile-de-France.

fertilisation, both the O/S and A/R/D scenarios follow current environmental regulations, external synthetic fertilisers will be used according to a yield objective for the former, while the latter banish the use of mineral fertilisers (and pesticides) replaced by biological nitrogen fixation and possibly by on-farm and recirculated external organic inputs. In both scenarios, the livestock species structure was kept identical to the current one (2004–2014) for the regions that already had livestock.

### 2.5. Uncertainty analysis

The GHG emission values calculated as described above result from complex calculations based on basic statistical data (fertilisation rates, livestock number, etc.) and a number of parameters (coefficient of statistical relationships, emission factors, etc.), both subject to a certain level of uncertainty. In order to assess how these uncertainties propagate to the final emission estimates, a bootstrap procedure was carried out, as developed by Le Noë et al. (2018), under Microsoft Excel and associated VBA macros. Shortly, after having stated the confidence interval of all primary data and parameters (typically 10–20% uncertainty was assumed), thousand independent estimations of the GHG emissions were computed with a random draw of each of these data and parameters according to a Monte Carlo sampling within a Gaussian distribution inside the confidence interval. The uncertainty on the final GHG emissions was calculated as the standard error of the mean of these thousand replicates.

## 3. Results


### 3.1. Long-term trends of the control variables of GHG emissions

#### 3.1.1. Temperature

A 1.5 °C increase of mean annual temperature was observed between the mid-19th century and the last decade for all of France, mainly from 1980 when the average temperature exceeded 10 °C (Fig. 3). The same trend was observed for the three selected regions, the Great South-West being closer to the overall average and the Great West showing the largest difference (-0.3 °C) (Fig. 3). The coldest and warmest regions are, respectively, Alpes (7 °C) and Grand Marseille (13 °C) in the South-East of France. Regionally, whereas the temperature increase was >1.7 °C in the East, South and South-West of France, with, respectively, a continental and Mediterranean climate, other mountainous regions (Jura, Savoie, Pyrénées) and oceanic temperate (Bretagne), showed an increase of 1.1 °C.

#### 3.1.2. Rainfall

Average rainfall over the period studied (1852–2014) was 805 mm yr<sup>-1</sup> for the whole of France, more or less 100 mm yr<sup>-1</sup> for



**Fig. 3.** Long-term evolution (1850–2014) for France and the three supra-regions selected in the annual average of a. temperature; b. rainfall. Long-term evolution of major indicators of agriculture in France and the three supra-regions selected for the 22 dates analysed; c. total exogenous N inputs; d. livestock size in terms of livestock units (LU), right axis for France LU.

the Great South-West ( $935 \text{ mm yr}^{-1}$ ) and Seine basin ( $703 \text{ mm yr}^{-1}$ ), respectively, whereas rainfall in the Great West averaged  $746 \text{ mm yr}^{-1}$  (Fig. 3). The wettest regions ( $>1000 \text{ mm yr}^{-1}$ ) were Savoie, Pyrénées Occidentales and Landes and the driest was Ile-de-France ( $\sim 600 \text{ mm yr}^{-1}$ ). No clear trend was observed over the long term, but wet and dry years alternated (Fig. 3).

### 3.1.3. Exogenous fertiliser

This accounts for total N inputs (synthetic, manure and deposition), excluding biological nitrogen fixation. From the 1850s until 1965, fertilisation increased from  $20$  to  $60 \text{ kgN ha}^{-1} \text{ yr}^{-1}$ , and more in the early 1970s when half of the regions received  $100 \text{ kgN ha}^{-1} \text{ yr}^{-1}$ . The inputs plateaued from the late 1980s at about  $135 \text{ kgN ha}^{-1} \text{ yr}^{-1}$  at the national scale. The Northern part of France firstly increased its fertilisation, especially for the fertile soil of Ile-de-France and Nord-Pas-de-Calais, with values exceeding  $150 \text{ kgN ha}^{-1} \text{ yr}^{-1}$  in the 1980s (Fig. 3). There were few regions where fertilisation remained below  $100 \text{ kgN ha}^{-1} \text{ yr}^{-1}$  until now (Savoie, Alpes, Côte d'Azur in South-East France, and Garonne, Pyrénées Orientales in the South-West), mostly mountainous areas. In these regions, mineral fertilisation was less than half the total exogenous N inputs. Before 1965, fertilisation was essentially based on manure.

The same trend was found for the three selected regions, the Seine Basin and Great West fertiliser trajectory being above the average, and the Great South-West below (Fig. 3).

### 3.1.4. Livestock

Livestock density, expressed in livestock units (LU, i.e. equivalent to an animal excreting  $85 \text{ kg N yr}^{-1}$ ) per ha of agricultural surface, increased slowly in all French regions until the mid-20th century. The increase was more pronounced in the 1950–1980 period, except in those regions specialising in stockless crop farming, such as the Seine Basin (Figs. 3, 4). The Great West, specialising in intensive livestock farming, dependent on feed import, showed the highest increase in livestock density (Le Noë et al., 2018). Differences among regions stabilised after the 1990s (Fig. 4).

## 3.2. Agricultural features

We chose three dates (1906 as a reference for traditional agriculture in France, 1970 characterising the beginning of modernisation and 2014

for evidencing the results of specialisation). These agricultural features are mapped for all 33 regions.

### 3.2.1. Cereal crop production and livestock density

The specificity of the French regions was already in place in the early 20th century with the northern half of France producing  $10$ – $25 \text{ kgN ha}^{-1} \text{ yr}^{-1}$  embedded in the harvested grain, similar to two other regions in the South (West: Garonne; East: Grand Marseille). Ile-de-France already distinguished itself as the most productive (Fig. 4a). At this time in 1906, the regions with the highest livestock density were generally associated with those with high cereal production, showing the importance of the crop–livestock connection at that time. Specialisation had already appeared in 1970, with an intensification of cereal production in the Seine Basin and a rise in livestock breeding most particularly in Bretagne and Loire Amont. In 2014, the decoupling between crop production and livestock is striking: one map is almost the opposite of the other (Fig. 4b).


### 3.2.2. Percentage of permanent grassland and forest

Except in the Seine Basin, grassland was present over the entire country in 1906, occupying up to  $60$ – $80\%$  of the total agricultural area in the South-Eastern quadrant of France (Fig. 4c), and in some other regions with (Manche) or without (Pyrénées) high livestock density. In the 1970s, this trend was accentuated, whereas in 2014 grassland was regressing everywhere. The overall forest area increased in proportion throughout the 20th century but more in the South-East of France and the South-Western border of the Massif Central (Fig. 4d).

## 3.3. Distribution of GHG emissions over the long term

### 3.3.1. $\text{N}_2\text{O}$

By construction,  $\text{N}_2\text{O}$  emissions by cropland, grassland and forested areas (the sum of their surface areas forms the rural area, in  $\text{km}^2$ ) reflect the long-term spatial and temporal variations in temperature, rainfall and fertilisation of agricultural land. The highest value ( $75$ – $125 \text{ kg N-N}_2\text{O km}^{-2} \text{ yr}^{-1}$ ) found for Ile-de-France in 1906 was typically related to its early use of mineral fertilisers (Fig. 5a), whereas the values for Côte d'Azur would be more a combination of fertilisation and temperature/rainfall. The same can be said for the North of France region on the one hand and the South and Center of France regions on the other hand, for the  $50$ – $75 \text{ kg N-N}_2\text{O km}^{-2} \text{ yr}^{-1}$  category. For a majority of regions,  $\text{N}_2\text{O}$  emissions showed a great increase in the 1970s, recently



**Fig. 4.** Maps of regional distributions in France for three dates (1906, 1970, 2014) representative of major time periods: a. cereal production; b. livestock density; c. percentage of permanent grassland in total agricultural area; d. percentage of forest in rural area (i.e. cropland + grassland + forest). UAA, utilised agriculture area. A: Alpes; Al: Alsace; AL: Aveyron-Lozère; AR: Ain-Rhône; B: Bourgogne; Br: Bretagne; C-A-Y: Champagne-Ardennes-Yonne; CC: Cantal-Corrèze; Cda: Côte d'Azur; CO: Calvados-Orne; DL: Dordogne-Lot; E: Eure; E&L: Eure-et-Loire; Gar: Garonne; Gd J: Grand Jura; Gd M: Grand Marseille; Gde L: Grande Lorraine; G-H: Gard-Hérault; Gir: Gironde; I-D-A: Isère-Drôme-Ardèche; IdF: Ile de France; L Am: Loire Aumont; L Av: Loire Aval; Lan: Landes; LC: Loire Centrale; M: Manche; N-PdC: Nord Pas-de-Calais; Pic: Picardie; Pocc: Pyrénées Occidentales; POR: Pyrénées Orientales; S: Savoie; VC: Vendée-Charentes.

reaching or exceeding  $250 \text{ kg N-N}_2\text{O km}^{-2} \text{ yr}^{-1}$  in 2014, i.e. doubling over the 20th century.

### 3.3.2. $\text{CH}_4$

The distribution of C- $\text{CH}_4$  emissions in time and space, even more than the livestock density map, clearly reflects the patterns of intensification and specialisation that persisted for the whole period studied (Fig. 5b). In brief, a large ring of high emissions was emerging in 1906 around the Seine Basin and was accentuated in 1970 and even more in 2014, when livestock density was reduced giving way to intensive cropping. From emission values generally lower than  $2000 \text{ kgC-CH}_4 \text{ km}^{-2} \text{ yr}^{-1}$  in 1906, they increased to above  $6000 \text{ kgC-CH}_4 \text{ km}^{-2} \text{ yr}^{-1}$  in the regions specialising in intensive livestock farming.


### 3.3.3. $\text{CO}_2$

The highest emissions of  $\text{CO}_2$  in 1906 are directly linked to the small amount of fossil fuels used for mineral fertilisers because no

mechanisation and no feed import were considered (Fig. 5c). The 1970 map resulted from the post Second World War (WWII) increase of all items taken into account in the calculation (fertiliser production, fossil fuels used for field work, machinery and feed import) the proportion of which was modulated from one region to another according to their rate of modernisation and specialisation. For example, in 1970 feed import concerned essentially Bretagne, Nord-Pas-de-Calais and Loire Centrale, also affected by high fertilisation and mechanisation. In 2014 the highest emissions in Bretagne typically related to increasing feed imports from South America (Billen et al., 2011; Le Noé et al., 2016).

### 3.4. Estimating GHGs under contrasting scenarios

For the O/S scenarios all GHG emissions would increase considerably (Fig. 6). The approximately 20% increase of  $\text{N}_2\text{O}$  emissions for the entire French territory would not be related to nitrogen synthetic fertilisers,



**Fig. 5.** Maps of regional distributions in France for three dates (1906, 1970, 2014) representative of major time periods of GHG emissions per km<sup>2</sup> of rural areas (cropland + grassland + forest) and per year: a. N<sub>2</sub>O; b. CH<sub>4</sub>; c. CO<sub>2</sub>. See legend of Fig. 4 for the names of the regions.

which are not increasing significantly owing to environmental regulations, but rather to manure application in livestock farming regions (Billen et al., 2018). The Seine Basin, as well as other regions such as Alsace and Landes, emptied of their livestock under extreme specialisation into stockless crop farming, would emit only very low CH<sub>4</sub> while the rest of France would show as high emissions as the most current emitting regions (>6000 kgC-CH<sub>4</sub> km<sup>-2</sup> yr<sup>-1</sup>) (Fig. 6) because they have reached the maximum authorised livestock density (European Nitrate Directive (91/676/CEE), i.e. 2 LU ha<sup>-1</sup> of agricultural area). CO<sub>2</sub> emissions would concomitantly increase mostly in breeding areas, due to feed imports, necessary after a loss in grassland and increased livestock density, in addition to C-CO<sub>2</sub> from fuel consumption and fertiliser production.

Under the A/R/D scenario, the reduction of GHG emissions would allow a “return” to the emissions before the heavy industrialisation of the 1970s and even 1955 (see Fig. 7) depending on the regions. The A/R/D scenario showed that (i) autonomy with respect to synthetic mineral fertilisers would significantly reduce N<sub>2</sub>O emissions, (ii) the decrease in animal loading to meet local feed autonomy would lower CH<sub>4</sub> emissions and (iii) CO<sub>2</sub> emissions would be also considerably lowered due to lack of both synthetic fertiliser use and feed imports (Fig. 6). This would allow France to achieve GHG emission targets committed to within the COP-21.


The results of these two scenarios for the three GHGs were put into perspective with their respective historical trajectory for the whole nation and for the three regions selected to distinguish the origin of emissions. We thus corroborate our choice of the three dates illustrating

three periods: 1906 for its traditional agriculture (until 1955), the 1970s for this period’s transitional status, and 2014, typical of intensive production and specialisation since the 1980s.

Whatever the date, croplands were the major N<sub>2</sub>O emitters compared to forests and grasslands, especially in the Seine Basin and the Great West (Fig. 7), whereas the O/S scenario would not significantly change the pattern and the level of N<sub>2</sub>O emissions for these two emblematic specialised regions. The Great South-West, assumed to develop high livestock density, hence imported feed, while optimising its crop production in this scenario, would not only increase N<sub>2</sub>O but all three GHG emissions. Concerning CH<sub>4</sub>, enteric fermentation dominated CH<sub>4</sub> emission, which overall accounted for 70% of the total, and manure (30%) increased in proportion in the O/S scenario for the Great West, traditionally livestock oriented, but also in the Great South-West, as shown by the fourfold increased CH<sub>4</sub> emissions. For CO<sub>2</sub>, concomitantly and consequently, imported feed was the dominant emitter sector in the Great West, which would remain so in the O/S scenario, differently from the Great South-West which, as for cropping, revealed its potential for more livestock, and imported feed, with more CO<sub>2</sub> emissions (Fig. 7).

### 3.5. Comparative agricultural GHG emissions in CO<sub>2</sub> equivalent

The GHG emission can be calculated in a single unit (CO<sub>2</sub> equivalent, CO<sub>2</sub> Eq) by multiplying the global warming potential (GWP) of the different sources with their respective emissions. CO<sub>2</sub> taken as a reference has a GWP of 1, whereas N<sub>2</sub>O and CH<sub>4</sub> GWPs are,



**Fig. 6.** Maps of regional distributions in France for present (2014) and for two prospective scenarios (O/S: opening and specialisation and A/R/D: autonomy, livestock reconnection and demitarian diet) of GHG emissions per km<sup>2</sup> and per year: a. N<sub>2</sub>O; b. CH<sub>4</sub>; c. CO<sub>2</sub>. See legend of Fig. 4 for the names of the regions.

respectively, 265 and 28 times the CO<sub>2</sub> GWP using the 100-year GWP (IPCC, 2014). Total agricultural emissions for France amounted to 113,939 ktons CO<sub>2</sub> Eq yr<sup>-1</sup> for the 2010s, with 49% CH<sub>4</sub>, 29% N<sub>2</sub>O and 22% CO<sub>2</sub> (Fig. 8a; Table 4). Regarding the three selected regions, similar in area, the Great West emissions were twice those of the other two, with the highest proportion in CH<sub>4</sub> (59%). N<sub>2</sub>O emission dominated in the intensively cropped Seine Basin (37%) and CH<sub>4</sub> in the Great South-East (48%). Whereas the O/S scenario would increase overall GHG emissions almost 1.5 times for France, the increase would be about 1.2 for the Seine Basin and the Great West, but emissions would explode for the Great South-West with a 2.5-fold increase. With no mineral fertilisation and extensive livestock breeding, the A/R/D scenario would lower emissions by 50% for France, 36% and 76% for the Seine Basin and the Great West, respectively, but emissions for the Great South-West would only decrease by 11% (Fig. 8b-d; Table 4). Note that, taking into account a 20-year GWP, i.e., 265 for N<sub>2</sub>O and 84 for CH<sub>4</sub> (IPCC, 2014), would double the total emissions at the scale of France, CH<sub>4</sub> then representing about 75% of the total for the 2010–2014 period, and up to 80% for both scenarios.

When expressed per km<sup>2</sup> of surface area, agricultural GHG emissions ranged from 148 to 408 tCO<sub>2</sub> eq km<sup>-2</sup> yr<sup>-1</sup> for the Great South West and Great West regions respectively, the GHG for the Seine Basin being close to the national value (Table 4). For comparison, agricultural GHG emissions were 3.4, 2.4, 1.3 times higher than the French ones for the Netherlands, Belgium and Germany, but 2.0, 1.8, 1.7, 1.6 times lower for Spain, Romania, Austria and Slovenia, respectively (EAA, 2018).

#### 4. Discussion

Greenhouse gas emission estimates are known to be highly uncertain, due to the complexity of their controlling factors at the local scale and the difficulty of measuring them on-site as well as due to the lack of uniform information on how highly diverse farming practices may impact net GHG balances. Consequently, the scientific community is encountering problems formalising mechanistic models to estimate GHG emissions (e.g. NOE, DNDC, Gu et al., 2014; Gilhespy et al., 2014; Zimmermann et al., 2018), for upscaling local emissions measurements to larger scales, from local areas to countries. Since the level of detail needed for a wide regional and long-term estimation could make the modelling unfeasible, the approach used here is more a budgeting one, based on activity data from agricultural census and emission factors determined on a rather detailed analysis of the controlling factors of N<sub>2</sub>O and CH<sub>4</sub>. For these two GHGs, our calculations allow linking plot-scale measurements or animal physiology to the regional and country scale. CO<sub>2</sub> emission estimates are still rough, although explicitly taking into account several contributions, mostly upstream of the production system, but in line with the Climagri approach for France (Doublet, 2011) and other studies (e.g. Canada, Dyer and Desjardins, 2009, and Spain, Aguilera et al., 2015).

##### 4.1. Controlling factors of GHG emissions

To explore strategies to reduce GHGs from agriculture, identifying, quantifying and understanding the factors which control their





Fig. 7. Long-term evolution (1852–2014) for France (a.) and the three selected supra-regions (b., c., d.) in annual average of  $N_2O$ ,  $CH_4$  and  $CO_2$  from top to bottom.  $N_2O$  is represented for the three major land uses (forest, grassland and cropland),  $CH_4$  for manure and enteric emissions, and  $CO_2$  as energy for imported feed, fertiliser production and other energy, including field work, machinery and livestock breeding. Error bars provide uncertainties as calculated by the Monte Carlo analysis.



**Fig. 8.** Long-term evolution (1852–2014) for France (a.) and the three selected supra-regions (b., c., d.) in annual average of N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub>, expressed in CO<sub>2</sub> equivalents per year. Error bars provide uncertainties as calculated by the Monte Carlo analysis.

producing processes, from biotic to abiotic, from microbial to industrial, are challenging.

#### 4.1.1.1. N<sub>2</sub>O

In temperate systems, denitrification is considered the major microbial process at the origin of N<sub>2</sub>O emissions in soils, mostly under temperate and highly humid conditions (Benoit et al., 2015b; Vilain et al.,

**Table 4**

Relative contributions of N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> and total GHG emissions from agriculture in France and in the three supra-regions considered, given in CO<sub>2</sub> equivalent. Specific GHG emissions (tCO<sub>2</sub> km<sup>-2</sup> yr<sup>-1</sup>) are provided for comparison.

CO <sub>2</sub> Eq		2000–2014	O/S scn	A/R/D scn
France 540,498 km <sup>2</sup>	% N <sub>2</sub> O	29	23	25
	% CH <sub>4</sub>	49	57	58
	% CO <sub>2</sub>	22	20	17
	ktCO <sub>2</sub> yr <sup>-1</sup>	113,939	165,010	56,678
	tCO <sub>2</sub> km <sup>-2</sup> yr <sup>-1</sup>	211	305	105
Seine Basin 69,713 km <sup>2</sup>	% N <sub>2</sub> O	37	42	20
	% CH <sub>4</sub>	32	21	62
	% CO <sub>2</sub>	31	37	18
	ktCO <sub>2</sub> yr <sup>-1</sup>	12,909	10,872	8246
	tCO <sub>2</sub> km <sup>-2</sup> yr <sup>-1</sup>	185	156	118
Great West 59,109 km <sup>2</sup>	% N <sub>2</sub> O	20	15	23
	% CH <sub>4</sub>	59	65	57
	% CO <sub>2</sub>	21	20	19
	ktCO <sub>2</sub> yr <sup>-1</sup>	24,123	29,471	6397
	tCO <sub>2</sub> km <sup>-2</sup> yr <sup>-1</sup>	408	499	108
Great South-West 65,437 km <sup>2</sup>	% N <sub>2</sub> O	29	23	25
	% CH <sub>4</sub>	48	60	59
	% CO <sub>2</sub>	23	17	17
	ktCO <sub>2</sub> yr <sup>-1</sup>	9662	23,937	8582
	tCO <sub>2</sub> km <sup>-2</sup> yr <sup>-1</sup>	148	366	131

2014; Gu et al., 2014). In contrast, nitrification plays an important role in Mediterranean and semi-arid regions (e.g., Sanz-Cobena et al., 2012; Aguilera et al., 2013). The factors affecting denitrification in soils have been largely reported in the literature (see Tiedje, 1988; Groffman, 1991; de Klein et al., 2001, for example). Following Saggari et al. (2013), who distinguished two types of controlling factors, namely (i) soils and plant factors (crops, soil mineral nitrogen, pH, water content and oxygen, carbon availability, C:N ratios) and (ii) environmental factors (temperature, rainfall, drying/wetting vs freezing/thawing), the variables we considered for our prediction relationships (N input, land use and climatic conditions [temperature and rainfall]) covered these two types of factors. Generally considered to be the major controlling factors of N<sub>2</sub>O production in soils, under both nitrifying and denitrifying conditions (e.g. Snyder et al., 2009; Smith, 2010; Kirschbaum et al., 2012; Sanz-Cobena et al., 2017), these variables are the best documented in the literature for achieving such a long-term study.

Our approach is a step further in the use of the Tier 1 emission factor (IPCC, 2008), which has been found too coarse for correctly representing N<sub>2</sub>O emissions possibly underestimated in wet areas (Lu et al., 2006 and reference herein) but overestimated in Mediterranean regions (Aguilera et al., 2013; Cayuela et al., 2017). The approach we developed here is advantageous for its low requirement in complex agronomic data, given that temperature and rainfall are routinely surveyed everywhere and fertilisation commonly quoted in agricultural statistics.

However, taking into account the annual scale of our study, pulses of N<sub>2</sub>O emissions cannot be represented as observed after fertiliser applications (Bouwman, 1996; Hénault et al., 1998; Laville et al., 2011; Plaza-Bonilla et al., 2014; Benoit et al., 2015a; Recio et al., 2018), rainy events (Zheng et al., 2000; Lu et al., 2006; Beare et al., 2009) and during freeze and thaw periods and rewetting in semi-arid regions (Vilain et al., 2010; Lu et al., 2015; Wertz et al., 2016; Sanchez-Martín et al., 2010). However, the N<sub>2</sub>O emissions data gathered here were selected

for their annual representativeness, thus already integrating seasonal variability. Spatial and long-term variations can be expected through the heterogeneous distribution in both rainfall and temperature in the country (cf. Fig. 1a) and fertiliser applications depending on soil characteristics, which condition the technical orientations of the farms (cf. Fig. 1b). N<sub>2</sub>O emissions related to manure management were not taken into account in the calculations, because they account for only a few percent of GHGs emitted from livestock, i.e., 2.94% in CO<sub>2</sub> Eq according to the review by (Zervas and Tsiplakou, 2012).

#### 4.1.2. CH<sub>4</sub>

Methanogenesis is a microbial process producing CH<sub>4</sub> mainly under anaerobic conditions. Enteric fermentation by grazing animals is the major process producing CH<sub>4</sub>, although losses from manure management is far from negligible, far above the net CH<sub>4</sub> flux in the soil. Although agricultural soils can be a sink for this GHG, as a result of the balance between methanogenesis and methanotrophy (microbial CH<sub>4</sub> consumption) (Dutaur and Verchot, 2007; Kirschbaum et al., 2012), this function of oxidation by soils has been reduced with the intensive use of fertilisers. Ammonium inputs, increasing nitrifying activity, would exclude methanotrophs from their ecological niche, which even when fertilisation ceases would not quickly recover (Lemke et al., 2007; Ball et al., 2002; Boeckx and Van Cleemput, 2001; Schnell and King, 1994). In addition, soil cultivation and change in soil structure would restrict diffusion sites, thus affecting CH<sub>4</sub> oxidation (Dobbie and Smith, 1996). Experimental measurements in the Seine Basin showed that CH<sub>4</sub> removal could account for 5.6% of the emissions by livestock (Garnier et al., 2013). The CH<sub>4</sub> depletion rate was the highest for grassland soils (1.27 mg C m<sup>-2</sup> d<sup>-1</sup>) followed by forest (0.7 mg C m<sup>-2</sup> d<sup>-1</sup>) and cropland (0.4 mg C m<sup>-2</sup> d<sup>-1</sup>). Interestingly, the riparian zones, which were alternatively a source or a sink, overall emitted 0.1 mg C m<sup>-2</sup> d<sup>-1</sup> methane.

Considering the low proportion of CH<sub>4</sub> removal by soils compared to the emissions from livestock, even when it was half as much on average than currently for the long 1850–1950 period, we can consider that neglecting soil sink would not lead to overestimation of CH<sub>4</sub> emission in the past. Further CH<sub>4</sub> sink would account for about 2% of N<sub>2</sub>O emissions from soils in CO<sub>2</sub> equivalents (Boeckx and Van Cleemput, 2001).

Enteric fermentation was estimated here to be responsible for 70% of CH<sub>4</sub> emissions in France, 30% originating from manure management, with a similar proportion over the time period studied, resulting in a proportion of enteric fermentation lower than the figure provided globally as being responsible for 83% of CH<sub>4</sub> emissions (Zervas and Tsiplakou, 2012). As reported by these authors, ruminant CH<sub>4</sub> production may depend on many factors including diet (fibrous content of the ration), enteric flora, while more generally CH<sub>4</sub> production is influenced by farm management and the farm production system.

In this study, we did not intend to finely characterise CH<sub>4</sub> production from livestock as reported in the study mentioned above (and references therein) but rather used the IPCC Tier 3 guidelines (IPCC, 1997), according to specific emissions (enteric and manure) for 19 categories of livestock and related to animal body weight over the period studied (see Material and methods section).

#### 4.1.3. CO<sub>2</sub>

An increasing greenhouse effect is currently dominated by the increase in CO<sub>2</sub> concentration derived from fossil-fuel consumption, clearly the compound responsible for the enhanced greenhouse effect (Forster et al., 2007). At the scale of the Seine Basin, CO<sub>2</sub> emissions account for 79% of the total emissions (agricultural and non-agricultural), 72% of which come from non-agricultural sectors (Marescaux et al., 2018). Similar figures were provided by CITEPA for France (i.e. 78% and 75%, respectively, in 2014 (<https://www.citepa.org/fr/activites/inventaires-des-emissions/secten>)). Regarding the three GHGs, the CITEPA figures showed an approximately 20% underestimation for the agricultural sector (90,000 ktons CO<sub>2</sub> eq yr<sup>-1</sup> versus

114,000 ktons CO<sub>2</sub> eq yr<sup>-1</sup> here), due to an underestimation of CO<sub>2</sub>, fertiliser production, and machinery manufacture, for example, accounted for in the industrial sector rather than the agricultural sector.

The coefficients from Doublet (2011) (Table 3) are quite comparable with other approaches (Bochu, 2006; Aguilera et al., 2015). For all of France, we showed that a large amount of CO<sub>2</sub> emissions (35%) is linked to the energy supply for production of commercial chemicals. CO<sub>2</sub> emissions from feed to livestock accounted for 10% for France, but 20% for the specialised Great West. These emissions are generally ignored in most studies, including national inventories. “Other energy” (i.e. fuel for farm fieldwork, electricity for on-farm operations and energy for machinery) appeared to be the largest emitter sector (55%). Farm machinery manufacturing energy, a significant part of total CO<sub>2</sub> emissions from agriculture (13% in 2014 with spatial differences) has been roughly taken into account in this paper, but could be included in future research with greater detail (taking into account for example the number of tractors and their power, etc.). This percentage is lower than the farm machinery percentage reported for US corn farms for all Canada, accounting for 15% of the direct CO<sub>2</sub> emissions in the US (Patzek, 2003) and 19% in Canada (Dyer and Desjardins, 2009).

#### 4.2. Past long-term changes in GHG emissions: practice and land use changes

Agriculture in France until the late 19th and early 20th centuries was characterised by a traditional family-based agriculture, with low external requirements (mechanisation, electrical power, etc.). Whereas the number of farms was around 5.5 million during the second-half of the 19th century, this number decreased to 2.3 million in 1955, with an increase in the size of farms, although small ones still dominated. During this period, only slow changes occurred in terms of the indicators selected, however (see Fig. 3, exogenous fertilisers and livestock including horses). Accordingly, N<sub>2</sub>O and CO<sub>2</sub> emissions remained rather stable while CH<sub>4</sub> increased slowly, due to a slow but general increase of livestock density within integrated crop–livestock farming systems.

A second period can be identified after WWII when indirect energy inputs in agriculture started, the war materials industry and chemicals being reoriented to agricultural goods (e.g., fertiliser production through the Haber-Bosch process and intensification of tractor manufacturing and other agricultural machinery). This initiated 3 decades of continuous growth with cheap fossil energy and expanding global food markets (Dyer and Desjardins, 2009). French farms decreased in number by a factor of five by 1980 (i.e. 1 million), intensified their production, with a strong dependence on mechanisation, industrial fertilisers, pesticides and imported feed. GHGs reached their maximum in the early 1990s, when the Rio conference (1992) alerted the world to the strong deterioration of the environment, including GHG emissions.

Despite continuing intensification and specialisation, the third period, from the mid-1990s to the present, showed a stabilisation in GHG emissions and even a tendency to decrease (N<sub>2</sub>O especially, see Figs. 7 and 8). Even before the Rio conference (1992), the use of mineral fertilisers – a major driver for N<sub>2</sub>O emissions – was reduced in France (see Lassaletta et al., 2014b), since the nitrate directive (91/676/CEE) was promulgated for water resource protection. At this time, the data show that ruminants were reduced by 10%, slightly more in the specialised Great West in livestock breeding (17%). During this period, the number of farms still decreased (664,000 in 2000, 452,000 in 2013), with small farms the most affected (Agreste, 2013). Between 1993 and 2014, we calculated a net loss of 1.2 million ha (1.53 million ha lost as permanent grassland and 0.33 million ha gained in cropland, with forested areas stable during this time).

Land use changes are reported to have modified the flux of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O through altered biogeochemical processes (Forster et al., 2007; Houghton et al., 2012; Kirschbaum et al., 2012; Tate, 2015). For example, according to Kim and Kirschbaum (2015), conversion from natural

forest to cropland or grassland would increase net emissions by  $7.3 \pm 0.6$  or  $5.9 \pm 0.3$  t CO<sub>2</sub> eq ha<sup>-1</sup> y<sup>-1</sup>, respectively, while conversion of cropland or grassland to secondary forest would decrease emissions by  $5.3 \pm 0.9$  or  $3.6 \pm 0.7$  t CO<sub>2</sub> eq ha<sup>-1</sup> y<sup>-1</sup>, i.e. figures on the same order of magnitude, although the delay in losses/recoveries might not be similar.

Between 1929 and 2014, 3.3 and 1.8 million ha of permanent grassland and cropland, respectively, have changed use whereas 7.7 million ha were gained as forested area, representing a total loss of 2.6 million ha of these total rural areas. We considered 1929 as the first date of French border stability after the Alsace and Lorraine regions were reattributed to France in 1918. Most of these land use changes were implicitly taken into account in our calculations. However, the possible loss of sink function by soils for CH<sub>4</sub> under increased fertilisation was not accounted for (Boeckx and Van Cleemput, 2001). We also did not consider wetlands as possible CH<sub>4</sub> emitters, which were lost after their conversion to agricultural and other land uses (Zedler and Kercher, 2005; Verhoeven and Setter, 2010).

#### 4.3. Contrasting scenarios for a possible future

The scenarios (Billen et al., 2018) explored here for GHG emissions were inspired from prospective documents elaborated at different regional, national and European levels (Poux et al., 2005; Solagro, 2014; Poux and Aubert, 2018) and also took into account other published results for the global scale (Schmitz et al., 2012; Erb et al., 2016; Muller et al., 2017).

The pursuit of intensification and specialisation in the O/S scenario, despite taking into account the current regulations, would, at a national scale, increase GHG emissions as a whole, but also each of the three GHGs considered separately. Some regional differences would appear however. Whereas increased N<sub>2</sub>O emissions would be generalized over the 33 regions, CO<sub>2</sub> and especially CH<sub>4</sub> would be lower in the regions dedicated to cereal production, with no livestock as predicted for the Seine Basin. In most other regions, livestock would increase CH<sub>4</sub> and its associated feed production would impact CO<sub>2</sub>, in accordance with the approach. This scenario, although credible taking into account the current trends of opening the agro-food system (Le Noé et al., 2016), would not be desirable for the environment, and specifically here for GHG warming, as stated by the last, recently published IPCC report (2018). This economic desire for growth, which is highly dependent on the international market, facilitating trade exchanges, is fully present in political discourse. This vision of the future of French agriculture devoted to cereal exportation driven by powerful food sectors, animal production oriented to milk powder export in industrial breeding farms and a human diet with a high ratio of animal proteins is in line with many documents coming from professional farmer organisations as well as Harbour Authorities (CRAN, 2006; DREAL HN, 2014; Benhalima, 2015; HAROPA, 2015). Furthermore, the emergence of the bio-fuel industries (mainly ethanol from sugar beet and biodiesel from rapeseed) is a paradigm that agriculture will have to face in future decades, even for energy consumption of the farm itself. It is worth mentioning that in this O/S scenario, the livestock structure was kept constant, since French livestock is traditionally dominated by cattle, despite a trend, at the global scale, to shift animal production from ruminants to non-ruminants, which could possibly reduce CH<sub>4</sub> emissions and CO<sub>2</sub> from feed (Westhoek et al., 2014).

Contrary to this vision of France fully involved in globalisation, the A/R/D scenario (Billen et al., 2018) amplifies the weak signals observed in recent years, showing the emergence of a new relationship to nature for a sustainable development, from which proactive policies could emerge. Due to several food crises (e.g. the mad cow problems in 1996, dioxin contamination in 1999, the *Escherichia coli* outbreak in 2011, among the most publicised events), increasing numbers of consumers are demanding healthy food products. At the same time, farmers exposed to

chemical products are concerned by health issues, victims either themselves or family members and neighbours.

In this context, the A/R/D scenario explores alternative agro-food systems which (i) generalise organic farming practices banning exogenous mineral fertilisers and pesticides, (ii) favour local supply and (iii) decrease animal protein consumption in the diet, while reducing food waste. Although not fully realistic over the short term, this scenario becomes desirable for the health of people and the environment. For this scenario total production of plant and animal products are reduced, but this reduction is compensated by a reduction in the demand associated with dietary changes (see Billen et al., 2018). The A/R/D scenario, despite requiring a deep structural change of the agro-food system, shows a significant reduction of GHGs, i.e. 50% for the French agricultural sector compared to the 2010–2014 period, more than one-third of the total reduction advocated. Interestingly, this scenario, beneficial in terms of GHGs, has also been shown to improve water quality and to prevent the risk of coastal eutrophication (Desmit et al., 2018; Garnier et al., 2018).

#### 4.4. Weaknesses and strengths of the approach

Agricultural statistics have been rather well documented in France since the mid-19th century, at the scale of the French “départements” (Poisvert et al., 2017; Le Noé et al., 2018; see also Garnier et al., 2014, 2016 for local studies), and hence the entire country (Harchaoui and Chatzimpiros, 2018), and provide realistic pictures of the changes occurring in the agricultural sectors for indicators such as fertiliser inputs, crop production, livestock units, etc.

The estimate of GHGs from these data is more ambitious given that no statistics exist for the past at this regional resolution. The task was the most difficult for N<sub>2</sub>O due to variability in the emission factors even for the present. Indeed, direct N<sub>2</sub>O emissions in agricultural fields occur essentially with great spatial and temporal variabilities related to many factors (soil parameters, crop species, agricultural practices, rainfall and water management, amount and type of fertilisation, etc.; Lu et al., 2006; Bouwman et al., 2013; Cayuela et al., 2017). Here, taking into account the accompanying N<sub>2</sub>O data in the literature, and the data available over the long term, previous N<sub>2</sub>O data mining (Garnier et al., 2009) was enlarged for the main land uses (forest, grassland and cropland), gathering temperature, rainfall and fertilisation, three major variables that were analysed for the past. Even though soil parameters were often documented in the papers examined, they were not included due to the lack of a homogeneous description in the studies considered. Because grassland studies, similar to cropland, reported fertiliser applications either through external inputs or grazing animals, N<sub>2</sub>O emissions for these two land uses were grouped under the same relationship. The relationship for forests was established with fewer data. Overall the evaluation of both relationships in terms of NRMSE and bias is quite good.

Regarding CH<sub>4</sub>, the uncertainties may be more related to changes in the size of the animals over time, modifying the figures documenting CH<sub>4</sub> emissions from enteric fermentation and manure management for the present. By applying a factor of change on the basis of a reconstruction of excretion for the major types of animals, we can consider that CH<sub>4</sub> emissions from livestock farming must be robust for the past. CH<sub>4</sub> emissions and oxidation from soils were not taken into account given that soils are known to be low emitters (Garnier et al., 2013; Kandel et al., 2018), which must be true for the period studied, because ecosystems such as wetlands, ponds, etc. had already been considerably reduced in the landscape, for health issues and for conversion of these lands into cultivated areas (Zedler and Kercher, 2005; Verhoeven and Setter, 2010).

The estimates for CO<sub>2</sub> are more indicative, because we only took into account its emissions related to field and on-farm work, fertiliser use, CO<sub>2</sub> emissions from machinery and CO<sub>2</sub> emissions linked to feed

imports, applying coefficients reported in the literature based on a full material flow analysis (Doublet, 2011; see also Aguilera et al., 2015).

However, the approach allows reflecting regional differences and, differently from an overall national estimate, provides a variable range of GHGs related to the details of specific characteristics and possible transitions for each region.

For the whole of France, our approach fits rather well with the national agricultural inventories (Doublet, 2011) reporting ~114,000 ktons CO<sub>2</sub> eq. yr<sup>-1</sup> (2010–2014) emitted from agriculture versus ~103,000 ktons CO<sub>2</sub> eq. yr<sup>-1</sup> (for 2006); the percentages we found for N<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> (29%, 49% and 22%, respectively) can be compared with the 39%, 44% and 17%, respectively, found in Doublet (2011). Interestingly, carbon sequestration by agricultural soils, amounting to 6200 ktons CO<sub>2</sub> eq. yr<sup>-1</sup> for France (Le Noë et al., 2019), accounts for only ~5% of our GHG emissions.

## 5. Conclusions

An analysis of the controlling factors of GHG emissions (land use, rainfall, temperature) together with the GRAFS approach gathering key variables of the agro-food system at any temporal and spatial resolution (fertilisers, livestock, crop production, etc.) made it possible to estimate N<sub>2</sub>O, CH<sub>4</sub>, CO<sub>2</sub> emissions at the scale of France's 33 agricultural regions.

The period studied, from 1852 to 2014, showed that the increase of GHGs can be divided into three major periods: (i) a long period from 1852 to 1955 with family mixed crop–livestock farming systems still in effect, small in size with very little mechanisation, and emissions increasing from ~30,000 to 54,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup>; (ii) a second period of modernisation of French agriculture after WWII, with an increasing dependence on mechanisation, industrial fertilisers, pesticides and imported feed, favoured by fossil energy at a relatively low monetary cost and expanding global food markets, a period corresponding to the maximum of GHG emissions, which reached 110,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup> in the 1990s and 120,000 CO<sub>2</sub> ktons Eq yr<sup>-1</sup> in the early 2000s; (iii) and then stabilisation (around 114,000 ktons CO<sub>2</sub> Eq yr<sup>-1</sup> for 2010–2014) related to the abandonment of government policies directly aimed at intensification, and their replacement by environmental regulations after the Rio conference (1992) and the following protocols (e.g., Kyoto in 1997), which prompted the countries of the world to control or decrease their GHG emissions to avoid the adverse effects of climate change. Progressive spatial specialisation was observed as early as 1906, with regions already characterised by cereal production in the Northern half of France, and in the South-West, others by livestock, such as Bretagne in the West, and still others with a typically low proportion of grasslands, such as in the Seine Basin. This specialisation had an influence on the distribution of emissions in terms of intensity and compounds.

The contrasting scenarios explored showed that a 50% reduction of GHG emissions in the agriculture sector could be achieved with a deep change in the structure of the agro-food system, whereas the pursuit of the present trends of specialisation and intensification associated with international trade, even applying the current environmental regulations, could increase the current GHG emissions by a factor of 1.5.

## Acknowledgements

The study was conducted within the framework of several scientific projects, C-Cascades (funded by Horizon 2020 - Research and Innovation Framework Programme under the Marie Skłodowska-Curie grant agreement No. 643052), HydroGES (financed by the Agency for the Environment and Mastery of Energy, ADEME) and Escapade (supported by the National Agency for Research, ANR). The PIREN-Seine programme and the Fédération Ile-de-France de Recherche pour l'Environnement (FIRE) are also acknowledged for their support. L. Lassaletta acknowledges support from MINECO, Spain, co-funded by the European

Commission ERDF (Ramon y Cajal fellowship, RYC-2016-20269) and from Programa Propio from Universidad Politécnica de Madrid.

## Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2019.01.048>.

## References

- Agreste, 2013. Agreste-Recensements agricoles 1988, 2000, 2010 et enquête structure 2013. <http://agreste.agriculture.gouv.fr/IMG/pdf/Gaf15p019-021.pdf>.
- Aguilera, E., Lassaletta, L., Sanz-Cobena, A., Garnier, J., Vallejo, A., 2013. The potential of organic fertilizers and water management to reduce N<sub>2</sub>O emissions in Mediterranean climate cropping systems. A review. *Agric. Ecosyst. Environ.* 164. <https://doi.org/10.1016/j.agee.2012.09.006>.
- Aguilera, E., Guzmán, G., Alonso, A., 2015. Greenhouse gas emissions from conventional and organic cropping systems in Spain. II. Fruit tree orchards. *Agron. Sustain. Dev.* 35, 725–737. <https://doi.org/10.1007/s13593-014-0265-y>.
- Ball, B.C., McTaggart, I.P., Watson, C.A., 2002. Influence of organic ley–arable management and afforestation in sandy loam to clay loam soils on fluxes of N<sub>2</sub>O and CH<sub>4</sub> in Scotland. *Agric. Ecosyst. Environ.* 90, 305–317. [https://doi.org/10.1016/S0167-8809\(01\)00207-9](https://doi.org/10.1016/S0167-8809(01)00207-9).
- Barles, S., 2010. Society, energy and materials: what are the contributions of industrial ecology, territorial ecology and urban metabolism to sustainable urban development issues? *J. Environ. Plan. Manag.* 53, 439–455. <https://doi.org/10.1080/09640561003703772>.
- Barles, S., 2017. Écologie territoriale et métabolisme urbain: quelques enjeux de la transition socio-écologique. *Revue d'Economie Régionale et Urbaine.* 5, pp. 819–836.
- Beare, M.H., Gregorich, E.G., St-Georges, P., 2009. Compaction effects on CO<sub>2</sub> and N<sub>2</sub>O production during drying and rewetting of soil. *Soil Biol. Biochem.* 41, 611–621. <https://doi.org/10.1016/j.soilbio.2008.12.024>.
- Benhalima, M., 2015. Analyse du système agro-alimentaire de la région Nord-Pas de Calais et ses enjeux sur l'eau. Etudes et Documents n° 125. Commissariat Général au Développement Durable. <http://www.developpement-durable.gouv.fr/IMG/pdf/ED125.pdf>.
- Benoit, M., Garnier, J., Billen, G., 2015a. Nitrous oxide production from nitrification and denitrification in agricultural soils: determination of temperature relationships in batch experiments. *Biochem. Process* 50, 79–85. <https://doi.org/10.1016/j.procbio.2014.10.013>.
- Benoit, M., Garnier, J., Billen, G., Tournebise, J., Gréhan, E., Mary, B., 2015b. Nitrous oxide emissions and nitrate leaching in an organic and a conventional cropping system (Seine basin, France). *Agric. Ecosyst. Environ.* 213, 131–141. <https://doi.org/10.1016/j.agee.2015.07.030>.
- Billen, G., Barles, S., Chatzimpiros, P., Garnier, J., 2011. Grain, meat and vegetables to feed Paris: where did and do they come from? Localising Paris food supply areas from the eighteenth to the twenty-first century. *Reg. Environ. Chang.* <https://doi.org/10.1007/s10113-011-0244-7>.
- Billen, G., Garnier, J., Lassaletta, L., 2013. Modelling the nitrogen cascade from watershed soils to the sea: from regional to global scales. *Philos. Trans. R. Soc. B* 368, 20130123. <https://doi.org/10.1098/rstb.2013.0123>.
- Billen, G., Lassaletta, L., Garnier, J., 2014. A biogeochemical view of the global agro-food system: nitrogen flows associated with protein production, consumption and trade. *Glob. Food Sec.* 3, 209–219. <https://doi.org/10.1016/j.gfs.2014.08.003>.
- Billen, G., Le Noë, J., Garnier, J., 2018. Two contrasted future scenarios for the French agro-food system. *Sci. Total Environ.* 637–638, 695–705. <https://doi.org/10.1016/j.scitotenv.2018.05.043>.
- Bochu, J.-L., 2006. Consommation et efficacité énergétique de différents systèmes de production agricoles avec la méthodologie PLANETE. *Fourrages* 186, 165–177.
- Boeckx, P., Van Cleemput, O., 2001. Estimates of N<sub>2</sub>O and CH<sub>4</sub> fluxes from agricultural lands in various regions in Europe. *Nutr. Cycl. Agroecosyst.* 60, 35–47.
- Bouwman, A.F., 1996. Direct emission of nitrous oxide from agricultural soils. *Nutr. Cycl. Agroecosyst.* 46, 53–70.
- Bouwman, A.F., Beusen, A.H.W., Griffioen, J., et al., 2013. Global trends and uncertainties in terrestrial denitrification and N<sub>2</sub>O emissions. *Philos. Trans. R. Soc. B* 368, 1621. <https://doi.org/10.1098/rstb.2013.0112>.
- Cayuela, M.L., Aguilera, E., Sanz-Cobena, A., Adams, D.C., Abalos, D., Barton, L., Ryals, R., Silver, W.L., Alfaro, M.A., Pappa, V.A., Smith, P., Garnier, J., Billen, G., Bouwman, L., Bondeau, A., Lassaletta, L., 2017. Direct nitrous oxide emissions in Mediterranean climate cropping systems: emission factors based on a meta-analysis of available. *Agric. Ecosyst. Environ.* <https://doi.org/10.1016/j.agee.2016.10.006>.
- Chatzimpiros, P., 2011. Les empreintes environnementales de l'approvisionnement alimentaire: Paris, ses viandes et lait, XIXe-XXIe siècles. (Ph-D). Architecture, aménagement de l'espace. Université Paris-Est (332pp + annexes).
- Cinotti, B., 1996. Evolution des surfaces boisées en France: proposition de reconstitution depuis le début du XIX<sup>e</sup> siècle. *Revue Forestière Française.* XLVIII – 6-547-562.
- CITEPA, 2012. Rapport national d'inventaire pour la France au titre de la convention cadre des nations unies sur le changement climatique et du protocole de Kyoto. Rapport CCNUCC/Mars 2012 (1364pp).
- Clayton, H., Arah, J.R.M., Smith, K.A., 1994. Measurement of nitrous oxide emissions from fertilised grassland using closed chambers. *J. Geophys. Res.* 99, 16599–16607.

- CRAN, Chambre Régionale d'Agriculture de Normandie, 2006. 2020: Que mangerons-nous? [www.normandie.chambagri.fr/eco\\_prosp.asp](http://www.normandie.chambagri.fr/eco_prosp.asp)
- Crutzen, P.J., Aselmann, I., Seiler, W., 1986. Methane production by domestic animals, wild ruminants, other herbivorous fauna, and humans. *Tellus* 38B, 271–284. <https://doi.org/10.3402/tellusb.v38i3-4.15135>.
- de Klein, C.A.M., Sherlock, R.R., Cameron, K.C., van der Weerden, T.J., 2001. Nitrous oxide emissions from agricultural soils in New Zealand – a review of current knowledge and directions for future research. *J. R. Soc. N. Z.* 31, 543–574. <https://doi.org/10.1080/03014223.2001.9517667>.
- Desmit, X., Thieu, V., Dulière, V., Ménesguen, A., Campuzano, F., Lassaletta, L., Sobrinho, J.L., Silvestre, M., Garnier, J., Neves, R., Billen, G., Lacroix, G., 2018. Reducing marine eutrophication may require a paradigmatic change. *Sci. Total Environ.* 635, 1444–1466. <https://doi.org/10.1016/j.scitotenv.2018.04.181>.
- Dobbie, K.E., Smith, K.A., 1996. Comparison of CH<sub>4</sub> oxidation rates in woodland, arable and set aside soils. *Soil Biol. Biochem.* 28, 1357–1365. [https://doi.org/10.1016/S0038-0717\(96\)00152-6](https://doi.org/10.1016/S0038-0717(96)00152-6).
- Doublet, S., 2011. CLIMAGRI: bilan énergies et GES des territoires ruraux, la ferme France en 2006 et 4 scénarios pour 2030. Rapport ADEME. Available at: <http://www.ademe.fr/sites/default/files/assets/documents/climagri-la-ferme-france-en-2006-et-4-scenarios-pour-2030.pdf>.
- DREAL HN, 2014. Scénarios prospectifs de développement de la Vallée de Seine à l'horizon 2040. Etude. <http://www.normandie.developpement-durable.gouv.fr/prospective-vallee-de-la-seine-2040-a250.html>.
- Dutaur, L., Verchot, L.V., 2007. A global inventory of the soil CH<sub>4</sub> sink. *Glob. Biogeochem. Cycles* 21, GB4013. <https://doi.org/10.1029/2006GB002734>.
- Dyer, J.A., Desjardins, R.L., 2009. A review and evaluation of fossil energy and carbon dioxide emissions in Canadian agriculture. *J. Sustain. Agric.* 33, 210–228. <https://doi.org/10.1080/10440040802660137>.
- Dyer, J.A., Kulshreshtha, S.N., McConkey, B.G., Desjardins, R.L., 2010. An assessment of fossil fuel energy use and CO<sub>2</sub> emissions from farm field operations using a regional level crop and land use database for Canada. *Energy* 35, 2261–2269. <https://doi.org/10.1016/j.energy.2010.02.013>.
- EEA, 2018. Annual European Union Greenhouse Gas Inventory 1990–2016 and Inventory Report 2018 Submission to the UNFCCC Secretariat (27 May 2018. 955pp).
- ENA, 2011. In: Sutton, et al. (Eds.), *European Nitrogen Assessment*. Cambridge University Press, London (612pp).
- Erb, K.-H., Lauk, C., Kastner, T., Mayer, A., Theurl, M.C., Haberl, H., 2016. Exploring the biophysical option space for feeding the world without deforestation. *Nat. Commun.* 7, 11382.
- Fischer-Kowalski, M., Haberl, J., 2007. Socioecological transitions and global change: trajectories of social metabolism and land use. *Advances in Ecological Economics Series*. Edward Elgar Pub (263pp).
- Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D.W., Haywood, J., Lean, J., Lowe, D.C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., Van Dorland, R., 2007. Changes in atmospheric constituents and in radiative forcing. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Garnier, J., Billen, G., Vilain, G., Martinez, A., Mounier, E., Silvestre, M., Toche, F., 2009. Nitrous oxide (N<sub>2</sub>O) in the Seine river and basin: observations and budgets. *Agric. Ecosyst. Environ.* 133, 223–233. <https://doi.org/10.1016/j.agee.2009.04.024>.
- Garnier, J., Vilain, G., Jehanno, S., Silvestre, M., Billen, G., Poirier, D., Martinez, A., Decuq, C., Cellier, P., Abril, G., 2013. Methane emissions from land use, livestock farming, and the river network of the Seine basin (France). *Biogeochemistry* 116, 199–214. <https://doi.org/10.1007/s10533-013-9845-1>.
- Garnier, J., Billen, G., Vilain, G., Benoit, M., Passy, P., Tallec, G., Tournebise, J., Anglade, J., Billy, C., Mercier, B., Ansart, P., Sebilo, M., Kao, C., 2014. Curative vs. preventive management of nitrogen transfers in rural areas: lessons from the case of the Orgeval watershed (Seine River basin, France). *J. Environ. Manag.* 144, 125–134. <https://doi.org/10.1016/j.jenvman.2014.04.030>.
- Garnier, J., Lassaletta, L., Billen, G., Romero, E., Grizzetti, B., Némery, J., Le, Q.L.P., Pistocchi, C., Aissa-Grouz, N., Luu, M.T.N., Vilmin, L., Dorioz, J.-M., 2015. Phosphorus budget in the water-agro-food system at nested scales in two contrasted regions of the world (ASEAN-8 and EU-27). *Glob. Biogeochem. Cycles* <https://doi.org/10.1002/2015GB005147>.
- Garnier, J., Anglade, J., Benoit, M., Billen, G., Puech, T., Ramarson, A., Passy, P., Silvestre, M., Lassaletta, L., Trommschlagel, J.-M., Schott, C., Tallec, G., 2016. Reconnecting crop and cattle farming to reduce nitrogen losses in river water of an intensive agricultural catchment (Seine basin, France). *Environ. Sci. Pol.* 63, 76–90. <https://doi.org/10.1016/j.envsci.2016.04.019>.
- Garnier, J., Ramarson, A., Billen, G., Théry, S., Thiéry, D., Thieu, V., Minaudo, C., Moatar, F., 2018. Nutrient inputs and hydrology together determine biogeochemical status of the Loire River (France): current situation and possible future scenarios. *Sci. Total Environ.* 637–638, 609–624. <https://doi.org/10.1016/j.scitotenv.2018.05.045>.
- Gilhespy, S.L., Anthony, S., Cardenas, L., Chadwick, D., del Prado, A., Li, C.S., Misselbrook, T., Rees, R.M., Salas, W., Sanz-Cobena, A., Smith, P., Tilston, E.L., Topp, C.F.E., Vetter, S., Yeluripati, J.B., 2014. First 20 years of DNDC (DeNitrification DeComposition): model evolution. *Ecol. Model.* 292, 51–62. <https://doi.org/10.1016/j.ecolmodel.2014.09.004>.
- Gingrich, S., Erb, K.-H., Krausmann, F., Gaube, V., Haberl, H., 2007. Long-term dynamics of terrestrial carbon stocks in Austria: a comprehensive assessment of the time period from 1830 to 2000. *Reg. Environ. Chang.* 7, 37–47. <https://doi.org/10.1007/s10113-007-0024-6>.
- Groffman, P., 1991. Ecology of nitrification and denitrification in soil evaluated at scales relevant to atmospheric chemistry. In: Rogers, J.E., Whitman, W.B. (Eds.), *Microbial Production and Consumption of Greenhouse Gases: Methane, Nitrogen Oxides and Halomethanes*. American Society of Microbiology, Washington, DC (USA), pp. 201–217.
- Gu, J., Loustau, D., Hénault, C., Rochette, P., Cellier, P., Nicoulaud, B., Gossel, A., Richard, G., 2014. Modeling nitrous oxide emissions from tile-drained winter wheat fields in Central France. *Nutr. Cycl. Agroecosyst.* 98, 27–40. <https://doi.org/10.1007/s10705-013-9593-6>.
- Harchaoui, S., Chatzimirois, P., 2018. Energy, nitrogen, and farm surplus transitions in agriculture from historical data modeling. France, 1882–2013. *J. Ind. Ecol.* <https://doi.org/10.1111/jieec.12760>.
- HAROPA, 2015. Projets stratégiques HAROPA 2030. [http://www.haropaports.com/projets\\_strategiques/](http://www.haropaports.com/projets_strategiques/).
- Hénault, C., Devis, X., Lucas, J.L., Germon, J.C., 1998. Influence of different agricultural practices (type of crop, form of N-fertilizer) on soil nitrous oxide emissions. *Biol. Fertil. Soils* 27, 299–306. <https://doi.org/10.1007/s003740050437>.
- Houghton, R.A., House, J.I., Pongratz, J., van der Werf, G.R., De Fries, R.S., Hansen, M.C., Le Quére, C., Ramankutty, N., 2012. Carbon emissions from land use and landcover change. *Biogeosciences* 9, 5125–5142. <https://doi.org/10.5194/bg-9-5125-2012>.
- IPCC, 1997. Revised 1996 IPCC guidelines for national greenhouse gas inventories. Reference manual. IPCC <http://www.ipcc-nggip.iges.or.jp/public/gl/invs6.html>, Accessed date: 27 October 2018.
- IPCC, 2008. 2006 IPCC guidelines for national greenhouse gas inventories – a primer, prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Miwa K., Srivastava N. and Tanabe K. (eds). IGES, Japan [https://www.ipcc-nggip.iges.or.jp/support/Primer\\_2006GLs.pdf](https://www.ipcc-nggip.iges.or.jp/support/Primer_2006GLs.pdf), Accessed date: 27 October 2018.
- IPCC, 2014. In: Pachauri, R.K., Meyer, L.A. (Eds.), *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland Core Writing Team. (151 pp).
- IPCC, 2018. Global warming of 1.5 °C summary for policymakers, approved at the First Joint Session of Working Groups I, II and III of the IPCC and accepted by the 48th Session of the IPCC, Incheon, Republic of Korea, 6 October 2018. Allen M. et al. 33pp. <https://www.ipcc.ch/>.
- Jouzel, J., Ouzeau, G., Déqué, M., Jouini, M., Planton, S., Vautard, R., 2014. Le climat de la France au XXI<sup>e</sup> siècle, Volume 4, Scénarios régionalisés: édition 2014 pour la métropole et les régions d'outre-mer. Ministère de l'Écologie, du Développement durable et de l'Énergie. 62pp. [https://www.ecologie-solidaire.gouv.fr/sites/default/files/ONERC\\_Climat\\_France\\_XXI\\_Volume\\_4\\_VF.pdf](https://www.ecologie-solidaire.gouv.fr/sites/default/files/ONERC_Climat_France_XXI_Volume_4_VF.pdf).
- Kandel, T.P., Lærke, P.E., Elsgaard, L., 2018. Annual emissions of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O from a temperate peat bog: comparison of an undrained and four drained sites under permanent grass and arable crop rotations with cereals and potato. *Agric. For. Meteorol.* 256–257, 470–481. <https://doi.org/10.1016/j.agrformet.2018.03.021>.
- Kim, D.-G., Kirschbaum, M.U.F., 2015. The effect of land-use change on the net exchange rates of greenhouse gases: a compilation of estimates. *Agric. Ecosyst. Environ.* 208, 114–126. <https://doi.org/10.1016/j.agee.2015.04.026>.
- Kirschbaum, M.U.F., Saggart, S., Tate, K.R., Giltrap, D.L., Ausseil, A.G.E., Greenhalgh, S., Whitehead, D., 2012. Comprehensive evaluation of the climate-change implications of shifting land use between forest and grassland: New Zealand as a case study. *Agric. Ecosyst. Environ.* 150, 123–138. <https://doi.org/10.1016/j.agee.2012.01.004>.
- Lassaletta, L., Billen, G., Romero, E., Garnier, J., Aguilera, E., 2014a. How changes in diet and trade patterns have shaped the N cycle at the national scale: Spain (1961–2009). *Reg. Environ. Chang.* 14, 785–797. <https://doi.org/10.1007/s10113-013-0536-1>.
- Lassaletta, L., Billen, G., Grizzetti, B., Anglade, J., Garnier, J., 2014b. 50 year trends in nitrogen use efficiency of world cropping systems: the relationship between yield and nitrogen input to cropland. *Environ. Res. Lett.* 9. <https://doi.org/10.1088/1748-9326/9/10/105011>.
- Laville, P., Lehuger, S., Loubet, B., Chaumartin, F., Cellier, P., 2011. Effect of management, climate and soil conditions on N<sub>2</sub>O and NO emissions from an arable crop rotation using high temporal resolution measurements. *Agric. For. Meteorol.* 151, 228–240. <https://doi.org/10.1016/j.agrformet.2010.10.008>.
- Le Noé, J., Billen, G., Lassaletta, L., Silvestre, M., Garnier, J., 2016. La place du transport de denrées agricoles dans le cycle biogéochimique de l'azote en France: un aspect de la spécialisation des territoires. *Cah. Agric.* 25, 15004. <https://doi.org/10.1051/cagri/2016002>.
- Le Noé, J., Billen, G., Garnier, J., 2017. Nitrogen, phosphorus and carbon fluxes through the French Agro-Food System: an application of the GRAFS approach at the territorial scale. *Sci. Total Environ.* 586, 42–55. <https://doi.org/10.1016/j.scitotenv.2017.02.040>.
- Le Noé, J., Billen, G., Esculier, F., Garnier, J., 2018. Long term socio-ecological trajectories of agro-food systems revealed by N and P flows: the case of French regions from 1850 to 2014. *Agric. Ecosyst. Environ.* 265, 132–143. <https://doi.org/10.1016/j.agee.2018.06.006>.
- Le Noé, J., Billen, G., Mary, B., Garnier, J., 2019. Drivers of long-term carbon dynamics in cropland: a bio-political history (France, 1852–2014). *Environ. Sci. Pol.* 93, 53–65. <https://doi.org/10.1016/j.envsci.2018.12.027>.
- Lemke, R.L., Zhong, Z., Campbell, C.A., Zentner, R., 2007. Can pulse crops play a role in mitigating greenhouse gases from North American agriculture? *Agron. J.* 99, 1719–1725. <https://doi.org/10.2134/agronj2006.0327s> (Symposium Papers).
- Lu, Y., Huang, Y., Zou, J., Zheng, X., 2006. An inventory of N<sub>2</sub>O emissions from agriculture in China using precipitation-rectified emission factor and background emission. *Chemosphere* 65, 1915–1924. <https://doi.org/10.1016/j.chemosphere.2006.07.035>.
- Lu, Z., Du, R., Du, P., Qin, S., Liang, Z., Li, Z., Wang, Y., Wang, Y., 2015. Influences of land use/cropland types on nitrous oxide emissions during freeze-thaw periods from waterlogged soils in Inner Mongolia. *PLoS ONE* 10 (9), e0139316. <https://doi.org/10.1371/journal.pone.0139316>.

- Marescaux, A., Thieu, V., Garnier, J., 2018. Carbon dioxide, methane and nitrous oxide emissions from the human-impacted Seine watershed in France. *Sci. Total Environ.* 643, 247–259. <https://doi.org/10.1016/j.scitotenv.2018.06.151>.
- Moss, A.R., Jouany, J.-P., Newbold, J., 2000. Methane production by ruminants: its global contribution to global warming. *Ann. Zootech.* 49, 231–253. <https://doi.org/10.1051/animres:2000119>.
- Muller, A., Schader, C., El-Hage Scialabba, N., Brüggemann, J., Isensee, A., Erb, K.-H., Smith, P., Klocke, P., Leiber, F., Stolze, M., Niggli, U., 2017. Strategies for feeding the world more sustainably with organic agriculture. *Nat. Commun.* 8, 1290. <https://doi.org/10.1038/s41467-017-01410-w>.
- Ojima, D.S., Valentine, D.W., Mosier, A.R., Parton, W.J., Schimel, D.S., 1993. Effect of land use change on methane oxidation in temperate forest and grassland soils. *Chemosphere* 26, 675–685. [https://doi.org/10.1016/0045-6535\(93\)90452-B](https://doi.org/10.1016/0045-6535(93)90452-B).
- Patzek, T.W., 2003. Energy required to produce corn. *Civil and Environmental Engineering, U.C. Berkeley*, March 12, 2003. <http://petroleum.berkeley.edu/patzek/ce24/ce24top.htm>.
- Plaza-Bonilla, D., Álvaro-Fuentes, J., Arrúe, J.L., Cantero-Martínez, C., 2014. Tillage and nitrogen fertilization effects on nitrous oxide yield-scaled emissions in a rainfed Mediterranean area. *Agric. Ecosyst. Environ.* 189, 43–52. <https://doi.org/10.1016/j.agee.2014.03.023>.
- Poisvert, C., Curie, F., Moatar, F., 2017. Annual agricultural N surplus in France over a 70-year period. *Nutr. Cycl. Agroecosyst.* <https://doi.org/10.1007/s10705-016-9814-x>.
- Poux, X., Aubert, P.-M., 2018. Une Europe agroécologique en 2050: une agriculture multifonctionnelle pour une alimentation saine. Enseignements d'une modélisation du système alimentaire européen, Study N°09/18. Iddri-AScA, Paris, France (78 p).
- Poux, X., Narcy, J.-B., Chenat, V., 2005. *Agriculture et environnement: 4 scénarios à l'horizon 2025*. Groupe de la Bussière. ASCA (123pp+annexes).
- Recio, J., Vallejo, A., Le-Noë, J., Garnier, J., García-Marco, S., Álvarez, J.M., Sanz-Cobena, A., 2018. The effect of nitrification inhibitors on NH<sub>3</sub> and N<sub>2</sub>O emissions in highly N fertilized irrigated Mediterranean cropping systems. *Sci. Total Environ.* 636, 427–436. <https://doi.org/10.1016/j.scitotenv.2018.04.294>.
- Saggar, S., Jha, N., Deslippe, J., Bolan, N.S., Luo, J., Giltrap, D.L., Kim, D.-G., Zaman, M., Tillman, R.W., 2013. Denitrification and N<sub>2</sub>O:N<sub>2</sub> production in temperate grasslands: processes, measurements, modelling and mitigating negative impacts. A review. *Sci. Total Environ.* 465, 173–195. <https://doi.org/10.1016/j.scitotenv.2012.11.050>.
- Sanchez-Martín, L., Meijide, A., García-Torres, L., Vallejo, A., 2010. Combination of drip irrigation and organic fertilizer for mitigating emissions of nitrogen oxides in semiarid climate. *Agric. Ecosyst. Environ.* 137, 99–107. <https://doi.org/10.1016/j.agee.2010.01.006>.
- Sanz-Cobena, A., Sánchez-Martín, L., García-Torres, L., Vallejo, A., 2012. Gaseous emissions of N<sub>2</sub>O and NO and NO<sub>3</sub>-leaching from urea applied with urease and nitrification inhibitors to a maize (*Zea mays*) crop. *Agric. Ecosyst. Environ.* 149, 64–73. <https://doi.org/10.1016/j.agee.2011.12.016>.
- Sanz-Cobena, A., Lassaletta, L., Aguilera, E., del Prado, A., Garnier, J., Billen, G., Iglesias, A., Sánchez, B., Guardia, G., Abalos, D., Plaza-Bonilla, D., Puigdueta, I., Moral, R., Galán, E., Arriaga, H., Merino, P., Infante-Amate, J., Meijide, A., Pardo, G., Alvaro-Fuentes, J., Gilsanz, C., Báez, D., Doltra, J., González, S., Cayuela, M.L., Menendez, S., Díaz-Pines, E., Le Noë, J., Quemada, M., Estellés, F., Calvet, S., van Grinsven, H., Yáñez, D., Westhoek, H., Sanz, M.J., Sánchez-Jimeno, B., Vallejo, A., Smith, P., 2017. Strategies for GHG mitigation in Mediterranean agricultural soils. A review. *Agric. Ecosyst. Environ.* 238, 5–24. <https://doi.org/10.1016/j.agee.2016.09.038>.
- Schmitz, C., Biewald, A., Lotze-Campen, H., Popp, A., Dietrich, J.P., Bodirsky, B., Krause, M., Weindl, I., 2012. Trading more food: implications for land use, greenhouse gas emissions, and the food system. *Glob. Environ. Chang.* 22, 189–209. <https://doi.org/10.1016/j.gloenvcha.2011.09.013>.
- Schnell, S., King, G.M., 1994. Mechanistic analysis of ammonium inhibition of atmospheric methane consumption in forest soils. *Appl. Environ. Microbiol.* 60, 3514–3521.
- Skiba, U., McTaggart, I.P., Smith, K.A., Hargreaves, K., Fowler, D., 1996. Estimates of nitrous oxide emissions from soil in the UK. *Energy Convers. Manag.* 37, 1303–1308. [https://doi.org/10.1016/0196-8904\(95\)00337-1](https://doi.org/10.1016/0196-8904(95)00337-1).
- Smith, K., 2010. *Nitrous Oxide and Climate Change*. Earthscan (240 pp).
- Smith, K.A., McTaggart, I.P., Tsuruta, H., 1997. Emissions of N<sub>2</sub>O and NO associated with nitrogen fertilization in intensive agriculture, and the potential for mitigation. *Soil Use Manag.* 13, 296–304. <https://doi.org/10.1016/j.agee.2009.04.021>.
- Snyder, C.S., Bruulsema, T.W., Jensen, T.L., Fixen, P.E., 2009. Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agric. Ecosyst. Environ.* 133, 247–266. <https://doi.org/10.1016/j.agee.2009.04.021>.
- Solagro, 2014. Afterres2050: Un scénario soutenable pour l'agriculture et l'utilisation des terres en France à l'horizon 2050. <http://www.solagro.org/site/393.html>.
- Springmann, M., Clark, M., Mason-D'Croz, D., Wiebe, K., Bodirsky, B.L., Lassaletta, L., de Vries, W., Vermeulen, S.J., Herrero, M., Carlson, K.M., Jonell, M., Troell, M., DeClerck, F., Gordon, L.J., Zurayk, R., Scarborough, P., Rayner, M., Loken, B., Fanzo, J., Godfray, H.C.J., Tilman, D., Rockström, J., Willett, W., 2018. Options for keeping the food system within environmental limits. *Nature* <https://doi.org/10.1038/s41586-018-0594-0>.
- Tate, K.R., 2015. Soil methane oxidation and land-use change – from process to mitigation. A review. *Soil Biol. Biochem.* 80, 260–272. <https://doi.org/10.1016/j.soilbio.2014.10.010>.
- Tiedje, J.M., 1988. Ecology of denitrification and dissimilatory nitrate reduction to ammonium. In: Zehnder, A.J.B. (Ed.), *Biology of Anaerobic Microorganisms*. Wiley & Sons, New York, pp. 179–244.
- Verhoeven, J.T.A., Setter, T.L., 2010. Agricultural use of wetlands: opportunities and limitations. *Ann. Bot.* 105, 155–164. <https://doi.org/10.1093/aob/mcp172>.
- Vermorel, M., Jouany, J., Eugène, M., Sauvart, D., Noblet, J., Durmad, J., 2008. Evaluation quantitative des émissions de méthane entérique par les animaux d'élevage en 2007 en France. *INRA. Prod. Anim.* 21, 403–418.
- Vilain, G., Garnier, J., Tallec, G., Cellier, P., 2010. Effect of slope position and land use on nitrous oxide (N<sub>2</sub>O) emissions (Seine Basin, France). *Agric. For. Meteorol.* 150, 1192–1202. <https://doi.org/10.1016/j.agrformet.2010.05.004>.
- Vilain, G., Garnier, J., Decuq, C., Lugnot, M., 2014. Nitrous oxide production by nitrification vs. denitrification in experimental conditions. *Nutr. Cycl. Agroecosyst.* <https://doi.org/10.1007/s10705-014-9604-2>.
- Wertz, S., Goyer, C., Zebbarth, B., Tatti, E., Burton, D.L., Chantigny, M.H., Filion, M., 2016. The amplitude of soil freeze-thaw cycles influences temporal dynamics of N<sub>2</sub>O emissions and denitrifier transcriptional activity and community composition. *Biol. Fertil. Soils* 52, 1149–1162. <https://doi.org/10.1007/s00374-016-1146-0>.
- Westhoek, H., Lesschen, J.P., Rood, T., Wagner, S., De Marco, A., Murphy-Bokern, D., Leip, A., van Grinsven, H., Sutton, M.A., Oenema, O., 2014. Food choices, health and environment: effects of cutting Europe's meat and dairy intake. *Glob. Environ. Chang.* 26, 196–205. <https://doi.org/10.1016/j.gloenvcha.2014.02.004>.
- Zedler, J.B., Kercher, S., 2005. Wetland resources: status, trends, ecosystem services, and restorability. *Annu. Rev. Environ. Resour.* 30, 39–74. <https://doi.org/10.1146/annurev.energy.30.050504.144248>.
- Zervas, G., Tsiplakou, E., 2012. An assessment of GHG emissions from small ruminants in comparison with GHG emissions from large ruminants and monogastric livestock. *Atmos. Environ.* 49, 13–23. <https://doi.org/10.1016/j.atmosenv.2011.11.039>.
- Zheng, X., Wang, M., Wang, Y., Shen, R., Gou, J., Li, J., Jin, J., Li, L., 2000. Impacts of soil moisture on nitrous oxide emission from croplands: a case study on the rice-based agroecosystems in Southeast China. *Chemosphere Global Change Sci.* 2, 207–224. [https://doi.org/10.1016/S1465-9972\(99\)00056-2](https://doi.org/10.1016/S1465-9972(99)00056-2).
- Zhou, J.B., Jiang, M.M., Chen, G.Q., 2007. Estimation of methane and nitrous oxide emission from livestock and poultry in China during 1949–2003. *Energy Policy* 35, 3759–3767. <https://doi.org/10.1016/j.enpol.2007.01.013>.
- Zimmermann, J., Carolan, R., Forrester, P., Hartly, M., Lanigan, G., Richards, K.G., Roche, L., Whitfield, M.G., Jones, M.B., 2018. Assessing the performance of three frequently used biogeochemical models when simulating N<sub>2</sub>O emissions from a range of soil types and fertiliser treatments. *Geoderma* 331, 53–69. <https://doi.org/10.1016/j.geoderma.2018.06.004>.