



HAL
open science

High-resolution quantification of earthworm calcite granules from western European loess sequences reveals stadial-interstadial climatic variability during the Last Glacial

Charlotte Prud'Homme, Olivier Moine, Jérôme Mathieu, Segolene Saulnier-Copard, Pierre Antoine

► To cite this version:

Charlotte Prud'Homme, Olivier Moine, Jérôme Mathieu, Segolene Saulnier-Copard, Pierre Antoine. High-resolution quantification of earthworm calcite granules from western European loess sequences reveals stadial-interstadial climatic variability during the Last Glacial. *Boreas*, 2019, 48 (1), pp.257-268. 10.1111/bor.12359 . hal-02147883

HAL Id: hal-02147883

<https://hal.sorbonne-universite.fr/hal-02147883v1>

Submitted on 15 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High-resolution quantification of earthworm calcite granules from western European loess sequences reveals stadial–interstadial climatic variability during the Last Glacial

CHARLOTTE PRUD'HOMME , OLIVIER MOINE, JEROME MATHIEU, SEGOLENE SAULNIER-COPARD AND PIERRE ANTOINE

BOREAS



Prud'homme, C., Moine, O., Mathieu, J., Saulnier-Copard, S. & Antoine, P.: High-resolution quantification of earthworm calcite granules from western European loess sequences reveals stadial–interstadial climatic variability during the Last Glacial. *Boreas*. <https://doi.org/10.1111/bor.12359>. ISSN 0300-9483.

High concentrations of calcite fossil granules produced by earthworms (ECG) have been identified in most of the stratigraphical units along the loess-palaeosol reference sequence of Nussloch (Germany). They are particularly abundant in interstadial brown soils and in tundra gley horizons, the latter reflecting short-term phases of aggradation then degradation of permafrost. These granules are characterized by a radial crystalline structure produced in the earthworms by specific bio-mineralization processes. In our study, we used this biological indicator combined with ¹⁴C and OSL dating, and sedimentological parameters to characterize millennial-time scale climatic variations recorded in loess sequences. The approach is based on high-resolution counts of ECG throughout a 17-m-thick loess sequence (332 samples). Strong increases in granule and mollusc concentrations suggest warmer climate conditions during palaeosol formation phases, associated with increasing biodiversity, biological activity and vegetation cover. Decreased granule concentrations occur within primary loess deposits, indicating a strong correlation with palaeoenvironmental conditions and demonstrating the reliability of ECG concentration variations as a new palaeoenvironmental proxy. Finally, this pattern is also recorded in loess sequences located about 600 km westward in northern France demonstrating the large-scale validity of this new palaeoclimatic proxy.

Charlotte Prud'homme (c.prudhomme@mpic.de), Laboratoire de Géographie Physique, Environnements Quaternaires et Actuels, (LGP), UMR8591 CNRS-Université Paris 1-UPEC, 1 Place Aristide Briand 92195 Meudon cedex, France and Research Group for Terrestrial Palaeoclimates, Max Planck Institute for Chemistry, Hahn-Meitner-Weg 1, Mainz 55128, Germany; Olivier Moine, Ségolène Saulnier-Copard and Pierre Antoine, Laboratoire de Géographie Physique, Environnements Quaternaires et Actuels, (LGP), UMR8591, CNRS-Université Paris 1-UPEC, 1 Place Aristide Briand 92195 Meudon cedex, France; Jerome Mathieu, Université Pierre-et-Marie-Curie (UPMC) - Institut d'Ecologie et des Sciences de l'Environnement de Paris, Institut de Recherche pour le Développement, Sorbonne Universités, 4 Place Jussieu, Paris 75005, France; received 31st May 2018, accepted 1st October 2018.

Covering almost all non-glaciated continental areas lower than 500 m between 42 and 52°N, the European Loess Belt represents the most extensive terrestrial archive of the Last Glacial (Pecsi 1990; Haase *et al.* 2007). Its western part is well situated to record the impact of the climatic variations triggered by fluctuations of both North Atlantic oceanic circulation and Fennoscandian sea-ice cover extension propagated eastward by the Westerlies (Renssen *et al.* 2007; Antoine *et al.* 2009). The loess sequences of western Europe provide very detailed and reliable records of climatic and environmental changes of the last 110 kyr in the European Plain (Remy 1968; Antoine *et al.* 2001, 2016; Meijs 2011; Haesaerts *et al.* 2016; Schirmer 2016). Despite the occurrence of some erosion gaps these sequences, made up of complex and cyclic successions of loess units (aeolian deposits) and of soil horizons, have recorded the impact of millennial-scale climatic variability during the Last Glacial (Antoine *et al.* 2009, 2016; Moine *et al.* 2017; Rousseau *et al.* 2017). They also exhibit high sedimentation rates of up to 1 mm per year especially between 30 and 20 ka during the Upper Pleniglacial (Meszner *et al.* 2011, 2013; Antoine *et al.* 2016). Previously based on pedostratigraphical studies, grain-size

analysis, terrestrial mollusc assemblages and OSL dating (Antoine *et al.* 2002, 2009; Rousseau *et al.* 2007; Moine *et al.* 2008; Meijs 2011; Schirmer 2012), correlations of palaeosols with Dansgaard–Oeschger events recorded in Greenland ice-cores have recently been improved by radiocarbon dating performed on earthworm calcite granules (ECG; Moine *et al.* 2017).

However, as pollen grains are generally not preserved in loess sediments due to their oxidation in this porous sediment, the impact of these millennial climate changes on palaeoenvironment has been until now mainly investigated through the analysis of terrestrial mollusc assemblages (Moine *et al.* 2005, 2008, 2011). The malacological analysis of Nussloch loess sequences revealed a vegetation cover developed on periglacial deposits (Moine 2008) and confirmed that the formation of palaeosols such as tundra gley horizons is associated with moisture increase. In parallel, a new proxy has been discovered in Quaternary deposits: ECG. Surprisingly, the presence of ECG in Last Glacial loess-palaeosol sequences has been found to be common in many regions from western-central Europe (Mazenot 1956; Kerney 1971; Meijer 1985; Becze-Deák *et al.* 1997; Barta 2014; Prud'homme *et al.* 2015; Moine *et al.* 2017).

ECG are composed of large rhombohedral calcite crystals organized in a radial crystalline structure (Canti 1998; Gago-Duport *et al.* 2008). They are produced by a biomineralization process within calciferous glands (also called Morren's glands) located in three pairs on each side of the oesophagus (Darwin 1881). This process starts in the four oesophagus pouches (segments XII and XI) with the secretion of a colloidal 'milky' fluid, which contains amorphous calcium carbonate (ACC) crystals of irregular shape and size ranging from 10 to 50 μm . Then, the milky fluid reaches the two oesophageal pouches in the front (segment X) where the ACC is precipitated into macroscopic crystals of calcium carbonate, which are finally released into the soil through the gut (Briones *et al.* 2008; Gago-Duport *et al.* 2008; Hodson *et al.* 2015). Currently, *Lumbricus* is identified as the most productive genus (Bräm 1956; Canti 1998; Canti & Pearce 2003). *In vitro* experiments showed that this genus releases granules principally at the soil surface in the litter (Canti & Pearce 2003). However, contrary to the species *Eisenia nordenskioldi* and *Dendrobaena octaedra* commonly found in permafrost environments, the *Lumbricus* genus is not considered as freeze-tolerant (Rasmussen & Holmstrup 2002; Holmstrup 2003; Calderon *et al.* 2009). Freeze-tolerant species can endure ice formation in extracellular body due to rapid glucose accumulation in cells. Moreover, *Eisenia nordenskioldi* and *Dendrobaena octaedra* belong to the epigeic class (i.e. living at the surface of the soil) whereas *Lumbricus terrestris* and *Lumbricus rubellus* belong to the anecic and epi-edogenic classes, respectively (i.e. living beneath the soil surface and able to dig burrows). Thus, their life cycle and behaviour cannot be comparable in the same context. *Lumbricus terrestris* and *Lumbricus rubellus* are found up to 60 and 65°N, respectively, in the central Russian Plain (Tiunov *et al.* 2006; Meshcheryakova & Berman 2014) and also in southwestern Finland and southern Sweden (Rundgren 1975; Nuutinen & Butt 2009). Although they are often absent from modern permafrost environments (Perel' 1979), adult individuals of both species can survive in soil temperatures as low as $-1\text{ }^{\circ}\text{C}$ (Nordström 1975; Rundgren 1975). Moreover, *L. terrestris* is able to overwinter in all life stages and its cocoons (i.e. envelopes protecting the eggs) can survive 1 month in frozen soil (Nuutinen & Butt 2009). Earthworm cocoons resist freezing by cryoprotective dehydration, characterized by a loss of 50% of the mass of their body fluids (Holmstrup 1994). Even though most earthworm species in western Europe were strongly affected by the Last Glacial (Mathieu & Davies 2014), the discovery of abundant granules in palaeosols developed during the Weichselian suggests that a few species have been able to survive in permafrost environment, perhaps even outside refuges. As *Lumbricus* species are currently very abundant in Germany (Jänsch *et al.* 2013) and in the absence of other common large taxa known to produce numerous granules in western Europe (Bräm 1956; Canti 1998), we

assume that *Lumbricus* individuals produced the large ($\leq 0.8\text{ mm}$) granules discovered in the Last Glacial loess sequence of Nussloch.

Previous works showed that ECG: (i) can constitute an ideal material for radiocarbon dating in loess contexts as demonstrated by the new chronology of the Nussloch sequence (Moine *et al.* 2017), (ii) allow the quantification of palaeoclimatic parameters (temperature and precipitation; Prud'homme *et al.* 2016, 2018) and (iii) can be used as a bio-indicator for palaeoenvironmental reconstructions (Prud'homme *et al.* 2015).

In this paper, we focus on the third point above by presenting the first continuous record of ECG concentration variations from the west European reference site of Nussloch. The 17-m-thick loess-palaeosol sequence encompasses the Weichselian Middle and Upper Pleniglacial from about 55 to 20 ka. The ECG counting profile is compared with high-resolution stratigraphy and grain-size analysis.

Material and methods

Stratigraphy

The Nussloch loess-palaeosol profiles have been studied in an active limestone quarry (latitude 49°18'59"N, longitude 8°43'54"E) located at about 10 km south-southeast of the city of Heidelberg on the eastern flank of the Upper Rhine valley (Fig. 1). The stratigraphy of the former four main profiles (P1 to P4) has already been published in great detail (Antoine *et al.* 2001, 2009). These profiles consist of four sub-sequences attributed to the main chronoclimatic phases of the Last Glacial, i.e. Early Glacial, Lower, Middle and Upper Pleniglacial (Antoine *et al.* 2009). Profile P8, located about 50 m from the reference profile P4 in the same loess greda (i.e. elongated loess ridge forms orientated parallel to the direction of winds; Leger 1990), revealed the same stratigraphical succession and can easily be correlated with the other profiles using specific palaeosol marker horizons. Moreover, a radiocarbon chronology based on 46 dates obtained from ECG has been produced for Nussloch profiles P3, P4 and P8 (Moine *et al.* 2017). This new chronology, combined with some of the previous OSL dates, highlights a very accurate time-framework for the deposition of the Nussloch loess sequence.

According to stratigraphy and sedimentology given in Antoine *et al.* (2009), the 17-m-thick P8 profile is made up of the succession of two main chronoclimatic phases of the Last Glacial: the lower 5 m allocated to the Middle Pleniglacial (from ~ 55 to 35 ka; MIS 3) and the upper 12 m representing the Upper Pleniglacial (from ~ 35 to 20 ka; end of MIS 3 and beginning of MIS 2). The lower part is composed of a succession of two thick boreal brown soils (Lower (GBL) and Upper (GBU) Gräselberger Boden) developed on silty aeolian sands, followed by loess and tundra gley horizons, topped by an arctic brown soil




Fig. 1. Map of the loess and sandy loess cover over western Europe with the location of the Nussloch site and of the northern France sequences (NFS) modified from Haase *et al.* (2007) and Antoine *et al.* (2016).

(Lohner Boden). This part of the record is characterized by an average sedimentation rate of 0.19 mm a^{-1} .

The upper part is characterized by a very high sedimentation rate (average: 0.93 mm a^{-1}) and by a cyclic alternation of typical calcareous loess and tundra gley horizons. Tundra gley horizons were formed by hydromorphic processes and correspond to active layers of former permafrost events. Thus, these horizons indicate the presence of a former permafrost. This interpretation is supported by observations from northwestern France plateau sections in which large ice-wedge cast networks are systematically connected with tundra gley horizons (Antoine *et al.* 2016). Furthermore, the new radiocarbon chronology of the Nussloch loess sequence (Moine *et al.* 2017) confirms the correlations of both brown soils and main tundra gley horizons with interstadials identified in Greenland ice-core $\delta^{18}\text{O}$ and dust records.

Sampling strategy

The 17-m-thick Nussloch profile P8 was sampled at high resolution (5-cm-thick) using the continuous column sampling protocol (Antoine *et al.* 2009). ~~Two continuous columns of 332 samples were collected in parallel: one 5 cm large producing 300 g samples for grain size analyses and one 50 cm large resulting in 10 L samples (10–12 kg) for malacological analyses from which granules have been extracted.~~

Grain-size analyses were performed using a Beckman Coulter LS-230 laser particle size analyser following the protocol presented in Antoine *et al.* (2009). Ten grams

of homogenized ~~subsamples~~ was first dispersed by addition of sodium hexametaphosphate (5‰), then processed overnight in a rotating agitator, and finally sieved at $160 \mu\text{m}$ in order to remove the coarse fraction (calcified roots, Fe and Mn concretions and fragments of mollusc shells). Each sample was measured with the Coulter at least three times in order to ~~obtain~~ good reproducibility. The limits of the grain-size classes applied to Nussloch were adjusted using a comparison between classical and laser grain-size analyses from a set of test samples: clay: $\leq 4.6 \mu\text{m}$; coarse silts: 26 to $52 \mu\text{m}$; fine sands: 52 to $160 \mu\text{m}$. A grain-size index (GSI) reflecting the variations in aeolian dynamics throughout the record has been defined as the ratio between the coarse silt fraction (%) and the fraction of fine silt plus clay (%) (Antoine *et al.* 2009).

ECG were extracted from the malacology samples. After wet sieving ($>0.425 \text{ mm}$) of the samples, ECG larger than 0.8 mm were extracted and counted under a binocular microscope to select mainly *Lumbricus* material (Canti 1998; Canti & Pearce 2003). According to Canti & Pearce (2003), this genus has its calciferous glands very well developed and produces the largest granules (with a diameter $>0.5 \text{ mm}$).

Results

Variation in ECG concentrations

Figure 2 shows that the variation in the ECG concentrations appears to be correlated with the stratigraphy

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57




Fig. 2. Pedostratigraphy of the Nussloch profile P8 associated with the earthworm granule concentration (in red), the clay fraction percentage (<4.6 μm , blue) and the grain-size index (GSI, brown). Detailed descriptions of the stratigraphy and lithology are available in Antoine *et al.* (2001, 2009). GBL = Lower Gräselberger Boden (boreal brown soil, Bw horizon); GBU = Upper Gräselberger Boden (boreal brown soil, Bw horizon); LB = Lohner Boden (arctic brown soil, Bw horizon); G = major tundra gley (gelic gleysol horizon); IG = incipient tundra gley; r = reworked.

and with both GSI and clay fraction variations. All the data are available in Table S1.

During the Middle Pleniglacial, the highest ECG concentrations are found in boreal and arctic brown soil (GBU and LB) with more than 600 granules per sample. However, the lower boreal brown soil (GBL) recorded a lower granule concentration (around 200), probably due to its higher coarse sand content. P8 is the first profile that provides evidence of the presence of two horizons in the GBL unit (P4-2b), with a stratigraphical limit found in the section at 16 m. The upper part is characterized by a humic horizon associated with a higher percentage of coarse sand (>160 μm) compared to the lower part. This observation is coherent with the double peak in the granule concentration. This palaeosol probably developed during two interstadials (GI 14 and 13). The maximal ECG concentrations occur between 15 and 30 cm below the surface of these palaeosols.

During the Upper Pleniglacial, the maximal concentrations are observed in tundra gley horizons with values between 130 and 400, whereas homogeneous loess units are associated with extremely low ECG concentrations (between 30 and 100 by 10⁻¹). The three laminated loess units are characterized by drastic decreases associated with very low concentrations of large granules (diameter >0.8 mm) between 0 and 10.

Grain-size distribution

The variations in ECG concentration were compared with clay contents and with GSI (ratio between coarse and fine silt+clay). High GSI values imply strong aeolian dynamics and occur within loess units, whereas low GSI values indicate a decrease in dust storms and are correlated within palaeosol formations. GSI highlights relative changes in the aeolian dynamics (Rousseau *et al.* 2007; Antoine *et al.* 2009). Throughout the profile P8, GSI varies from 0.5 to 1.9 (Fig. 2). Palaeosols are characterized by the lowest values whereas the highest ones are associated with loess units.

The clay percentage is affected by two processes: (i) weathering of loess leading to soil formation and (ii) variations in aeolian dynamics (highest values corresponding to phases of low sedimentation rates). The clay fraction varies between 6.3 and 17.2% (Fig. 2). The lowest value is found in unit 13, whereas the highest value is found in the Lohner Boden (unit 20). During the Upper Pleniglacial, the tundra gleys are characterized by higher clay percentages and lower GSI values compared to loess units. The grain-size analyses of profile P8 confirm the cyclic pattern of the loess-palaeosol sequence pedostratigraphy as already shown by Antoine *et al.* (2009) and Rousseau *et al.* (2007). Moreover, the comparison between the variations in ECG concentration and in clay percentage shows a medium correlation ($R = 0.6$) as illustrated by Fig. 2.


Discussion

Survival strategy and seasonal activity of earthworms during the Last Glacial

The discovery of high ECG concentration in palaeosols formed during the Last Glacial and the good correlation with the pedostratigraphy demonstrate the presence and the survival of earthworm species during this period in western Europe, preferentially during interstadials, as already observed in northern France loess sequences (Prud'homme *et al.* 2015). In this study, a more complete record of ECG concentration variations has been obtained between ~55 and 20 ka due to the almost continuous sedimentation and high sedimentation rate of the Nussloch sequence, especially for the Upper Pleniglacial (12 m). High large granule concentration in the tundra gley horizons (active layer of a former permafrost) suggests the presence of the earthworm genus *Lumbricus* (*terrestris* and *rubellus*) in permafrost environment. By contrast, the very low concentration or even the lack of large ECG in loess units indicates a drastic decrease in earthworm activity, but not a total absence of earthworms as a few small granules (0.5 to 0.8 mm) can be found.

According to the literature, *Lumbricus* species are not considered as freeze-tolerant, and they are not found in modern permafrost (Perel' 1979; Holmstrup 2003; Calderon *et al.* 2009). However, as already described in the Introduction, *Lumbricus* is the only earthworm genus able to produce a high quantity of large granules (Bräm 1956; Canti 1998; Canti & Pearce 2003), and this genus is found in Russia and in Scandinavia (Tiunov *et al.* 2006; Meshcheryakova & Berman 2014). The activity of earthworms and the production of granules are directly linked to both soil temperature and humidity and are thus influenced by seasonal climate fluctuations. Nowadays, the maximum activity of earthworms is observed during spring and autumn and can stop the rest of the year (Satchell 1967). Experimental studies on natural environment indicate that *Lumbricus terrestris* are inactive when the soil is frozen (Nordström 1975; Rundgren 1975); however, adult individuals are able to survive and their cocoons (egg matrix) are cryo-resistant (Holmstrup & Zachariassen 1996; Meshcheryakova & Berman 2014). In Fig. 3, a conceptual model is proposed to explain how earthworms were able to survive during the Last Glacial in a permafrost environment in function of the seasons:

- Spring is associated with the thawing of the upper horizon of the permafrost (active layer), development of vegetation and increases in biological activity. Earthworm populations are growing and their activity is basically concentrated at the surface (feeding). In addition, they are confined to the upper part of the active layer (~15–30 cm thick), which is coherent with the thickness of the tundra gley horizons on the field.



41 Fig. 3. Schematic model of the different earthworm activity phases in a permafrost environment depending on the season. See on the text for the
42 description of the figure.

(French 2007; Burn 2012). The conditions are favourable for the production of granules.

- 45 • In summer, although the conditions are drier, *Lumbricus terrestris* are still active (Nordström 1975). The earthworms are mainly found in the upper 5 cm of the soil due to the high proportion of juveniles, which are not able to dig in the deeper layers (Rundgren 1975).
- 46 • In autumn, granule production becomes lower, and the earthworms start to overwinter. The cryo-resistant cocoons are already produced (Holmstrup & Zachariassen 1996) and are mainly distributed in the upper 7.5 cm of the soil (Gerard 1967).
- 47 • In winter, the biological activity stops and the earthworms are in diapause. The snow cover should be thick enough to protect the soil from extreme temperatures (French 2007). The earthworms are confined to the interface between the thick snow cover and the topsoil. The cocoons protect the eggs until they hatch during the next spring, when the conditions become milder (Tiunov *et al.* 2006; Meshcheryakova & Berman 2014).

The earthworms are thus confined to the active layer of the permafrost, greatly reducing the mixing of the granules through depth, allowing an accurate chronological approach of these horizons using ECG ¹⁴C dating (Moine *et al.* 2017). Furthermore, field observations and thin sections indicate an absence of galleries and hiber-

nation chambers in these horizons as the earthworms remain mainly at the surface (Rundgren 1975). Moreover, the rate of active horizontal movements (due to locomotion) varies amongst species, but is limited to 14 m a⁻¹ in the most favourable conditions, and is generally around 4 m a⁻¹ (Eijsackers 2011; Caro *et al.* 2013). The effect of dryness on earthworm movements has not been studied so far, yet it is indeed an interesting question. However, it has been shown for a long time that earthworms go into quiescence, a form of resistance, when the soil dries out (Satchell 1967; Bouché 1972). Bearing in mind that dispersal is very costly in terms of survival, due to predation and other risks, it is more likely that earthworms enter into such a form of resistance rather than dispersing during unfavourable conditions.

Comparison between ECG concentration, pedomatography and grain-size distribution

The good correlation between ECG concentration, clay percentage variations and the stratigraphy (Fig. 2) confirms that earthworms produced the granules during the formation of palaeosols and not after. Granules are thus contemporaneous with the horizons in which they are found. Additionally, the variations in mollusc concentration in the profile P3 of Nussloch (Moine *et al.*

2008) show a similar pattern to the ECG variations, especially during the Middle Pleniglacial. The boreal (GBU) and arctic (LB) brown soils as well as the major tundra gleys are associated with the highest ECG abundances (up to 700) and with high molluscan species diversity and individual concentration. ~~As already observed in the counting of the ECG in P8,~~ the boreal brown soil (GBL) is characterized by lower molluscan abundances than GBU and LB in P3. The increase in molluscan species diversity in these palaeosols indicates a densification of the vegetation cover from steppe to a herb/shrub tundra environment during the Lower–Middle Pleniglacial transition (Moine *et al.* 2005).

The vertical distribution of ECG through the palaeosols between about 55 and 35 ka differs from that in the present-day soils (Fig. 4). In the brown soils and tundra gley horizons, the maximum number of ECG is not found at the surface of the soil but 15 to 30 cm below. However, in present-day soils, earthworms (especially *Lumbricus terrestris* and *L. rubellus*) need a land surface to feed and they produce a high concentration of granules within the litter layer (Canti & Pearce 2003; Canti 2007). This pattern of the ECG concentration suggests that boreal and arctic brown soils result from accretionary soil processes (upbuilding soils; Nikiforoff 1949; Almond & Tonkin 1999) associated with a low aeolian sedimentation, which is contemporaneous with weathering processes. This hypothesis is also confirmed by ~~the distribution of the mollusc concentration through the Middle Pleniglacial palaeosols in Nussloch~~ (Moine *et al.* 2008, 2011). Indeed, most of the mollusc species found in Nussloch are surface species and do not bury more than 5 cm in depth to hibernate (Moine *et al.* 2008). Therefore, loessic sedimentation did not stop completely but only reduced during the palaeosol formation.

Major and incipient tundra gley horizons were developed during the Upper Pleniglacial (Fig. 2). The incipient tundra gley horizons described in the Nussloch loess profiles are associated with high ECG concentrations comparable to major tundra gleys and with an increase in the clay content, especially for IG5, IG6 and IG8a (Fig. 2, units 30a–b, 33a–c, 37a). This pattern is not recorded in the number of individual molluscs, although a decrease in the diversity and a slight increase from 20 to 30% in the proportion of *S. oblonga* is observed. The comparison between the stratigraphical observations and the variations in ECG concentration suggests that incipient tundra gleys might correspond to rapid and very short climate improvements (≤ 0.3 ka), less intense than the interstadial phases. Based on recent ECG radiocarbon chronology, incipient tundra gleys appear to be correlated with significant decreases in Greenland ice-core dust records (Moine *et al.* 2017). Moreover, a slight shift between pedostratigraphical limits, ECG concentrations and clay percentage is observed very clearly for the incipient tundra gleys IG5, IG6 and IG8 and for the major tundra gleys G2, G3 and G4 (Fig. 2).

Thus, the increases in ECG concentration and clay percentage seem to record the beginning of a warming phase, associated with an increase in edaphic humidity and a reduction in the sedimentation rate, slightly below the stratigraphical limits, i.e. preceding the development of the tundra gley horizons.

By contrast, the low concentration of large granules and the absence of molluscs in loess units indicate that the conditions were too extreme for earthworm and mollusc activities. The homogeneous loess units (especially units 22 and 23b) are associated with low ECG concentrations (50–150 granules), but not as low as the laminated loess units, in which they are close to zero ~~and not significant~~ (0–10 in 10 L samples).

This pattern suggests that the climatic conditions during the deposition of homogeneous loess units were not extreme enough (cold, dry and windy) to affect the activity of the earthworm population, in contrast to the mollusc population. Earthworms were thus probably more resistant to extreme conditions than molluscs during the extreme cold phases of the Last Glacial. Indeed, whereas terrestrial molluscs live mainly at the surface of the soil, earthworms live in the soil and feed at the surface. This difference can explain the two patterns of these two ~~proxies~~, especially during the Upper Pleniglacial. Besides, the deposition of the laminated loess units (units 24, 31–32 and 34) is characterized by a high sedimentation rate (≥ 1 mm a⁻¹) and by the occurrence of thin sand beds resulting from dust storms able to rework sand grains from the alluvial plain of the Rhine and niveo-aeolian processes (Antoine *et al.* 2009). An abrupt decrease in ECG concentration is systematically observed in these facies in the Nussloch loess sequence ~~(0 to 10 granules)~~ associated with a decrease in the percentage of clay content and an increase in the GSI. These observations highlight an enhancement of the wind and aeolian dynamics coupled to extreme climatic conditions, which increase the desiccation of the surface. The correlation between the pedostratigraphy of the loess sequence and the NGRIP dust data suggests that these particular horizons are contemporaneous with the Heinrich Stadials HS 3 (around 30 ka) and HS 2 (between 26 and 24 ka) (Rasmussen *et al.* 2014; Moine *et al.* 2017). Heinrich Events, primary ~~highlight~~ in the Atlantic marine core (Heinrich 1988; Bond & Lotti 1995), are characterized by a cold and arid climate on the continent (Fletcher *et al.* 2010; Harrison & Sanchez Goñi 2010; Meniel *et al.* 2014). The comparison between the Heinrich layer in the Atlantic marine core and the Greenland ice-core suggests that the Heinrich Events are contemporaneous with dust peaks in the Greenland ice-core (Fig. 5). Thus, variations in ECG concentrations through loess-palaeosol sequences can be considered as a new proxy of relative variations in aridity. Moreover, ~~the impact of extreme cold events and wind intensity can also be recorded when the ECG concentration is close to 0 during Heinrich Events~~. The same signal (very low to




Fig. 4. Distribution of the ECG concentration through (A) present-day experimental soil and (B) three types of palaeosols from the Nussloch loess profile. The distribution of granules through the present-day experimental soil is associated with the distribution of the number of *Lumbricus terrestris* as a function of depth. The granules were extracted from 50 g of the soil and around 10 000 granules were found in the litter sample (Canti & Pearce 2003). The ECG were extracted from 10 L of sediment every 5 cm through the tundra gley horizons and arctic and boreal brown soils.

zero ECG concentration) is recorded 600 km away in two northern France loess sequences (Fig. 1) for the same time-span (around 30 ka), suggesting that variation

in ECG concentrations can be used to evidence the response of west European environments to large-scale rapid climatic changes.

Comparison with Greenland records

The radiocarbon chronology performed on the ECG confirmed that the two brown soils formed during the Middle Pleniglacial (GBU and LB) can be correlated with Greenland interstadials (GI) 12 and GI 8/7c, and that major tundra gleys G1, G2, G3, G4 and G7 can be correlated with GI 6, GI 5, GI 4, GI 3 and GI 2, respectively (Moine *et al.* 2017). During this time period, the Greenland interstadials GI 12 and GI 8 are the longest interstadials (2700 and 1700 years, respectively), whereas the duration of the GIs correlated with the development of tundra gley horizons are shorter, between 600 and 200 years. The temperature reconstructions performed on NGRIP indicated an increase between

8.5 °C (GI 2) and 15.5 °C (GI 7), with no particular difference between long and short interstadials in this arctic area (Landais *et al.* 2004; Capron *et al.* 2010; Kindler *et al.* 2014). Moreover, palaeoclimate parameters (temperature and precipitation, Fig. 5) have been estimated from oxygen and carbon stable isotope analyses performed on ECG (Prud'homme *et al.* 2016, 2018). During the formation of GBU, the air temperature for the five warmest months was 13.0 ± 4.1 °C and the mean palaeoprecipitation was $287 + 223 / - 161$ mm a^{-1} , whereas during the formation of the Lohner Boden, the temperature of the warmest season was 11.7 ± 3.6 °C and the mean rainfall was $322 + 236 / - 171$ mm a^{-1} (Prud'homme *et al.* 2016, 2018). The palaeoenvironmental reconstructions from the coastal Atlantic marine




Fig. 5. Correlation between ECG concentration and quantitative palaeoclimate parameters (temperature and precipitation; Prud'homme *et al.* 2016, 2018) of the Nussloch loess sequence with the Greenland ice-core record (Rasmussen *et al.* 2014) and the Heinrich events (Rasmussen *et al.* 2016) based on the ECG radiocarbon chronology of the loess sequence (Moine *et al.* 2017).

core MD04 2845 next to the French coast and located above 45°N indicate that GI 12 is characterized by a mild and moderately humid environment with mainly *Pinus* forest whereas GI 8 is still mild but less humid with a vegetation represented by grassland and dry shrubland (Fletcher *et al.* 2010).

The difference in humidity between GI 12 and GI 8 recorded in Nussloch loess sequence is also confirmed by the pedostratigraphy. The LB is an arctic brown soil with the presence of cryostructures (platy structure indicating ice-lensing processes), which indicate more humid conditions than in the boreal brown soils (GBU) (Antoine *et al.* 2009). Furthermore, during the Upper Pleniglacial, palaeoclimate data indicate that tundra gley horizon G2a/b was formed during mild and humid conditions (Fig. 5). The air temperature of the warmest season was 9.4 ± 4.3 °C and the mean annual precipitation was $425 + 273 / - 201$ mm a⁻¹. These climate conditions are supported by (i) the very high proportion (more than 50%) of the hygrophilous mollusc taxon *Succinella oblonga* (Moine *et al.* 2005, 2008) and by (ii) more intense hydromorphic features, such as stronger reduction, greenish colour and more developed iron oxide precipitation around root tracks (Antoine *et al.* 2009). The two horizons G2a and G2b of tundra gley G2 are respectively correlated with GI 5.2 and GI 5.1 (Moine *et al.* 2017). This interstadial (600 years) is characterized by an increase of +12.5 °C prior to the Heinrich Event H3 (Kindler *et al.* 2014; Rasmussen *et al.* 2014). Wetter conditions are also recorded on the laminated Eifel maar sediments in Germany, where four flooding events were discovered around 30 to 31 ka, suggesting intense spring snow-melt events (Sirocko *et al.* 2015). This period (around 31–30 ka) corresponds to the first step of the maximal expansion of the Fennoscandian ice sheet, implying an increase in moisture conditions in NW Europe (Hardt *et al.* 2016; Hughes *et al.* 2016). This time-span is also characterized by the first considerable increase in typical loess deposition throughout north-western Europe (Antoine *et al.* 2016).

Conclusions

The first continuous record of the variations in large ECG concentration through the Nussloch loess sequence led to the discovery of very high granule concentrations in boreal and arctic brown soils and tundra gleys horizons, whereas loess deposits are distinguished by very low values that are even close to zero in laminated loess units. Generally, the variation in ECG concentration is well correlated with stratigraphical boundaries and variations in clay percentages, suggesting a link with palaeoclimate and palaeoenvironmental changes. Nevertheless, the maximal abundance of ECG is not found at the surface (as in the present-day soil profile) but 15 to 30 cm deeper. This observation suggests that loess sedimenta-

tion was not null during the formation of boreal and arctic brown soils and tundra gleys horizons. Moreover, the high concentration of large granules in the incipient tundra gley horizons indicates that these units may be associated with rapid and short climate improvement but less intense than that during the interstadial phases, which is favourable to earthworm activity. However, the absence of terrestrial molluscs in these horizons suggests that earthworms are more resistant to periods of marked climatic stress (cold and arid conditions) as they can protect themselves by burrowing into the soil, especially in very exposed conditions such as at the top of the Nussloch loess gredas.

Furthermore, the good correlation between the ECG concentration and the clay content in tundra gley horizons suggests that the beginning of the warming associated with the increase in humidity is anterior to the formation of the stratigraphical units and thus to the functioning of the active layer.

The comparison between the high concentration of ECG in palaeosols and the radiocarbon dating confirmed that boreal and arctic brown soils and tundra gley horizons were formed under milder climatic conditions (interstadial events) associated with the development of biodiversity.

Finally, the three drastic decreases in ECG concentration suggest that they are contemporaneous with peaks of dust recorded in the Greenland ice-core associated with Heinrich Events (H3 and H2). Thus, the absence of large ECG can be interpreted as a new indicator of aridity through the loessic sequence of Nussloch, even though the role of potentially very low temperatures and the increase in wind intensity during these three cold stages should also be considered. This approach could therefore highlight in the future, climatic oscillations that are hardly identifiable in other contexts. Moreover, the same pattern is observed in two loess sequences of northern France (600 km to the west) in the same period (30 ka), suggesting a global aridity signal over western Europe. In the future, an investigation of variation in the ECG concentration at the top and at the sides of a greda in loess-palaeosol sequences formed in gentle topography could provide information about the migration of earthworms and their possible refuges during the Last Glacial.

The comparison between ECG concentration, grain-size variations, palaeoclimate data and radiocarbon dating confirms that ECG can be used as a new palaeoenvironmental and palaeoclimate proxy of the Last Glacial in continental environments. Therefore, it can help to better understand the response of terrestrial archives to the millennial climate variations highlighted in ice-cores.

Acknowledgements. – This work received financial support from the LabEx DynamiTe (ANR-11-LABX-0046), as part of the 'Investissements d'Avenir' program. The authors thank the two anonymous reviewers and Jan A. Piotrowski for their constructive comments that helped us to improve the scientific content of this study.

References

- Almond, P. C. & Tonkin, P. J. 1999: Pedogenesis by upbuilding in an extreme leaching and weathering environment, and slow loess accretion, south Westland, New Zealand. *Geoderma* 92, 1–36.
- Antoine, P., Coutard, S., Guerin, G., Deschodt, L., Goval, E., Loch, J. L. & Paris, C. 2016: Upper Pleistocene loess-palaeosol records from Northern France in the European context: environmental background and dating of the Middle Palaeolithic. *Quaternary International* 411, 4–24.
- Antoine, P., Rousseau, D. D., Hatté, C., Zöller, L., Lang, A., Fontugne, M. & Moine, O. 2002: Événements éoliens rapides dans les loess du pléistocène supérieur weichselien : L'exemple de la séquence de Nussloch Vallée du Rhin-Allemagne. *Quaternaire* 13, 199–208.
- Antoine, P., Rousseau, D. D., Moine, O., Kunesch, S., Hatté, C., Lang, A., Tissoux, H. & Zöller, L. 2009: Rapid and cyclic aeolian deposition during the Last Glacial in European loess: a high-resolution record from Nussloch, Germany. *Quaternary Science Reviews* 28, 2955–2973.
- Antoine, P., Rousseau, D. D., Zöller, L., Lang, A., Munaut, H., Hatté, C. & Fontugne, M. 2001: High-resolution record of the last Interglacial glacial cycle in the Nussloch loess palaeosol sequences, Upper Rhine Area, Germany. *Quaternary International* 77, 211–229.
- Barta, G. 2014: Palaeoenvironmental reconstruction based on the morphology and distribution of secondary carbonates of the loess-palaeosol sequence at Süttő, Hungary. *Quaternary International* 319, 64–75.
- Becze-Deák, J., Langohr, R. & Verrecchia, E. P. 1997: Small scale secondary CaCO₃ accumulations in selected sections of the European loess belt. *Geoderma* 76, 221–252.
- Bond, G. C. & Lotti, R. 1995: Iceberg discharges into the north Atlantic on millennial time scales during the last glaciation. *Science* 267, 1005–1010.
- Bouché, M. B. 1972: *Lombriciens de France. Ecologie et Systématique*. 671 pp. INRA, Paris.
- Bräm, H. 1956: Was sind Arion kinkelinii Wenz und A. hochheimensis Wenz. *Ecologiae Geologicae Helveticae* 49, 593–598.
- Briones, M. J. I., López, E., Méndez, J., Rodríguez, J. B. & Gago-Duport, L. 2008: Biological control over the formation and storage of amorphous calcium carbonate by earthworms. *Mineralogical Magazine* 72, 227–231.
- Burn, C. 2012: Permafrost distribution and stability. In French, H. & Slaymaker, O. (eds.): *Changing Cold Environments: A Canadian Perspective*, 126–146. Wiley, Chichester.
- Calderon, S., Holmstrup, M., Westh, P. & Overgaard, J. 2009: Dual roles of glucose in the freeze-tolerant earthworm *Dendrobaena octaedra*: cryoprotection and fuel for metabolism. *Journal of Experimental Biology* 212, 859–866.
- Canti, M. G. 1998: Origin of calcium carbonate granules found in buried soils and Quaternary deposits. *Boreas* 27, 275–288.
- Canti, M. G. 2007: Deposition and taphonomy of earthworm granules in relation to their interpretative potential in Quaternary stratigraphy. *Journal of Quaternary Science* 22, 111–118.
- Canti, M. G. & Pearce, T. G. 2003: Morphology and dynamics of calcium carbonate granules produced by different earthworm species. *Pedobiologia* 47, 511–521.
- Capron, E., Landais, A., Chappellaz, J., Schilt, A., Buiron, D., Dahl-Jensen, D., Johnsen, S. J., Jouzel, J., Lemieux-Dudon, B., Loulergue, L., Leuenberger, M., Masson-Delmotte, V., Mayer, H., Oerter, H. & Stenni, B. 2010: Millennial and sub-millennial scale climatic variations recorded in polar ice cores over the last glacial period. *Climate of the Past Discussions* 6, 135–183.
- Caro, G., Decaëns, T., Lecarpentier, C. & Mathieu, J. 2013: Are dispersal behaviours of earthworms related to their functional group? *Soil Biology and Biochemistry* 58, 181–187.
- Darwin, C. 1881: The formation of vegetable mould through the action of worms, with observations on their habits. *Science* 3, 26–26.
- Eijsackers, H. 2011: Earthworms as colonizers of natural and cultivated soil environments. *Applied Soil Ecology* 50, 1–13.
- Fletcher, W. J., Sánchez-Goni, M. F., Allen, J. R. M., Cheddadi, R., Combourieu-Nebout, N., Huntley, B., Lawson, I., Londeix, L., Magri, D., Margari, V., Müller, U. C., Naughton, F., Novenko, E., Roucoux, K. & Tzedakis, P. C. 2010: Millennial-scale variability during the last glacial in vegetation records from Europe. *Quaternary Science Reviews* 29, 2839–2864.
- French, H. M. 2007: *The Periglacial Environment*. 478 pp. Wiley, Chichester.
- Gago-Duport, L., Briones, M. J. I., Rodríguez, J. B. & Covelo, B. 2008: Amorphous calcium carbonate biomineralization in the earthworm's calciferous gland: pathways to the formation of crystalline phases. *Journal of Structural Biology* 162, 422–435.
- Gerard, B. M. 1967: Factors affecting earthworms in pastures. *Journal of Animal Ecology* 36, 235–252.
- Haase, D., Fink, J., Haase, G., Ruske, R., Pécsi, M., Richter, H., Altermann, M. & Jäger, K. D. 2007: Loess in Europe—its spatial distribution based on a European Loess Map, scale 1:2,500,000. *Quaternary Science Reviews* 26, 1301–1312.
- Haesaerts, P., Damblon, F., Gerasimenko, N., Spagna, P. & Pirson, S. 2016: The Late Pleistocene loess-palaeosol sequence of Middle Belgium. *Quaternary International* 411, 25–43.
- Hardt, J., Lüthgens, C., Hebenstreit, R. & Böse, M. 2016: Geochronological OSL: and geomorphological investigations at the presumed Frankfurt ice marginal position in northeast Germany. *Quaternary Science Reviews* 154, 85–99.
- Harrison, S. P. & Sanchez Goñi, M. F. 2010: Global patterns of vegetation response to millennial-scale variability and rapid climate change during the last glacial period. *Quaternary Science Reviews* 29, 2957–2980.
- Heinrich, H. 1988: Origin and consequences of cyclic ice rafting in the Northeast Atlantic Ocean during the past 130,000 years. *Quaternary Research* 29, 142–152.
- Hodson, M. E., Benning, L. G., Demarchi, B., Penkman, K. E. H., Rodriguez-Blanco, J. D., Schofield, P. F. & Versteegh, E. A. A. 2015: Biomineralisation by earthworms – an investigation into the stability and distribution of amorphous calcium carbonate. *Geochemical Transactions* 16, 1–16.
- Holmstrup, M. 1994: Physiology of cold hardiness in cocoons of five earthworm taxa Lumbricidae: Oligochaeta. *Journal of Comparative Physiology B* 164, 222–228.
- Holmstrup, M. 2003: Overwintering adaptations in earthworms. *Pedobiologia* 47, 504–510.
- Holmstrup, M. & Zachariassen, K. E. 1996: Physiology of cold hardiness in earthworms. *Comparative Biochemistry and Physiology - A Physiology* 115, 91–101.
- Hughes, A. L. C., Gyllencreutz, R., Lohne, Ø. S., Mangerud, J. & Svendsen, J. I. 2016: The last Eurasian ice sheets - a chronological database and time-slice reconstruction, DATED-1. *Boreas* 45, 1–45.
- Jänsch, S., Steffens, L., Höfer, H., Horak, F., Roß-Nickoll, M., Russell, D., Toschki, A. & Römbke, J. 2013: State of knowledge of earthworm communities in German soils as a basis for biological soil quality assessment. *Soil Organisms* 85, 215–233.
- Kerney, M. P. 1971: A Middle Weichselian deposit at Halling, Kent. *Proceedings of the Geologists' Association* 82, 1–11.
- Kindler, P., Guillevic, M., Baumgartner, M., Schwander, J., Landais, A. & Leuenberger, M. 2014: Temperature reconstruction from 10 to 120 kyr b2k from the NGRIP ice core. *Climate of the Past* 10, 887–902.
- Landais, A., Caillon, N., Goujon, C., Grachev, A. M., Barnola, J. M., Chappellaz, J., Jouzel, J., Masson-Delmotte, V. & Leuenberger, M. 2004: Quantification of rapid temperature change during DO event 12 and phasing with methane inferred from air isotopic measurements. *Earth and Planetary Science Letters* 225, 221–232.
- Leger, M. 1990: Loess landforms. *Quaternary International* 7–8, 53–61.
- Mathieu, J. & Davies, T. J. 2014: Glaciation as an historical filter of below-ground biodiversity. *Journal of Biogeography* 41, 1204–1214.
- Mazenot, G. 1956: Recherches sur les faunes malacologiques du loess récent wurmien et de divers limons terrestres holocènes dans le sud-est de la France. *Société Linnéenne de Lyon* 25, 9–24.
- Meijer, T. 1985: The pre-Weichselian nonmarine molluscan fauna from Maastricht-Belvédère Southern Limburge, the Netherlands. *Analecta Praehistorica Leidensia* 18, 75–103.
- Meijs, E. P. M. 2011: The Veldwezelt site province of Limburg, Belgium: environmental and stratigraphical interpretations. *Geologie en Mijnbouw/Netherlands Journal of Geosciences* 90, 73–94.

- Menviel, L., Timmermann, A., Friedrich, T. & England, M. H. 2014: Hindcasting the continuum of Dansgaard-Oeschger variability: mechanisms, patterns and timing. *Climate of the Past* 10, 63–77.
- Meshcheryakova, E. N. & Berman, D. I. 2014: Cold hardness and geographic distribution of earthworms Oligochaeta, Lumbricidae, Moniligastridae. *Entomological Review* 94, 486–497.
- Meszner, S., Fuchs, M. & Faust, D. 2011: Loess-Palaeosol-Sequences from the loess area of Saxony Germany. *E&G Quaternary Science Journal* 60, 47–65.
- Meszner, S., Kreutzer, S., Fuchs, M. & Faust, D. 2013: Late Pleistocene landscape dynamics in Saxony, Germany: paleoenvironmental reconstruction using loess-paleosol sequences. *Quaternary International* 296, 94–107.
- Moine, O. 2008: West-European malacofauna from loess deposits of the Weichselian Upper Pleniglacial: compilation and preliminary analysis of the database. *Quaternaire* 19, 11–29.
- Moine, O., Antoine, P., Deschodt, L. & Sellier-Segard, N. 2011: Enregistrements malacologiques à haute résolution dans les loess et les gleys de toundra du pléniglaciaire weichselien supérieur: premiers exemples du nord de la France. *Quaternaire* 22, 307–325.
- Moine, O., Antoine, P., Hatté, C., Landais, A., Mathieu, J., Prud'homme, C. & Rousseau, D. D. 2017: The impact of Last Glacial climate variability in west-European loess revealed by radiocarbon dating of fossil earthworm granules. *Proceedings of the National Academy of Sciences of the United States of America* 114, 6209–6214.
- Moine, O., Rousseau, D. D. & Antoine, P. 2005: Terrestrial molluscan records of Weichselian Lower to Middle Pleniglacial climatic changes from the Nussloch loess series Rhine Valley, Germany. The impact of local factors. *Boreas* 34, 363–380.
- Moine, O., Rousseau, D. D. & Antoine, P. 2008: The impact of Dansgaard-Oeschger cycles on the loessic environment and malacofauna of Nussloch Germany: during the Upper Weichselian. *Quaternary Research* 70, 91–104.
- Nikiforoff, C. C. 1949: Weathering and soil evolution. *Soil Science* 67, 219–230.
- Nordström, S. 1975: Seasonal activity of lumbricids in Southern Sweden. *Oikos* 26, 307–315.
- Nuutinen, V. & Butt, K. R. 2009: Worms from the cold: lumbricid life stages in boreal clay during frost. *Soil Biology and Biochemistry* 41, 1580–1582.
- Pecsi, M. 1990: Loess is not the accumulation of dust. *Quaternary International* 7–8, 1–21.
- Perel', T. S. 1979: *Rasprostranenie i zakonmernosti raspredeleniä doždevyh červejfauny SSSR s opredelitel'nymi tablicami Lumbricidae i drugih Megadrili*. Nauka, Moskva.
- Prud'homme, C., Antoine, P., Moine, O., Turpin, E., Huguenard, L., Robert, V. & Degeai, J.-P. 2015: Earthworm calcite granules: a new tracker of millennial-timescale environmental changes in Last Glacial loess deposits. *Journal of Quaternary Science* 30, 529–536.
- Prud'homme, C., Lécuyer, C., Antoine, P., Moine, O., Hatté, C., Fourel, F., Amiot, R., Martineau, F. & Rousseau, D. D. 2018: $\delta^{13}\text{C}$ signal of earthworm calcite granules: a new proxy for palaeoprecipitation reconstructions during the Last Glacial in Western Europe. *Quaternary Science Reviews* 179, 158–166.
- Prud'homme, C., Lécuyer, C., Antoine, P., Moine, O., Hatté, C., Fourel, F., Martineau, F. & Rousseau, D. D. 2016: Palaeotemperature reconstruction during the Last Glacial from $\delta^{18}\text{O}$ of earthworm calcite granules from Nussloch loess sequence, Germany. *Earth and Planetary Science Letters* 442, 13–20.
- Rasmussen, L. M. & Holmstrup, M. 2002: Geographic variation of freeze-tolerance in the earthworm *Dendrobaena octaedra*. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology* 172, 691–698.
- Rasmussen, S. O., Bigler, M., Blockley, S. P., Blunier, T., Buchardt, S. L., Clausen, H. B., Cvijanovic, I., Dahl-Jensen, D., Johnsen, S. J., Fischer, H., Gkinis, V., Guillevic, M., Hoek, W. Z., Lowe, J. J., Pedro, J. B., Popp, T., Seierstad, I. K., Steffensen, J. P., Svensson, A. M., Vallenga, P., Vinther, B. M., Walker, M. J. C., Wheatley, J. J. & Winstrup, M. 2014: A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy. *Quaternary Science Reviews* 106, 14–28.
- Rasmussen, T. L., Thomsen, E. & Moros, M. 2016: North Atlantic warming during Dansgaard-Oeschger events synchronous with Antarctic warming and out-of-phase with Greenland climate. *Scientific Reports* 6, 20535.
- Remy, H. 1968: Zur Stratigraphie und Klimaentwicklung des jüngeren Pleistozäns in Mittel- und Westeuropa unter besonderer Berücksichtigung des Lösses. *Decheniana* 121, 121–145.
- Renssen, H., Kasse, C., Vandenbergh, J. & Lorenz, S. J. 2007: Weichselian Late Pleniglacial surface winds over northwest and central Europe: a model-data comparison. *Journal of Quaternary Science* 22, 281–293.
- Rousseau, D. D., Boers, N., Sima, A., Svensson, A., Bigler, M., Lagroix, F., Taylor, S. & Antoine, P. 2017: MIS3 & 2: millennial oscillations in Greenland dust and Eurasian aeolian records – a paleosol perspective. *Quaternary Science Reviews* 139, 99–113.
- Rousseau, D. D., Sima, A., Antoine, P., Hatté, C., Lang, A. & Zöller, L. 2007: Link between European and North Atlantic abrupt climate changes over the last glaciation. *Geophysical Research Letters* 34, L22713.
- Rundgren, A. S. 1975: Vertical distribution of Lumbricids in Southern Sweden. *Oikos* 26, 299–306.
- Satchell, J. 1967: Lumbricidae. In Burges, A. & Raw, F. (eds.): *Soil Biology*, 259–322. Academic Press, London.
- Schirmer, W. 2012: Rhine loess at Schwalbenberg II – MIS 4 and 3. *E&G Quaternary Science Journal* 61, 32–47.
- Schirmer, W. 2016: Late Pleistocene loess of the Lower Rhine. *Quaternary International* 411, 44–61.
- Sirocko, F., Knapp, H., Dreher, F., Förster, M. W., Albert, J., Brunck, H., Veres, D., Dietrich, S., Zech, M., Hambach, U., Röhrner, M., Rudert, S., Schwibus, K., Adams, C. & Sigl, P. 2015: The ELSA-vegetation-stack: reconstruction of landscape evolution zones LEZ: from laminated Eifel maar sediments of the last 60,000 years. *Global and Planetary Change* 142, 108–135.
- Tiunov, A. V., Hale, C. M., Holdsworth, A. R. & Vsevolodova-Perel, T. S. 2006: Invasion patterns of Lumbricidae into the previously earthworm-free areas of northeastern Europe and the western Great Lakes region of North America. *Biological Invasions Belowground: Earthworms as Invasive Species* 8, 1223–1234.

Supporting Information

Additional Supporting Information may be found in the online version of this article at <http://www.boreas.dk>.

Table S1. Numerical data of granule counting and granulometry analysis (clay content <4.6 μm , coarse sand content >160 μm) of Nussloch profile P8.