



New Expectation from DNP-Enhanced SS-NMR to figure out the Role of the Organic-Silica Interfaces: the Case of Diatom Frustules and Marine Siliceous Sponge Spicules

Sylvie Masse, Guillaume Laurent, Thibaud Coradin, Andrzej Pisera

► To cite this version:

Sylvie Masse, Guillaume Laurent, Thibaud Coradin, Andrzej Pisera. New Expectation from DNP-Enhanced SS-NMR to figure out the Role of the Organic-Silica Interfaces: the Case of Diatom Frustules and Marine Siliceous Sponge Spicules. Magnetism and Magnetic Resonance: Magnetic Resonance, Understanding, Measurements and Modeling, Jun 2019, Strasbourg, France. . hal-02156116

HAL Id: hal-02156116

<https://hal.sorbonne-universite.fr/hal-02156116>

Submitted on 18 Jun 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

New Expectation from DNP-Enhanced SS-NMR to figure out the Role of the Organic-Silica Interfaces: the Case of Diatom Frustules and Marine Siliceous Sponge Spicules



Sylvie MASSE ¹, **Guillaume LAURENT** ¹, **Thibaud CORADIN** ¹ and **Andrzej PISERA** ²
¹: Lab. Chimie de la Matière Condensée de Paris, Sorbonne Université, 4 place Jussieu, 75005 Paris, France
²: Institute of Paleobiology, Polish Academy of Sciences, ul. Twarda 51/55, 00-818 Warszawa, Poland






Microalgal culture at ISOMER- in collaboration with Dr Véronique Martin-Jézéquel and Dr Benoit Tesson Faculté des Sciences et Techniques, Nantes, France




NMR Facility-Sorbonne Université Campus P. et M. Curie -T 32-33 SB





Living sponge collection: Pr A. PISERA on board of the IRD ship for an expedition in New Caledonia (Financially supported by National Science Centre, Grant No. 2016/21/B/ST10/02332)




The two-valves silica cell wall (« Frustule ») of diatoms: the silaffin-mediated formation of a silica shell embedded with organic matter.



Frustule and associated organic matter, from Hatté et al, Mar. Chem., 2008




¹H MAS and ³¹P HPDEC-MAS NMR spectra of the Whole-cell, SDS-treated and H₂O₂-treated diatom frustule samples isotopically enriched in ²⁹Si, ¹³C/²⁹Si/¹⁵N and ¹³C/¹⁵N, resp. (Ref.1). A signal broadening and a loss in intensity are observed after chemical treatment. While SDS/EDTA is used first to clean the frustule, further H₂O₂ treatment seems to be much more aggressive, probably leading to partial dissolution-recrystallization





SEM pictures of a series of sponge spicules

(a,b) Petrosid demosponge PETRO: (a) untreated, scale bar = 500 mm, (b) after organics removal, scale bar = 200 mm; (c,d) Geodia sp demosponge GEODIA: (c) untreated, scale bar = 1mm, (d) untreated, scale bar = 20 mm; (e,f) Aphrocallistes sp. hexactinellid sponge APHRO: (e) untreated, scale bar = 1 mm, (f) after organics removal, scale bar = 500 mm; (g,h) Laocoetis perion hexactinellid LAOCO: (g) after organics removal, scale bar = 2 mm, (h) after organics removal, scale bar = 200 mm.




The silica skeleton of sponges: the silicatein-mediated formation of a silica shell around an axial filament, mainly composed of proteins.




Variability in ¹³C CP-MAS NMR response depending on species and history: nature of the taxon, aging, conservation, chemical treatment... (Ref.2).

Sensitivity limit



Peak Signal-to-Noise Ratio (PSNR) and Root Mean Square (rms) for the ¹³C CP-MAS SS-NMR spectrum of the ThalaE (diatom frustule) and Laocoetis Perion (sponge spicule) samples. Signal and noise regions are highlighted with dotted vertical red and green lines, resp.; critical (L_c), detection (L_d) and quantitative (L_q) limits with dotted horizontal red, orange and green lines, resp. (according to ref. 3). In both cases, signal is detected but must be amplified.

(¹H)-²⁹Si-¹³C double-CP MAS NMR spectrum at 9.4T (AV400) of a triply-enriched (¹³C,²⁹Si,¹⁵N) Thalassiosira Pseudonana frustule sample (d1=3s, tcp1=3ms, tcp2=40ms, NS=9.600, ro= 5kHz and LB=100Hz).



Conclusion

While a lot of work is needed to figure out the organic-silica interfaces in natural materials such as diatom frustules or marine siliceous sponge spicules, Solid-State NMR appears to be a powerful toolbox with several nuclei and methods to carry out. Nevertheless, natural abundance in ¹³C as well as a too poor C-content in the cleaned specimen do not allow nor 2D correlations neither well-resolved 1D spectra, that are necessary to go further on species proximity and connectivity assessment. Conjugating DNP to SS-NMR appears to be a promising solution to enhance the signal.

References

- Tesson B., Masse S., Laurent G., Maquet J., Livage J., Martin-Jézéquel V. and Coradin T. « Contribution of multi-nuclear solid state NMR to the characterization of the Thalassiosira pseudonana diatom cell wall », *Anal Bioanal Chem*, 390, 1889–1898 (2008), doi 10.1007/s00216-008-1908-0
- Masse S., Pisera A., Laurent G., Coradin T. « A Solid-State NMR Investigation of Recent Marine Siliceous Sponge Spicules », *Minerals*, 6, 21-30 (2016), doi: 10.3390/min6010021
- Laurent G., Woelfel W., Barret-Vivien V., Gouillart E., Bonhomme C. « Denoising applied to spectroscopies – part I : concept and limits », *Appl. Spectrosc. Rev.*, doi: 10.1080/05704928.2018.1523183

3rd Thematic School: Magnetism and Magnetic Resonance

Magnetic Resonance, Understanding, Measurements and Modeling – June 2-6, 2019, Strasbourg, France