



HAL
open science

Cohomology classes of strata of differentials

Adrien Sauvaget

► **To cite this version:**

Adrien Sauvaget. Cohomology classes of strata of differentials. *Geometry & Topology Monographs*, 2019, Cohomology classes of strata of differentials, 23 (3), pp.1085-1171. 10.2140/gt.2019.23.1085 . hal-02170857v2

HAL Id: hal-02170857

<https://hal.sorbonne-universite.fr/hal-02170857v2>

Submitted on 8 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cohomology classes of strata of differentials

ADRIEN SAUVAGET

We introduce a space of stable meromorphic differentials with poles of prescribed orders and define its tautological cohomology ring. This space, just as the space of holomorphic differentials, is stratified according to the set of multiplicities of zeros of the differential. The main goal of this paper is to compute the Poincaré-dual cohomology classes of all strata. We prove that all these classes are tautological and give an algorithm to compute them.

In the second part of the paper we study the Picard group of the strata. We use the tools introduced in the first part to deduce several relations in these Picard groups.

14C17, 14H10, 30F30, 32G15

1. Introduction	1085
2. Stable differentials	1098
3. Stratification of spaces of stable differentials	1118
4. Boundary components of strata of stable differentials	1133
5. Computation of classes of strata	1147
6. Examples of computation	1160
7. Relations in the Picard group of the strata	1162
References	1169

1 Introduction

1.1 Stratification of the Hodge bundle

Let $g \geq 1$. Let \mathcal{M}_g be the space of smooth curves of genus g . The *Hodge bundle*

$$\mathcal{H}_g \rightarrow \mathcal{M}_g$$

is the vector bundle whose fiber over a point $[C]$ of \mathcal{M}_g is the space of holomorphic differentials on C . A point of \mathcal{H}_g is then a pair $([C], \alpha)$, where C is a curve and α a differential on C . We will denote by $\mathbb{P}\mathcal{H}_g \rightarrow \mathcal{M}_g$ the projectivization of the Hodge bundle.

Notation 1.1 Let Z (for zeros) be a vector (k_1, k_2, \dots, k_n) of positive integers satisfying

$$\sum_{i=1}^n k_i = 2g - 2.$$

We will denote by $\mathbb{P}\mathcal{H}_g(Z)$ the subspace of $\mathbb{P}\mathcal{H}_g$ composed of pairs $([C], \alpha)$ such that α is a differential (defined up to a multiplicative constant) with zeros of orders k_1, \dots, k_n .

The locus $\mathbb{P}\mathcal{H}_g(Z)$ is a smooth orbifold (or a Deligne–Mumford stack); see for instance Polishchuk [32]. However, neither $\mathbb{P}\mathcal{H}_g$ nor the strata $\mathbb{P}\mathcal{H}_g(Z)$ are compact.

The Hodge bundle has a natural extension to the space of stable curves:

$$\overline{\mathcal{H}}_g \rightarrow \overline{\mathcal{M}}_g.$$

We recall that abelian differentials over a nodal curve are allowed to have simple poles at the nodes with opposite residues on the two branches.

The space $\mathbb{P}\overline{\mathcal{H}}_g$ is compact and smooth, and we can consider the closures $\mathbb{P}\overline{\mathcal{H}}_g(Z)$ of the strata inside this space. Computing the Poincaré-dual cohomology classes of these strata is our motivating problem. In this paper we solve this problem and present a more general computation in the case of meromorphic differentials.

1.2 Stable differentials

On a fixed smooth curve C with one marked point x consider a family of meromorphic differentials with one pole of order p at x such that the leading coefficient of the differential at the pole tends to 0. In order to construct a compact moduli space of meromorphic differentials we need to decide what the limit of a family like that should be. One natural idea is to include differentials with poles of orders less than p in the moduli space. It turns out, however, that a more convenient way to represent the limit is to allow the underlying curve to bubble at x ; in other words, to allow differentials defined on semistable curves.

The first uses of semistable objects to compactify moduli problems can be found in the work of Gieseker for the moduli space of stable bundles (see [20]), or in Caporaso's construction of a universal Picard variety over the moduli space of curves (see [6]).

A *semistable curve* is a nodal curve with smooth marked points such that every genus 0 component of its normalization contains at least two marked points and preimages of

nodes (instead of at least three for stable curves). In the example above, the limit of the family would be a meromorphic differential defined on a semistable curve with one unstable component and one marked point x on it. The curve maps to C under the contraction of the unstable component. The meromorphic differential still has a pole of order exactly p at x .

Definition 1.2 Let $n, m \in \mathbb{N}$ and let P (for poles) be a vector (p_1, p_2, \dots, p_m) of positive integers. A *stable differential* of type (g, n, P) is a tuple $(C, x_1, \dots, x_{n+m}, \alpha)$, where (C, x_1, \dots, x_{n+m}) is a semistable curve with $n + m$ marked points and α is a meromorphic differential on C , such that

- the differential α has no poles outside the m last marked points and nodes;
- the poles at the nodes are at most simple and have opposite residues on the two branches;
- if $p_i > 1$ then the pole at the marked point x_{n+i} is of order exactly p_i ; if $p_i = 1$ then x_i can be a simple pole, a regular point or a zero of any order;
- the group of isomorphisms of C preserving α and the marked points is finite.

Definition 1.3 A *family of stable differentials* is a tuple $(C \rightarrow B, \sigma_1, \dots, \sigma_n, \alpha)$ where $(C \rightarrow B, \sigma_1, \dots, \sigma_n)$ is a family of marked semistable curves and α is a meromorphic section of the relative dualizing line bundle $\omega_{C/B}$ such that for each geometric point b of B , the tuple $(C_b, \sigma_1(b), \dots, \sigma_n(b), \alpha|_{C_b})$ is a stable differential.

The stack $\overline{\mathfrak{H}}_{g,n,P}$ of stable differentials of type (g, n, P) is the category of families of stable differentials of type (g, n, P) , fibered over the category of \mathbb{C} -schemes.

Proposition 1.4 The moduli space $\overline{\mathfrak{H}}_{g,n,P}$ is a smooth Deligne–Mumford (DM) stack. It is of dimension $4g - 4 + \sum p_i$ if P is nonempty and $4g - 3$ otherwise.

The space $\overline{\mathfrak{H}}_{g,n,P}$ carries a natural \mathbb{C}^* -action given by the multiplication of the differential by nonzero scalars. Further, there exists a forgetful map $\overline{\mathfrak{H}}_{g,n,P} \rightarrow \overline{\mathcal{M}}_{g,n+m}$ that maps a family stable differentials to the stabilization of its underlying family of semistable curves. However, the space $\overline{\mathfrak{H}}_{g,n,P}$ does not have a natural vector bundle structure in general because there is no natural definition of the sum of two differentials with fixed orders of poles.

We will construct a partial coarsification of $\overline{\mathfrak{H}}_{g,n,P}$ that has the structure of an orbifold cone over $\overline{\mathcal{M}}_{g,n+m}$.

Proposition 1.5 *There exists a unique DM stack $\overline{\mathcal{H}}_{g,n,P}$ fitting in the commutative triangle*

$$\begin{array}{ccc}
 \overline{\mathfrak{H}}_{g,n,P} & \longrightarrow & \overline{\mathcal{H}}_{g,n,P} \\
 & \searrow & \downarrow \pi \\
 & & \overline{\mathcal{M}}_{g,n+m}
 \end{array}$$

and satisfying:

- The morphism π is schematic, ie for any \mathbb{C} -scheme U with a morphism $U \rightarrow \overline{\mathcal{M}}_{g,n+m}$, the pullback $\overline{\mathcal{H}}_{g,n,P} \times_{\overline{\mathcal{M}}_{g,n+m}} U$ is representable by a \mathbb{C} -scheme.
- For any such $U \rightarrow \overline{\mathcal{M}}_{g,n+m}$, the scheme $\overline{\mathcal{H}}_{g,n,P} \times_{\overline{\mathcal{M}}_{g,n+m}} U$ is the coarse space of $\overline{\mathfrak{H}}_{g,n,P} \times_{\overline{\mathcal{M}}_{g,n+m}} U$.

Definition 1.6 The space $\overline{\mathcal{H}}_{g,n,P}$ is called the *space of stable differentials*.

Proposition 1.7 *The space of stable differentials is an orbifold cone over $\overline{\mathcal{M}}_{g,n+m}$. Further, the space $\overline{\mathcal{H}}_{g,n,P}$ and its projectivization are normal.*

We prove these propositions in Section 2, where we will also give a definition of an orbifold cone. At present it suffices to note that the cone structure on $\overline{\mathcal{H}}_{g,n,P}$ allows one to define a projectivization $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$, a line bundle $\mathcal{O}(1)$ over the projectivization, and the Segre class. Further, the morphism $\overline{\mathfrak{H}}_{g,n,P} \rightarrow \overline{\mathcal{H}}_{g,n,P}$ is \mathbb{C}^* -equivariant.

Remark 1.8 The stack $\overline{\mathfrak{H}}_{g,n,P}$ can be endowed with the structure of an orbifold cone over a different moduli space $\overline{\mathcal{M}}_{g,n,P}$. The space $\overline{\mathcal{M}}_{g,n,P}$ is a $(\prod_{i=1}^m \mathbb{Z}/(p_i-1)\mathbb{Z})$ -gerbe over $\overline{\mathcal{M}}_{g,n+m}$. The fibers of $\overline{\mathfrak{H}}_{g,n,P} \rightarrow \overline{\mathcal{M}}_{g,n,P}$ are vector spaces, but the \mathbb{C}^* -action on these spaces has nontrivial weights.

One can define the projectivization of $\overline{\mathfrak{H}}_{g,n,P}$ and the tautological line bundle over this projectivization. Then we have a map $\mathbb{P}\overline{\mathfrak{H}}_{g,n,P} \rightarrow \mathbb{P}\overline{\mathcal{H}}_{g,n,P}$ which is a bijection between the geometric points of these two stacks. Therefore, we have natural isomorphisms $H^*(\mathbb{P}\overline{\mathfrak{H}}_{g,n,P}, \mathbb{Q}) \simeq H^*(\mathbb{P}\overline{\mathcal{H}}_{g,n,P}, \mathbb{Q})$ and $A^*(\mathbb{P}\overline{\mathfrak{H}}_{g,n,P}, \mathbb{Q}) \simeq A^*(\mathbb{P}\overline{\mathcal{H}}_{g,n,P}, \mathbb{Q})$. Thus, all the results of this text are valid for both spaces.

While the space $\overline{\mathfrak{H}}_{g,n,P}$ is the more natural choice for the moduli space of differentials, in this paper we prefer to work with $\overline{\mathcal{H}}_{g,n,P}$ in order to have $\overline{\mathcal{M}}_{g,n+m}$ as the base of our cone.

Notation 1.9 Let $P = (p_1, \dots, p_m)$ be a vector of positive integers and $Z = (k_1, \dots, k_n)$ a vector of nonnegative integers. We denote by $A_{g,Z,P} \subset \overline{\mathcal{H}}_{g,n,P}$ the locus of stable differentials $(C, x_1, \dots, x_{n+m}, \alpha)$ such that C is smooth and α has zeros exactly of orders prescribed by Z at the first n marked points. The locus $A_{g,Z,P}$ is invariant under the \mathbb{C}^* -action. We denote by $\mathbb{P}A_{g,Z,P}$ the projectivization of $A_{g,Z,P}$. Moreover, we denote by $\overline{A}_{g,Z,P}$ (resp. $\mathbb{P}\overline{A}_{g,Z,P}$) the closures of $A_{g,Z,P}$ (resp. $\mathbb{P}A_{g,Z,P}$) in the space $\overline{\mathcal{H}}_{g,n,P}$ (resp. in $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$).

1.3 The tautological ring of $\overline{\mathcal{M}}_{g,n}$

Let g and n be nonnegative integers satisfying $2g - 2 + n > 0$. Let $\overline{\mathcal{M}}_{g,n}$ be the space of stable curves of genus g with n marked points. Define the following cohomology classes:

- $\psi_i = c_1(\mathcal{L}_i) \in H^2(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$, where \mathcal{L}_i is the cotangent line bundle at the i^{th} marked point.
- $\kappa_m = \pi_*(\psi_{n+1}^{m+1}) \in H^{2m}(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$, where $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ is the forgetful map.
- $\lambda_k = c_k(\overline{\mathcal{H}}_{g,n}) \in H^{2k}(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ for $k = 1, \dots, g$.

Definition 1.10 A *stable graph* is the datum of

$$\Gamma = (V, H, g: V \rightarrow \mathbb{N}, a: H \rightarrow V, i: H \rightarrow H, E, L)$$

satisfying the following properties:

- V is a vertex set with a genus function g .
- H is a half-edge set equipped with a vertex assignment a and an involution i .
- E , the edge set, is defined as the set of length 2 orbits of i in H (self-edges at vertices are permitted).
- (V, E) defines a connected graph.
- L is the set of fixed points of i called *legs*.
- For each vertex v , the stability condition holds: $2g(v) - 2 + n(v) > 0$, where $n(v) = \#(a^{-1}(v))$ (the cardinality of $a^{-1}(v)$).

The genus of Γ is defined by $\sum g(v) + \#(E) - \#(V) + 1$.

Let $v(\Gamma)$, $e(\Gamma)$ and $n(\Gamma)$ denote the cardinalities of V, E and L , respectively. A boundary stratum of the moduli space of curves naturally determines a stable graph of genus g with n legs by considering the dual graph of a generic pointed curve parametrized by the stratum. Thus, the boundary strata of $\overline{\mathcal{M}}_{g,n}$ are in one-to-one correspondence with stable graphs.

Let Γ be a stable graph. Define the moduli space $\overline{\mathcal{M}}_\Gamma$ by the product

$$\overline{\mathcal{M}}_\Gamma = \prod_{v \in V} \overline{\mathcal{M}}_{g(v),n(v)},$$

and let $\zeta_\Gamma: \overline{\mathcal{M}}_\Gamma \rightarrow \overline{\mathcal{M}}_{g,n}$ be the natural morphism.

Definition 1.11 A *tautological class* is a linear combination of classes β of the form

$$\beta = \zeta_\Gamma * \left(\prod_{v \in V} P_v \right),$$

where Γ is a stable graph and P_v is a polynomial in κ, λ and ψ classes on $\overline{\mathcal{M}}_{g(v),n(v)}$.

Proposition/Definition 1.12 Let $RH^*(\overline{\mathcal{M}}_{g,n}) \subset H^*(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ be the vector subspace spanned by tautological classes. This subspace is a subring, called the **tautological ring** of $\overline{\mathcal{M}}_{g,n}$.

See Graber and Pandharipande [21] for the description of the product of tautological classes.

Remark 1.13 Actually, the classes α as above that do not involve λ -classes span the tautological ring. However, it will be more convenient for us to use this larger set of generators.

1.4 The tautological ring of $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$

Let P be a vector of positive integers. From now on, unless specified otherwise, we will denote by $\pi: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ the forgetful map and by $p: \overline{\mathcal{H}}_{g,n,P} \rightarrow \overline{\mathcal{M}}_{g,n+m}$ the projection from the space of stable differentials to $\overline{\mathcal{M}}_{g,n}$. Moreover, we use the same notation $p: \mathbb{P}\overline{\mathcal{H}}_{g,n,P} \rightarrow \overline{\mathcal{M}}_{g,n+m}$ for the projectivized cone. Let

$$\mathcal{L} = \mathcal{O}(1) \rightarrow \mathbb{P}\overline{\mathcal{H}}_{g,n,P}$$

be the tautological line bundle of $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$, and let $\xi = c_1(\mathcal{L})$.

Definition 1.14 The tautological ring of $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$ is the subring of the cohomology ring $H^*(\mathbb{P}\overline{\mathcal{H}}_{g,n,P}, \mathbb{Q})$ generated by ξ and the pullback of $RH^*(\overline{\mathcal{M}}_{g,n+m})$ under p . We denote it by $RH^*(\mathbb{P}\overline{\mathcal{H}}_{g,n,P})$.

Remark 1.15 We have $\xi^d = 0$ for $d > \dim(\mathbb{P}\overline{\mathcal{H}}_{g,n,P})$. Therefore, the tautological ring of $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$ is a finite extension of the tautological ring of $\overline{\mathcal{M}}_{g,n+m}$.

Example 1.16 In the absence of poles, the Hodge bundle is a vector bundle and we have

$$RH^*(\mathbb{P}\overline{\mathcal{H}}_{g,n}) = RH^*(\overline{\mathcal{M}}_{g,n})[\xi]/(\xi^g + \lambda_1 \xi^{g-1} + \dots + \lambda_g).$$

Proposition 1.17 The Segre class of the cone $\overline{\mathcal{H}}_{g,n,P} \rightarrow \overline{\mathcal{M}}_{g,n+m}$ equals

$$\prod_{i=1}^m \frac{(p_i - 1)^{p_i - 1}}{(p_i - 1)!} \cdot \frac{1 - \lambda_1 + \dots + (-1)^g \lambda_g}{\prod_{i=1}^m (1 - (p_i - 1)\psi_i)}.$$

This proposition will be proved in Section 2. An important corollary of this proposition is that the pushforward of a tautological class from $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$ to $\overline{\mathcal{M}}_{g,n+m}$ is tautological.

1.5 Statement of the results

Now, we have all elements to state the main theorems of this article.

Theorem 1 For any vectors Z and P , the class $[\mathbb{P}\overline{A}_{g,Z,P}]$ introduced in Notation 1.9 lies in the tautological ring of $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}$ and is explicitly computable.

The main ingredient to prove this theorem will be the induction formula of Theorem 5 (see page 1150).

Definition 1.18 Let V be a vector; in this article we will denote by $|V|$ the sum of elements of V and by $\ell(V)$ the length of V .

Given g and P , we will say that Z is complete if it satisfies $|Z| - |P| = 2g - 2$. If Z is complete, we denote by $Z - P$ the vector $(k_1, \dots, k_n, -p_1, \dots, -p_m)$.

Restricting ourselves to the holomorphic case and applying the forgetful map of the marked points we obtain the following corollary:

Theorem 2 For any complete vector Z , the class $[\mathbb{P}\overline{\mathcal{H}}_g(Z)]$ introduced in Notation 1.1 lies in the tautological ring of $\mathbb{P}\overline{\mathcal{H}}_g$ and is explicitly computable.

Remark 1.19 As a guideline for the reader, it will be important to understand that the holomorphic case in Theorem 1 cannot be proved without using strictly meromorphic differentials. Thus, Theorem 2 is a consequence of a specific case of Theorem 1 but one cannot avoid proving Theorem 1 in its full generality.

The second important corollary is obtained by forgetting the differential instead of the marked points. Let $P = (p_1, \dots, p_m)$ be a vector of poles and $Z = (k_1, \dots, k_n)$ be a complete vector of zeros. We define $\mathcal{M}_g(Z - P) \subset \mathcal{M}_{g,n+m}$ as the locus of points (C, x_1, \dots, x_n) that satisfy

$$\omega_C \left(-\sum_{i=1}^n k_i x_i + \sum_{j=1}^m p_j x_{n+j} \right) \simeq \mathcal{O}_C.$$

We denote by $\overline{\mathcal{M}}_g(Z - P)$ the closure of $\mathcal{M}_g(Z - P)$ in $\overline{\mathcal{M}}_{g,n+m}$.

Theorem 3 For any vectors Z and P , the class $[\overline{\mathcal{M}}_g(Z - P)]$ lies in the tautological ring of $\overline{\mathcal{M}}_{g,n+m}$ and is explicitly computable.

Remark 1.20 Theorems 1, 2 and 3 are stated for the Poincaré-dual rational cohomology classes. However, all identities of this paper are actually valid in the Chow groups.

In the second part of the text (Section 7) we will consider the rational Picard group of the space $\overline{\mathcal{M}}_g(Z - P)$. We will define several natural classes in this Picard group and apply the tools developed in the first part of the paper to deduce a series of relations between these classes (see Theorem 6).

1.6 An example

Here we illustrate the general method used in this article by computing the class of differentials with a double zero $[\mathbb{P}\overline{\mathcal{H}}_g(2, 1, \dots, 1)]$. This computation was carried out by D Zvonkine in an unpublished note [35] and was the starting point of the present work.

We begin by marking a point, ie we study the space $\mathbb{P}\overline{\mathcal{H}}_{g,1}$ of triples $(C, x_1, [\alpha])$ composed of a stable curve C with one marked point x_1 and an abelian differential α modulo a multiplicative constant. Recall that $\mathbb{P}\overline{\mathcal{A}}_{g,(2)} \subset \mathbb{P}\overline{\mathcal{H}}_{g,1}$ is the closure of the locus of smooth curves with a double zero at the marked point. In order to compute $[\mathbb{P}\overline{\mathcal{A}}_{g,(2)}]$, we consider the line bundle

$$\mathcal{L} \otimes \mathcal{L}_1 \simeq \text{Hom}(\mathcal{L}^\vee, \mathcal{L}_1)$$

over $\mathbb{P}\overline{\mathcal{H}}_{g,1}$. (Recall that \mathcal{L}^\vee is the dual tautological line bundle of the projectivization $\mathbb{P}\overline{\mathcal{H}}_{g,1}$ and \mathcal{L}_1 is the cotangent line bundle at the marked point x_1 .) We construct a natural section s_1 of this line bundle,

$$s_1: \mathcal{L}^\vee \rightarrow \mathcal{L}_1, \quad \alpha \mapsto \alpha(x_1).$$

Namely, an element of \mathcal{L}^\vee is an abelian differential on C , and we take its restriction to the marked point.

The section s_1 vanishes if and only if the marked point is a zero of the abelian differential. Thus, we have the following identity in $H^2(\mathbb{P}\overline{\mathcal{H}}_{g,1})$:

$$[\mathbb{P}\overline{A}_{g,(1)}] = [\{s_1 = 0\}] = c_1(\mathcal{L} \otimes \mathcal{L}_1) = \xi + \psi_1.$$

Now we restrict ourselves to the locus $\{s_1 = 0\}$ and consider the line bundle

$$\mathcal{L} \otimes \mathcal{L}_1^{\otimes 2}.$$

We build a section s_2 of this new line bundle. An element of $\mathcal{L}^\vee|_{\{s_1=0\}}$ is an abelian differential with at least a simple zero at the marked point x_1 . Its first derivative at x_1 is then an element of $\mathcal{L}_1^{\otimes 2}$ (we can verify this assertion using a local coordinate at x_1).

As before, s_2 is equal to zero if and only if the marked point is at least a double zero of the abelian differential. However, $\{s_2 = 0\}$ is composed of three components:

- $\mathbb{P}\overline{A}_{g,(2)}$.
- The locus a_e where the marked point lies on an elliptic component attached to the rest of the stable curve at exactly one point and the abelian differential vanishes identically on the elliptic component.
- The locus a_r where the marked point lies on a “rational bridge”, that is, a rational component attached to two components of the stable curve that are not connected except by this rational component (in this case the abelian differential automatically vanishes on the rational bridge).

We deduce the following formula for $[\mathbb{P}\overline{A}_{g,(2)}]$:

$$\begin{aligned} [\mathbb{P}\overline{A}_{g,(2)}] &= [\{s_2 = 0\}] - [a_e] - [a_r] \\ &= (\xi + \psi_1)(\xi + 2\psi_1) - [a_e] - [a_r] \\ &= \xi^2 + 3\psi_1\xi + 2\psi_1^2 - [a_e] - [a_r]. \end{aligned}$$

Remark 1.21 We make a series of remarks on this result:

- To transform the above considerations into an actual proof we need to check that the vanishing multiplicity of s_2 along all three components equals 1. We will prove this assertion and its generalization in Section 3.
- Denote by $\pi: \mathbb{P}\bar{\mathcal{H}}_{g,1} \rightarrow \mathbb{P}\bar{\mathcal{H}}_g$ the forgetful map, by δ_{sep} the boundary divisor composed of curves with a separating node, and δ_{nonsep} the boundary divisor of curves with a nonseparating node. Let us apply the pushforward by π to the above expression of $[\mathbb{P}\bar{A}_{g,(2)}]$:
 - The term $\pi_*(\xi^2)$ vanishes by the projection formula, since it is a pushforward of a pullback.
 - The term $\pi_*(3\xi\psi_1)$ gives $3\kappa_0\xi = (6g - 6)\xi$ by the projection formula.
 - The term $\pi_*(2\psi_1^2)$ gives $2\kappa_1$.
 - The term $\pi_*([a_e])$ vanishes, because the geometric image of a_e is of codimension 2 in $\mathbb{P}\bar{\mathcal{H}}_g$.
 - The term $\pi_*([a_r])$ gives δ_{sep} since π induces a degree 1 map from a_r onto δ_{sep} .

Thus, we get

$$[\mathbb{P}\bar{\mathcal{H}}(2, 1, \dots, 1)] = \pi_*[\mathbb{P}\bar{A}_{g,(2)}] = (6g - 6)\xi + 2\kappa_1 - \delta_{\text{sep}}.$$

Using the relation $\kappa_1 = 12\lambda_1 - \delta_{\text{sep}} - \delta_{\text{nonsep}}$ on $\bar{\mathcal{M}}_g$ (see for example Arbarello, Cornalba and Griffiths [1, Chapter 17]), we have

$$[\mathbb{P}\bar{\mathcal{H}}(2, 1, \dots, 1)] = (6g - 6)\xi + 24\lambda_1 - 3\delta_{\text{sep}} - 2\delta_{\text{nonsep}}.$$

This formula was first proved by Korotkin and Zograf in 2011 using an analysis of the Bergman tau function [26]. Dawei Chen gave another proof of this result in 2013 using test curves [9].

- In general, to prove Theorem 1 we will work by induction. Let P and $Z = (k_1, k_2, \dots, k_n)$ be vectors of positive integers. Let $Z' = (k_1, \dots, k_i + 1, \dots, k_n)$. Then we will show that

$$[\mathbb{P}\bar{A}_{g,Z',P}] = (\xi + (k_i + 1)\psi_i)[\mathbb{P}\bar{A}_{g,Z,P}] - \text{boundary terms}.$$

The computation of these boundary terms is the crucial part of the proof.

1.7 Applications and related work

Classes in the Picard group of $\overline{\mathcal{M}}_g$ Scott Mullane and Dawei Chen gave a closed formula for the class of $\pi_*[\overline{\mathcal{M}}_g(Z)]$ in the rational Picard group of $\overline{\mathcal{M}}_g$ for all Z of length $g - 2$ (see [8; 27]). They used test curves and linear series to compute this formula. This result has the advantage of giving explicit expressions; however, it has the drawback of not keeping track of the positions of the zeros and of being restricted to the vectors Z of length $g - 2$ (see Section 6.2 for an example of computation).

Incidence variety compactification The problem of the compactification of the strata is extensively studied from different approaches in joint work of Bainbridge, Chen, Gendron, Grushevsky and Möller (see [2; 18]). Their compactification (called *incidence variety compactification*) is slightly different from the one that we use here. We will recall their definitions in Section 4.2 since we will make use of some of their results.

Moduli space of twisted canonical divisors In [15], Farkas and Pandharipande proposed another compactification of the strata. Let g, n and m be nonintegers such that $2g - 2 + n + m > 0$. Let P be a vector of positive integers of length m and let Z be vector of nonnegative integers of length n that is complete for g and P . We recall that $\mathcal{M}_g(Z - P) \subset \mathcal{M}_{g,n+m}$ is the locus of smooth curves such that $\omega_C(-k_1x_1 - \dots - k_nx_n + p_1x_{n+1} + \dots + p_mx_{n+m}) \simeq \mathcal{O}_C$ and that we denote by $\overline{\mathcal{M}}_g(Z - P)$ its closure in $\overline{\mathcal{M}}_{g,n+m}$. Farkas and Pandharipande [15] defined the space of *twisted canonical divisors* denoted by $\widetilde{\mathcal{M}}(Z - P)$. The space of twisted canonical divisors is a singular closed subspace of $\overline{\mathcal{M}}_{g,n+m}$ such that $\overline{\mathcal{M}}(Z - P)$ is one of the irreducible components of $\widetilde{\mathcal{M}}(Z - P)$.

We assume that $m \geq 1$. In the appendix of [15], Farkas and Pandharipande defined a class $H_g(Z - P)$ in $A_g(\overline{\mathcal{M}}_{g,n+m})$ (or $H^{2g}(\overline{\mathcal{M}}_{g,n+m})$): this class is a weighted sum over the classes of irreducible components.

Conjectural expression of $H_g(Z - P)$ Let r be a positive integer and

$$(C, x_1, \dots, x_{n+m})$$

be a smooth curve with markings. An r -spin structure is a line bundle L such that $L^{\otimes r} \simeq \omega_C(-k_1x_1 - \dots - k_nx_n + p_1x_{n+1} + \dots + p_mx_{n+m})$. We denote the moduli space of r -spin structures by $\mathcal{M}_{g,Z-P}^{1/r}$. This space admits a standard compactification $\overline{\mathcal{M}}_{g,Z-P}^{1/r}$ by twisted r -spin structures. We denote by $\pi: \overline{\mathcal{C}}_{g,Z-P}^{1/r} \rightarrow \overline{\mathcal{M}}_{g,Z-P}^{1/r}$ the

universal curves and by $\mathcal{L} \rightarrow \bar{\mathcal{C}}_{g,Z-P}^{1/r}$ the universal line bundle. The moduli space of twisted r -spin structures has a natural forgetful map $\epsilon: \bar{\mathcal{M}}_{g,Z-P}^{1/r} \rightarrow \bar{\mathcal{M}}_{g,n+m}$; the map ϵ is finite of degree r^{2g-1} . We consider $R\pi_*(\mathcal{L})$, the image of \mathcal{L} in the derived category of $\bar{\mathcal{M}}_{g,Z-P}^{1/r}$.

If $m \geq 1$, then we consider the class $c_g^r(Z - P) \stackrel{\text{def}}{=} c_g(R\pi_*\mathcal{L}) \in A_g(\bar{\mathcal{M}}_{g,Z-P}^{1/r})$. If $m = 0$, then we consider a different class, namely Witten’s class, $c_W^r(Z) \in A_{g-1}(\bar{\mathcal{M}}_{g,Z}^{1/r})$. There are several equivalent definitions of Witten’s class, all of which require several technical tools that we will not describe here (see Polishchuk and Vaintrob [33], Chiodo [11] or Chang, Li and Li [7]).

We consider the two functions

$$\begin{aligned}
 P_{g,Z-P}: \mathbb{N}^* &\rightarrow A_*(\bar{\mathcal{M}}_{g,n+m}), & r &\mapsto r\epsilon_*(c_g^r(Z - P)), \\
 P_{g,Z}^W: \mathbb{N}^* &\rightarrow A_*(\bar{\mathcal{M}}_{g,n+m}), & r &\mapsto r^{g-1}\epsilon_*(c_W^r(Z)).
 \end{aligned}$$

Both $P_{g,Z-P}$ and $P_{g,Z}^W$ are polynomials for large values of r (this result is due to Aaron Pixton; see Jana, Pandharipande, Pixton and Zvonkine [23; 31]). We denote by $\tilde{P}_{g,Z-P}$ and $\tilde{P}_{g,Z}^W$ the asymptotic polynomials. The two following conjectures have been proposed:

Conjecture A (see [15]) *If $m \geq 1$ then the equality $H_g(Z - P) = \tilde{P}_{g,Z-P}(0)$ holds in $A_g(\bar{\mathcal{M}}_{g,n+m})$.*

Conjecture B (see [30]) *If $m = 0$ then the equality $[\bar{\mathcal{M}}_g(Z)] = (-1)^g \tilde{P}_{g,Z}^W(0)$ holds in $A_{g-1}(\bar{\mathcal{M}}_{g,n})$.*

As a consequence of Theorem 3, we know that the classes $H_g(Z - P)$ and $[\bar{\mathcal{M}}_g(Z)]$ are tautological and we have an algorithm to check the validity of the conjectures case by case (see Section 6.2 for examples of computations).

These two conjectures are the analogues for differentials of the formula for the so-called double-ramification cycles (DR cycles): the DR cycle is a natural extension to $\bar{\mathcal{M}}_{g,n}$ of the cycle in $\mathcal{M}_{g,n}$ defined as the locus of marked curves (C, x_1, \dots, x_n) such that

$$\sum_{i=1}^n a_i(x_i) \simeq \mathcal{O}_C$$

for any fixed vector of integers $(a_i)_{1 \leq i \leq n}$ such that $\sum a_i = 0$ (see [23]).

Compactification via log geometry Jérémy Guéré constructed a moduli space of “rubber” differentials using log geometry. He proved that this space is endowed with a perfect obstruction theory. Moreover, if $m \geq 1$, this moduli space surjects onto the moduli space of twisted canonical divisors and the class $H_g(Z - P)$ is the pushforward of the virtual fundamental cycle (see [22]).

If $m = 0$ has only positive values, Dawei Chen and Qile Chen have also used log geometry to define a compactification of the strata $\mathcal{H}_g(Z)$ (see [10]).

Induction formula for singularities in families The central result of the present work is the induction formula of Section 5. A similar formula has been proved by Kazarian, Lando and Zvonkine for classes of singularities in families of genus 0 stable maps (see [24]). Their formula contains only the genus 0 part of our induction formula.

They gave an interpretation of the induction formula in genus 0 as a generalization of the completed cycle formula of Okounkov and Pandharipande (see [29]). For now, it is not clear if this generalized completed cycle formula has an extension to higher genera.

This type of induction formula had been previously introduced by Gathmann in the context of genus 0 relative Gromov–Witten invariants (see [17]) and has been recently adapted to the genus 0 quasimap invariants (see Battistella and Nabijou [3]).

Computation of the Lyapunov exponents of strata Strata of differentials are endowed with the structure of a dynamical system. Several numerical invariants have been introduced to characterize the dynamics of the strata: volumes, Siegel–Veech constants, Lyapunov exponents. Some relations exist between these invariants. These relations come in general from relations in the cohomology of the strata.

Our computation of cohomology classes of strata of differentials could be useful to compute these numerical invariants. This idea is developed for example in [26; 8] based on the work of Eskin, Kontsevich and Zorich [13] (see Section 7.3.3). This has been explored in our subsequent paper [34].

Plan of the paper

In Section 2 we construct the space of stable differentials and compute its Segre class. Then we generalize the definition of stable differentials for disconnected curves and for unstable irreducible curves. In the last subsection we present the tautological rings

of spaces of stable differentials in this most general setting (with possible disconnected and semistable curves).

In Section 3 we introduce the stratification of the interior of spaces of stable differentials according to the orders of zeros and we study the geometry of the strata: local parameters, dimension, neighborhood in the space of differentials. Then, in Section 4 we describe the boundary components of the Zariski closure of strata.

Theorems 1, 2 and 3 are proved in Section 5. The main tool involved in their proof is the induction formula for the Poincaré-dual classes of strata of differentials with prescribed orders of zeros (see Theorem 5).

In Section 6 we present two examples of explicit computations.

Finally, in Section 7 we introduce several classes in the Picard group of strata of differentials and prove several relations between these classes by using the induction formula.

Acknowledgements I am very grateful to my PhD advisor Dimitri Zvonkine, who proposed to me this interesting problem, supervised my work with great involvement and brought many beneficial comments.

I would also like to thank Jérémy Guéré, Charles Fougeron, Samuel Grushevsky, Rahul Pandharipande, Dawei Chen, Qile Chen and Anton Zorich for helpful conversations. They have enriched my understanding of the subject in many ways. I am grateful to Felix Janda for the computations that he made and for the long discussions we had regarding the present work. Moreover, I would like to thank the organizers of the conference *Dynamics in the Teichmüller space* at the CIRM (Marseille, France) for their invitation as well as the people I met for the first time at this conference: Pascal Hubert, Erwan Lanneau, Samuel Lelièvre, Anton Zorich, Quentin Guendron and Martin Möller.

Finally, I am very grateful to the anonymous referee for *Geometry & Topology*, whose dedicated work significantly contributed to the final shape of the paper.

2 Stable differentials

In this section, we construct the space of stable differentials and compute its Segre class. We also define stable differentials on disconnected and/or unstable curves. Finally, we define and describe the tautological rings in this generalized setup.

2.1 The cone of generalized principal parts

2.1.1 Orbifold cones We follow here the approach of [12]. Let X be a projective DM stack.

Definition 2.1 An *orbifold cone* is a finitely generated sheaf of graded \mathcal{O}_X -algebras $S = S^0 \oplus S^1 \oplus S^2 \oplus \dots$ such that $S^0 = \mathcal{O}_X$.

Remark 2.2 This definition of cone is weaker than the classical definition of Fulton (see [16]) because we do not ask that S be generated by S^1 . In the classical definition, a cone is a subvariety of a vector bundle (the dual of S^1) given by homogeneous equations. Its projectivization is a subvariety of a bundle of projective spaces. In the orbifold case, the cone is, again, a suborbifold of a vector bundle, but is now given by quasihomogeneous equations. Its projectivization is a suborbifold of the corresponding bundle of weighted projective spaces, which carries a tautological line bundle $\mathcal{O}(1)$ in the orbifold sense (called canonical line bundle in [16]). Thus, the projectivization $\mathbb{P}\mathcal{C}$ of a cone is an orbifold and carries a natural orbifold line bundle $\mathcal{O}(1)$, the tautological line bundle. We write $p: \mathbb{P}\mathcal{C} = \text{Proj}(S) \rightarrow X$ and $\xi = c_1(\mathcal{O}(1))$. Let $\mathcal{C} \rightarrow X$ be a pure-dimensional cone and r the rank of the cone, defined as $\dim(\mathcal{C}) - \dim(X)$. The i^{th} Segre class of \mathcal{C} is defined as

$$s_i = p_*(\xi^{r+i-1}) \in H^{2i}(X, \mathbb{Q}).$$

Example 2.3 Let us consider the graded algebra $\mathbb{C}[x, y, z]$ such that x is an element of weight 2, y is an element of weight 3 and z is an element of weight 1. This graded algebra is not generated by its degree 1 elements. The associated projectivized cone over a point is the weighted projective space $\mathbb{P}(2, 3)$ which is the quotient of $(\mathbb{C}^3)^*$ by \mathbb{C}^* with the action

$$\lambda \cdot (x, y, z) = (\lambda^2 x, \lambda^3 y, \lambda z).$$

Example 2.4 More generally, consider a sheaf of algebras of the form $\mathcal{O}_X \otimes_{\mathbb{C}} S$, where S is a graded algebra over \mathbb{C} . The projective spectrum of this sheaf is a direct product of X with $\text{Proj}(S)$. We call this a *trivial orbifold cone*.

2.1.2 Cone of generalized principal parts

Definition 2.5 Let p be an integer greater than 1. A *principal part* of order p at a smooth point of a curve is an equivalence class of germs of meromorphic differentials with a pole of order p ; two germs f_1 and f_2 are equivalent if $f_1 - f_2$ is a meromorphic differential with at most a simple pole.

First, we parametrize the space of principal parts at a point. Let z be a local coordinate at $0 \in \mathbb{C}$. A principal part at 0 of order p is given by

$$\left[\left(\frac{u}{z}\right)^{p-1} + a_1 \left(\frac{u}{z}\right)^{p-2} + \cdots + a_{p-2} \left(\frac{u}{z}\right) \right] \frac{dz}{z}$$

with $u \neq 0$. However, given a principal part, the choice of (u, a_1, \dots, a_{p-2}) is not unique. Indeed, there are $p-1$ choices for u given by the $\zeta^l \cdot u$ (with $\zeta^l = \exp(2i\pi \cdot l/(p-1))$ for $0 \leq l < p-1$) and, once the value of u is chosen, the a_i are determined uniquely. Therefore, the coordinates (u, a_1, \dots, a_{p-2}) parametrize a degree $p-1$ covering of the space of principal parts. This motivates the following definition:

Definition 2.6 Assign to u the weight $1/(p-1)$ and to a_j the weight $j/(p-1)$. The graded algebra $S \subset \mathbb{C}[u, a_1, \dots, a_{p-2}]$ spanned by the monomials of integral weights is called the *algebra of generalized principal parts* and $\mathcal{P} = \text{Spec}(S)$ is the *space of generalized principal parts*.

The space \mathcal{P} is the quotient of \mathbb{C}^{p-1} by the group $\mathbb{Z}/(p-1)\mathbb{Z}$, which, from now on, we will denote by \mathbb{Z}_{p-1} for brevity. An element $\zeta \in \mathbb{Z}_{p-1}$ acts by

$$\zeta \cdot (u, a_1, \dots, a_{p-2}) = (\zeta u, \zeta a_1, \dots, \zeta^{p-2} a_{p-2}).$$

Moreover, the natural action of \mathbb{C}^* on \mathcal{P} is given by

$$\lambda \cdot (u, a_1, \dots, a_{p-2}) = (\lambda^{1/(p-1)} u, \lambda^{1/(p-1)} a_1, \dots, \lambda^{(p-2)/(p-1)} a_{p-2}).$$

Note that this action is not well defined on the covering space \mathbb{C}^{p-1} , but is well defined on its \mathbb{Z}_{p-1} quotient \mathcal{P} .

Notation 2.7 Denote by $I_u \subset S$ the ideal of polynomials divisible by u . Denote by $\mathcal{A} \subset \mathcal{P}$ the suborbifold defined by I_u .

The suborbifold $\mathcal{A} \subset \mathcal{P}$ is the Weil divisor obtained as the image of the Cartier divisor $\{u = 0\} \subset \mathbb{C}^{p-1}$ under the quotient of \mathbb{C}^{p-1} by the action of \mathbb{Z}_{p-1} . The divisor $(p-1)\mathcal{A}$ is the Cartier divisor given by the equation $u^{p-1} = 0$. (Note that u^{p-1} lies in S while u does not.) The space of principal parts embeds into \mathcal{P} as the complement of \mathcal{A} .

Lemma 2.8 A change of local coordinate z induces an isomorphism of S that preserves the grading and acts trivially on the quotient algebra S/I_u .

Proof Let $z = f(w) = \alpha_1 w + \alpha_2 w^2 + \dots$ be a local coordinates change. We denote by $(u', a'_1, \dots, a'_{p-2})$ the parameters of the presentation of principal parts in coordinate w . We have the transformation

$$\begin{aligned} u &\mapsto \alpha_1 u, \\ a_1 &\mapsto a_1 + \gamma_{1,1} u, \\ a_2 &\mapsto a_2 + \gamma_{2,1} u a_1 + \gamma_{2,2} u^2, \\ &\vdots \end{aligned}$$

where the $\gamma_{i,j}$ are polynomials in $\alpha_1, \alpha_2, \dots$ depending only on the order of the principal part. By taking u to be 0, we see that the coordinates (a_1, \dots, a_{p-2}) of \mathcal{A} are independent of the choice of local coordinate. □

Remark 2.9 In Section 2.2 we will see that the locus \mathcal{A} corresponds to the appearance of a semistable bubble of the underlying curve C at the i^{th} marked point. The coordinate on the bubble is $w = u/z$.

Remark 2.10 The cone of principal parts of differentials differs from the cone of principal parts of functions of [12] only by the coefficients $\gamma_{i,j}$.

Now, let g and n be nonnegative integers such that $2g - 2 + n > 0$. Let $i \in \llbracket 1, n \rrbracket$ and $p_i \geq 2$. We denote by \mathbb{P}_i the following sheaf of graded algebras over $\overline{\mathcal{M}}_{g,n}$:

Pick an open chart $U \subset \overline{\mathcal{M}}_{g,n}$ together with a trivialization of a tubular neighborhood of the i^{th} section σ_i of the universal curve over U . In other words, denoting by Δ the unit disc, we choose an embedding

$$U \times \Delta \hookrightarrow \overline{\mathcal{C}}_{g,n}$$

commuting with $U \hookrightarrow \overline{\mathcal{M}}_{g,n}$ and such that $U \times \{0\}$ is the i^{th} section of the universal curve. The sheaf \mathbb{P}_i over U is given by $\mathbb{P}_i(U) = \mathcal{O}_U \otimes S$.

Now, given two overlapping charts U and V we need to define the gluing map between the sheaves on their intersection. To do that, denote by z the coordinate on Δ in the product $U \times \Delta$ and by w the coordinate on Δ in the product $V \times \Delta$. Over the intersection $U \cap V$ we get a change of local coordinates $z(w)$. We use this change of local coordinate and the constants $\gamma_{i,j}$ from Lemma 2.8 to construct an identification between the two algebras $\mathbb{P}_i(U)|_{U \cap V}$ and $\mathbb{P}_i(V)|_{U \cap V}$.

Note that the sheaf of ideals I_u is well defined and the quotients S/I_u are identified with each other in a canonical way that does not depend on the local coordinates z and w .

We denote by $\mathcal{P}_i = \text{Spec}(\mathbb{P}_i)$ the spectrum of \mathbb{P}_i and by $\mathcal{A}_i = \text{Spec}(\mathbb{P}_i/I_u)$ the spectrum of the quotient. The latter is a trivial cone over $\overline{\mathcal{M}}_{g,n}$.

Proposition 2.11 *The cone \mathcal{P}_i and its projectivization are normal.*

Proof Indeed, the space $\overline{\mathcal{M}}_{g,n}$ is smooth and the sheaf of fractions of the algebra \mathbb{P}_i is the same as the sheaf of fractions of \mathbb{P}_i^1 . □

Lemma 2.12 *The cone \mathcal{A}_i is the product of $\overline{\mathcal{M}}_{g,n}$ with the weighted projective space with weights $(1/(p_i - 1), \dots, (p_i - 2)/(p_i - 1))$ quotiented by the action of \mathbb{Z}_{p_i-1} . Moreover, the Segre classes of \mathcal{A}_i and \mathcal{P}_i are given by*

$$s(\mathcal{A}_i) = \frac{(p_i - 1)^{p_i-2}}{(p_i - 1)!}, \quad s(\mathcal{P}_i) = \frac{(p_i - 1)^{p_i-1}}{(p_i - 1)!} \cdot \frac{1}{1 - (p_i - 1)\psi_i}.$$

Proof The proof is based on the same arguments as for the cone of principal parts of functions. The section u^{p_i-1} is a section of the line bundle $\mathcal{L}_i^{-\otimes(p_i-1)}$, which vanishes with multiplicity $p_i - 1$ along \mathcal{A}_i . □

2.1.3 Stack of generalized principal parts In the above paragraph we defined the cone of generalized principal parts, which is a normal scheme over $\overline{\mathcal{M}}_{g,n}$. We introduce here another approach to the quotient by the \mathbb{Z}_{p_i-1} -action. Let $\tilde{\mathbb{P}}_i$ be the sheaf of algebra defined locally by

$$\tilde{\mathbb{P}}_i(U) = \mathcal{O}_U[u, a_1, \dots, a_{p_i-2}],$$

where U is a chart with a trivialization of a tubular neighborhood of the i^{th} section of the universal curve and the coordinates $(u, a_1, \dots, a_{p_i-2})$ are defined as above.

Definition 2.13 *The stack of generalized principal parts \mathfrak{P}_i is the stack quotient*

$$\text{Spec}(\tilde{\mathbb{P}}_i)/\mathbb{Z}_{p_i-1}.$$

By construction we have the following proposition:

Proposition 2.14 *For all schemes U with a map $U \rightarrow \overline{\mathcal{M}}_{g,n}$, the scheme $U \times_{\overline{\mathcal{M}}_{g,n}} \mathcal{P}_i$ is the coarse space of $U \times_{\overline{\mathcal{M}}_{g,n}} \mathfrak{P}_i$.*

Proposition 2.15 *The stack of generalized principal parts is a smooth DM stack.*

Proof The space $\overline{\mathcal{M}}_{g,n}$ is a smooth DM stack and \mathfrak{P}_i is locally the quotient of an affine smooth scheme over $\overline{\mathcal{M}}_{g,n}$ by a finite group. □

2.1.4 Cones of generalized principal parts and jet bundles From now on in the text, unless otherwise mentioned, for any family of semistable curves $C \rightarrow S$ we denote by ω the relative cotangent line bundle $\omega_{C/S}$.

Definition 2.16 Let $\pi: \overline{\mathcal{C}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ be the universal curve and $(\sigma_i)_{1 \leq i \leq n}: \overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{C}}_{g,n}$ the global sections of marked points. Let $1 \leq i \leq n$ and $p_i \geq 1$. The vector bundle $J_i \rightarrow \overline{\mathcal{M}}_{g,n}$ of polar jets of order p_i at the i^{th} marked point is defined as the quotient

$$J_i = R^0 \pi_*(\omega(p_i \sigma_i)) / R^0 \pi_*(\omega(\sigma_i)).$$

We fix $1 \leq i \leq n$ and $p_i > 0$. The bundle of polar jet of order p_i is a vector bundle of rank $p_i - 1$. As before, we consider an open chart U of $\overline{\mathcal{M}}_{g,n}$ with a trivialization z_i of a tubular neighborhood of the section σ_i . Over the chart U the jet bundle is trivial. Indeed, an element of J_i over U is given by

$$\left[\frac{b_0}{z_i^{p_i-1}} + \dots + \frac{b_{p_i-2}}{z_i^{p_i-2}} \right] \frac{dz_i}{z_i}.$$

Thus, the jet bundle J_i restricted to U is given by $\text{Spec}(\mathcal{O}_U[b_0^i, \dots, b_{p_i-2}^i])$. Recall that, using the trivialization z_i , we have defined coordinates u, a_1, \dots, a_{p_i-2} such that $\mathbb{P}_i(U)$ is the subalgebra of

$$\mathcal{O}_U[u, a_1, \dots, a_{p_i-2}]$$

generated by monomials with integral weights. We define the following morphism of graded algebras over \mathcal{O}_U :

$$\phi_i(U): \text{Sym}^*(J_i^\vee)(U) \rightarrow \mathbb{P}_i(U), \quad \begin{aligned} b_0 &\mapsto u^{p_i-1}, \\ b_j &\mapsto u^{p_i-1-j} a_j \quad \text{for } 1 \leq j \leq p_i - 2. \end{aligned}$$

The morphism $\phi_i(U)$ is defined for a chart U with a choice of trivialization z_i . We can easily check that the $\phi_i(U)$ can be glued into a morphism of sheaves of graded algebras. Thus, we have constructed a morphism of cones

$$\phi_i: \mathcal{P}_i \rightarrow J_i.$$

It is important to note that for $p_i \geq 3$ the morphism ϕ_i is neither surjective nor injective.

Lemma 2.17 We define the two spaces

$$\begin{aligned} \mathcal{P}_i \supset \tilde{\mathcal{P}}_i &= (\mathcal{P}_i \setminus \mathcal{A}_i) \cup \text{the zero section,} \\ J_i \supset \tilde{J}_i &= (J_i \setminus \{b_0 = 0\}) \cup \text{the zero section.} \end{aligned}$$

The image of the morphism ϕ_i is the space \tilde{J}_i . Moreover, the morphism ϕ_i restricted to $\tilde{\mathcal{P}}_i$ induces an isomorphism from $\tilde{\mathcal{P}}_i$ to \tilde{J}_i .

The proof is a simple check.

Remark 2.18 In particular, the morphism ϕ_i does not define a morphism of projectivized cones. Indeed, certain points outside of the zero section of \mathcal{P}_i are mapped to the zero section of J_i .

2.2 The space of stable differentials

Let g, n and m be nonnegative integers satisfying $2g - 2 + n + m > 0$. Let $P = (p_1, p_2, \dots, p_m)$ be a vector of positive integers. For all $1 \leq i \leq m$, we denote by \mathcal{P}_{n+i} (respectively \mathfrak{P}_{n+i} and J_{n+i}) the cone of principal parts (respectively the stack of principal parts and the vector bundle of polar jets) of order p_i at the $(n+i)^{\text{th}}$ marked point. Let $p: \bar{\mathcal{H}}_{g,n,P} \rightarrow \bar{\mathcal{M}}_{g,n+m}$ be the space of stable differentials of Definition 1.2 together with the forgetful map.

We recall that $\pi: \bar{\mathcal{C}}_{g,n+m} \rightarrow \bar{\mathcal{M}}_{g,n+m}$ is the universal curve and the

$$(\sigma_i)_{1 \leq i \leq n+m}: \bar{\mathcal{M}}_{g,n+m} \rightarrow \bar{\mathcal{C}}_{g,n+m}$$

are the global sections corresponding to marked points.

Notation 2.19 Let $K\bar{\mathcal{M}}_{g,n}(P) \rightarrow \bar{\mathcal{M}}_{g,n+m}$ be the vector bundle

$$R^0 \pi_* \left(\omega \left(\sum_{i=1}^m p_i \sigma_{n+i} \right) \right) \rightarrow \bar{\mathcal{M}}_{g,n+m}.$$

It is a vector bundle of rank $g - 1 + \sum p_i$ if P is not empty.

We have the exact sequence of vector bundles over $\bar{\mathcal{M}}_{g,n+m}$

$$(1) \quad 0 \rightarrow R^0 \pi_* \left(\omega \left(\sum_{i=1}^m \sigma_{n+i} \right) \right) \rightarrow K\bar{\mathcal{M}}_{g,n}(P) \rightarrow \bigoplus_{i=1}^m J^{n+i} \rightarrow 0.$$

This exact sequence is simply the long exact sequence obtained from the residue exact sequence.

Proposition 2.20 *The stack $\overline{\mathfrak{H}}_{g,n,P}$ is isomorphic to the fiber product of $K\overline{\mathcal{M}}_{g,n}(P)$ and $\bigoplus_{i=1}^m \mathfrak{P}_{n+i}$ over $\bigoplus_{i=1}^m J_{n+i}$, where the map $\mathfrak{P}_{n+i} \rightarrow J_{n+i}$ is the composition of maps $\mathfrak{P}_{n+i} \rightarrow \mathcal{P}_{n+i} \xrightarrow{\phi_i} J_{n+i}$.*

Proof We denote by $\tilde{\mathfrak{H}}_{g,n,P}$ the fiber product

$$(2) \quad \begin{array}{ccc} \tilde{\mathfrak{H}}_{g,n,P} & \longrightarrow & \bigoplus_{i=1}^m \mathfrak{P}_{n+i} \\ \downarrow & & \downarrow \\ K\overline{\mathcal{M}}_{g,n}(P) & \longrightarrow & \bigoplus_{i=1}^m J_{n+i} \end{array}$$

We construct the two directions of the isomorphism $\tilde{\mathfrak{H}}_{g,n,P} \simeq \overline{\mathfrak{H}}_{g,n,P}$ separately.

From $\overline{\mathfrak{H}}_{g,n,P}$ to $\tilde{\mathfrak{H}}_{g,n,P}$ To construct a morphism $F_1: \overline{\mathfrak{H}}_{g,n,P} \rightarrow \tilde{\mathfrak{H}}_{g,n,P}$ we define morphisms $\Phi_i: \overline{\mathfrak{H}}_{g,n,P} \rightarrow \mathfrak{P}_{n+i}$ for all $1 \leq i \leq m$ and $\chi: \overline{\mathfrak{H}}_{g,n,P} \rightarrow K\overline{\mathcal{M}}_{g,n}(P)$ fitting in diagram (2).

Let $(C \rightarrow S, \sigma_1, \dots, \sigma_{n+m}, \alpha)$ be a family of stable differentials. Let $s \rightarrow S$ be a geometric point of S and $(C_s, x_1, \dots, x_{n+m}, \alpha_s)$ be the stable differential determined by s . The element $\Phi_i(\alpha_s)$ is determined as follows:

- If x_{n+i} does not belong to a rational component then $\Phi_i(s)$ is the principal part at the marked point. It belongs to $\mathfrak{P}^{n+i} \setminus \{u = 0\}$.
- If x_{n+i} belongs to a rational component, let w_{n+i} be a global coordinate of the rational component such that: x_{n+i} is at infinity, the node is at 0 and the term of α in front of $w_{n+i}^{p_i-2} dw_{n+i}$ is -1 . Then α_s is of the form

$$-(w_{n+i}^{p_i-1} + a_1 w_{n+i}^{p_i-2} + \dots + a_{p_i-2} w_{n+i} + \text{res}_{\sigma_{n+i}}(\alpha)) \frac{dw_{n+i}}{w_{n+i}}$$

and we set $\Phi_i(s) = (0, a_1, \dots, a_{p_i-2})$. Indeed, the substack $\{u = 0\}$ is the quotient of a trivial vector bundle by \mathbb{Z}_{p_i-1} and the a_i are the global coordinates of this vector bundle.

We will prove that the map Φ_i depends holomorphically on s . If s is a point of the first type, this is an obvious statement. If s is a point of the second type, let U be an open neighborhood of s in S with a trivialization z_{n+i} of a tubular neighborhood of σ_{n+i} in C (see the previous section). Let C' be the stabilization of C . The differential α restricted to C' is a differential with poles of order at most p_i at $z_{n+i} = 0$. The

differential α in this coordinate is given by

$$(3) \quad \alpha = \left(\left(\frac{u}{z_{n+i}} \right)^{p_i-1} + \dots + a_{p_i-2}^i \frac{u}{z_{n+i}} + \text{res}_{\sigma_{n+i}}(\alpha) + \underset{z_{n+i} \rightarrow 0}{O}(z_{n+i}) \right) \frac{dz_{n+i}}{z_{n+i}}$$

The value u^{p_i-1} depends holomorphically on s and up to a choice of smaller U we can fix a choice of $(p_i + 1)^{\text{st}}$ root u . The function u depends holomorphically on s .

Now we use the function u and the local trivialization z_{n+i} to construct the family of semistable curves $C'' \subset C' \times \mathbb{P}^1$ defined by the equation $z_{n+i}w = u$ (where, as previously, w is the global coordinate of the rational component and the pole is located at $w = \infty$). This family of curves is isomorphic to C (the stabilizations of C'' and C' are isomorphic and each fiber of these two families has the same dual graph). In particular, α is a meromorphic differential on C'' with constant order of pole at $w = \infty$. Further, in the chart w the highest-order coefficient of α is given by 1. In particular, the coordinate w is equal to the coordinate w_{n+i} on the unstable rational component of the fiber of s . In the chart w , the meromorphic differential α is given by

$$(4) \quad \alpha = -(w^{p_i-1} + a_1 w^{p_i-2} + \dots + a_{p_i-2} w + \text{res}_{\sigma_{n+i}}(\alpha) + \underset{w \rightarrow 0}{O}(w)) \frac{dw}{w},$$

where the a_i depend holomorphically on s . Therefore, Φ depends holomorphically on s .

Now we construct the map $\chi: \bar{\mathfrak{H}}_{g,n,P} \rightarrow K\bar{\mathcal{M}}_{g,n}(P)$. Let $(C \rightarrow S, \sigma_1, \dots, \sigma_{n+m}, \alpha)$ be a family of stable differentials. We denote by $\tilde{C} \rightarrow S$ the stabilization of C and write $\tilde{\alpha} = \alpha|_{\tilde{C}}$. The family $(\tilde{C} \rightarrow S, \sigma_1, \dots, \sigma_{n+m}, \tilde{\alpha})$ is a section of $\omega_{C/S}(\sum p_i \sigma_{n+i})$, and thus a map $S \rightarrow K\bar{\mathcal{M}}_{g,n}(P)$. By construction, the morphisms χ and the $(\Phi_i)_{i=1, \dots, m}$ fit in diagram (2).

From $\tilde{\mathfrak{H}}_{g,n,P}$ to $\bar{\mathfrak{H}}_{g,n,P}$ Let S be a \mathbb{C} -scheme and let $S \rightarrow \tilde{\mathfrak{H}}_{g,n,P}$ be a morphism. By composition with the morphism $\tilde{\mathfrak{H}}_{g,n,P} \rightarrow K\bar{\mathcal{M}}_{g,n}(P)$, we get a family of stable curves $C \rightarrow S$ with $n + m$ sections σ_i and a section α of $\omega_{C/S}(\sum p_i \sigma_{n+i})$. The family $S \rightarrow \tilde{\mathfrak{H}}_{g,n,P}$ determines also families of generalized principal parts. From the family of meromorphic differentials α and the principal parts we will construct a family of stable differentials.

Let z_{n+i} be local trivializations of the tubular neighborhoods of the sections σ_{n+i} of the curve C/S for $1 \leq i \leq m$. Let w_{n+i} be global coordinates of the complex plane. We denote by $(u^i, a_1^i, \dots, a_{p_i-2}^i)$ the standard coordinates of the principal parts \mathfrak{P}_{n+i} obtained from the trivializations z_{n+i} . We construct a family of semistable curves

$\tilde{C} \rightarrow S$ defined by the equation $z_{n+i}w_{n+i} = u^i$. On the curve \tilde{C} we construct a differential $\tilde{\alpha}$. This differential is given by the expression (4) in coordinate w_{n+i} and by the expression (3) in coordinate z_{n+i} . The tuple $(\tilde{C}, \sigma_1, \dots, \sigma_{n+m}, \tilde{\alpha})$ is a family of stable differentials over S .

Therefore, we have determined a morphism $F_2: \tilde{\mathfrak{H}}_{g,n,P} \rightarrow \bar{\mathfrak{H}}_{g,n,P}$. By construction it is the inverse of F_1 previously defined. □

The following proposition finishes the proof of Proposition 1.4 and thus completes Definition 1.3.

Proposition 2.21 *We denote by $\bar{\mathcal{H}}_{g,n,P}$ the following fiber product (in the category of cones over $\bar{\mathcal{M}}_{g,n+m}$ or in the category of DM stacks):*

$$(5) \quad \begin{array}{ccc} \bar{\mathcal{H}}_{g,n,P} & \longrightarrow & \bigoplus_{i=1}^m \mathcal{P}_{n+i} \\ \downarrow & & \downarrow \bigoplus \phi_i \\ K\bar{\mathcal{M}}_{g,n}(P) & \longrightarrow & \bigoplus_{i=1}^m J_{n+i} \end{array}$$

Then space $\bar{\mathcal{H}}_{g,n,P}$ is the unique space that satisfies the properties of Proposition 1.5.

Proof The fact that $\bar{\mathcal{H}}_{g,n,P}$ satisfies the properties of Proposition 1.5 is a direct consequence of Propositions 2.14 and 2.20. The uniqueness of this stack follows from the uniqueness of coarse spaces. □

From now on we will denote by $\text{stab}: \bar{\mathcal{H}}_{g,n,P} \rightarrow K\bar{\mathcal{M}}_{g,n}(P)$ the vertical projection in diagram (5).

2.3 Properties of spaces of stable differentials

We keep the notation g, n, m and P of the previous sections. We state here several general properties of $\bar{\mathfrak{H}}_{g,n,P}$ and $\bar{\mathcal{H}}_{g,n,P}$ that will be needed further in the text.

Proposition 2.22 *Suppose that P is not empty. Then the spaces $\bar{\mathfrak{H}}_{g,n,P}$ and $\bar{\mathcal{H}}_{g,n,P}$ are irreducible DM stacks of pure dimension $4g - 4 + \sum p_i$ and $\mathbb{P}\bar{\mathcal{H}}_{g,n,P}$ is a proper DM stack (of dimension one less). The space $\bar{\mathcal{H}}_{g,n,P}$ and its projectivization are normal. The space $\bar{\mathfrak{H}}_{g,n,P}$ is a smooth DM stack.*

If P is empty then both $\bar{\mathfrak{H}}_{g,n,P}$ and $\bar{\mathcal{H}}_{g,n,P}$ are isomorphic to the Hodge bundle, which is a smooth DM stack of dimension $4g - 3$.

Proof The first part of the proposition follows from Propositions 2.15, 2.20 and 2.11. The second part is straightforward. \square

We consider the following two maps: on the one hand the inclusion of vector bundles $R^0\pi_*(\omega(\sum_{i=1}^m \sigma_{n+i})) \rightarrow K\overline{\mathcal{M}}_{g,n}(P)$, and on the other hand the zero map $K\overline{\mathcal{M}}_{g,n,P} \rightarrow \bigoplus \mathcal{P}^{n+i}$. Then we get an embedding $R^0\pi_*(\omega(\sum_{i=1}^m \sigma_{n+i})) \rightarrow \overline{\mathcal{H}}_{g,n,P}$ by the universal property of the cartesian diagram (5).

Proposition 2.23 *For all g, n and P , we have the exact sequence of cones (in the sense of [16, Proposition 4.1.6])*

$$0 \rightarrow R^0\pi_*\left(\omega\left(\sum_{i=1}^m \sigma_{n+i}\right)\right) \rightarrow \overline{\mathcal{H}}_{g,n,P} \rightarrow \bigoplus_{i=1}^m \mathcal{P}_{n+i} \rightarrow 0.$$

Proof By construction, the sheaf of algebras defining $\overline{\mathcal{H}}_{g,n,P}$ is locally the tensor product of the sheaves of algebras $\text{Sym}^\vee(R^0\pi_*(\omega(\sum_{i=1}^m \sigma_{n+i})))$ and the \mathbb{P}_{n+i} . \square

The action of \mathbb{C}^* on the space $\overline{\mathcal{H}}_{g,n,P}$ is determined by multiplication of the differential by a scalar. Let us give a description of the \mathbb{C}^* -fixed locus, ie the locus of points that are invariant under the action of \mathbb{C}^* .

Let (C, x_1, \dots, x_{n+m}) be a curve in $\overline{\mathcal{M}}_{g,n+m}$. We denote by m' the number of entries of P greater than 1. From C we construct a semistable curve \tilde{C} as follows. The curve \tilde{C} has $m'+1$ irreducible components: one main component isomorphic to C and m' rational components attached to C at the marked points corresponding to poles of order greater than 1. We mark points (x'_1, \dots, x'_{n+m}) on \tilde{C} . The first n marked points and the points corresponding to poles of order at most 1 are on the main component and satisfy $x_i = x'_i$. The poles of orders greater than 1 are carried by the rational components.

Now we define a meromorphic differential α on \tilde{C} by:

- The differential α vanishes identically on the main component.
- On an exterior rational component, if we assume that the marked point is at 0 and the node at ∞ , then α is given by dz/z^{p_i} .

The tuple $(\tilde{C}, x'_1, \dots, x'_{n+m}, \alpha)$ is a stable differential invariant under the action of \mathbb{C}^* . Indeed, let λ be a scalar in \mathbb{C}^* ; the differential $\lambda\alpha$ vanishes on the main component and $\lambda dz/z^{p_i}$ is equal to $d w/w^{p_i}$ if we use the change of coordinate $z = w/\lambda^{1/p_i}$ for any $(p_i)^{\text{th}}$ root of λ .

Conversely, any \mathbb{C}^* -invariant point of $\bar{\mathcal{H}}_{g,n,P}$ is of this type. Indeed, $\bar{\mathcal{H}}_{g,n,P}$ is a cone; thus, the locus of \mathbb{C}^* -invariant points is a section of this cone and we have constructed this section here.

2.4 Residues

Let g, n, m and P be as in the previous sections.

Definition 2.24 Let \mathfrak{R} be the vector subspace of \mathbb{C}^m defined by

$$\mathfrak{R} = \{(r_1, r_2, \dots, r_m) \mid r_1 + r_2 + \dots + r_m = 0\}.$$

The vector space \mathfrak{R} will be called the *space of residues*. The *residue map* is the map of cones over $\bar{\mathcal{M}}_{g,n+m}$

$$\text{res}: \bar{\mathcal{H}}_{g,n,P} \rightarrow \mathfrak{R}, \quad \alpha \mapsto (\text{res}_{x_{n+1}}(\alpha), \text{res}_{x_{n+2}}(\alpha), \dots, \text{res}_{x_{n+m}}(\alpha)),$$

where \mathfrak{R} stands for the trivial cone. We use the same notation for the residue map $\text{res}: K\bar{\mathcal{M}}_{g,n}(P) \rightarrow \mathfrak{R}$. In this case it is a morphism of vector bundles.

These two residue maps fit in the commutative triangle

$$(6) \quad \begin{array}{ccc} \bar{\mathcal{H}}_{g,n,P} & \xrightarrow{\text{stab}} & K\bar{\mathcal{M}}_{g,n}(P) \\ & \searrow \text{res} & \downarrow \text{res} \\ & & \mathfrak{R} \end{array}$$

Let $\bar{\mathcal{H}}_{g,n,P}^0 \subset \bar{\mathcal{H}}_{g,n,P}$ (resp. $K\bar{\mathcal{M}}_{g,n}^0(P) \subset K\bar{\mathcal{M}}_{g,n}(P)$) be the subcone (resp. sub-vector bundle) of differentials without residues.

We recall that the Hodge bundle is by definition equal to $\bar{\mathcal{H}}_{g,n+m} = R^0\pi_*\omega$. The following sequence of vector bundles over $\bar{\mathcal{M}}_{g,n+m}$ is exact:

$$(7) \quad 0 \rightarrow \bar{\mathcal{H}}_{g,n+m} \rightarrow R^0\pi_*\left(\omega\left(\sum_{i=1}^m \sigma_{n+i}\right)\right) \xrightarrow{\text{res}} \mathfrak{R} \rightarrow 0$$

(this is the exact sequence obtained from the residue exact sequence $0 \rightarrow \omega_C(\sum x_i) \rightarrow \omega_C \rightarrow \mathbb{C}^m \rightarrow 0$). The vector bundle $K\bar{\mathcal{M}}_{g,n}^0(P)$ fits into the following commutative diagram of vector bundles over $\bar{\mathcal{M}}_{g,n+m}$:

$$(8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & K\bar{\mathcal{M}}_{g,n}^0(P) & \longrightarrow & K\bar{\mathcal{M}}_{g,n}(P) & \xrightarrow{\text{res}} & \mathfrak{R} \longrightarrow 0 \\ & & \uparrow & & \uparrow & \nearrow & \\ \bar{\mathcal{H}}_{g,n+m} & \longrightarrow & R^0\pi_*\left(\omega\left(\sum_{i=1}^m \sigma_{n+i}\right)\right) & & & & \end{array}$$

where the central square is cartesian. The first line of diagram (8) is exact by exactness of the sequence (7). Therefore, the cone structure of $\overline{\mathcal{H}}_{g,n,P}^0$ can be defined equivalently from the cone structure of $\overline{\mathcal{H}}_{g,n,P}$ or by saying that $\overline{\mathcal{H}}_{g,n,P}^0$ is the fiber product

$$\begin{array}{ccc} \overline{\mathcal{H}}_{g,n,P}^0 & \longrightarrow & \bigoplus P_{n+i} \\ \downarrow & & \downarrow \\ K\overline{\mathcal{M}}_{g,n}^0(P) & \longrightarrow & \bigoplus J_{n+i} \end{array}$$

We have the exact sequence of cones

$$0 \rightarrow \overline{\mathcal{H}}_{g,n+m} \rightarrow \overline{\mathcal{H}}_{g,n,P}^0 \rightarrow \bigoplus P_{n+i} \rightarrow 0.$$

Remark 2.25 We cannot say that the sequence

$$0 \rightarrow \overline{\mathcal{H}}_{g,n,P}^0 \rightarrow \overline{\mathcal{H}}_{g,n,P} \rightarrow \mathfrak{X} \rightarrow 0$$

is exact because exactness for morphisms of cones is ill defined if the first term is not a vector bundle.

More generally we define the following:

Definition 2.26 Let R be a vector subspace of \mathfrak{X} . Let $\overline{\mathcal{H}}_{g,n,P}^R \subset \overline{\mathcal{H}}_{g,n,P}$ (resp. $K\overline{\mathcal{M}}_{g,n}^R(P) \subset K\overline{\mathcal{M}}_{g,n}(P)$) be the subcone (resp. subvector bundle) of differentials with a vector of residues lying in R . We will call R a *space of residue conditions*.

Lemma 2.27 Let $R \subset \mathfrak{X}$ be a vector subspace.

- The space $\overline{\mathcal{H}}_{g,n,P}^R$ is a closed subcone of $\overline{\mathcal{H}}_{g,n,P}$ of codimension $\dim(\mathfrak{X}/R)$ (where we set $\dim(\mathfrak{X}/R) = 0$ if P is empty).
- The Segre classes of $\overline{\mathcal{H}}_{g,n,P}^R$ and $\overline{\mathcal{H}}_{g,n,P}$ are equal.
- The Poincaré-dual class of $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}^R$ in $H^*(\mathbb{P}\overline{\mathcal{H}}_{g,n,P}, \mathbb{Q})$ is given by

$$[\mathbb{P}\overline{\mathcal{H}}_{g,n,P}^R] = \xi^{\dim(\mathfrak{X}/R)}.$$

Proof Let us denote by res_R the composition of morphisms $\overline{\mathcal{H}}_{g,n,P} \rightarrow \mathfrak{X} \rightarrow \mathfrak{X}/R$ (we use the same notation for its alter ego for $K\overline{\mathcal{M}}_{g,n}(P)$). We denote by $\overline{\mathcal{H}}_{g,n+m}^R$ the kernel of the morphism

$$R^0 \pi_* \left(\omega \left(\sum_{i=1}^m \sigma_{n+i} \right) \right) \xrightarrow{\text{res}_R} \mathfrak{X}/R \rightarrow 0.$$

It is a vector bundle of rank $g + \dim(R)$. By repeating the above argument, we have the exact sequence of cones

$$0 \rightarrow \overline{\mathcal{H}}_{g,n+m}^R \rightarrow \overline{\mathcal{H}}_{g,n,P}^R \rightarrow \mathfrak{R}/R.$$

We deduce from this exact sequence that

- the codimension of $\overline{\mathcal{H}}_{g,n,P}^R$ in $\overline{\mathcal{H}}_{g,n,P}$ is $\dim(\mathfrak{R}/R)$;
- the Segre class of $\overline{\mathcal{H}}_{g,n,P}^R$ is given by

$$c_*(\overline{\mathcal{H}}_{g,n+m}^R) \cdot s_*\left(\bigoplus \mathcal{P}_{n+i}\right)$$

(see [16, Proposition 4.1.6]).

Further, the vector bundle \mathfrak{R}/R is trivial; thus,

$$c_*(\overline{\mathcal{H}}_{g,n+m}^R) = c_*\left(R^0 \pi_*\left(\omega\left(\sum_{i=1}^m \sigma_{n+i}\right)\right)\right)$$

and the Segre class of $\overline{\mathcal{H}}_{g,n,P}^R$ does not depend on the choice of R .

To prove the last statement, we study the vector bundle $\mathcal{O}(1) \otimes p^*(\mathfrak{R}/R) \rightarrow \mathbb{P}\overline{\mathcal{H}}_{g,n,P}$, where we recall that $p: \mathbb{P}\overline{\mathcal{H}}_{g,n,P} \rightarrow \overline{\mathcal{M}}_{g,n+m}$ is the forgetful map. We have

$$\mathcal{O}(1) \otimes p^*(\mathfrak{R}/R) \simeq \text{Hom}(\mathcal{O}(-1), p^*(\mathfrak{R}/R)).$$

A section of this vector bundle is given by

$$s: \alpha \mapsto \text{res}_R(\alpha).$$

The vanishing locus of s is $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}^R$, which is of codimension $\dim(\mathfrak{R}/R)$ and irreducible. Thus, the Poincaré-dual class of $\mathbb{P}\overline{\mathcal{H}}_{g,n,P}^R$ in $H^*(\mathbb{P}\overline{\mathcal{H}}_{g,n,P}, \mathbb{Q})$ is given by

$$d \cdot c_{\text{top}}(\mathcal{O}(1) \otimes p^*(\mathfrak{R}/R)) = d \cdot \xi^{\dim(\mathfrak{R}/R)},$$

where d is a rational number. Further, the cones $\overline{\mathcal{H}}_{g,n,P}^R$ and $\overline{\mathcal{H}}_{g,n,P}$ have the same Segre class, and thus

$$s_0 = p_*(\xi^{\text{rk}(\overline{\mathcal{H}}_{g,n,P})-1}) = p_*([\mathbb{P}\overline{\mathcal{H}}_{g,n,P}^R]_{\xi^{\text{rk}(\overline{\mathcal{H}}_{g,n,P})-1}}) = ds_0,$$

and the coefficient d is equal to 1. □

Proposition 2.28 *The Segre class of $\overline{\mathcal{H}}_{g,n,P}$ is given by*

$$\prod_{i=1}^m \frac{(p_i - 1)^{p_i - 1}}{(p_i - 1)!} \cdot \frac{1 - \lambda_1 + \dots + (-1)^g \lambda_g}{\prod_{i=1}^m (1 - (p_i - 1)\psi_i)}.$$

Proof From the above lemma, we have

$$\begin{aligned}
 s_*(\overline{\mathcal{H}}_{g,n,P}) &= s_*(\overline{\mathcal{H}}_{g,n,P}^0) \\
 &= c_*(\overline{\mathcal{H}}_{g,n+m})^{-1} \cdot s_*\left(\bigoplus_{i=n+1}^m \mathcal{P}_{n+i}\right) \\
 &= c_*(\overline{\mathcal{H}}_{g,n+m}^\vee) \cdot s_*\left(\bigoplus_{i=n+1}^m \mathcal{P}_{n+i}\right) \\
 &= \prod_{i=1}^m \frac{(p_i - 1)^{p_i - 1}}{(p_i - 1)!} \cdot \frac{1 - \lambda_1 + \dots + (-1)^g \lambda_g}{\prod_{i=1}^m (1 - (p_i - 1)\psi_i)}.
 \end{aligned}$$

From the third line to the fourth we have used the fact that $c(\overline{\mathcal{H}}_g)^{-1} = c(\overline{\mathcal{H}}_g^\vee)$ (see [28]). □

2.5 Unstable base

Here we extend the definition of the spaces of stable differentials to differentials supported on an unstable base.

Definition 2.29 A triple (g, n, P) composed of a nonnegative integers g and n and a vector P of positive integers is *semistable* if either

- $2g - 2 + n + \ell(P) > 0$ (in which case we also say that (g, n, P) is *stable*);
- $g = 0, n = 1$ and $P = (p)$ with $p > 1$; or
- $g = 0, n = 0$ and $P = (1, p)$ with $p > 1$.

We want to define the space $\overline{\mathcal{H}}_{g,n,P}$ for all semistable

triples. However, the space $\overline{\mathcal{M}}_{0,2}$ is empty; thus, we cannot define the spaces $\overline{\mathcal{H}}_{0,1+1,(p)}$ and $\overline{\mathcal{H}}_{0,2,(1,p)}$ as cones over a moduli space of curves. Still, we can define the cone structure of these two spaces over $\text{Spec}(\mathbb{C})$.

The space $\overline{\mathcal{H}}_{0,1+1,(p)}$ is defined as the complement of $\{u = 0\}$ in the space of generalized principal parts defined in Section 2.1. In other words $\overline{\mathcal{H}}_{0,1+1,(p)}$ is the spectrum of the graded subalgebra of $\mathbb{C}[a_1, \dots, a_{p-2}]$ generated by monomials with integral weights (where the weight of a_j is $j/(p - 1)$).

The space $\overline{\mathcal{H}}_{0,2,(1,p)}$ is the spectrum of the graded subalgebra of $\mathbb{C}[a_1, \dots, a_{p-2}, r]$ generated by monomials with integral weights, where r (for residue) has weight 1.

2.6 Stable differentials on disconnected curves

In the paper, we will need stable differentials supported on disconnected curves. Let q be a positive integer, and

$$\mathbf{g} = (g_1, g_2, \dots, g_q), \quad \mathbf{n} = (n_1, n_2, \dots, n_q), \quad \mathbf{m} = (m_1, m_2, \dots, m_q)$$

be lists of nonnegative integers, and let

$$\mathbf{P} = (P_j)_{1 \leq j \leq q} = (P_{j,i})_{1 \leq j \leq q, 1 \leq i \leq m_j}$$

be a list of vectors of positive integers of length m_j .

Definition 2.30 The triple $(\mathbf{g}, \mathbf{n}, \mathbf{P})$ is *stable* (or *semistable*) if the triple (g_j, n_j, P_j) is stable (or semistable) for all $1 \leq j \leq q$ (see Definition 2.29).

Unless otherwise stated, we assume from now on that $(\mathbf{g}, \mathbf{n}, \mathbf{P})$ is semistable.

Definition 2.31 The *space of stable differentials of type $(\mathbf{g}, \mathbf{n}, \mathbf{P})$* is the space

$$\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}} = \prod_{i=1}^q \overline{\mathcal{H}}_{g_i, n_i, P_i}.$$

We define the *interior* of $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}$ as the open substack $\mathcal{H}_{\mathbf{g}, \mathbf{n}, \mathbf{P}} \subset \overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}$ of differentials supported on smooth curves.

Definition 2.32 The *reduced base of type $(\mathbf{g}, \mathbf{n}, \mathbf{P})$* (or of type $(\mathbf{g}, \mathbf{n}, \mathbf{m})$) is the space

$$\overline{\mathcal{M}}_{\mathbf{g}, \mathbf{n}, \mathbf{m}}^{\text{red}} = \prod_j \overline{\mathcal{M}}_{g_j, n_j + m_j} \\ 2g_j - 2 + n_j + m_j > 0$$

if the product is nonempty and $\text{Spec}(\mathbb{C})$ otherwise.

Proposition 2.33 *The space of stable differentials of type \mathbf{P} is a cone over $\overline{\mathcal{M}}_{\mathbf{g}, \mathbf{n}, \mathbf{m}}$. If the triple $(\mathbf{g}, \mathbf{n}, \mathbf{P})$ is stable then the Segre class is given by*

$$s(\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}) = \prod_{j=1}^q s(\overline{\mathcal{H}}_{g_j, n_j + m_j, P_j}),$$

where $s(\overline{\mathcal{H}}_{g_j, n_j, P_j})$ is the pullback of the Segre class of $\overline{\mathcal{H}}_{g_j, n_j, P_j}$ to the product $\prod_{j=1}^q \overline{\mathcal{M}}_{g_j, n_j + m_j}$ under the j^{th} projection.

Proof The proof is straightforward because the space $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}$ is a product of cones. \square

To handle the residues, we extend the definition of the space of residues \mathfrak{R} :

$$(9) \quad \mathfrak{R} = \bigoplus_{j=1}^q \mathfrak{R}_j = \left\{ (r_{j,i})_{j,i} \mid \sum_{i=1}^{m_j} r_{j,i} = 0 \text{ for all } j \in [1, q] \right\} \subset \mathbb{C}^{m_1 + \dots + m_q}.$$

Definition 2.34 Let R be a vector subspace of \mathfrak{R} . The space $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}^R$ is the space of stable differentials with residues lying in R .

Lemma 2.35 Let R be a linear subspace of \mathfrak{R} . The space $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}^R$ is a subcone of $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}^R$ of codimension $\dim(\mathfrak{R}) - \dim(R)$ and we have:

- The cones $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}$ and $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}^R$ have the same Segre class.
- The Poincaré-dual class of $[\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}^R]$ in $H^*(\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}, \mathbb{Q})$ is given by

$$\xi^{\dim(\mathfrak{R}) - \dim(R)}.$$

Proof The proof of Lemma 2.27 can be adapted immediately to the general case. \square

Definition 2.36 Let $p: \overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}} \rightarrow \overline{\mathcal{M}}_{\mathbf{g}, \mathbf{n}, \mathbf{m}}^{\text{red}}$ be the projection to the base. The tautological ring of $\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}$ is the subring of $H^*(\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}})$ generated by $\xi = c_1(\mathcal{O}(1))$ and pullbacks by p of tautological classes from the base $\overline{\mathcal{M}}_{\mathbf{g}, \mathbf{n}, \mathbf{m}}^{\text{red}}$. We denote this ring by $RH^*(\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}})$.

2.7 Semistable graphs

Let $\mathbf{g}, \mathbf{n}, \mathbf{m}$ and \mathbf{P} be lists of genera, numbers of marked points without poles, numbers of marked poles and vectors of positive integers indexed by $j \in \llbracket 1, q \rrbracket$ as in the previous section. We assume that $(\mathbf{g}, \mathbf{n}, \mathbf{P})$ is semistable.

In this section we define a combinatorial object called semistable graphs. We show here that the space $\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}$ has a natural stratification according to semistable graphs and that semigraphs allow to define some tautological classes.

Definition 2.37 A semistable graph of type $(\mathbf{g}, \mathbf{n}, \mathbf{P})$ is given by the data

$$\left(V, H, g: V \rightarrow \mathbb{N}, a: H \rightarrow V, i: H \rightarrow H, E, \pi^0(V, E) \simeq \llbracket 1, q \rrbracket, L \simeq \bigcup_{j=1}^q \llbracket 1, n_j + m_j \rrbracket \right),$$

satisfying the following properties:

- V is a vertex set with a genus function g .

- H is a half-edge set equipped with a vertex assignment a and an involution i .
- The edge set E is defined as the set of length 2 orbits of i in H (self-edges at vertices are permitted).
- The graph (V, E) has q labeled connected components.
- For all $1 \leq j \leq q$, the genus of the connected component labeled by j is defined by $\sum g(v) + \#(E_j) - \#(V_j) + 1$ and is equal to g_j .
- L is the set of fixed points of i , called *legs*.
- For all $1 \leq j \leq q$, there are $n_j + m_j$ legs on the j^{th} connected component and this set of legs is identified with the set $\llbracket 1, n_j + m_j \rrbracket$.
- For each vertex v in V belonging to the j^{th} component:
 - Let $n(v)$ be the number of legs adjacent to v with label at most n_j .
 - Let $m(v)$ be the number of legs adjacent to v with label at least $n_j + 1$.
 - Let $P'(v) = (P_{j,m-n_j})_{m \mapsto v, m > n_j}$: it is the vector obtained from P_j by keeping only the entries associated to the legs of the second type adjacent to v . We denote by $P(v)$ the concatenation of $P'(V)$ with the vector $(1, \dots, 1)$ of length equal to number of half-edges adjacent to v that are not legs.
- For each vertex v , the triple $(g(v), n(v), P(v))$ is semistable.

We define the following lists indexed by the vertices of Γ :

$$g_\Gamma = (g(v))_{v \in V}, \quad n_\Gamma = (n(v))_{v \in V}, \quad m_\Gamma = (m(v))_{v \in V}, \quad P_\Gamma = (P(v))_{v \in V}.$$

The triple $(g_\Gamma, n_\Gamma, P_\Gamma)$ is semistable (it is implied by the last condition of the definition of a semistable graph). We consider the space $\overline{\mathcal{H}}_{g_\Gamma, n_\Gamma, P_\Gamma}$. We denote by \mathfrak{R}_Γ the space of residues of $\overline{\mathcal{H}}_{g_\Gamma, n_\Gamma, P_\Gamma}$. We define the subspace $R_\Gamma \subset \mathfrak{R}$ by the equations

$$r_h + r_{h'} = 0$$

for all edges $e = (h, h')$.

Notation 2.38 Let Γ be a semistable graph; we denote by $\overline{\mathcal{H}}_\Gamma$ the moduli space $\overline{\mathcal{H}}_{g_\Gamma, n_\Gamma, P_\Gamma}^{R_\Gamma}$ and by

$$\zeta_\Gamma^\# : \overline{\mathcal{H}}_\Gamma \rightarrow \overline{\mathcal{H}}_{g, n, P}$$

the natural closed morphism.

Proposition 2.39 *The set of semistable graphs is finite and the space $\overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}}$ is stratified according to the semistable graphs; ie for all x in $\overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}}$ there exists a unique graph Γ such that $x \in \zeta_\Gamma(\mathcal{H}_\Gamma)$.*

Proof If we fix the datum $(\mathbf{g}, \mathbf{n}, \mathbf{P})$, then there are finitely many semistable graphs Γ for $(\mathbf{g}, \mathbf{n}, \mathbf{P})$ such that the graph Γ is stable. Indeed, there are finitely many stabilization of Γ and then the graph Γ is determined by the choice of which set of marked points is on an unstable rational component (we recall that unstable rational bridges between components are not permitted because the triple $(0, 0, (1, 1))$ is not semistable).

Now, for all semistable graphs the only possible unstable vertices are vertices of genus 0 with two marked points: a leg and a half-edge. Therefore, for all stable graphs Γ of type $(\mathbf{g}, \mathbf{n}, \mathbf{P})$, there are finitely many semistable graphs Γ' such that the stabilization of Γ' is equal to Γ . Therefore, there are finitely many semistable graphs.

Now, if x is a point in $\overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}}$ then, if we denote by Γ the dual graph of the underlying curve of x then x lies in $\zeta_\Gamma(\mathcal{H}_\Gamma)$. This graph is uniquely determined. □

The space $\mathbb{P}\overline{\mathcal{H}}_\Gamma$ is a cone; thus, it has a tautological line bundle $\mathcal{O}(1)$. This line bundle is the pullback by $\zeta_\Gamma^\#$ of the tautological line bundle of $\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}}$. By abuse of notation we will write ξ for the first Chern class of the tautological line bundle for both spaces. We have the following important proposition:

Proposition 2.40 *Let Γ be semistable graph. The morphism $\zeta_{\Gamma,*}^\#: H^*(\mathbb{P}\overline{\mathcal{H}}_\Gamma, \mathbb{Q}) \rightarrow H^*(\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}}, \mathbb{Q})$ maps tautological classes to tautological classes.*

Proof Let Γ be a semistable graph. Let $k \geq 0$ and $\beta \in \overline{\mathcal{M}}_\Gamma^{\text{red}}$. We need to prove that the class $\zeta_{\Gamma,*}^\#(\xi^k p^*(\beta))$ is tautological. We will prove this statement in three steps.


Stable graphs We suppose first that Γ is a stable graph. We recall that in this case we have defined a map $\zeta_\Gamma: \overline{\mathcal{M}}_\Gamma \rightarrow \overline{\mathcal{M}}_{\mathbf{g},\mathbf{n},\mathbf{m}}$. Then $\overline{\mathcal{H}}_\Gamma$ is the fiber product

$$\begin{array}{ccc} \overline{\mathcal{H}}_\Gamma & \xrightarrow{\zeta_\Gamma^\#} & \overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}} \\ p_\Gamma \downarrow & & \downarrow p \\ \overline{\mathcal{M}}_\Gamma & \xrightarrow{\zeta_\Gamma} & \overline{\mathcal{M}}_{\mathbf{g},\mathbf{n},\mathbf{m}} \end{array}$$

Let β be a cohomology class in $H^*(\overline{\mathcal{M}}_\Gamma, \mathbb{Q})$. We use the projection formula and the fact that $\overline{\mathcal{H}}_\Gamma$ is a fiber product to get $\zeta_{\Gamma,*}^\#(\xi^k \cdot p_\Gamma^*(\beta)) = \xi^k p^*(\zeta_{\Gamma,*}(\beta))$. Therefore, if the

class β belongs to the tautological ring $RH^*(\overline{\mathcal{M}}_\Gamma, \mathbb{Q})$, then the class $\zeta_{\Gamma^*}^\#(\xi^k \cdot p_\Gamma^*(\beta))$ belongs to the tautological ring of $\overline{\mathcal{H}}_{g,n,P}$.

Graph with one main vertex We no longer assume that Γ is stable. Let $1 \leq j \leq q$ and $1 \leq i \leq m_j$. Let p_i be the i^{th} entry of P_j . Assume that Γ is the graph




(we take the trivial graph for all the other connected components). We will prove that the class $\zeta_{\Gamma^*}^\#(1)$ lies in $RH^*(\mathbb{P}\overline{\mathcal{H}}_{g,n,m,P})$. We use the parametrization of the cone of principal parts at x

$$\left[\left(\frac{u}{z}\right)^{p_i-1} + a_1 \left(\frac{u}{z}\right)^{p_i-2} + \dots + a_{p_i-2} \left(\frac{u}{z}\right) \right] \frac{dz}{z}.$$

The stratum defined by Γ is the vanishing locus of u . We have seen that u^{p_i-1} is a section of the line bundle $\text{Hom}(\mathcal{O}(-1), \mathcal{L}_i^{p_i-1})$. Therefore, the vanishing locus of u has Poincaré-dual class given by

$$[u = 0] = \frac{1}{p_i-1} \xi - \psi_i.$$

By the same argument, if Γ is the graph



where the set $\{i_k\}$ is a set of indices in $\llbracket 1, m_j \rrbracket$, then we have

$$\zeta_{\Gamma^*}^\#(1) = \prod_k \left(\frac{1}{p_{i_k}-1} \xi - \psi_{i_k} \right).$$

And, more generally, for a class β in $RH^*(\overline{\mathcal{M}}_{g,n,m,P}^{\text{red}})$ and $k \in \mathbb{N}$, we have

$$\zeta_{\Gamma^*}^\#(\xi^k \beta) = \xi^k \beta \cdot \prod_k \left(\frac{1}{p_{i_k}-1} \xi - \psi_{i_k} \right) \in RH^*(\overline{\mathcal{H}}_{g,n,m,P}).$$

General unstable graph We combine the two previous arguments. Let Γ be a general semistable graph. Let $\tilde{\Gamma}$ be the graph obtained by contracting all edges between stable vertices. We have $\overline{\mathcal{M}}_{\mathbf{g},\mathbf{n},\mathbf{m}}^{\text{red}} = \overline{\mathcal{M}}_{\tilde{\Gamma}}^{\text{red}}$. The space $\overline{\mathcal{H}}_{\Gamma}$ is the fiber product

$$\begin{array}{ccccc} \overline{\mathcal{H}}_{\Gamma} & \longrightarrow & \overline{\mathcal{H}}_{\tilde{\Gamma}}^{\text{red}} & \xrightarrow{\zeta_{\tilde{\Gamma}}^{\#}} & \overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{m}} \\ p_{\Gamma} \downarrow & & \downarrow p_{\tilde{\Gamma}} & & \\ \overline{\mathcal{M}}_{\Gamma} & \xrightarrow{\zeta_{\Gamma}} & \overline{\mathcal{M}}_{\mathbf{g},\mathbf{n},\mathbf{m}}^{\text{red}} & & \end{array}$$

Thus, $\zeta_{\Gamma,*}^{\#}(\xi^k p_{\Gamma}^* \beta) = \zeta_{\tilde{\Gamma},*}^{\#}(\xi^k p_{\tilde{\Gamma}}^*(\zeta_{\Gamma,*} \beta))$. Now $\tilde{\Gamma}$ has one stable vertex, and $\zeta_{\tilde{\Gamma},*} \beta \in RH^*(\overline{\mathcal{M}}_{\mathbf{g},\mathbf{n},\mathbf{m}}^{\text{red}})$; thus, the class $\zeta_{\Gamma,*}^{\#}(\xi^k p_{\Gamma}^* \beta)$ is tautological. \square

3 Stratification of spaces of stable differentials

The interior of space of stable differentials is stratified according to the orders of the zeros of the differential. In this section we study the local parametrization of these strata and compute their dimension.

3.1 Definitions and notation

In the paper we will often consider the following setup:

Assumption 3.1 The quadruple $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$ is of the following type:

- $\mathbf{g} = (g_1, \dots, g_q)$, $\mathbf{Z} = (Z_1, \dots, Z_q)$ and $\mathbf{P} = (P_1, \dots, P_q)$ are lists of the same length $q \geq 1$.
- For all $1 \leq j \leq q$, g_j is a positive integer, Z_j is a vector of nonnegative integers of length n_j and P_j is a vector of positive integers of length m_j .
- We write $\mathbf{n} = (n_1, \dots, n_q)$ and $\mathbf{m} = (m_1, \dots, m_q)$.
- The triple $(\mathbf{g}, \mathbf{n}, \mathbf{P})$ is semistable (in the sense of Definition 2.30).
- R is a linear subspace of $\mathfrak{X} = \bigoplus_{j=1}^q \mathfrak{X}_j \simeq \bigoplus_{j=1}^q \mathbb{C}^{m_j-1}$ (defined as in (9)).

Let $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$ be a quadruple satisfying Assumption 3.1.

Notation 3.2 We denote by

$$A_{\mathbf{g},\mathbf{Z},\mathbf{P}}^R \subset \mathcal{H}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$$

the locus of points $(C, (x_{j,i})_{1 \leq j \leq q, 1 \leq i \leq n_j + m_j}, \alpha) \in \mathcal{H}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ such that C is smooth

and α is nonzero on each connected component and has a zero of order exactly $k_{j,i}$ at the i^{th} point of the j^{th} connected component for all $1 \leq j \leq q$ and $1 \leq i \leq n_j$.

If there is no condition on the residues, we will simply denote it by $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}$.

Definition 3.3 We say that \mathbf{Z} is *complete* for (\mathbf{g}, \mathbf{P}) if Z_j is complete for (g_j, P_j) for all $1 \leq j \leq q$.

3.2 Standard coordinates

In this section we describe how to parametrize differentials with prescribed singularities. We use the notation $\Delta_\rho = \{z \in \mathbb{C} : |z| < \rho\}$ for the disks of radius $\rho \in \mathbb{R}^+$ and $A_{\rho_1, \rho_2} = \{z \in \mathbb{C} : \rho_1 < |z| < \rho_2\}$ for the annulus of parameters $0 < \rho_1 < \rho_2$.

3.2.1 Standard coordinates Let α be a meromorphic differential on a small disk $\Delta_\rho \subset \mathbb{C}$. We denote by r the residue of α at 0. Then, there exists a conformal map $\varphi: \Delta_{\rho'} \rightarrow \Delta_\rho$ for ρ' small enough such that $\varphi(0) = 0$ and

$$\varphi^*(\alpha) = \begin{cases} d(z^k) & \text{if } 0 \text{ is a zero of order } k - 1, \\ r dz/z & \text{if } 0 \text{ is a pole of order } 1, \\ d(1/z^k) + r dz/z & \text{if } 0 \text{ is a pole of order } k + 1. \end{cases}$$

The map φ is unique up to multiplication of the coordinate z by a k^{th} root of unity when 0 is a zero of order $k - 1$ or a pole of order $k + 1$. The coordinate z will be called the *standard coordinate*.

More generally, if U is an open neighborhood of 0 in \mathbb{C}^n and α_u is a holomorphic family of differentials on Δ_ρ such that the order of α_u at 0 is constant, then there exists a holomorphic map $\varphi: \tilde{U} \times \Delta_{\rho'} \rightarrow \Delta_\rho$ such that $\varphi(u, \cdot)^*(\alpha_u)$ is in the standard form for some neighborhood \tilde{U} of 0. Once again the map φ is unique up to multiplication of the standard coordinate by a root of unity.

Now the following classical lemma describes the deformations of $d(z^k)$ (see [25] for a proof):

Lemma 3.4 Let $\rho > 0$ and $U \subset \mathbb{C}^n$ be a domain containing 0. Let α_u be a family of holomorphic differentials on Δ_ρ such that α_0 has a zero of order $k - 1$ at the origin. Then, there exists $\rho' > 0$, a neighborhood \mathcal{Z} of 0 in \mathbb{C}^{k-2} and a conformal map

$$\varphi: U \times \Delta_{\rho'} \rightarrow \Delta_\rho \times \mathcal{Z}$$

such that $\varphi(u, \cdot)^*(\alpha_u) = d(z^k + a_{k-2}z^{k-2} + \dots + a_1z)$. The map φ is unique up to multiplication of z by a k^{th} root of unity.

The locus $z = 0$ determines a section of the projection $U \times \Delta_\rho$ that does not depend on the choice of k^{th} root of unity. This section is called the *local center of mass of zeros*.

Now we would like to generalize the above lemma to deformations of poles of order 1.

Definition 3.5 Let $\rho > 0$ and $U \subset \mathbb{C}^n$ be a domain containing 0. Let α be a differential on Δ_ρ in the standard form $d(z^k)$. A *standard deformation* of α is defined by a holomorphic function $\beta: U \times \Delta_\rho \rightarrow \mathbb{C}$ satisfying $\beta(0, z) = 0$. A standard deformation associated to β is the family of differentials on Δ_ρ parametrized by U

$$\alpha_u = d(z^k) + \frac{\beta(u, z)}{z} dz.$$

In general, there exists no standard coordinate for a standard deformation. However, the following proposition has been proved in [2] (see Theorem 4.3):

Proposition 3.6 We consider the annulus A_{ρ_1, ρ_2} for any choice of $0 < \rho_1 < \rho_2 < \rho$. Choose a point $p \in A_{\rho_1, \rho_2}$ and $\zeta^l = \exp(2i\pi l/k)$ a k^{th} root of unity. Choose a map $\sigma: U \rightarrow \Delta_\rho$ such that $\sigma(0) = \zeta^l p$. Then there exists a neighborhood \tilde{U} of 0 in U and a holomorphic map $\varphi: \tilde{U} \times A_{\rho_1, \rho_2} \rightarrow \Delta_R$ such that

$$\varphi_u^*(\alpha_u) = d(z^k) + \frac{\beta(u, 0)}{z} dz,$$

and $\varphi(0, z) = \zeta^l z$ and $\varphi(u, p) = \sigma(u)$ for all $u \in \tilde{U}$ and $z \in A_{\rho_1, \rho_2}$. For \tilde{U} small enough, the map φ is unique.

3.2.2 Neighborhood of strata Let $(g, \mathbf{Z}, \mathbf{P}, R)$ be a quadruple satisfying Assumption 3.1.

Lemma 3.7 There exists a neighborhood V of $A_{g, \mathbf{Z}, \mathbf{P}}$ in $\mathcal{H}_{g, \mathbf{n}, \mathbf{P}}$ and a holomorphic retraction $\eta: V \rightarrow A_{g, \mathbf{Z}, \mathbf{P}}$ such that η preserves the residues at the poles.

Proof The general statement follows immediately from the connected case. Indeed, $A_{g, \mathbf{Z}, \mathbf{P}}$ is locally isomorphic to $\prod_{j=1}^q A_{g_j, \mathbf{Z}_j, \mathbf{P}_j}$; therefore, we can define the neighborhood V and the retraction η as the products of the V_j and η_j for all $1 \leq j \leq q$. Therefore, we will assume that $q = 1$.

Let $y_0 = (C_0, x_1, \dots, x_{n+m}, \alpha_0)$ be a point in $A_{g, \mathbf{Z}, \mathbf{P}}$. Let n' be the number of zeros of α distinct from the marked points. We choose an ordering of these zeros $(\tilde{x}_1, \dots, \tilde{x}_{n'})$ and we denote by \tilde{k}_i the order of α at \tilde{x}_i for all $1 \leq i \leq n'$.

We denote by $d = \dim(A_{g,Z,P})$ and by $d' = \dim(\mathcal{H}_{g,n,P})$. A neighborhood of y_0 in $A_{g,Z,P}$ is of the form $U/\text{Aut}(y_0)$ where U is a contractible domain of \mathbb{C}^d . A neighborhood of $U/\text{Aut}(y_0)$ in $\mathcal{H}_{g,n,P}$ is of the form $W/\text{Aut}(y_0)$ where W is a contractible domain of $\mathbb{C}^{d'}$.

For all $y = (C, \alpha, (x_{j,i}))$ in U we denote by $P(y) \subset C$ the set of poles of α and by $Z(y)$ the set of zeros (marked or not). For all y , the form α determines a class in the relative cohomology group $H^1(C \setminus P(y), Z(y), \mathbb{C})$. Further, we have a canonical identification of $H^1(C \setminus P(y), Z(y), \mathbb{C})$ with $H^1(C_0 \setminus P(y_0), Z(y_0), \mathbb{C})$ (this is the Gauss–Manin connection); therefore, we have a holomorphic map

$$\Phi_U: U \rightarrow H^1(C_0 \setminus P(y_0), Z(y_0), \mathbb{C}).$$

This map can be described as follows. Let $(\gamma_1, \dots, \gamma_d)$ be simple closed curves of $C_0 \setminus (P(y_0) \cup Z(y_0))$ that form a basis of the relative homology group

$$H_1(C_0 \setminus P(y_0), Z(y_0), \mathbb{Z}).$$

Then the map Φ_U is defined by

$$\Phi_U: U \rightarrow H^1(C_0 \setminus P(y_0), Z(y_0), \mathbb{C}), \quad (C, \alpha, (x_i)) \mapsto \left(\gamma \mapsto \int_{\gamma_i} \alpha \right),$$

where the cycles on $C_0 \setminus (P(y_0) \cup Z(y_0))$ are identified with cycles on $C \setminus (P(y) \cup Z(y))$ by the Gauss–Manin connection. The map Φ_U is a local biholomorphism (see [4] for example). We call the map Φ_U a *period coordinates* chart.

Now we will construct the holomorphic maps

$$\begin{aligned} \Phi^1: W &\rightarrow H^1(C_0 \setminus P(y_0), Z(y_0), \mathbb{C}), \\ \Phi^{2,i}: W &\rightarrow \mathcal{Z}^{k_i} \quad \text{for all } 1 \leq i \leq n, \\ \Phi^{3,i}: W &\rightarrow \tilde{\mathcal{Z}}^{k_i} \quad \text{for all } 1 \leq i \leq n', \end{aligned}$$

where \mathcal{Z}_i is a domain of \mathbb{C}^{k_i} containing of 0 for all $1 \leq i \leq n$ and $\tilde{\mathcal{Z}}_i$ is a domain of $\mathbb{C}^{\tilde{k}_i-1}$ containing of 0 for all $1 \leq i \leq n'$:

- For all $1 \leq i \leq n$, the map $\Phi^{2,i}$ is determined by a slight modification of Lemma 3.4 for marked differentials. We consider a tubular neighborhood $W \times \Delta_\rho \rightarrow C_W$ around the i^{th} section of the universal curve. There exists a $\rho' > 0$ and a neighborhood \mathcal{Z}_i of $0 \in \mathbb{C}^{k_i}$ with coordinates $(a_{i,1}, \dots, a_{i,k_i})$ and a map $\varphi: W \times \Delta_\rho \rightarrow \Delta_{\rho'} \times \mathcal{Z}_i$ such that the marked point is at $z_i = 0$ and

$$\alpha_s = d(z_i^{k_i+1} + a_{i,k_i} z_i^{k_i} + \dots + a_{i,1} z_i)$$

for each point s of W . The map φ is unique up to a multiplication of z_i by a $(k_i + 1)^{\text{st}}$ root of unity. Thus, we have defined a map from W to \mathcal{Z}_i given by $\alpha_s \mapsto (a_{i,1}, \dots, a_{i,k_i})$.

- For all $1 \leq i \leq n'$, the map $\Phi^{2,i}$ is determined by Lemma 3.4. We consider a tubular neighborhood $W \times \Delta_\rho \rightarrow C_W$ around the i^{th} section of the universal curve. There exists a $\rho' > 0$ and a neighborhood $\tilde{\mathcal{Z}}_i$ of $0 \in \mathbb{C}^{\tilde{k}_i - 1}$ with coordinates $(a_{i,1}, \dots, a_{i,\tilde{k}_i - 1})$ and a map $\varphi: W \times \Delta_\rho \rightarrow \Delta_{\rho'} \times \mathcal{Z}_i$ such that

$$\alpha_y = d(z_i^{\tilde{k}_i + 1} + \dots + a_{i,1}z_i)$$

for each point y of W . The map φ is once again unique up to a multiplication of z_i by a $(\tilde{k}_i + 1)^{\text{st}}$ root of unity. Thus, we have defined a map from W to \mathcal{Z}_i given by $\alpha_y \mapsto (a_{i,1}, \dots, a_{i,\tilde{k}_i - 1})$.

Further, the point $z_i = 0$ is called the center of mass of the differential. It does not depend on the choice of a root of unity; therefore, we have a uniquely determined point $\tilde{x}_i \in C$ for all s .

- The map Φ^1 is defined as Φ_U by the Gauss–Manin connection. For a point $y = (C, \alpha, x_1, \dots, x_{n+m})$ in W we let $Z(y) = \{x_1, \dots, x_n\} \cup \{\tilde{x}_1, \dots, \tilde{x}_{n'}\}$ (the union of the marked points with the center of masses defined above). Then the differential α defines a point in $H^1(C \setminus P(y), Z(y))$, which is once again canonically identified with $H^1(C_0 \setminus P(y_0), Z(y_0))$.

We will prove that the map

$$\Phi = \Phi^1 \times \left(\prod_{i=1}^n \Phi^{2,i} \right) \times \left(\prod_{i=1}^{n'} \Phi^{3,i} \right)$$

is a local biholomorphism (see [25, Section 5.2] in the holomorphic case). The source and the target have the same dimension; therefore, we only need to check that the differential of each component of Φ is surjective. For Φ^1 this is obvious because $\Phi^1|_U = \Phi_U$ is a local biholomorphism.

Let $1 \leq i \leq n$ and let Δ_ρ be a disk in C_0 around x_i such that $\alpha = d(w^{k_i + 1})$. Up to a choice of a smaller \mathcal{Z}_i , for all $(a_{i,1}, \dots, a_{i,k_i - 1}) \in \mathcal{Z}_i$ we have $(z^{k_i + 1} + \dots + a_{i,1}z) \neq 0$ for all $\frac{1}{2}\rho < |z| < \rho$. Then we construct a family of curves $C_i \rightarrow \mathcal{Z}_i$ by gluing the two families of curves $(C_0 \setminus \Delta_{\rho/2}) \times \mathcal{Z}_i$ with $\Delta_\rho \times \mathcal{Z}_i$ along the identification

$$w = (z^{k_i + 1} + \dots + a_{i,1}z)^{1/(k_i + 1)}$$

(this family depends on the choice of a $(k_i + 1)^{\text{st}}$ root). Now, the differential α on C_i is determined by α_0 on $(C_0 \setminus \Delta_{\rho/2}) \times \mathcal{Z}_i$ and by $d(z^{k_i+1} + \dots + a_{i,1}z)$ on $\Delta_{\rho} \times \mathcal{Z}_i$. The two differentials agree by construction of the complex structure. Therefore, the differential of $\Phi^{2,i}$ is surjective. The same argument holds for $\Phi^{3,i}$ for $1 \leq i \leq n'$.

Now we set $\eta_W = \Phi_U^{-1} \circ \Phi^1$. This retraction does not depend on the choice of the root of unity nor on the choice of ordering of the nonmarked zeros. Indeed, it is defined by the inverse procedure of patching $d(w^{k_i+1})$ instead of $d(z^{k_i+1} + \dots + a_{i,1}z)$ for all $1 \leq i \leq n$ (and for nonmarked zeros). Therefore, it does not depend on the local identification of the relative homology group. Thus, if we consider two maps η_W and $\eta_{W'}$ (for neighborhoods of points y_0 and y'_0) then these two maps agree on $W \cap W'$.

Finally, the residues are preserved by η . Indeed, for any choice of y_0 , we can choose a basis $(\gamma_1, \dots, \gamma_d)$ of $H_1(C_0 \setminus P(y_0), \mathbb{Z}(y_0), \mathbb{Z})$ such that γ_i is a small loop around the $(n+i)^{\text{th}}$ marked point for all $1 \leq i \leq m-1$. The period of α around this loop is the residue of α at the i^{th} pole and is preserved by η . □

Corollary 3.8 *The residue map restricted to $A_{g,Z,P}^R \rightarrow R$ is a submersion.*

Proof Let $(C, x_1, \dots, x_{n+m}, \alpha)$ be a point of $A_{g,Z,P}^R$. Let $r = (r_1, \dots, r_m)$ be a vector in R . There exists a meromorphic differential φ on C with at most simple poles at the m last marked points with residues prescribed by r . Let Δ be a disk of \mathbb{C} centered at 0 and parametrized by ϵ . Let η be the retraction map of Lemma 3.7. The residues of $\eta(\alpha + \epsilon\varphi)$ at the poles are given by

$$\text{res}_{x_{n+i}}(\alpha) + \epsilon r_i.$$

Thus, the vector r belongs to the image of the tangent space of $A_{g,Z,P}^R$ under the differential of the map res . □

Remark 3.9 Recently, Gendron and Tahar studied the surjectivity of the residue maps for open strata in the space of meromorphic differentials (and also of higher-order differentials — see [19]). Our statement that the residue map is a submersion does not imply surjectivity. However, the image of an algebraic submersion is always a Zariski open set. Thus, we can claim that the residue map is surjective on the *closure* of every nonempty stratum.

3.2.3 Neighborhood of strata with appearance of residues We consider a slightly more general setup. Let $q \geq 2$ and g, n, n' and m be lists of nonnegative integers of length q . Let $P = (P_1, \dots, P_q)$ be a list of vectors of positive integers such that

$\text{length}(P_j) = m_j$ for all $1 \leq j \leq q$ and let $Z = (Z_1, \dots, Z_q)$ be a list of vectors of nonnegative integers such that $\text{length}(Z_j) = n_j + n'_j$. We assume that the triple $(\mathbf{g}, \mathbf{n} + \mathbf{n}', \mathbf{P})$ is semistable (in the sense of Definition 2.30).

For all $1 \leq j \leq q$, we denote by $P'_j = (p_1, \dots, p_{m_j}, 1, \dots, 1)$ the vector obtained from p by appending n'_j 1s and by $Z'_j = (k_1, \dots, k_n)$ the vector obtained by erasing the last n' entries of Z .

The space $\mathcal{H}_{\mathbf{g}, \mathbf{n} + \mathbf{n}', \mathbf{P}}$ is embedded in $\mathcal{H}_{\mathbf{g}, \mathbf{n}, \mathbf{P}'}$. We denote by \mathfrak{X} and \mathfrak{X}' the vector spaces of residues of $\mathcal{H}_{\mathbf{g}, \mathbf{n} + \mathbf{n}', \mathbf{P}}$ and $\mathcal{H}_{\mathbf{g}, \mathbf{n}, \mathbf{P}'}$. Let R' be a vector subspace of \mathfrak{X}' . The vector space \mathfrak{X} is a vector subspace of \mathfrak{X}' , and we let $R = \mathfrak{X} \cap R'$. We have the series of embeddings

$$A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R \hookrightarrow A_{\mathbf{g}, \mathbf{Z}', \mathbf{P}}^R \hookrightarrow A_{\mathbf{g}, \mathbf{Z}', \mathbf{P}'}^{R'}$$

Proposition 3.10 *Let y_0 be a point in $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$. Let U be a neighborhood of y_0 in $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$. There exists a neighborhood V of y_0 in $A_{\mathbf{g}, \mathbf{Z}', \mathbf{P}'}^{R'}$ and a map*

$$\phi: V \xrightarrow{\sim} U \times \left(\prod_{\substack{1 \leq j \leq q \\ 1 \leq i \leq n'_j}} \mathcal{Z}_{j,i} \right) \times \mathcal{Z},$$

where:

- $\mathcal{Z}_{j,i}$ is a neighborhood of 0 in $\mathbb{C}^{k_j \cdot n_j + i}$ for all $1 \leq j \leq q$ and $1 \leq i \leq n'_j$, and \mathcal{Z} is a neighborhood of 0 in R'/R .
- If Δ_ρ is a disk and $s: U \times \Delta_\rho \rightarrow \left(\prod \mathcal{Z}_{j,i} \right) \times \mathcal{Z}$ is a holomorphic map such that $s(u, 0) = 0$, then the family of differentials

$$\tilde{s}: U \times \Delta_\rho \rightarrow V, \quad (u, \epsilon) \mapsto \phi^{-1}(u, s(u, \epsilon)),$$

is a standard deformation of $d(z^{k_j \cdot n_j + i + 1})$ for all $1 \leq j \leq q$ and $1 \leq i \leq n'_j$.

Proof We have seen that a neighborhood of U in $A_{\mathbf{g}, \mathbf{Z}', \mathbf{P}}^R$ is isomorphic to the space $U \times \prod_{j=1}^q \prod_{i=1}^{n_j + n'_j} \mathcal{Z}_{j,i}$. For all $1 \leq j \leq q$ and $1 \leq i \leq n'_j$, the differential at the marked point $x_{j, n_j + i}$ is given by $d(z^{k_j \cdot n_j + i} + a_1 z^{k_j \cdot n_j + i} + \dots)$ (Lemma 3.4).

Now, for all $1 \leq j \leq q$ and $1 \leq i \leq n_j$, we choose a meromorphic differential $\varphi_{j,i}$ with simple poles at the marked points in such a way that the vectors of residues $\mathbf{r}_{j,i}$ of $\varphi_{j,i}$ form a basis of R'/R . The residue map $A_{\mathbf{g}, \mathbf{Z}', \mathbf{P}'}^{R'} \rightarrow R'$ is a submersion (Corollary 3.8). Thus, a neighborhood of $U \times \prod \mathcal{Z}_{j,i}$ in $A_{\mathbf{g}, \mathbf{Z}', \mathbf{P}'}^{R'}$ is naturally identified with some $U \times \left(\prod \mathcal{Z}_{j,i} \right) \times \mathcal{Z}$ with \mathcal{Z} a neighborhood of 0 in R'/R . The identification is given by adding a linear combination of the $\varphi_{j,i}$.

Both the deformations of U into $U \times \prod \mathcal{Z}_{j,i}$ and the deformations of $U \times \prod \mathcal{Z}_{j,i}$ into $U \times (\prod \mathcal{Z}_{j,i}) \times \mathcal{Z}$ are standard deformations at the marked point x_{j,n_j+i} for all $1 \leq j \leq q$ and $1 \leq i \leq n_j$. \square

The isomorphism ϕ is not unique. Our construction depends on the choice of standard coordinates at the x_{j,n_j+i} for all $1 \leq j \leq q$ and $1 \leq i \leq n_j$ and on the choice of the differentials $\varphi_{j,i}$ with simple poles. However, Proposition 3.10 implies the following corollary:

Corollary 3.11 *If a morphism ϕ satisfies the conditions of Proposition 3.10, then it defines a local retraction $\eta: V \rightarrow U$ such that $\eta \circ \tilde{s} = \text{Id}_U$ for any holomorphic section $s: U \times \Delta_\rho \rightarrow (\prod \mathcal{Z}_{j,i}) \times \mathcal{Z}$.*

3.3 Dimension of the strata

Let (g, Z, P, R) be quadruple satisfying Assumption 3.1.

Definition 3.12 A completion of Z is a list of q vectors of nonnegative integers $Z'_1 = (k'_{1,1}, \dots, k'_{1,n'_1}), \dots, Z'_q = (k'_{q,1}, \dots, k'_{q,n'_q})$ such that

- $n'_j \geq n_j$ for all $1 \leq j \leq q$;
- $k'_{j,i} \geq k_{j,i}$ for all $1 \leq j \leq q$ and $1 \leq i \leq n_j$;
- Z' is complete for (g, P) .

We will say that the completion Z' is exterior if $k'_i = k_i$ for all j and all $1 \leq i \leq n_j$. Finally, we will denote by Z_m the maximal completion, ie the exterior completion of Z that satisfies $k'_{j,i} = 1$ for all j and $n_j + 1 \leq i \leq n'_j$.

If Z' is a completion of Z , we denote by $\pi: A^R_{g,Z',P} \rightarrow A^R_{g,Z,P}$ the forgetful map of marked point that are not accounted for by Z , ie the restriction of the forgetful map of marked points $\pi: \mathcal{H}_{g,n',P} \rightarrow \mathcal{H}_{g,n,P}$ to $A^R_{g,Z',P}$. We have the following straightforward lemma:

Lemma 3.13 We have

$$A^R_{g,Z,P} = \bigcup_{Z'} \pi(A^R_{g,Z',P}),$$

where the union is over all exterior completions of Z .

Lemma 3.14 *If $q = 1$ and the vector Z is complete for g and P , then the forgetful map of the differential $p: A^R_{g,Z,P} \rightarrow p(A^R_{g,Z,P}) \subset \mathcal{M}_{g,n+m}$ is a line bundle minus the zero section. In particular, $\mathbb{P}A^R_{g,Z,P}$ is isomorphic to its image.*

Proof Let (C, x_1, \dots, x_{n+m}) be a point of $\text{Im}(p)$. The curve C is smooth and the divisor $\omega_C - \sum_{i=1}^n k_i(x_i) + \sum_{j=1}^m p_j(x_{n+j})$ is a principal divisor of degree 0. Therefore, the fiber of p over (C, x_1, \dots, x_{n+m}) is given by the nonzero multiples of one differential with fixed orders of zeros and poles. \square

Proposition 3.15 *The space $A_{g,Z,P}^R$ is either empty or of codimension*

$$\sum_{j=1}^q |Z_j| + \dim(\mathfrak{R}/R)$$

in $\mathcal{H}_{g,n,P}$.

Proof First we assume that $q = 1$ (connected case), Z is complete and $R = \mathfrak{R}$ (no residue condition). The dimension of $\mathbb{P}A_{g,Z,P}$ is equal to the dimension of its image in the moduli space of curves. Then the image of $\mathbb{P}A_{g,Z,P}$ is of dimension $2g - 2 + n$ if P is empty (see [32]) and $2g - 3 + n + m$ otherwise (see [15]). By a simple count of dimension we can check that the proposition is valid in this specific case.

We no longer assume that $q = 1$ (but we still assume that Z is complete and $R = \mathfrak{R}$). Then the space $A_{g,Z,P}$ is birationally equivalent to $\prod_j A_{g_j,Z_j,P_j}$. Thus, $\dim(A_{g,Z,P}) = \sum \dim(A_{g_j,Z_j,P_j})$ and once again, the proposition holds by a simple count of dimensions

Now, we still assume that Z is complete; however, we no longer assume that $R = \mathfrak{R}$. We have seen that the residue map $A_{g,Z,P}^R \rightarrow R$ is a submersion; therefore, the dimension of $A_{g,Z,P}^R$ is equal to the dimension of R plus the dimension of the fiber of the residue map at any point. If we consider the case $R = \mathfrak{R}$, then we see that the dimension of the fiber at any point is $\dim A_{g,Z,P} - \dim \mathfrak{R}$. Therefore, the dimension of $A_{g,Z,P}^R$ is equal to $\dim A_{g,Z,P} - (m - 1) + \dim(R)$. Thus, the proposition is valid for all choices of R .

Now let Z be any vector. Let Z' be an exterior completion of Z . The map

$$\pi: A_{g,Z',P}^R \rightarrow A_{g,Z,P}^R$$

is quasifinite. Indeed, the preimage of a point $(C, x_1, \dots, x_{n+m}, \alpha)$ is finite: the points in the preimage correspond to the different orderings of the zeros that are not accounted for by Z .

The proof of Lemma 3.7 implies that if $A_{g,Z',P}$ is not empty for some exterior completion then $A_{g,Z_m,P}$ is not empty: indeed, we can always perturb a differential

to “break up” a zero of order greater than 1. By counting the dimensions, we have $\dim(A_{g, Z_m, P}^R) > \dim(A_{g, Z', P}^R)$ for all exterior completions $Z' \neq Z_m$. Therefore, $\dim(A_{g, Z_m, P}^R) = \dim(A_{g, Z, P}^R)$ and the proposition is proved. \square

3.4 Fibers of the map $p: A_{g, Z, P}^R \rightarrow \mathcal{M}_{g, n, m}$

Let $(g, Z, P, R \subset \mathfrak{R})$ be a quadruple satisfying Assumption 3.1. Throughout this section we assume that the triple (g, n, P) is *stable*.

If the context is clear, we denote by the same letter the map $p: \mathcal{H}_{g, n, P} \rightarrow \mathcal{M}_{g, n, m}$ and its restriction $p: A_{g, Z, P}^R \rightarrow p(A_{g, Z, P}^R)$. We denote by $\text{Im}(p) = p(A_{g, Z, P}^R) \subset \mathcal{M}_{g, Z, P}$ its image.

We recall that, by definition, $\mathfrak{R} = \bigoplus_{j=1}^q \mathfrak{R}_j \simeq \bigoplus_{j=1}^q \mathbb{C}^{m_j-1}$ (see Section 2.6).

Notation 3.16 Let $1 \leq j \leq q$; we denote by $\text{pr}_j: \mathfrak{R} \rightarrow \mathfrak{R}_j$ the projection onto \mathfrak{R}_j along $\bigoplus_{j' \neq j} \mathfrak{R}_{j'}$. We denote by R_j the space $\text{pr}_j(R)$.

Remark 3.17 The linear relations that define the space R may involve residues at poles of different connected components. Thus, in general we have $R \cap \mathfrak{R}_j \subsetneq R_j$.

Let $1 \leq j \leq q$. We denote by p_j the map from $A_{g_j, Z_j, P_j}^{R_j}$ to $\mathcal{M}_{g_j, n_j + m_j}$. Finally, we denote by $\text{Im}(p_j)$ the image of p_j . We have a natural embedding of $A_{g, Z, P}^R$ into $\prod_{j=1}^q A_{g_j, Z_j, P_j}^{R_j}$ and of $\text{Im}(p)$ into $\prod_{i=1}^n \text{Im}(p_j)$.

The purpose of this section is to state the condition $(\star\star)$ (see Notation 3.23) that ensures that the projectivized morphism $p: \mathbb{P} A_{g, Z, P}^R \rightarrow \text{Im}(p)$ is birational. This will be needed in Section 4.4 to describe the boundary divisors of the stratum $A_{g, Z, P}^R$. We will proceed in two steps: first we consider the case that Z is complete and then a general Z .

Complete case For now we assume that Z is complete for g and P .

We have seen that the fact that Z_j is complete for all $1 \leq j \leq q$ implies that $A_{g_j, Z_j, P_j}^{R_j} \rightarrow \text{Im}(p_j)$ is a line bundle minus the zero section. We denote by L_j the pullback of this line bundle to $\text{Im}(p)$.

We define the j^{th} *evaluation map of residues* $\text{ev}_j: L_j \rightarrow \mathfrak{R}_j$ as the morphism of vector bundles over $\text{Im}(p)$ given by the evaluation of the residues at the j^{th} connected component. We define the *evaluation of residues* as the morphism of vector bundles $\text{ev} = (\bigoplus_{j=1}^q \text{ev}_j): \bigoplus_{j=1}^q L_j \rightarrow \mathfrak{R}$.

Remark 3.18 The evaluation map (ev) and the residue map (res) are not defined on the same spaces. The first one is a morphism of vector bundles on the space $\text{Im}(p)$ while the second one is defined as a morphism of vector bundles over $\mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$. If $q = 1$, then $\mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$ is isomorphic to its image and the two morphisms are equal.

Proposition 3.19 *Suppose that \mathbf{Z} is complete. Then the families*

$$p: A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R \rightarrow \text{Im}(p)$$

and

$$\tilde{p}: \text{ev}^{-1}(R) \cap \left(\prod_{j=1}^q L_j^* \right) \rightarrow \text{Im}(p)$$

are isomorphic. If $q \geq 2$, the fiber of p over a point is of dimension 1 if and only if ev is injective and $R \cap \text{ev}(\bigoplus_j L_j)$ is of dimension 1.

Proof The proposition is straightforward for $q = 1$. We suppose from now on that $q \geq 2$.

For a point $x \in \text{Im}(p)$, the fiber of p can be described as follows: it is the choice of a nonzero differential for each connected component such that the residues at the poles define a vector in R . Therefore, the fiber over x is the subset of points of $\prod L_j^*$ with residues in R . This fiber is given by $\text{ev}^{-1}(R) \cap \prod_{j=1}^q L_j^*$.

The fiber of $\text{ev}^{-1}(R) \cap \prod_{j=1}^q L_j^*$ over $x \in \text{Im}(p)$ is not empty. Indeed, suppose that for some $1 \leq j \leq q$ the space $\text{ev}^{-1}(R)$ is contained in $\{0\} \times \bigoplus_{j' \neq j} L_{j'}$; then the residue condition R imposes that the differential on one of the components is zero, in which case x is not a point of $\text{Im}(p)$. Therefore, the dimension of $\text{ev}^{-1}(R) \cap \prod_{j=1}^q L_j^*$ is the same as the dimension of $\text{ev}^{-1}(R) \cap \bigoplus_{j=1}^q L_j$.

The only point that remains to prove is: if the map ev is not injective then the fiber of p is of dimension greater than 1. We assume that the map ev is not injective. Then one of the L_j is mapped to zero for some $1 \leq j \leq q$: indeed, for all $1 \leq j \leq q$, the j^{th} component of ev is the composition of $\text{ev}_j: L_j \rightarrow \mathfrak{R}_j$ with the inclusion of $\mathfrak{R}_j \rightarrow \mathfrak{R}$; thus, if a vector in $\bigoplus L_j$ with a nonzero j^{th} entry is mapped to zero in \mathfrak{R} then the generator of L_j is mapped to zero in \mathfrak{R}_j and L_j is mapped to zero in \mathfrak{R} .

Therefore,

$$\text{ev}^{-1}(R) \cap \bigoplus_{j=1}^q L_j = L_j \oplus \left(\text{ev}^{-1}(R) \cap \bigoplus_{j' \neq j} L_{j'} \right).$$

We have seen that $\text{ev}^{-1}(R)$ cannot be contained in $L_j \times \{0\}$; thus, the second summands is of positive dimension and $\text{ev}^{-1}(R) \cap \bigoplus_{j=1}^q L_j$ is of dimension greater than 1. \square

Let Σ be the union of the vector subspaces $R \cap \ker(\text{pr}_j)$ for $1 \leq i \leq q$. If R is of positive dimension, we denote by $\mathbb{P}\Sigma$ the image of Σ in $\mathbb{P}R$. This is the locus of vectors of residues that vanish on at least one connected component. Suppose that all R_j are of positive dimension; then $\Sigma \subsetneq R$ and there is a natural map $\rho: \mathbb{P}R \setminus \mathbb{P}\Sigma \rightarrow \prod_{j=1}^q \mathbb{P}R_j$ defined as the projection on each factor.

Notation 3.20 We will say that the tuple of residue vector spaces $(\mathfrak{R}, R, (\mathfrak{R}_j)_{1 \leq j \leq q})$ satisfies the condition (\star) if either $q = 1$ or the two following conditions hold:

- The spaces R and the R_j are of positive dimension.
- There exists an open and dense set U in $\mathbb{P}R$ such that the restriction of the natural map $\rho: \mathbb{P}R \setminus \mathbb{P}\Sigma \rightarrow \prod_{i=1}^q \mathbb{P}R_j$ to U is finite.

Proposition 3.21 *Suppose that Z is complete and that q is at least 2. Then the fiber of p over a generic point of $\text{Im}(p)$ is of dimension 1 if and only if $(\mathfrak{R}, R, (\mathfrak{R}_j)_{1 \leq j \leq q})$ satisfy the condition (\star) .*

Proof We have already seen that if R_j is reduced to the trivial space, then the map $\text{ev}: \bigcup_{j=1}^q L_j \rightarrow \mathfrak{R}$ is not injective and the fibers of p are all of dimension greater than 1 (see the proof of Proposition 3.19). We assume that all R_j are nontrivial. For all j , we denote by $A_j^0 \subset A_{g,Z,P}^R$ to be the locus of differentials with zero residues on the j^{th} component. The image of A_j^0 by the residue map lies in $R \cap \ker(\text{pr}_j)$ which is of positive codimension in R . Further, the residue map is a submersion; thus, $\dim(A_j^0) < \dim(A_{g,Z,P}^R)$. We will write

$$A' = A_{g,Z,P}^R \setminus \bigcup_{j=1}^q A_j^0.$$

The locus A' is dense in $A_{g,Z,P}^R$. If we assume that the fibers of p are generically of dimension 1, then $p(A')$ is also dense in $\text{Im}(p)$. Therefore, we only need to prove that a generic point of $p(A')$ has fibers of dimension 1 if and only if condition (\star) is satisfied.

It is easy to check that the residue map sends A' to $R \setminus \Sigma$. Therefore, the locus $p(A')$ is the locus of points such that the map ev defined in the proof of Proposition 3.19 is injective. Thus, a point of $p(A')$ has fibers of dimension 1 by p if and only if $R \cap \text{ev}(\bigoplus_j L_j)$ is of dimension 1. Now, $R \cap \text{ev}(\bigoplus_j L_j)$ is of dimension 1 if and only if the preimage under ρ of the point $(L_1, \dots, L_q) \in \prod_{j=1}^q \mathbb{P}R_j$ is composed of a unique point.

Now, the residue map is a submersion from $A_{g,Z,P}^R$ to R . Therefore, the map ρ is finite on a dense open subset of $\mathbb{P}R \setminus \mathbb{P}\Sigma$ if and only if the fiber of p is of dimension 1 on a dense open set of $\text{Im}(p)$. \square

3.4.1 General case We no longer assume that Z is complete. We denote by $Z_m = (Z_{1,m}, \dots, Z_{q,m})$ the maximal completion of Z . Further, we denote by

$$p_m: A_{g,Z_m,P}^R \rightarrow \text{Im}(p_m)$$

the forgetful map of the differential.

Proposition 3.22 *We suppose that $(\mathfrak{X}, R, (\mathfrak{X}_j)_{1 \leq j \leq q})$ satisfies the condition (\star) . Then we have $\dim(\text{Im}(p_m)) = \dim(\text{Im}(p))$ if and only if for all $1 \leq j \leq q$ we have $\dim(A_{g_j,Z_j,P_j}^{R_j}) - 1 \leq \dim(\mathcal{M}_{g_j,n_j+m_j})$.*

Proof We proceed in two steps: first we assume that the base is connected and then we consider the general case.

Connected case We assume that $q = 1$. In this case, the “only if” is trivial. Indeed, $\mathbb{P}A_{g,Z,P}^R = \dim(\text{Im})(p_m)$ and $\dim(\text{Im})(p) \leq \mathcal{M}_{g,n+m}$.

We assume that the dimension of $\mathbb{P}A_{g,Z,P}^R$ is less than or equal to the dimension of $\mathcal{M}_{g,n+m}$. We have the commutative diagram

$$\begin{array}{ccc} A_{g,Z_m,P}^R & \longrightarrow & A_{g,Z,P}^R \\ p_m \downarrow & & \downarrow p \\ \text{Im}(p_m) & \longrightarrow & \text{Im}(p) \end{array}$$

where the horizontal arrows are the forgetful map of the zeros that are not accounted for by Z . We have seen that the image of $A_{g,Z_m,P}^R$ is dense in $A_{g,Z,P}^R$. Therefore, the image of $\text{Im}(p_m)$ under the forgetful map of the points that are not accounted for by Z is dense in $\text{Im}(p)$. Then we have $\dim(\text{Im}(p_m)) \geq \dim(\text{Im}(p))$. Now we will prove that $\dim(\mathbb{P}A_{g,Z,P}^R) \leq \dim(\text{Im}(p))$.

We consider the two vector bundles over the moduli space of curves $\mathcal{M}_{g,n+m}$

$$K\mathcal{M}_{g,n}(P) = R^0\pi_*\left(\omega_C\left(\sum_{i=1}^m p_i\sigma_{n+i}\right)\right), \quad E = \mathfrak{X}/R \oplus \left(\bigoplus_{i=1}^n J_{i,k_i}^{\text{hol}}\right),$$

where J_{i,k_i}^{hol} is the vector space of holomorphic jets of order k_i at the marked point x_i , ie

$$J_{i,k_i}^{\text{hol}} = R^0 \pi_*(\omega(-k_i x_i)/\omega)$$

(beware the vector space of jets here is not the vector space of polar jets used in Section 2.2). We have a morphism $e: K\mathcal{M}_{g,n}(P) \rightarrow E$. The rank of $K\mathcal{M}_{g,n}(P)$ is $r_1 = g - 1 + \sum p_i$ if P is not empty and $r_1 = g$ otherwise. The rank of E is $r_2 = \dim(\mathfrak{R}/R) + \sum k_i$. By assumption, we have

$$\dim(\mathbb{P}A_{g,Z,P}^R) = \dim(\mathcal{M}_{g,n+m}) + r_1 - r_2 - 1 \leq \dim(\mathcal{M}_{g,n+m}).$$

Let $\mathcal{E} \subset \mathcal{M}_{g,n+m}$ be the locus where e is not injective. We have $r_1 \leq r_2 + 1$; thus, the locus \mathcal{E} is of codimension at most $r_2 - r_1 + 1$ because it is the vanishing locus of $r_2 - r_1 + 1$ minors of the map e . Therefore, the locus \mathcal{E} is of dimension greater than or equal to $\dim(\mathbb{P}A_{g,Z,P}^R) = \dim(\text{Im}(p_m))$.

To complete the proof, we show that $\text{Im}(p)$ is open and dense in \mathcal{E} . Let P' be a vector of m positive integers such that $P' \leq P$. Let Z' be a vector of n nonnegative integers such that $Z' \geq Z$. The image of $\mathbb{P}A_{g,Z',P'}^R$ lies in \mathcal{E} . Conversely, the locus \mathcal{E} is the union of all the $\text{Im}(p')$ where p' is the map from $\mathbb{P}A_{g,Z',P'}^R$ to $\mathcal{M}_{g,n+m}$ for $P' \leq P$ and $Z' \geq Z$. We have $\dim(\mathbb{P}A_{g,Z',P'}^R) < \dim(\mathbb{P}A_{g,Z,P}^R) \leq \dim(\mathcal{E})$ if $P' < P$ or $Z' > Z$. Therefore, all irreducible components of $\text{Im}(p)$ have the same dimension as \mathcal{E} and $\dim(\text{Im}(p)) = \dim(\text{Im}(p_m))$.

Disconnected case Suppose that there exists $1 \leq j \leq q$ such that $\dim(\mathbb{P}A_{g_j,Z_j,P_j}^{R_j}) > \dim(\mathcal{M}_{g_j,n_j+m_j})$. Then the fibers of the map $\text{Im}(p_{j,m}) \rightarrow \text{Im}(p_j)$ are of positive dimension. Thus, for all points in $\text{Im}(p)$ the fibers of the map $A_{g,Z,P} \rightarrow \text{Im}(p)$ are of positive dimension.

Conversely, suppose that $\dim(\mathbb{P}A_{g_j,Z_j,P_j}^{R_j}) \leq \dim(\mathcal{M}_{g_j,n_j+m_j})$ for all $1 \leq j \leq q$. Thus, for all $1 \leq j \leq q$, we have $\dim(\text{Im}(p_j)) = \dim(\text{Im}(p_{j,m})) = \dim(\mathbb{P}A_{g_j,Z_j,P_j}^{R_j})$. Therefore, there exists a dense open subset $U_j \subset \mathbb{P}A_{g_j,Z_j,P_j}^{R_j}$ such that the morphism $\text{Im}(p_{j,m}) \rightarrow \text{Im}(p_j)$ is finite over its image. Further, the map $\mathbb{P}A_{g,Z,P}^R \rightarrow \mathbb{P}R$ and the maps $\mathbb{P}A_{g_j,Z_j,P_j}^{R_j} \rightarrow \mathbb{P}R_j$ are submersions. Thus, for all $1 \leq j \leq q$, the image of U_j under the residue map is an open subset of $\mathbb{P}R_j$, which we denote by $\tilde{U}_j \subset \mathbb{P}R_j$.

Now we consider the morphism $\rho: \mathbb{P}R \setminus \mathbb{P}\Sigma \rightarrow \prod_j U_j$. We claim that the preimage of $\prod_j U_j$ under ρ is a nonempty open subset in $\mathbb{P}R$. Indeed, if we suppose that $\rho^{-1}(\prod_j U_j)$ is empty, then the image of $\mathbb{P}R \setminus \mathbb{P}\Sigma$ under ρ is contained in a finite

union of closed subsets of the form $(\mathbb{P}R_j \setminus \tilde{U}_j) \times \prod_{j' \neq j} \mathbb{P}R_{j'}$ for some $1 \leq j \leq q$. However, the space $\mathbb{P}R \setminus \mathbb{P}\Sigma$ is irreducible; thus, its image under ρ is contained in one such subspace. This would imply that the image of $\mathbb{P}R \setminus \mathbb{P}\Sigma \rightarrow \mathbb{P}R_j$ is contained in a closed subspace and this is not possible (because R_j is the image of the projection of R onto \mathfrak{R}_j).

Putting everything together, the preimage of $\prod_j \tilde{U}_j$ under the composition of morphisms $\mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R \rightarrow \mathbb{P}R \setminus \mathbb{P}\Sigma \rightarrow \prod_j R_j$ is an open and dense subspace U :

$$\begin{array}{ccc}
 U \subset \mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R & \longrightarrow & \text{Im}(p) \\
 \downarrow & & \downarrow \\
 \prod_{j=1}^q U_j \subset \prod_{j=1}^q \mathbb{P}A_{\mathbf{g}_j, \mathbf{Z}_j, \mathbf{P}_j}^{R_j} & \longrightarrow & \prod_{j=1}^q \text{Im}(p_j)
 \end{array}$$

The lower arrow is finite from $\prod_j U_j$ to its image. By construction, the subspace U is embedded in $\prod U_j$. Therefore, the map $U \rightarrow \text{Im}(p)$ is finite over its image and $\dim(\text{Im}(p)) = \dim(\mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R) = \dim(\text{Im}(p_m))$. □

Notation 3.23 We will say that $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$ satisfies condition $(\star\star)$ if and only if the two following conditions are satisfied:

- The vector spaces $(\mathfrak{R}, R, (\mathfrak{R}_j)_{1 \leq j \leq q})$ satisfy the condition (\star) .
- For all $1 \leq j \leq q$, we have $\dim(A_{\mathbf{g}_j, \mathbf{Z}_j, \mathbf{P}_j}^{R_j}) - 1 \leq \dim(\mathcal{M}_{\mathbf{g}_j, n_j + m_j})$.

Proposition 3.24 *The morphism $p: \mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R \rightarrow \text{Im}(p)$ is birational if and only if $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$ satisfies the condition $(\star\star)$.*

Proof Proposition 3.21 implies that $\dim(\text{Im}(p)) = \dim(\mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R)$ if and only if $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$ satisfies the condition $(\star\star)$. Therefore, if $p: \mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R \rightarrow \text{Im}(p)$ is birational then the condition $(\star\star)$ is satisfied.

Conversely, if $(\star\star)$ is satisfied, then there exists a dense open subspace U in $\text{Im}(p)$ such that for any point in U , the fiber of p over this point is finite. Suppose that there are at least two points in the preimage of a marked curve $(C, (x_{j,i})_{j,i}) \in \text{Im}(p)$. Then there exist two nonproportional meromorphic differentials α and α' supported on C with orders of zeros and poles prescribed by \mathbf{Z} and \mathbf{P} and with the same residues at the poles. Any nonzero linear combination of these two differentials is in $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$ and in the preimage of $(C, x_{j,i})$. This is a contradiction with the finiteness of the fibers of p over U . □

4 Boundary components of strata of stable differentials

Let $(g, Z, P, R \subset \mathfrak{R})$ be a quadruple satisfying Assumption 3.1.

Notation 4.1 We denote by $\overline{A}_{g,Z,P}^R$ and $\mathbb{P}\overline{A}_{g,Z,P}^R$ the Zariski closures of $A_{g,Z,P}^R$ and $\mathbb{P}A_{g,Z,P}^R$ in $\overline{\mathcal{H}}_{g,Z,P}$ and $\mathbb{P}\overline{\mathcal{H}}_{g,Z,P}$, respectively.

In this section we describe the boundary components of $\overline{A}_{g,Z,P}^R$. We will see that these can be described with combinatorial objects called P -admissible graphs. We also describe the subset of boundary divisors among these boundary components.

4.1 Twisted graphs with level structures

We introduce P -admissible graphs here and in the subsequent section, we explain how they correspond to strata of $\overline{A}_{g,Z,P}^R$.

Let Γ be a semistable graph of type (g, n, P) . We denote by H_e the set of half-edges of Γ which are not legs.

Definition 4.2 A *twist* on Γ is a function

$$I: H_e \rightarrow \mathbb{Z}$$

satisfying the following conditions:

- If h and h' form an edge, then $I(h) + I(h') = 0$.
- Let v and v' be two vertices, and $\{(h_1, h'_1), \dots, (h_n, h'_n)\}$ be the set of edges from v to v' . Then either $I(h_j) = 0$ for all $1 \leq j \leq n$, or $I(h_j) > 0$ for all $1 \leq j \leq n$, or $I(h_j) < 0$ for all $1 \leq j \leq n$. We say that $v = v'$, or $v > v'$, or $v < v'$, depending on the above inequalities.
- The relation \leq thus defined on vertices is transitive.

For brevity, a semistable graph endowed with a twist function will be called a *twisted graph*. If (Γ, I) is a twisted graph, the above conditions define a partial order on the set of vertices of Γ .

Definition 4.3 A *level structure* on a twisted graph is a function

$$l: \text{Vertices} \rightarrow \mathbb{Z}^-,$$

compatible with the partial order induced by the twist; ie for all vertices v and v' ,

$$v = v' \implies l(v) = l(v'), \quad v < v' \implies l(v) < l(v').$$

We impose that the image of l is an interval containing all integers from 0 to $-d$ and we call d the *depth* of the twisted graph. We will denote by V^i the set of vertices of level i .

Definition 4.4 An edge between vertices of the same level will be called a *horizontal edge*.

Definition 4.5 A twisted graph with level structure is called P -*admissible* if all marked poles of order at least 2 belong to vertices of level 0. For brevity we will call such graphs *admissible graphs*.

This definition of P -admissibility implies in particular that unstable vertices can only be present at the level 0. In the sequel, we will see that P -admissible graphs represents loci in $\mathcal{H}_{g,n,P}$, where the differential vanishes identically on the components of negative levels. As explained in the introduction, the appearance of unstable components on the level 0 ensures that the poles remains of fixed order.

Remark 4.6 The reader should keep in mind that a stable differential cannot vanish identically on an unstable component. Indeed, otherwise there would be infinitely many automorphisms of the curve preserving the differential; this would contradict the stability condition (see Definition 1.2).

Example 4.7 We represent in Figure 1 an example of admissible graph. Each vertex v is represented by a circle containing the integer g_v . The marked poles and zeros are represented by legs. A leg corresponding to a pole (respectively a zero) of order k is marked by $-k$ (respectively $+k$). The twists are indicated on each edge.

Definition 4.8 Let (Γ, I, l) be a semistable graph with a twist and a level structure. We say that (Γ, I, l) is a *twisted stable graph* if Γ is a stable graph (in the sense of Definition 1.10).

Definition 4.9 Let (Γ, I, l) be a semistable graph with a twist and a level structure. We say that (Γ, I, l) is *realizable* if for all vertices v of Γ we have

$$(10) \quad \sum_{(j,i) \mapsto v} k_{j,i} - \sum_{(j,n_j+i) \mapsto v} p_{j,i} + \sum_{h \mapsto v} I(h) - 1 \leq 2g(v) - 2,$$




Figure 1: An example of admissible graph of genus 7 for the vectors $Z = (2, 4, 4, 7)$ and $P = (-3, -2)$

where the sums are respectively over marked points corresponding to zeros, marked points corresponding to poles and half-edges adjacent to v .


The following lemma will be needed later to compare the space of stable differentials and the incidence variety:

Lemma 4.10 *If Z is complete, then there exists a bijection between the set of realizable and admissible graphs and the set of realizable and twisted stable graphs.*

Proof To an admissible graph we assign its stabilization. The twists and levels on this graph are obtained by restriction of the former twists and level functions.

From a twisted stable graph, we construct an admissible graph by adding an unstable vertex for each marked point corresponding to a pole of order p greater than 1 and adjacent to a vertex of level < 0 . This new vertex is of level 0 and the new edge between this vertex and the rest of the curve has twists given by $+p - 1$ and $-p + 1$. \square

Example 4.11 Here is the stabilization of the admissible graph of Figure 1:



4.2 Boundary strata associated to admissible graphs

Let $(g, \mathbf{Z}, \mathbf{P}, R \subset \mathfrak{A})$ be a quadruple satisfying Assumption 3.1. Let (Γ, I, l) be an admissible graph. In this subsection, we assign to this admissible graph a stratum of abelian differentials $A_{\Gamma, I, l} \subset \overline{\mathcal{H}}_{g, n, \mathbf{P}}$ that lies in the closure of $A_{g, \mathbf{Z}, \mathbf{P}}^R$. We build this stratum level by level.

To every level 0 vertex we assign a substack of the corresponding space of differentials. To every vertex of negative levels we assign a substack of the corresponding moduli space of curves. The product of these cycles will give us a substack of the space $\overline{\mathcal{H}}_{\Gamma}$ by putting an identically vanishing differential on every component of the curve of negative level. Thus, our input is (\mathbf{Z}, R) and an admissible graph (Γ, I, l) of type (g, n, \mathbf{P}) ; our output is a collection of subspaces of the spaces of differentials (for level 0 vertices) and of the spaces of curves (for vertices of negative levels).

Levels 0 and -1 We respectively denote by q_0 and q_1 the numbers of vertices of levels 0 and -1. Further, we denote by \mathbf{g}_0 and \mathbf{g}_1 the lists of genera of vertices of levels 0 and -1. We determine orders of zeros and poles as follows:

- For all $1 \leq j \leq q_0$, we construct the vector P_j^0 by taking the entries of \mathbf{P} for all marked poles on the j^{th} component and a -1 for each horizontal half-edge; we construct the vector Z_j^0 by taking the entries of \mathbf{Z} for all marked zeros carried by the j^{th} component and $I(h) - 1$ for each half-edge h to a deeper level.
- For all $1 \leq j \leq q_1$, we construct the vector P_j^1 by taking $I(h) + 1$ for all half-edges to levels 0 and 1 for all horizontal half-edges adjacent to the j^{th} component; we construct the vector Z_j^1 by taking the entries of \mathbf{Z} for all marked zeros carried by the j^{th} component and $I(h) - 1$ for each half-edge to a deeper level.
- We write $\mathbf{Z}_i = (Z_1^i, \dots, Z_{q_i}^i)$ and $\mathbf{P}_i = (P_1^i, \dots, P_{q_i}^i)$ for $i = 0, 1$.

Now we define the residue conditions as follows:

- We let hor_0 be the number of horizontal half-edges of level 0 and let $\mathfrak{R}^H = \mathbb{C}^{\text{hor}_0}$. We denote by \mathfrak{R}^1 the space of residues of the space of stable differentials $\overline{\mathcal{H}}_{g_1, n_1, \mathbf{P}_1}$ (where n_1 is determined by the length of entries of \mathbf{Z}_1).
- We define

$$\text{proj}: \mathfrak{R} \oplus \mathfrak{R}^H \oplus \mathfrak{R}^1 \rightarrow \mathfrak{R}^1$$

as the projection along $\mathfrak{R} \oplus \mathfrak{R}^H$.

- We consider the vector subspace $R' = R \oplus \mathfrak{R}^H \oplus \mathfrak{R}^1$, and we define the vector subspace \tilde{R} of R' by the following linear relations:
 - $r_h + r'_h = 0$ for all horizontal edges (h, h') .
 - For all vertex of level 0, we have

$$\sum_{p \rightarrow v} r_p + \sum_{\substack{h \text{ horizontal} \\ h \rightarrow v}} r_h + \sum_{\substack{h \text{ to level } -1 \\ h \rightarrow v}} -r_h = 0,$$

where the first sum is over marked poles adjacent to v , the second is over horizontal edges and the last one is over the edges to level -1 (in this last sum r_h is the value of the residue at the corresponding half-edge of level -1).

- Finally, we let $R^0 = \ker(\text{proj}) \cap \tilde{R}$ and $R^1 = \text{proj}(\tilde{R})$.


With these data, we define the level 0 and -1 strata as

$$A_{\Gamma, I, l}^0 = A_{g_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0} \subset \overline{\mathcal{H}}_{g_0, n_0, \mathbf{P}_0},$$

$$A_{\Gamma, I, l}^1 = p(A_{g_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1}) \subset \overline{\mathcal{M}}_{g_1, n_1, m_1} = \prod_{v \in V^1} \overline{\mathcal{M}}_{g_v, n_v + m_v},$$

where $p: \overline{\mathcal{H}}_{g_1, n_1, \mathbf{P}_1} \rightarrow \overline{\mathcal{M}}_{g_1, n_1, m_1}$ is the forgetful map.

Example 4.12 To illustrate the definition of R^0 and R^1 , we compute all vector spaces for the following two graphs:



On these two examples we have not represented the genera of the vertices and we have only represented the legs with poles (thus at level 0). In the first case, $R = \mathfrak{R} = \{0\}$ (there are no poles). In the second case, we assume that $R = \mathfrak{R} \simeq \mathbb{C}$ (we impose no condition on the residues).

All letters stand for the value of the residue, ie for a coordinate in $\tilde{\mathfrak{R}} \oplus \mathfrak{R}^1$ corresponding either to a half-edge or to a marked pole. In the following table we give the dimensions and equations of all subvector spaces of $\tilde{\mathfrak{R}}$ and a presentation of $\tilde{\mathfrak{R}}^1$ and R^1 :


Vector space	Left example	Right example
$\mathfrak{R} \oplus \mathfrak{R}^H \oplus \mathfrak{R}^1$	$\{0\} \oplus \{0\} \oplus \mathbb{C}^2$	$\mathbb{C} \oplus \{0\} \oplus \mathbb{C}^2$
R'	$\mathfrak{R} \oplus \mathfrak{R}^H \oplus \mathfrak{R}^1$	$\mathfrak{R} \oplus \mathfrak{R}^H \oplus \mathfrak{R}^1$
relations from edges	none	none
relations from vertices	$\{a + b = 0\}$	$\{c - a - b = 0\}$
\tilde{R}	$\{a = \epsilon, b = -\epsilon : \epsilon \in \mathbb{C}\}$	$\{a = \epsilon_1, b = \epsilon_2, c = -\epsilon_1 - \epsilon_2 : (\epsilon_1, \epsilon_2) \in \mathbb{C}^2\}$
R^1	$\{a = \epsilon, b = -\epsilon : \epsilon \in \mathbb{C}\}$	$\{a = \epsilon_1, b = \epsilon_2 : (\epsilon_1, \epsilon_2) \in \mathbb{C}^2\}$
R^0	$\{0\}$	$\{0\}$

Level $-\ell$ Let (Γ', I', l') be the graph obtained from Γ by contracting edges between vertices of levels 0 through $-\ell + 1$. The twist on Γ restricts to Γ' and the level structure is shifted. Vertices of levels 0 to $-\ell + 1$ merge to level 0, level $-\ell$ vertices become level -1 vertices, and so on. Therefore, we have the natural identification

$$\prod_{\substack{v \in V(\Gamma) \\ l(v) = -\ell}} \bar{\mathcal{M}}_{g(v), n(v)} = \prod_{\substack{v \in V(\Gamma') \\ l(v) = -1}} \bar{\mathcal{M}}_{g(v), n(v)}$$

and we define $A_{\Gamma, I, l}^\ell$ as $A_{\Gamma', I', l'}^1$.

Example 4.13 The contraction of levels 0 and -1 of the admissible graph of Figure 1 gives the following admissible graph with two levels:



If we assume here that $R = \mathfrak{R} \simeq \mathbb{C}$, then we have $R^0 = \{0\}$ while $R^{0'} = R$.

Notation 4.14 Now that we have defined the $A_{\Gamma, I, l}^\ell$ for all levels, we let

$$A_{\Gamma, I, l} = \prod_{\ell \in \mathbb{Z}^-} A_{\Gamma, I, l}^\ell.$$

We have a natural morphism of $A_{\Gamma, I, l} \hookrightarrow \overline{\mathcal{H}}_{g, n, P}$: the differential is nonzero only on the level 0 vertices and vanishes identically everywhere else. We will call $A_{\Gamma, I, l}$ the *boundary stratum of type (g, Z, P, R) associated to (Γ, I, l)* .

Remark 4.15 The stratum $A_{\Gamma, I, l}$ is constructed from an admissible graph (Γ, I, l) of type (g, n, P) , a space of residues $R \subset \mathfrak{R}$ and q vectors of zeros Z . However, for simplicity, R and Z do not explicitly appear in the notation.

Remark 4.16 If $R = \mathfrak{R}$, then the construction of the space of residues is the translation of the global residue condition of [2]. For every level $-\ell$ and every vertex v of level greater than $-\ell$ that does not contain a pole, the following condition holds: Let h_1, \dots, h_k denote the half-edges adjacent to v and part of an edge to a vertex of level $-\ell$. Then the sum of residues assigned to this set of half-edges is zero.

Our definition of the R^i is more complicated to state because we need to take into account any vector subspace R of \mathfrak{R} .

4.3 Stratification of $\bar{A}_{g,Z,P}^R$

Let $(g, Z, P, R \subset \mathfrak{R})$ be a quadruple satisfying Assumption 3.1.

Lemma 4.17 *Let (Γ, I, l) be an admissible graph of type (g, Z, P, R) . The locus $A_{\Gamma,I,l}$ lies in the closure of $A_{g,Z,P}^R$. Conversely, if y is a point of $\bar{A}_{g,Z,P}^R$ then there exists an exterior completion Z' of Z and an admissible graph (Γ, I, l) of type (g, Z', P, R) such that y lies in $\pi(A_{\Gamma,I,l})$, where $\pi: A_{g,Z',P}^R \rightarrow A_{g,Z,P}^R$ is the forgetful map of the marked zeros that are not accounted for by Z .*

Remark 4.18 The set of admissible and realizable graphs (see Definition 4.9) is finite. Further, if (Γ, I, l) is an admissible graph, then the locus $A_{\Gamma,I,l}$ is empty if (Γ, I, l) is not realizable. Thus, Lemma 4.17 asserts that $\bar{A}_{g,Z,P}^R$ is stratified by finitely many strata corresponding to admissible graphs.

Before proving it we will introduce the incidence variety compactification of [2].

Notation 4.19 We suppose that $2g_j - 2 + n_j + m_j > 0$ for all $1 \leq j \leq q$. Then we denote by $K\bar{\mathcal{M}}_{g,n}(P)$ the vector bundle

$$R^0 \pi_* \left(\omega \left(\sum_{j=1}^q \sum_{i=1}^{m_j} p_{j,i} \sigma_{j,n_j+i} \right) \right),$$

where $\pi: \bar{\mathcal{C}}_{g,n,m} \rightarrow \bar{\mathcal{M}}_{g,n,m}$ is the forgetful map, ω is the relative cotangent bundle and the $\sigma_{j,i}$ are the sections of the universal curve (this generalizes Notation 2.19 to the disconnected case).

As in Section 2, there exists a natural morphism of cones

$$\text{stab}: \bar{\mathcal{H}}_{g,n,P} \rightarrow K\bar{\mathcal{M}}_{g,n}(P).$$

4.3.1 The image of $\bar{A}_{g,n,P}^R$ under the morphism stab

Definition 4.20 We denote by $\Omega\mathcal{M}_g^{\text{inc}}(Z, P)^R \subset K\bar{\mathcal{M}}_{g,n}(P)$ the image of $A_{g,n,P}^R$ under the morphism stab . The *incidence variety* for the tuple (g, Z, P, R) is the closure of $\Omega\mathcal{M}_g^{\text{inc}}(Z, P)^R$ in $K\bar{\mathcal{M}}_{g,n}(P)$.

The morphism stab induces a map from $A_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ to $\overline{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$. We will use the same notation for the morphism stab and its restriction

$$\text{stab}: \overline{A}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R \rightarrow \overline{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R.$$

Proposition 4.21 *If \mathbf{Z} is complete, then the map $\text{stab}: \overline{A}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R \rightarrow \overline{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$ is an isomorphism.*

Remark 4.22 Beware that this statement is valid only under the hypothesis that \mathbf{Z} is complete. Otherwise, the map stab may have fibers of positive dimension and/or may not be surjective.

Proof In Section 2 we proved that the following square is cartesian:

$$\begin{array}{ccc} \overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}} & \xrightarrow{\Phi_{j,i}} & \bigoplus_{\substack{1 \leq j \leq q \\ 1 \leq i \leq m_j}} \mathcal{P}_{j,n_j+i} \\ \downarrow & & \downarrow \\ K\overline{\mathcal{M}}_{\mathbf{g},\mathbf{n}}(\mathbf{P}) & \xrightarrow{\text{proj}_{j,i}} & \bigoplus_{\substack{1 \leq j \leq q \\ 1 \leq i \leq m_j}} J_{j,n_j+i} \end{array}$$

where \mathcal{P}_{j,n_j+i} is the cone of principal parts of order $p_{j,i}$ at the i^{th} marked point of the j^{th} connected component and J_{j,n_j+i} is the vector bundle of polar jets of order $p_{j,i}$. We recall that we have defined the spaces

$$\begin{aligned} \tilde{\mathcal{P}}_{j,n_j+i} &= (\mathcal{P}_{j,n_j+i} \setminus \mathcal{A}_{j,n_j+i}) \cup \text{the zero section,} \\ \tilde{J}_{j,n_j+i} &= (J_{j,n_j+i} \setminus \{\text{leading term} = 0\}) \cup \text{the zero section.} \end{aligned}$$

We have seen that the map $\Phi_{j,i}$ maps \mathcal{P}_{n_j+i} to \tilde{J}_{n_j+i} and that the restriction of $\phi_{i,j}$ to $\tilde{\mathcal{P}}_{j,n_j+i} \rightarrow \tilde{J}_{j,n_j+i}$ is an isomorphism (see Lemma 2.17). Thus, the morphism $\overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}} \rightarrow K\overline{\mathcal{M}}_{\mathbf{g},\mathbf{n}}(\mathbf{P})$ is an isomorphism from the preimage of $\bigoplus \tilde{\mathcal{P}}_{j,n_j+i}$ to the preimage of $\bigoplus \tilde{J}_{j,n_j+i}$.

The spaces $\overline{A}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ and $\overline{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$ are defined as the Zariski closures of open subspaces of $\overline{\mathcal{H}}_{\mathbf{g},\mathbf{n},\mathbf{P}}$ and $K\overline{\mathcal{M}}_{\mathbf{g},\mathbf{n}}(\mathbf{P})$. Therefore, we will prove that for all $1 \leq j \leq i$ and $1 \leq i \leq m_j$, the image of $\overline{A}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ (resp. $\overline{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$) under $\Phi_{j,i}$ (resp. $\text{proj}_{j,i}$) is included in $\tilde{\mathcal{P}}_{j,n_j+i}$ (resp. \tilde{J}_{j,n_j+i}) to deduce the proposition.

Let us consider a differential (C, α) in $\overline{A}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ and one of the marked points x_{j,n_j+i} corresponding to a pole. There are two possibilities:

- The point x_{j,n_j+i} belongs to a stable irreducible component of level 0, in which case the principal part belongs to $\mathcal{P}_{n_j+i} \setminus \mathcal{A}_{n_j+i}$.
- The point x_{j,n_j+i} belongs to an unstable rational component. In this case the differential restricted to this rational component is necessarily given by $dw/w^{p_{j,i}}$ (the marked point is at 0 and the node at ∞). Indeed, this follows from the assumption that \mathbf{Z} is complete: suppose that α has a zero outside the node; then let $B \rightarrow \bar{A}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ be a irreducible family of differentials with a special point $b_0 \in B$ whose image is the class $[(C, \alpha)]$ and the image of $B \setminus \{b_0\}$ lies in $A_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$. Then there exists a neighborhood U of b_0 such that the differential parametrized by U has an unmarked zero (this follows from Lemma 3.4). This is contradictory with the assumption that \mathbf{Z} is complete (all zeros of differentials in $A_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ are at marked points). Therefore, the principal part is equal to 0.

Therefore, the image of $\bar{A}_{\mathbf{g},\mathbf{n},\mathbf{P}}^R$ under $\Phi_{j,i}$ is included in $\tilde{\mathcal{P}}^{j,n_j+i}$. Now, let us consider a differential in $\bar{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$, and one of the marked points x_{j,n_j+i} corresponding to a pole. Once again, there are two possibilities:

- The point x_{j,n_j+i} belongs to an irreducible component of level 0. In this case the differential has a pole of order exactly $p_{j,i}$ at this marked point and the jet at x_{j,n_j+i} is in $\tilde{\mathcal{J}}_{j,n_j+i}$.
- The point x_{j,n_j+i} belongs to an irreducible component of level $-\ell < 0$. Then the differential vanishes identically on this component and the jet at x_{j,n_j+i} is 0.

Therefore, the image of $\bar{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$ under $\text{proj}_{j,i}$ is included in $\tilde{\mathcal{J}}_{j,n_j+i}$. This completes the proof. □

4.3.2 The image of the $A_{\Gamma,I,l}$ under the morphism stab To complete the description of the map stab we describe the image of the strata defined by admissible graphs.

Notation 4.23 Suppose that \mathbf{Z} is complete and (Γ, I, l) is a realizable stable twisted graph. Let (Γ', I', l') be the corresponding admissible graph. We denote by $\Omega\mathcal{M}_{\Gamma',I',l'}^{\text{inc}}$ the locus $\text{stab}(A_{\Gamma,I,l}) \subset K\bar{\mathcal{M}}_{\mathbf{g},\mathbf{n}}(\mathbf{P})$.

4.3.3 Stratification of $\bar{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$ Recall the main result of [2]:

Lemma 4.24 [2, Theorem 1.3] *Suppose that \mathbf{Z} is complete and that the triple (g_j, n_j, P_j) is stable for all $1 \leq j \leq q$. Let (Γ, I, l) be a stable graph. The locus*

$\Omega\mathcal{M}_{\Gamma,I,l}^{\text{inc}}$ lies in the closure of $\Omega\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$. Conversely, the space $\overline{\Omega}\mathcal{M}_{\mathbf{g}}^{\text{inc}}(\mathbf{Z}, \mathbf{P})^R$ is the union of the $\Omega\mathcal{M}_{\Gamma,I,l}^{\text{inc}}$ for all stable graphs (Γ, I, l) .

Remark 4.25 The statement here is slightly more general than [2, Theorem 1.3]. Indeed, it takes into account possibly disconnected bases and general choices of vector subspace $R \subset \mathfrak{R}$. However, all arguments in the proof of [2] can be adapted mutatis mutandis to get the general statement above.

Proof of Lemma 4.17 Suppose that \mathbf{Z} is complete and that the triple (g_j, n_j, P_j) is stable for all $1 \leq j \leq q$. Then, using Lemma 4.24 and Proposition 4.21 we automatically get

$$\overline{A}_{\mathbf{g},\mathbf{Z},\mathbf{P}}^R = \bigcup A_{\Gamma,I,l},$$

where the union is taken over all admissible graphs. Therefore, we only need to prove that the statement of Lemma 4.17 is still valid if we allow unstable base curves and noncomplete lists of vectors \mathbf{Z} .

Unstable basis We assume that \mathbf{Z} is complete but we no longer impose that the base curves are stable. Then, on a rational component with two points, the only possible configuration is $P = (p)$ and $Z = (p - 2)$. This is a closed point in $\overline{\mathcal{H}}_{0,1,(p)}$. Thus, the statement of Lemma 4.17 is still valid if we consider an unstable basis.

Noncomplete \mathbf{Z} We no longer impose that \mathbf{Z} is complete. The space $A_{\mathbf{g},\mathbf{Z},\mathbf{P}}^R$ is the union of the $\pi(A_{\mathbf{g},\mathbf{Z}',\mathbf{P}}^R)$ for all exterior completions \mathbf{Z}' of \mathbf{Z} (π being the forgetful map of the zeros which are accounted for by \mathbf{Z}). Therefore, we have

$$\overline{A}_{\mathbf{g},\mathbf{Z},\mathbf{P}}^R = \bigcup \pi(\overline{A}_{\mathbf{g},\mathbf{Z}',\mathbf{P}}^R) = \bigcup \pi(A_{\Gamma,I,l}),$$

where the last union is over all possible completions and admissible graphs. □

4.4 Description of boundary divisors

Let $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R \subset \mathfrak{R})$ be a quadruple satisfying Assumption 3.1. In the proof of the main theorem, we will be interested in the vanishing loci of sections of certain line bundles over $\overline{A}_{\mathbf{g},\mathbf{Z},\mathbf{P}}^R$. That is why we need to understand the boundary divisors of $\overline{A}_{\mathbf{g},\mathbf{Z},\mathbf{P}}^R$. The purpose of this section is to determine the set of admissible graphs which are associated to strata of codimension 1, ie to divisors.

4.4.1 Bicolored graphs

Lemma 4.26 Let (Γ, I, l) be an admissible graph. The codimension of $\overline{A}_{\Gamma,I,l}$ in $\overline{A}_{\mathbf{g},\mathbf{Z},\mathbf{P}}^R$ is greater than or equal to the depth of the level structure l .

Proof Let (Γ, I, l) be an admissible graph of depth d . Let (Γ', I', l') be the admissible graph obtained by merging the levels 0 and -1 . The locus $A_{\Gamma, I, l}$ lies in the closure of $A_{\Gamma', I', l'}$. Indeed, this follows from Lemma 4.17 applied to the stratum $A_{\Gamma', I', l'}^0$: the subgraph of (Γ, I, l) obtained by keeping only vertices of levels 0 and -1 determines a boundary stratum of $A_{\Gamma', I', l'}^0$. Thus, $A_{\Gamma, I, l}$ is of dimension at most $\dim(A_{\Gamma, I, l}) - 1$. Therefore, every time we merge two levels we decrease the codimension at least by 1. \square

Lemma 4.27 *Let (Γ, I, l) be an admissible graph of depth 1. The codimension of $\bar{A}_{\Gamma, I, l}$ in $\bar{A}_{g, Z, P}^R$ is greater than the number of horizontal edges.*

Proof We can independently merge vertices along horizontal edges (see ‘‘classical plumbing’’ in [2]). At every merging, we decrease the codimension by at least 1. \square

It follows from Lemmas 4.26 and 4.27 that a nontrivial admissible graph corresponding to a divisor of $A_{g, Z, P}^R$ is necessarily of depth at most 1. Moreover, if it is of depth 1 then it has no horizontal edges.

We recall from Section 4.2 that the boundary stratum associated to a graph of depth 1 is equal to $p(A_{g_1, Z_1, P_1}^{R^1}) \times A_{g_0, Z_0, P_0}^{R^0}$, where p is the map from $A_{g_1, Z_1, P_1}^{R^1}$ to the moduli space of curves $\bar{\mathcal{M}}_{g_1, n_1, m_1}$.

Notation 4.28 We denote by $\text{Bic}(g, Z, P, R)$ the set of realizable and admissible graphs with two levels and no horizontal edges. We will call such graphs *bicolored graphs*.

We say that a bicolored graph (Γ, I, l) satisfies condition $(\star\star)$ if (g_1, Z_1, P_1, R^1) satisfies the condition $(\star\star)$ (see Notation 3.23). We denote by $\text{Div}(g, Z, P, R)$ the set of bicolored graphs satisfying condition $(\star\star)$.

Remark 4.29 Elements of $\text{Bic}(g, Z, P, R)$ are twisted graphs with level structures. However, the level structure of a bicolored graph is completely determined by the twists. This is why we will denote by (Γ, I) the elements of $\text{Bic}(g, Z, P, R)$.

Proposition 4.30 *Let (Γ, I) be a bicolored graph in $\text{Bic}(g, Z, P, R)$. The locus $A_{\Gamma, I}$ is of codimension 1 in $\bar{A}_{g, Z, P}^R$ if and only if (Γ, I) belongs to $\text{Div}(g, Z, P, R)$.*

Proof Let $(\Gamma, I) \in \text{Bic}(g, Z, P, R)$. The proposition follows easily from the equation

$$(11) \quad \dim(A_{g_0, Z_0, P_0}^{R^0}) + \dim(A_{g_1, Z_1, P_1}^{R^1}) = \dim(A_{g, Z, P}^R).$$

Indeed, $A_{\Gamma, I}$ is of codimension 1 in $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$ if and only if $\dim(p(A_{\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1})) = \dim(\mathbb{P} A_{\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1})$, ie if and only if $(\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1, R^1)$ satisfy condition $(\star\star)$ (see Proposition 3.22).

Let us prove (11). We assume first that \mathbf{Z} is complete for (\mathbf{g}, \mathbf{P}) and the dimension of $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$ is given by $(\sum_{j=1}^q (2g_j - 1 + n_j)) + \dim(R)$. Therefore, we have

$$\begin{aligned} & \dim(A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R) - \dim(A_{\mathbf{g}_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0}) - \dim(A_{\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1}) \\ &= \left(\sum_{j=1}^q (2g_j - 1 + n_j) \right) + \dim(R) - \dim(R^1 \oplus R^0) \\ & \qquad \qquad \qquad - \left(\sum_{v \in V^0} (2g_v - 1 + n_v) + \sum_{v \in V^1} (2g_v - 1 + n_v) \right) \\ &= 2h^1(\Gamma) - q + \text{Card}(V(\Gamma)) - \text{Card}(E(\Gamma)) + \dim(R) - \dim(R^1 \oplus R^0) \\ &= h^1(\Gamma) + \dim(R) - \dim(R^1 \oplus R^0). \end{aligned}$$

Thus, we will prove that $\dim(R^1 \oplus R^0) = \dim(R) + h^1(\Gamma)$.

Let us recall the construction of R^0 and R^1 . In the absence of horizontal edges, we consider the vector space $\mathfrak{R} \oplus \mathfrak{R}^1$ and the projection $\text{proj}: \mathfrak{R} \oplus \mathfrak{R}^1 \rightarrow \mathfrak{R}^1$ along \mathfrak{R} . We also consider the vector subspace $\tilde{R} \subset R \oplus \mathfrak{R}^1 \subset \mathfrak{R} \oplus \mathfrak{R}^1$ defined by the linear relations

$$\sum_{h \in H(\Gamma), h \mapsto v} r_h = 0$$

for all vertices v of level 0 (the sum is over all residues at half-edges adjacent to v). We defined $R^0 = \ker(\text{proj}) \cap \tilde{R}$ and $R^1 = \text{proj}(\tilde{R})$. Thus, $\dim(R^0) + \dim(R^1) = \dim(\tilde{R})$. Therefore, we need to prove that $\dim(\tilde{R}) = \dim(R) + h^1(\Gamma)$.

To prove this equality we use the graph Γ' obtained from Γ by adding one vertex per marked pole and one edge between this vertex and the vertex that carries the marked pole. We consider the spaces $C_0 = \mathbb{C}^{V(\Gamma')}$ and $C_1 = \mathbb{C}^{E(\Gamma')}$. We have the chain complex $d: C_1 \rightarrow C_0$.

The space R is a subspace of C_0 : indeed, the space \mathfrak{R} is a subspace of the subspace of \mathfrak{R} spanned by the vertices in $V(\Gamma') \setminus V(\Gamma)$. The space \tilde{R} is naturally identified with $d^{-1}(R)$. Therefore, $\dim(\tilde{R}) = \dim(R) + \dim(\ker(d)) = \dim(R) + h^1(\Gamma)$.

If Z is not complete, then we consider Z_m the maximal completion of Z . Then (11) still holds by

$$\begin{aligned} \dim(A_{g,Z,P}^R) &= \dim(A_{g,Z_m,P}^R) = \dim(A_{g_0,Z_{0,m},P_0}^{R^0}) + \dim(A_{g_1,Z_{1,m},P_1}^{R^1}) \\ &= \dim(A_{g_0,Z_0,P_0}^{R^0}) + \dim(A_{g_1,Z_1,P_1}^{R^1}). \quad \square \end{aligned}$$

4.4.2 Classification of boundary divisors

Notation 4.31 Let $1 \leq j \leq q$ and $1 \leq i \leq \ell(Z_j)$. We denote by $Z_{j,i}$ the list of vectors obtained from Z by increasing the i^{th} coordinate of Z_j by one.

Proposition 4.32 Let Z' be a completion of Z and let (Γ, I, l) be an admissible graph such that $D = \pi(\bar{A}_{\Gamma,I})$ is a divisor of $\bar{A}_{g,Z,P}^R$ (where π is the forgetful map of the points), then D is necessarily of one of four kinds:

- (1) the stratum $\bar{A}_{\Gamma,I}$ for $(\Gamma, I) \in \text{Div}(g, Z, P, R)$;
- (2) the locus $\bar{A}_{g,Z_{j,i},P}^R$ for some label (j, i) corresponding to a marked point which is not a pole;
- (3) the locus $\bar{A}_{\Gamma,I,l}$ for a P -admissible graph of depth 0 with a unique horizontal edge;
- (4) the locus $\bar{A}_{g,Z,P}^{R'}$ for the vector subspace $R' \subset R$ defined by the condition $\text{res}_{x_{j,n_j+i}} = 0$ for a choice of j and i such that the point x_{j,n_j+i} corresponds to a pole of order at most -1 .

Proof Let Z' be a completion of Z . If Z' is not the maximal completion then $\dim(A_{g,Z',P}^R) < \dim(A_{g,Z,P}^R)$. The only possible admissible graph is the trivial and we obtain a divisor of type 2.

We suppose now that $Z' = Z_m$; then (Γ, I, l) is of depth less than or equal to 1 by Lemma 4.26. If (Γ, I, l) is of depth 0 then (Γ, I, l) has at most one horizontal edge (type 3). If (Γ, I, l) is of depth 1 then either all or none of the edges of (Γ, I, l) are contracted under the forgetful map of the marked points which are not accounted for by Z (otherwise this graph does not satisfy condition $(\star\star)$). If none of the edges are contracted, then D is a divisor of type 1. If all edges are contracted then we get a divisor of type 2 or 4 (depending on whether there is a leg corresponding to a pole of order 1 on a level -1 vertex or not). □

Proposition 4.33 *Let D_1 and D_2 be two divisors obtained from an admissible graph as in Proposition 4.32. Then D_1 and D_2 have no common irreducible components.*

Proof The divisors D_1 and D_2 can be of one of the four types described in Proposition 4.32. We will prove this proposition by considering every possible case:

Type 1 Let (Γ, I) and (Γ', I') in $\text{Div}(\mathfrak{g}, \mathbf{Z}, \mathbf{P}, R)$ be such that $A_{\Gamma, I}$ and $A_{\Gamma', I'}$ have a common irreducible component D . The component D determines a semistable graph by taking the dual graph of a any point of $D \cap A_{\Gamma, I}$; therefore, $\Gamma = \Gamma'$. Moreover, the vertices of Γ with identically zero differentials are the vertices of level -1 . Therefore, the level structure (or more precisely the signs of the twists) are the same for (Γ, I) and (Γ', I') . Now the twist at an edge is determined by the vanishing order of the differential at the corresponding node on the component of level 0 for any point in $D \cap A_{\Gamma, I}$. Therefore, $(\Gamma, I) = (\Gamma', I')$. Thus, divisors of type 1 have no common irreducible components.

Types 2 and 4 The underlying generic curve of the divisors of type 2 or 4 is a curve without singularities; therefore, divisors of type 2 or 4 do not intersect divisors of type 1 or type 3. Now the differentials of the generic differentials of two divisors of type 2 have different vanishing order at two of the marked points (either a marked zero or a marked pole of order -1).

Type 3 Two divisors of type 3 are distinguished by the topological types of a generic curve. Further, a divisor of type 3 is distinguished from a divisor of type 1 because none of the components carries a vanishing differential in a divisor of type 3. \square

5 Computation of classes of strata

Let $(\mathfrak{g}, \mathbf{Z}, \mathbf{P}, R \subset \mathfrak{A})$ be a quadruple satisfying Assumption 3.1. The purpose of this section is to prove the following generalization of Theorem 1 stated in the introduction.

Theorem 4 *Let $(\mathfrak{g}, \mathbf{Z}, \mathbf{P}, R)$ be a quadruple satisfying Assumption 3.1. The Poincaré-dual class of $\mathbb{P}\bar{A}_{\mathfrak{g}, \mathbf{Z}, \mathbf{P}}^R \in H^*(\mathbb{P}\bar{\mathcal{H}}_{\mathfrak{g}, n, \mathbf{P}}, \mathbb{Q})$ is tautological (in the sense of Definition 2.36) and is explicitly computable.*

Theorems 1, 2 and 3 will be deduced from Theorem 4 at the end of the section. The most technical result involved in the proof of Theorem 4 is the induction formula for the classes $[\mathbb{P}\bar{A}_{\mathfrak{g}, \mathbf{Z}, \mathbf{P}}^R]$ (see Section 5.2).

5.1 A meromorphic function on $\bar{A}_{g,Z,P}^R$

Let $1 \leq j \leq q$ and $1 \leq i \leq n_j$. Let $k_{i,j}$ be the i^{th} entry of Z_j . We consider the line bundle

$$\mathcal{O}(-1) \otimes \mathcal{L}_{j,i}^{k_{j,i}+1} \Big|_{A_{g,P,Z}^R} \simeq \text{Hom}(\mathcal{O}(-1), \mathcal{L}_{j,i}^{k_{j,i}+1}) \Big|_{A_{g,P,Z}^R},$$

where $\mathcal{L}_{j,i}$ is the cotangent line bundle to the i^{th} marked point of the j^{th} connected component. Let $s_{j,i}$ be the holomorphic section of the line bundle

$$\text{Hom}(\mathcal{O}(-1), \mathcal{L}_{j,i}^{k_{j,i}+1}) \Big|_{A_{g,P,Z}^R}$$

that maps a differential to its $(k_{j,i}+1)^{\text{st}}$ -order term at the i^{th} marked point of the j^{th} connected component.

Lemma 5.1 *The section $s_{j,i}$ vanishes with multiplicity 1 along $\mathbb{P}\bar{A}_{g,Z_j,i,P}^R$.*

Proof Let $y_0 = (C, \alpha, Z(y_0) \cup P(y_0))$ be a point of $A_{g,Z_j,i,P}^R$, where we denote by $P(y_0) \subset C$ be the set of poles of α and $Z(y_0) \subset C$ be the set of marked zeros of C . Further, we denote by $Z'(y_0) \subset C$ be the set of nonmarked zeros.

Let $W/\text{Aut}(y_0)$ be a contractible neighborhood of y_0 . Up to a choice of a smaller W , in the proof of Lemma 3.7, we constructed the three maps

$$\begin{aligned} \Phi^1: W &\rightarrow H^1(C \setminus P(y_0), Z(y_0) \cup Z'(y_0), \mathbb{C}), \\ \Phi^{2,x}: W &\rightarrow \mathcal{Z}^{k_x} \quad \text{for all } x \in Z(y_0), \\ \Phi^{3,x}: W &\rightarrow \tilde{\mathcal{Z}}^{k_x-1} \quad \text{for all } x \in Z'(y_0), \end{aligned}$$

where k_x is the order of α at x (be it a marked or nonmarked zero) and where \mathcal{Z}^k is a domain in \mathbb{C}^k containing 0. These maps are not uniquely determined; however, we saw in the proof of Lemma 3.7 that the map $\Phi_1 \times \prod_{x \in Z(y_0)} \Phi^{2,x} \times \prod_{x \in Z'(y_0)} \Phi^{3,x}$ is a local biholomorphism.

Now we consider the marked point $x_{j,i}$. We let

$$\Phi^{(j,i)} = \Phi^{2,x_{j,i}} \quad \text{and} \quad \tilde{\Phi}^{(j,i)} = \prod_{x \in Z_{y_0} \setminus \{x_{j,i}\}} \Phi^{2,x}.$$

We recall that the map $\Phi^{(j,i)}$ is defined as follows: for all points s in a neighborhood of y_0 , the differential representing y is given in neighborhood of the marked point $x_{j,i}(y)$ by

$$\alpha = (z^{k_{j,i}+1} + a_{k_{j,i}} z^{k_{j,i}} + \dots + a_0) dz$$

(the marked point being at $z = 0$); then we define $\Phi_2(y) = (a_0, \dots, a_{k_{j,i}}) \in \mathbb{Z}^{k_{j,i}+1}$ (this definition is unique up to choice of a $(k_{j,i} + 2)^{\text{nd}}$ root of unity).

Then, with this parametrization we have

$$W \cap A_{\mathbf{g}, \mathbf{Z}_{j,i}, \mathbf{P}}^R = (\Phi^{j,i} \times \tilde{\Phi}^{j,i})^{-1} \left(\prod_{x \in Z(y_0)} \{0\} \right),$$

$$W \cap A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R = (\Phi^{j,i} \times \tilde{\Phi}^{j,i})^{-1} \left((0, \dots, 0, \epsilon) \times \prod_{x \in Z(y_0) \setminus \{x_{j,i}\}} \{0\} \right).$$

In other words, the coordinate $a_{k_{j,i}}$ is a transverse parameter to the divisor $A_{\mathbf{g}, \mathbf{Z}_{j,i}, \mathbf{P}}^R$ in $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$. We obviously have $s_{j,i}(y) = a_{k_{j,i}}$. Therefore, the vanishing order of $s_{j,i}$ along $\mathbb{P}A_{\mathbf{g}, \mathbf{Z}_{j,i}, \mathbf{P}}^R$ is equal to 1. □

Notation 5.2 We denote by $\text{Bic}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i} \subset \text{Bic}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)$ the subset of bicolored graphs such that the i^{th} marked point of the j^{th} connected component belongs to a level -1 vertex and we denote by $\text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}$ the intersection of $\text{Bic}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}$ and $\text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)$.

Lemma 5.3 *The divisors contained in the vanishing locus of $s_{j,i}$ are exactly the divisors corresponding to admissible graphs in $\text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}$ and the divisor $\mathbb{P}\bar{A}_{\mathbf{g}, \mathbf{Z}_{j,i}, \mathbf{P}}^R$. No two of these divisors have a common irreducible component.*

Proof This is a consequence of Propositions 4.32 and 4.33. □

5.2 Induction formula

Let $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R \subset \mathfrak{R})$ be a quadruple satisfying Assumption 3.1. Let $1 \leq j \leq q$ and $1 \leq i \leq n_j$. We recall that we denote by $\mathbf{Z}_{j,i}$ the list of vectors obtained from \mathbf{Z} by increasing $k_{j,i}$ by 1. Further, as in the previous section, we denote by $\mathcal{L}_{j,i}$ the cotangent line to the i^{th} marked point on the j^{th} connected component of the curve and let $\psi_{j,i} = c_1(\mathcal{L}_{j,i}) \in H^2(\mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}, \mathbb{Q})$.

5.2.1 Multiplicity of (Γ, I)

Definition 5.4 Let $(\Gamma, I) \in \text{Bic}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)$. The *multiplicity* of (Γ, I) is defined as

$$m(I) = \prod_{h \rightarrow V^0} I(h),$$

where the product runs over the half-edges which are not legs, pointing to vertices of level 0. The *least common multiple* and the *group of roots* of the twist are

$$L(I) = \text{LCM}(\{I(h)\}_{h \rightarrow V^0}), \quad G_I = \left(\prod_{h \rightarrow V^0} \mathbb{Z}_{I(h)} \right) / \mathbb{Z}_{L(I)}.$$

5.2.2 Locus of generic points Let $(\Gamma, I) \in \text{Div}(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$. We recall that

$$A_{\Gamma, I} = p(A_{\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1}) \times A_{\mathbf{g}_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0},$$

where $p: A_{\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1} \rightarrow \overline{\mathcal{M}}_{\mathbf{g}_1, \mathbf{n}_1, \mathbf{m}_1}$ is the forgetful map. The condition $(\star\star)$ ensures that there exists an open dense locus $A_1^{\text{gen}} \subset A_{\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1}$ such that the map $p: A_1^{\text{gen}} \rightarrow p(A_1^{\text{gen}})$ has fibers of dimension 1 (see Proposition 3.24). Then we set

$$A_{\Gamma, I}^{\text{gen}} = A_1^{\text{gen}} \times A_{\mathbf{g}_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0}.$$

This open locus of generic points will be important for us because the map

$$p: A_1^{\text{gen}} \times A_{\mathbf{g}_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0} \rightarrow A_{\Gamma, I}^{\text{gen}} = p(A_1^{\text{gen}}) \times A_{\mathbf{g}_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0}$$

is a line bundle minus the zero section.

Notation 5.5 We denote by $p: \mathcal{N}_{\Gamma, I} \rightarrow A_{\Gamma, I}^{\text{gen}}$ this line bundle.

5.2.3 Induction formula We finally have all elements to state the main result of the paper.

Theorem 5 In $H^*(\mathbb{P}\overline{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}, \mathbb{Q})$ we have

$$(12) \quad [\mathbb{P}\overline{A}_{\mathbf{g}, \mathbf{Z}_{j,i}, \mathbf{P}}^R] = (\xi + (k_{j,i} + 1)\psi_{j,i}) \cdot [\mathbb{P}\overline{A}_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R] - \sum_{(\Gamma, I) \in \text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}} m(I) [\mathbb{P}\overline{A}_{\Gamma, I}]$$

if $2g_j - 2 + n_j + m_j > 0$, or

$$(13) \quad [\mathbb{P}\overline{A}_{\mathbf{g}, \mathbf{Z}_{j,1}, \mathbf{P}}^R] = \frac{p-k-2}{p-1} \xi \cdot [\mathbb{P}\overline{A}_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R]$$

if $g_j = 0$, $Z_j = (k)$ and $P_j = (p)$.

Proof of (12) As in Section 5.1, we consider the line bundle $\text{Hom}(\mathcal{O}(-1), \mathcal{L}_{j,i}^{k_{j,i}+1}) \rightarrow \mathbb{P}\overline{A}_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$. Its first Chern class is equal to $\xi + (k_{j,i} + 1)\psi_{j,i}$. Moreover, this line bundle has a global section $s_{j,i}$ which maps a differential to its $(k_{j,i} + 1)^{\text{st}}$ -order term at the

marked point (j, i) . In Lemma 5.1 we showed that $s_{j,i}$ vanishes along $\mathbb{P}\bar{A}_{\mathbf{g}, \mathbf{Z}, i, \mathbf{P}}^R$ with multiplicity 1. In Lemma 5.3 we showed that the remaining vanishing loci of $s_{j,i}$ are supported on the $\mathbb{P}\bar{A}_{\Gamma, I}$ for (Γ, I) of $\text{Div}(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)_{j,i}$. Therefore, we deduce that

$$(\xi + (k_{j,i} + 1)\psi_{j,i}) \cdot [\mathbb{P}\bar{A}_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R] = [\mathbb{P}\bar{A}_{\mathbf{g}, \mathbf{Z}, i, \mathbf{P}}^R] + \mathbb{Z},$$

where \mathbb{Z} is a cycle supported on the union of $\mathbb{P}\bar{A}_{\Gamma, I}$ for $(\Gamma, I) \in \text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}$.

Now we claim that the vanishing order of $s_{j,i}$ along the locus $\mathbb{P}\bar{A}_{\Gamma, I}$ is equal to $m(I)$ (see Definition 5.4). Lemma 5.6 below implies this statement and thus equation (12). \square

Lemma 5.6 *Let (Γ, I) be a divisor graph in $\text{Div}(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)_{j,i}$. Let $y_0 \in \mathbb{P}A_{\Gamma, I}^{\text{gen}}$. Let Δ be an open disk in \mathbb{C} containing 0 and parametrized by ϵ . There exists an open neighborhood U of y_0 in $\mathbb{P}A_{\Gamma, I}^{\text{gen}}$ together with a map $\iota: U \times \Delta \times G_I \rightarrow \mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{m}, \mathbf{P}}$ satisfying:*

- *The restriction $\iota|_{U \times 0 \times g}$ is the identity on U for all $g \in G_I$.*
- *The image of the restriction $\iota|_{\epsilon \neq 0}$ lies in the open stratum $\mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$.*
- *For all $g \in G_I$, the section $s_{j,i}$ restricted to $\iota(U \times \Delta \times g)$ vanishes along $\iota(U \times 0 \times g)$ with multiplicity $L(I)$.*
- *The map $\iota: U \times \Delta \times G_I \rightarrow \mathbb{P}\bar{A}_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$ is a degree 1 parametrization of a neighborhood of U in $\mathbb{P}\bar{A}_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$.*

The proof of Theorem 5 immediately follows from Lemma 5.6 because the vanishing order of $s_{j,i}$ along $\mathbb{P}\bar{A}_{\Gamma, I}$ is equal to

$$L(I) \cdot \text{Card}(G_I) = m(I).$$

Proof of Lemma 5.6 We prove the lemma in two steps: first we will prove the first three points of the lemma and then we will prove that ι is a parametrization of degree 1 of a neighborhood of U in $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$.

Proof of the first three points For the sake of clarity we will successively prove the first three points at three levels of generality: first for a divisor graph with one edge, then for divisor graph with $R^1 = \{0\}$ and finally in full generality.

Bicolored graph with one edge For the moment we place ourselves in the simplest case: (Γ, I) is an admissible graph with two vertices, one at level 0 and one at level -1 .

We suppose that there is only one edge with a twist given by $k > 0$. Let y_0 be a point of $\mathbb{P}A_{\Gamma,I}^{\text{gen}}$. Let U be an open neighborhood of y_0 in $\mathbb{P}A_{\Gamma,I}^{\text{gen}}$. A point y of U is given by

$$([C^0], [C^1], \bar{x}^0, \bar{x}^1, [\alpha^0]),$$

where C^0 and C^1 are the curves corresponding to the two vertices of the graph, \bar{x}^0 and \bar{x}^1 are their marked point sets, α^0 is a differential on the curve C^0 and $[\alpha^0]$ its equivalence class under the \mathbb{C}^* -action. More precisely, we denote by $\alpha^0(y)$ a nonvanishing section of the line bundle $\mathcal{O}(-1)$ over U . (Also recall that on C^1 the differential vanishes identically.)

The condition that $y \in A_{\Gamma,I}^{\text{gen}}$ implies that the curve C^1 carries a *unique* meromorphic differential α^1 with zeros and poles of prescribed multiplicities at the marked points, up to a scalar factor. Let $\alpha^1(y)$ be a nonvanishing section of the line bundle $\mathcal{N}_{\Gamma,I}$, ie a choice of the scalar factor for each point y .

At the neighborhood of the node, the curves C^1 and C^0 have standard coordinates z and w such that $\alpha^0 = d(z^k)$ and $\alpha^1 = d(1/w^k)$. The local coordinates z and w are unique up to multiplication by a k^{th} root of unity. We fix one such choice in a uniform way over U . We define a family of curves $C(y, \epsilon)$ over $U \times \Delta$ by smoothing the node between C^0 and C^1 via the equation $zw = \epsilon$, where ϵ is the coordinate on the disc Δ , and z and w are as above. The differentials α^0 and $\epsilon^k \alpha^1$ automatically glue together into a differential on $C(y, \epsilon)$.

The deformation that we have constructed does not depend on the choice of standard coordinates z and w . For instance, if we multiply z by a k^{th} root of unity ζ , the equation of the deformation becomes $zw = \zeta \epsilon$, which is isomorphic to the original deformation under a rotation of the disc Δ .

The section $s_{j,i}$ vanishes with multiplicity k along the locus defined by $\epsilon = 0$: indeed, we have explicitly

$$s_{j,i}(y, \epsilon) = \epsilon^k \cdot \alpha_1(y).$$

Bicolored graph (Γ, I) with $R^1 = \{0\}$ We suppose now that the space R^1 is trivial (residues at the nodes between vertices of levels 0 and -1 are equal to 0). A point y in U still determines

$$([C^0], [C^1], \bar{x}^0, \bar{x}^1, [\alpha^0], [\alpha^1]),$$

where α^0 and α^1 are sections of $\mathcal{O}(-1)$ and $\mathcal{N}_{\Gamma,I}$ as in the previous paragraph.

Let e be an edge of Γ . We denote by k_e the positive integer equal to $|I(h)|$ for any of the two half-edges of e . Let z_e and w_e be choices of standard coordinates in a neighborhood of the node corresponding to e , ie $\alpha^0 = d(z_e^{k_e})$ and $\alpha^1 = d(1/w_e^{k_e})$. This choice of standard coordinates being fixed for all edges, we choose, on top of that, a $(k_e)^{\text{th}}$ root of unity ζ_e for each edge e .

We define a family of curves $C(y, \epsilon)$ over $U \times \Delta$ by smoothing the node corresponding to an edge e of Γ via the equation $z_e w_e = (\zeta_e \epsilon)^{L(I)/k_e}$, where ϵ is the coordinate on the disc Δ . The differentials defined by α^0 and by $\epsilon^{L(I)} \alpha_1$ automatically glue together into a differential on $C(y, \epsilon)$.

A multiplication of ϵ by a $L(I)^{\text{th}}$ root of unity ζ gives an isomorphic deformation. Thus, two choices of roots $(\zeta_e)_{e \in \text{Edges}}$ and $(\zeta'_e)_{e \in \text{Edges}}$ give isomorphic deformations if $\zeta'_e = \zeta^{L(I)/k_e} \zeta_e$ for all edges. The vanishing multiplicity of $s_{j,i}$ along the locus defined by $\epsilon = 0$ is equal to $L(I)$.

General bicolored graph (Γ, I) We no longer impose restrictions on R^1 . We still define

$$([C^0], [C^1], \bar{x}^0, \bar{x}^1, \alpha^0, \alpha^1),$$

as above. Moreover, we define the section r by

$$r(y) = (r_e(y))_{e \in \text{Edges}},$$

where $r_e(y)$ is the residue of α_1 at the node of C^1 corresponding to the edge e . For every edge e , we fix a choice of standard coordinates of z_e and w_e in a neighborhood of the node corresponding to e , ie coordinates satisfying $\alpha^0 = d(z_e^{k_e})$ and $\alpha^1 = d(1/w_e^{k_e}) + r_e(y) dw_e/w_e$.

Using Proposition 3.10, we get a family of differentials $(\tilde{C}^0, \bar{x}^0, \tilde{\alpha}^0)$ parametrized by $U \times \Delta$ such that

- when $\epsilon = 0$, we have $(C^0, \bar{x}^0, \alpha^0) = (\tilde{C}^0, \bar{x}^0, \tilde{\alpha}^0)$;
- the zeros of the differential which are not at the marked points corresponding to nodes are of fixed orders;
- the differential $\tilde{\alpha}^0$ has at most simple poles at the nodes of \tilde{C}^0 and the residue at the node corresponding to the edge e is equal to $-\epsilon^{L(I)} r_e(y)$;
- the vector of residues at the poles of $\tilde{\alpha}^0$ lies in R ;
- for each node corresponding to an edge e with a twist k_e , the family of differentials defined by $U \times \Delta$ is a standard deformation of $d(z_e^{k_e})$ (see Definition 3.5).

We use the fact that the family parametrized by $U \times \Delta$ is a standard deformation of $d(z_e^{k_e})$ to apply Proposition 3.6. At each node e the differential $\tilde{\alpha}_0$ can be written in the form $d(z_e^{k_e}) - \epsilon^{L(I)} r(u) dz_e/z_e$ in any annulus contained in a neighborhood of the node. Therefore, we can still glue the two components together along this annulus with the identification $z_e w_e = \zeta_e \epsilon^{L(I)/k_e}$ for any choice of the $(k_e)^{\text{th}}$ root of unity ζ_e . The end of the proof is the same as for divisor graphs with trivial residue conditions.

Proof of the fourth point Now we will prove that the map $\iota: U \times \Delta \times G_I \rightarrow \mathbb{P} \bar{A}_{g, \mathbf{Z}, \mathbf{P}}^R$ is a degree 1 parametrization of a neighborhood of U in $\mathbb{P} \bar{A}_{g, \mathbf{Z}, \mathbf{P}}^R$.

First we prove that the image $\iota(U \times \Delta \times G_I)$ covers *entirely* a neighborhood of U in $A_{g, \mathbf{Z}, \mathbf{P}}^R$. Let $y_0 = (C = C_0 \cup C_1, \bar{x}_0, \bar{x}_1, \alpha_0)$ be a point in $A_{\Gamma, I}^{\text{gen}}$. Let $\tilde{\iota}: \Delta \rightarrow \bar{A}_{g, \mathbf{Z}, \mathbf{P}}$ be a family of differentials such that $\tilde{\iota}(0) = y_0$ and $\tilde{\iota}(\epsilon) \in A_{g, \mathbf{Z}, \mathbf{P}}^R$ for $\epsilon \neq 0$. We denote by $\pi: \mathcal{C} \rightarrow \Delta$ the induced family of curves and by α the induced family of differentials on the fibers of $\mathcal{C} \rightarrow \Delta$.

Let e be a node of C with a twist of order k_e . Let γ_e be a simple loop in the curve C_0 around the node e . Let W_e be a neighborhood of γ_e in \mathcal{C} such that $W_e \cap \pi^{-1}(\epsilon)$ is an annulus for any ϵ small enough. Now, the differential α_0 is given by $d(z_e^{k_e})$ in a standard coordinate. Thus, the differential $\alpha|_{\pi^{-1}(\epsilon)}$ is given by $d(z_e^{k_e}) + \phi(\epsilon, z_e) dz_e$ and we denote by $r_e(\epsilon)$ the integral of $\phi(\epsilon, z_e) dz_e$ along γ_e . We consider the differential $\alpha_e(\epsilon) = dz_e + \phi(\epsilon, z_e) dz_e - r_e(\epsilon) dz_e/z_e$. We fix a point p in the annulus $W_e \cap \pi^{-1}(\epsilon)$, the function $f: z \mapsto \left(\int_p^z \alpha_e\right)^{1/k_e}$ is uniquely determined for small values of ϵ . This determines a coordinate (that we will still denote z_e) such that $\alpha_0 = z_e^{k_e} dz_e - \varphi(\epsilon, z_e) dz_e/z_e$ with φ holomorphic and thus a standard deformation of α^0 . Proposition 3.6 implies that there exists a coordinate z_e on this annulus such that $\alpha|_{\pi^{-1}(\epsilon)} = d(z_e^{k_e}) + r_e(\epsilon) dz_e/z_e$.

We fix ϵ small enough that the coordinates z_e are defined for all edges e . We cut the curve $\pi^{-1}(\epsilon)$ along simple loops contained in W_e . This gives two (possibly disconnected) curves with boundary C_0^{open} and C_1^{open} . We “plug” the holes of C_0^{open} with disks parametrized by the coordinate z_e and the holes of C_1^{open} with disks with coordinate $1/z_e$. This determines two curves $C_0(\epsilon)$ and $C_1(\epsilon)$. On both C_0 and C_1 , the local chart used to “plug” the holes allow us to define differentials $\alpha_0(\epsilon)$ and $\alpha_1(\epsilon)$.

The differential $\alpha_1(\epsilon)$ has a pole of order $k_e + 1$ at $w_e = 0$; thus, $(C_1, \bar{x}_1, \alpha_1)(\epsilon)$ is an element of $A_{g_1, \mathbf{Z}_1, \mathbf{P}_1}^R$. Now, at the level 0, we use Corollary 3.11: in a neighborhood of y_0 we can apply the retraction η . The point $\eta((C_0, \bar{x}_0, \alpha_0)(\epsilon))$ is a point

of $A_{\mathbf{g}_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0}$. Therefore, we define

$$y(\epsilon) = (\eta(C_0, \bar{x}_0, \alpha_0), (C_1, \bar{x}_1, \alpha_1))(\epsilon) \in A_{\Gamma, I}^{\text{gen}}$$

For all ϵ in a neighborhood of 0, the point $\tilde{\iota}(\epsilon)$ lies in the deformation of $y(\epsilon)$ by the family ι restricted to $y(\epsilon) \times \Delta \times g$ for some $g \in G_I$ (in fact here $g = 1$ because of the choices of the parameters around y_0 that we have fixed).

To finish the proof of the fourth point, we need to prove that the parametrization is of degree 1. For this, we once again use the retraction η defined in Corollary 3.11. We have $\eta \circ \iota = \text{Id}_U$; thus, we only need to prove that for all $y \in U$, the family ι restricted to $y \times \Delta \times G_I$ is of degree 1. We consider this family in the moduli space of curves, ie let

$$\iota': \Delta \times G_I \rightarrow \bar{\mathcal{M}}_{\mathbf{g}, \mathbf{n}, \mathbf{m}}, \quad \epsilon \times g \mapsto p(\iota(y, \epsilon, G_I)).$$

This family is of degree 1. Indeed, the stack $\bar{\mathcal{M}}_{\Gamma}$ is regularly embedded in $\bar{\mathcal{M}}_{\mathbf{g}, \mathbf{n}, \mathbf{m}}$ and its normal bundle is the direct sum of the $T_h \otimes T_{h'}$ for all edges $e = (h, h')$ of Γ . Thus, the family ι' is given by the family

$$\iota': \Delta \times G_I \rightarrow \bigoplus_{(h, h') \in \text{Edges}} T_h \otimes T_{h'}, \quad (\epsilon, (\zeta_e)_{e \in \text{Edges}}) \mapsto (\zeta_e \in L(I)/k_e)_{e \in \text{Edges}},$$

which is of degree 1. □

Proof of formula (13) We have seen that the space of differentials on an unstable component is a weighted projective space parametrized by

$$[w^{p-1} + a_1 w^{p-2} + \dots + a_{p-2} w] \frac{dw}{w},$$

where the weight of a_j is $j/(p-1)$. The fact that the order of the point x is $k_{j,i}$ is equivalent to the vanishing of the terms $a_{p-2}, \dots, a_{p-k_{j,i}-3}$. Therefore, the class of $[\mathbb{P} \bar{A}_{\mathbf{g}, \mathbf{Z}_{j,i}, \mathbf{P}}^R]$ is the closure of the vanishing locus $a_{p-k_{j,i}-2}$. Moreover, we can easily check that $a_{p-k_{j,i}+1}^{p-1}$ is a global section of $\mathcal{O}(-1)^{p-k_{j,i}+1}$. □

5.3 Class of a boundary divisor

Let $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R \subset \mathfrak{R})$ be a quadruple satisfying Assumption 3.1. We want to compute the Poincaré-dual class of the locus associated to an element of $\text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)$.

5.3.1 Decomposition of the morphism $A_{\Gamma,I} \rightarrow \overline{\mathcal{H}}_{g,Z,P}$ Let (Γ, I) be an admissible graph in $\text{Bic}(g, P, Z, R)$ (this graph may be a divisor or not). We recall that the semistable graph Γ determines a stratum

$$\zeta_{\Gamma}^{\#}: \overline{\mathcal{H}}_{\Gamma} = \overline{\mathcal{H}}_{g_{\Gamma},n_{\Gamma},P_{\Gamma}}^{R_{\Gamma}} \rightarrow \overline{\mathcal{H}}_{g,n,P}$$

(see Section 2.7).

Remark 5.7 Beware that $\overline{\mathcal{H}}_{\Gamma}$ is purely determined by Γ (and not I). However, the twist I determines the components that are of level -1 and the P -admissibility condition implies that these components do not carry marked poles. Therefore, the poles on these components are of order at most -1 and only at the marked points that will be mapped to the branches of nodes.

We define the linear subspace $R_{\Gamma}^{\dagger} \subset R_{\Gamma}$, as the space of vectors in R_{Γ} defined by the condition: all residues of poles on components of level -1 (or equivalently at the nodes) vanish.

Now, we have seen that (Γ, I) defines the space

$$A_{\Gamma,I} \simeq A_{g_0,Z_0,P_0}^{R_0} \times p(A_{g_1,Z_1,P_1}^{R_1}) \hookrightarrow \overline{\mathcal{H}}_{g_0,n_0,P_0} \times \overline{\mathcal{M}}_{g_1,n_1,m_1}$$

(see Notation 4.14 for the definitions of g_i, Z_i, P_i and R^i). We denote by $\tilde{A}_{\Gamma,I}$ the space on the right-hand side.

With this notation, we have the isomorphism

$$\overline{\mathcal{H}}_{g_{\Gamma},n_{\Gamma},m_{\Gamma},P_{\Gamma}}^{R_{\Gamma}^{\dagger}} \simeq \overline{\mathcal{H}}_{g_0,n_0,P_0} \times \left(\prod_{v \in V^1} \overline{\mathcal{H}}_{g_v,n_v+m_v} \right),$$

where, in the second product, n_v and m_v are the lengths of Z_v and P_v , respectively, and we recall that $p_v: \overline{\mathcal{H}}_{g_v,n_v+m_v} \rightarrow \overline{\mathcal{M}}_{g_v,n_v+m_v}$ is the Hodge bundle. Indeed, a differential in $\overline{\mathcal{H}}_{g_{\Gamma},n_{\Gamma},m_{\Gamma},P_{\Gamma}}^{R_{\Gamma}^{\dagger}}$ is a differential on the normalization of a curve in $\zeta_{\Gamma}^{\#}(\overline{\mathcal{H}}_{\Gamma})$ such that the differential has no residues (thus no poles) at the branches of a node. Therefore, the restriction of this differential to components of level 0 is a point in $\overline{\mathcal{H}}_{g_0,n_0,P_0}$ (without residue condition at the marked poles), and its restriction to the component of level -1 is an holomorphic differential (thus a point in the product of the Hodge bundles).

All in all we have the sequence of embeddings

$$A_{\Gamma,I} \hookrightarrow \tilde{A}_{\Gamma,I} \hookrightarrow \overline{\mathcal{H}}_{g_{\Gamma},n_{\Gamma},m_{\Gamma},P_{\Gamma}}^{R_{\Gamma}^{\dagger}} \hookrightarrow \overline{\mathcal{H}}_{\Gamma},$$

where the second one is given by the zero section embedding of $\overline{\mathcal{M}}_{g_1,n_1,m_1}$ in the

Hodge bundle. All these embeddings are compatible with the \mathbb{C}^* -action; therefore, we get the sequence of embeddings

$$\mathbb{P}A_{\Gamma,I} \hookrightarrow \mathbb{P}\tilde{A}_{\Gamma,I} \hookrightarrow \mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}_{\Gamma},\mathbf{n}_{\Gamma},\mathbf{m}_{\Gamma},\mathbf{P}_{\Gamma}}^{R_{\Gamma}^{\dagger}} \hookrightarrow \mathbb{P}\bar{\mathcal{H}}_{\Gamma}.$$

From here, we will compute the Poincaré-dual cohomology class of the element $\mathbb{P}\tilde{A}_{\Gamma,I}$ in $H^*(\mathbb{P}\bar{\mathcal{H}}_{\Gamma}, \mathbb{Q})$ by computing successively the class of each of these substacks in $H^*(\mathbb{P}\bar{\mathcal{H}}_{\Gamma}, \mathbb{Q})$.

5.3.2 The Poincaré-dual class of $\mathbb{P}\tilde{A}_{\Gamma,I}$ We let

$$d_{\Gamma} = \dim(R_{\Gamma}) - \dim(R_{\Gamma}^{\dagger}).$$

The Poincaré-dual class of $\mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}_{\Gamma},\mathbf{n}_{\Gamma},\mathbf{P}_{\Gamma}}^{R_{\Gamma}^{\dagger}}$ in $H^*(\mathbb{P}\bar{\mathcal{H}}_{\Gamma}, \mathbb{Q})$ is equal to $\xi^{d_{\Gamma}}$ (see Lemma 2.27).

Now we consider the morphism $\psi: \mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}_{\Gamma},\mathbf{n}_{\Gamma},\mathbf{P}_{\Gamma}}^{R_{\Gamma}^{\dagger}} \rightarrow \bar{\mathcal{M}}_{\mathbf{g}_1,\mathbf{n}_1,\mathbf{m}_1}$ (it is the composition of a projection the forgetful map of the differential). The restriction of a differential in $\mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}_{\Gamma},\mathbf{n}_{\Gamma},\mathbf{P}_{\Gamma}}^{R_{\Gamma}^{\dagger}}$ gives rise to a morphism of vector bundles

$$\begin{array}{ccc} \Psi: \mathcal{O}(-1) & \longrightarrow & \psi^*\left(\bigoplus_{v \in V_1} \bar{\mathcal{H}}_{g_v, n_v + m_v}\right) \\ \downarrow & & \swarrow \\ \mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}_{\Gamma},\mathbf{n}_{\Gamma},\mathbf{m}_{\Gamma},\mathbf{P}_{\Gamma}}^{R_{\Gamma}^{\dagger}} & & \end{array}$$

The morphism Ψ can equivalently be seen as a section of

$$\mathcal{O}(1) \otimes \psi^*\left(\bigoplus_{v \in V_1} \bar{\mathcal{H}}_{g_v, n_v + m_v}\right).$$

The vanishing locus of Ψ is the locus of differentials whose restriction to level -1 components is identically zero, ie $\mathbb{P}\tilde{A}_{\Gamma,I}$ (with the reduced closed substack structure). The Poincaré-dual class of this locus in $H^*(\mathbb{P}\bar{\mathcal{H}}_{\Gamma}, \mathbb{Q})$ is then given by

$$\xi^{d_{\Gamma}} \cdot \prod_{v \in V_1} (\xi^{g_v} + \lambda_1 \xi^{g_v - 1} + \dots + \lambda_{g_v}).$$

5.3.3 The Poincaré-dual class of $\mathbb{P}\tilde{A}_{\Gamma,I}$ We have the natural isomorphism

$$\mathbb{P}\tilde{A}_{\Gamma,I} \simeq \mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}_0,\mathbf{n}_0,\mathbf{P}_0} \times \bar{\mathcal{M}}_{\mathbf{g}_1,\mathbf{n}_1,\mathbf{m}_1}.$$

We denote by Φ_0 and Φ_1 the projections on both factors.

Definition 5.8 The class $a_{\Gamma, I} \in H^*(\mathbb{P}\overline{\mathcal{H}}_{g, n, P}, \mathbb{Q})$ is defined by

$$\frac{1}{|\text{Aut}(\Gamma, I)|} \zeta_{\Gamma}^{\#} \left(\xi^{d_{\Gamma}} \cdot \Phi_1^* (p_* [\mathbb{P}\overline{A}_{g_1, Z_1, P_1}^{R^1}]) \cdot \Phi_0^* [\mathbb{P}\overline{A}_{g_0, Z_0, P_0}^{R^0}] \right. \\ \left. \times \prod_{v \in V^1} (\xi^{g_v} + \lambda_1 \xi^{g_v - 1} + \dots + \lambda_{g_v}) \right),$$

where $\text{Aut}(\Gamma, I)$ is the group of automorphisms of Γ preserving the twists at the edges.

Proposition 5.9 Let $(\Gamma, I) \in \text{Bic}(g, P, Z, R)$. We have:

- (1) If (Γ, I) is divisor graph then $a_{\Gamma, I} = [\mathbb{P}\overline{A}_{\Gamma, I}]$.
- (2) If (Γ, I) is not a divisor graph then $a_{\Gamma, I} = 0$.
- (3) If $[\mathbb{P}\overline{A}_{g_0, Z_0, P_0}^{R^0}]$ and $[\mathbb{P}\overline{A}_{g_1, Z_1, P_1}^{R^1}]$ are tautological and can be explicitly computed, then the same is true of $a_{\Gamma, I}$.

Proof of the first and second points If (Γ, I) is a divisor graph then

$$p: \mathbb{P}A_{g_1, Z_1, P_1}^{R^1} \rightarrow \text{Im}(p)$$

is of degree 1; thus, $p_*[\mathbb{P}A_{g_1, Z_1, P_1}^{R^1}] = [p(\mathbb{P}A_{g_1, Z_1, P_1}^{R^1})]$. Therefore, by construction, $a_{\Gamma, I}$ is the Poincaré-dual class of $\mathbb{P}\overline{A}_{\Gamma, I}$.

If (Γ, I) belongs to $\text{Bic}(g, P, Z, R) \setminus \text{Div}(g, P, Z, R)$ then the fibers of

$$p: \mathbb{P}A_{g_1, Z_1, P_1}^{R^1} \rightarrow \text{Im}(p)$$

are of positive dimension and $p_*[\mathbb{P}\overline{A}_{g_1, Z_1, P_1}^{R^1}] = 0$. □

Proof of the third point We assume that $[\mathbb{P}\overline{A}_{g_0, Z_0, P_0}^{R^0}]$ and $[\mathbb{P}\overline{A}_{g_1, Z_1, P_1}^{R^1}]$ are tautological and can be explicitly computed.

The projections Φ_1 is equal to the composition of the forgetful map from $\overline{\mathcal{H}}_{\Gamma}$ to $\overline{\mathcal{M}}_{\Gamma}^{\text{red}}$ with the projection to the vertices of level -1 . Thus, by definition, if β is a tautological class of $\overline{\mathcal{M}}_{g_1, Z_1, m_1}$ then $\Phi_1^* \beta$ is a tautological class of $H^*(\mathbb{P}\overline{\mathcal{H}}_{g, Z, P}, \mathbb{Q})$. Further, if $[\mathbb{P}\overline{A}_{g_1, Z_1, P_1}^{R^1}]$ is tautological and be explicitly computed, then so is $p_*[\mathbb{P}\overline{A}_{g_1, Z_1, P_1}^{R^1}]$: indeed, the Segre class of $\overline{\mathcal{H}}_{g_1, n_1, P_1}$ is a tautological class of $\overline{\mathcal{M}}_{g_1, m_1, P_1}$.

The map Φ_1 is equivariant with respect to the \mathbb{C}^* -action; thus, $\Phi_1^{-1}(c_1(\mathcal{O}(1))) = c_1(\mathcal{O}(1))$. Further, the following diagram commutes:

$$\begin{array}{ccc}
 \mathbb{P}\tilde{A}_{\Gamma, I} & \xrightarrow{\Phi_0} & \mathbb{P}\bar{A}_{\mathbf{g}_0, \mathbf{n}_0, \mathbf{P}_0} \\
 \downarrow & & \downarrow p \\
 \bar{\mathcal{M}}_{\Gamma}^{\text{red}} & \longrightarrow & \bar{\mathcal{M}}_{\mathbf{g}_0, \mathbf{n}_0, \mathbf{m}_0}^{\text{red}}
 \end{array}$$

Thus, if β is a tautological class of $\bar{\mathcal{M}}_{\mathbf{g}_0, \mathbf{n}_0, \mathbf{m}_0}^{\text{red}}$, then the class $\Phi_0^*(p^*(\beta))$ is a tautological class of $\mathbb{P}\bar{\mathcal{H}}_{\Gamma}$ and thus a tautological class of $H^*(\mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}, \mathbb{Q})$. \square

We can already remark that Proposition 5.9 implies the following:

Corollary 5.10 *The following equality holds:*

$$\sum_{(\Gamma, I) \in \text{Bic}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}} m(I)a_{\Gamma, I} = \sum_{(\Gamma, I) \in \text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}} m(I)a_{\Gamma, I}.$$

Proof This follows from the fact that if $(\Gamma, I) \in \text{Bic}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i} \setminus \text{Div}(\mathbf{g}, \mathbf{P}, \mathbf{Z}, R)_{j,i}$ then $a_{\Gamma, I} = 0$. \square

5.4 Proof of Theorems 1, 2 and 3

We have all the ingredients to prove Theorem 4 (see the beginning of the section).

Proof of Theorem 4 For a list $\mathbf{Z} = (Z_1, \dots, Z_q)$ of vectors of nonnegative integers, we write $|\mathbf{Z}| = \sum_{j=1}^q |Z_j|$. We prove Theorem 4 by induction on $|\mathbf{Z}|$.

Base of the induction: $|\mathbf{Z}| = 0$ Let $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$ be a quadruple that satisfies Assumption 3.1 and is such that $|\mathbf{Z}| = 0$ is trivial; then $A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R$ is dense in $\bar{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}$. Therefore,

$$[\mathbb{P}A_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R] = [\mathbb{P}\bar{\mathcal{H}}_{\mathbf{g}, \mathbf{n}, \mathbf{P}}^R] = \xi^{\dim(\mathfrak{A}) - \dim(R)},$$

by Lemma 2.27.

Induction Now, let $(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$ be a quadruple satisfying Assumption 3.1 and such that $|\mathbf{Z}| > 0$. The induction formulas (12) and (13) of Theorem 5 express the class $[\mathbb{P}\bar{A}_{\mathbf{g}, \mathbf{Z}, \mathbf{P}}^R]$ in terms of a class with a smaller sum of the order of zeros and a sum over all bicolored graph (by Corollary 5.10). We only need to prove that the class $a_{\Gamma, I}$ is tautological for any $(\Gamma, I) \in \text{Div}(\mathbf{g}, \mathbf{Z}, \mathbf{P}, R)$.

The vectors of zeros \mathbf{Z}_0 and \mathbf{Z}_1 of the levels 0 and -1 satisfy $|\mathbf{Z}_i| < |\mathbf{Z}|$. Therefore, the classes $[\mathbb{P}\bar{A}_{\mathbf{g}_1, \mathbf{Z}_1, \mathbf{P}_1}^{R^1}]$ and $[\mathbb{P}\bar{A}_{\mathbf{g}_0, \mathbf{Z}_0, \mathbf{P}_0}^{R^0}]$ can be computed and are tautological. Using Proposition 5.9, this implies that the class $a_{\Gamma, I}$ is tautological and can be computed. \square

Theorems 1, 2 and 3, stated in Section 1.5, are straightforward corollaries of Theorem 4:

Proof of Theorems 1, 2 and 3 Theorem 1 is the special case of Theorem 4 for connected and stable curves. Theorem 3 is a consequence of Theorem 1 and Proposition 1.4 (the Segre class of the spaces of stable differential is tautological).

To prove Theorem 2, we recall that we denote by $\tilde{\pi}_n: \mathbb{P}\bar{\mathcal{H}}_{g,n} \rightarrow \mathbb{P}\bar{\mathcal{H}}_g$ the forgetful map of points. The bundle $\bar{\mathcal{H}}_{g,n}$ is the pullback of $\bar{\mathcal{H}}_g$ by π_n ; then $\xi \in H^*(\mathbb{P}\bar{\mathcal{H}}_{g,n}, \mathbb{Q})$ is the pullback of $\xi \in H^*(\mathbb{P}\bar{\mathcal{H}}_g, \mathbb{Q})$. Therefore, the pushforward of a tautological class of $RH^*(\mathbb{P}\bar{\mathcal{H}}_{g,n}, \mathbb{Q})$ by π_n is in $RH^*(\mathbb{P}\bar{\mathcal{H}}_g, \mathbb{Q})$ and can be explicitly computed.

If $Z = (k_1, \dots, k_n)$ is complete, the map $\tilde{\pi}_n$ restricted to $\mathbb{P}A_{g,Z}$ is finite of degree $\text{Aut}(Z)$ onto $\mathbb{P}\mathcal{H}[Z]$. We have

$$[\mathbb{P}\bar{\mathcal{H}}[Z]] = \frac{1}{\text{Aut}(Z)} \cdot \tilde{\pi}_{n*}[\mathbb{P}\bar{A}_{g,Z}],$$

and the class $[\mathbb{P}\bar{\mathcal{H}}[Z]]$ is tautological and can be explicitly computed. □

6 Examples of computation

We give two examples of computation: the first one is a computation in the projectivized Hodge bundle (we forget the marked points); the second is a computation in the moduli space of curves (we forget the differential).

6.1 The class $[\mathbb{P}\bar{\mathcal{H}}_g(3)]$

We consider here $g > 2$ and $Z = (3, 1, \dots, 1)$. We have seen in the introduction the computation of $[\mathbb{P}A_{g,(2)}]$. Therefore, in order to compute $[\mathbb{P}A_{g,(3)}]$ we need to list the divisor graphs contributing to $[\mathbb{P}\bar{A}_{g,(3)}] - (\xi + 3\psi_1)[\mathbb{P}\bar{A}_{g,(2)}]$. See Figure 2.




Figure 2: List of boundary terms in $[\mathbb{P}\bar{A}_{g,(3)}] - (\xi + 3\psi_1)[\mathbb{P}\bar{A}_{g,(2)}]$

We explain the notation of the above expression. The legs on the graphs stands for the only marked point. We have decorated graph with classes P_v in $\overline{\mathcal{M}}_{g(v),n(v)}$ for each vertex. These classes are either ψ_1 (for the marked point), ψ_e for an half-edge or λ_1 for a vertex.

We recall that $\mathcal{H}_3(4)$ has two connected components (hyperelliptic and odd). In this case one can compute $[\overline{\mathcal{M}}_3(4)^{\text{hyp}}]$ by using the work of Faber and Pandharipande (see [14]). This way one can also compute $[\overline{\mathcal{M}}_3(4)^{\text{odd}}] = [\overline{\mathcal{M}}_3(4)] - [\overline{\mathcal{M}}_3(4)^{\text{hyp}}]$. In general, it is possible to compute the class of the hyperelliptic component but we do not know how to compute separately the classes of odd and even components for $g \geq 4$.

Felix Janda has compared this expression with the expression of Conjecture B. The two expressions agree modulo tautological relations (see Section 1.7 for presentation of the conjecture).

If we forget the marked point, then we get a class in $\text{Pic}(\overline{\mathcal{M}}_g) \otimes \mathbb{Q}$. Using the string and dilaton equations and Mumford’s formula, for κ_1 we get

$$\begin{aligned} \pi_*[\overline{\mathcal{M}}_3(4)] &= 0 - 10 \times 4\lambda_1 + 35\kappa_1 - 5\delta_{\text{nonsep}} - 0 + 6 \cdot 0 + 0 + 6 \cdot 0 - 34\delta_{\text{sep}} - 11\delta_{\text{sep}} \\ &\quad + 0 - 10 \times 3\delta_{\text{sep}} - \delta_{\text{sep}} \\ &= 380\lambda_1 - 40\delta_{\text{nonsep}} - 100\delta_{\text{sep}}. \end{aligned}$$

The expression agrees with the formula of Scott Mullane (see [27]).

7 Relations in the Picard group of the strata

We fix the notation for this section. Let $g, n, m \geq 0$ such that $2g - 2 + n + m > 0$. Let $Z = (k_1, \dots, k_n)$ and $P = (p_1, \dots, p_m)$ be vectors of positive integers such that $|Z| - |P| = 2g - 2$. In this section we consider the space $\overline{\mathcal{M}}_g(Z - P) \subset \overline{\mathcal{M}}_{g,n+m}$ (see Section 1.5 for definitions). The purpose is to define several natural classes in $\text{Pic}(\overline{\mathcal{M}}_g(Z - P)) \otimes \mathbb{Q}$ and to compute relations between these elements. Namely, there are two types of classes which arise naturally:

- divisors associated to admissible graphs (see Sections 4.2 and 4.4);
- intersections of $\overline{\mathcal{M}}_g(Z - P)$ with the tautological classes of $A_1(\overline{\mathcal{M}}_{g,n})$.

7.1 Classes defined by admissible graphs

We consider the moduli space of stable differentials $\overline{\mathcal{H}}_{g,n,P}$ and the locus $\overline{A}_{g,Z,P} \subset \overline{\mathcal{H}}_{g,n,P}$. We recall that $p: \overline{\mathcal{H}}_{g,n,P} \rightarrow \overline{\mathcal{M}}_{g,n+m}$ is the forgetful map. We have seen

that $\overline{A}_{g,Z,P}$ admits a stratification indexed by admissible graphs (see Lemma 4.17). Here, we will describe the set of admissible graphs (Γ, I, l) such that $p(\overline{A}_{\Gamma,I,l})$ is a divisor in $\overline{\mathcal{M}}_g(Z - P) = p(\overline{A}_{g,Z,P})$.

The map $p: \mathbb{P}A_{g,Z,P} \rightarrow \mathcal{M}_g(Z - P)$ is an isomorphism (see Lemma 3.14). Thus, if $p(\mathbb{P}\overline{A}_{\Gamma,I,l})$ is a divisor in $\overline{\mathcal{M}}_g(Z - P)$ then $\mathbb{P}\overline{A}_{\Gamma,I,l}$ is a divisor in $\mathbb{P}\overline{A}_{g,Z,P}$. We saw that an admissible graph (Γ, I, l) defines to a divisor of $\overline{A}_{g,Z,P}$ if and only if it is of one of the three following types (see Section 4.4):

- (1) the admissible graph of depth 0 with one vertex and one edge;
- (2) an admissible graph of depth 0 with two vertices and one edge;
- (3) a bicolored graph that satisfies the condition $(\star\star)$.

Proposition 7.1 *Let (Γ, I, l) be an admissible graph. The locus $p(\mathbb{P}\overline{A}_{\Gamma,I,l})$ is a divisor of $\overline{\mathcal{M}}_g(Z - P)$ if and only if*

- (Γ, I, l) is of the type 1 above; or
- (Γ, I, l) is a bicolored graph with one vertex of level -1 , one stable vertex of level 0 and possibly other semistable vertices of level 0.

We call an *irreducible divisor* the divisor of $\overline{\mathcal{M}}_g(Z - P)$ of the first type. We denote this divisor by D_0 (with the reduced structure).

In the second case, the stabilization of the graph Γ determines a unique stable twisted graph of depth 1, (Γ', I') (we no longer write the level structure, which is uniquely determined by I). Conversely, a twisted stable graph of depth 1 with two vertices, we can uniquely determine an admissible graph satisfying the condition of Proposition 7.1 by putting all the poles on the component of level -1 on unstable rational components of level 0 (see Lemma 4.10 and Example 7.3 below).

Definition 7.2 *A simple bicolored graph is a twisted stable graph of depth 1 with two vertices. We denote by $SB(Z, P)$ the set of simple bicolored graphs. If (Γ, I) is a simple bicolored graph, we denote by $D_{\Gamma,I}$ the corresponding divisor in $\overline{\mathcal{M}}_g(Z - P)$ (with the reduced structure) and by $a_{\Gamma,I}$ its class in $\text{Pic}(\overline{\mathcal{M}}_g(Z - P)) \otimes \mathbb{Q}$.*

The class $i_*a_{\Gamma,I}$ (where i is the closed immersion of $p(\overline{A}_{\Gamma,I})$ in $\overline{\mathcal{M}}_{g,n+m}$) in the moduli space of curves is simply given by

$$\zeta_{\Gamma*}([\overline{\mathcal{M}}_{g_0}(Z_0 - P_0)], [\overline{\mathcal{M}}_{g_1}(Z_1 - P_1)]),$$




Figure 3: Example of the correspondence between admissible and stable graphs

where g_0 and g_1 are the genera of the vertices of levels 0 and -1 and the vectors Z_0, P_0, Z_1 and P_1 are the vectors encoding the orders of zeros and poles at the marked points and half-edges induced by Z, P and the twist I .

Example 7.3 We illustrate this correspondence between simple bicolored graphs and boundary divisors. We consider $g = 3, Z = (2, 6)$ and $P = (-2, -2)$ and the admissible graph in Figure 3 (in this example we take the twists equal to 1 on all edges). In this example, the class $i_*a_{\Gamma, I}$ in the moduli space of curves will be given by

$$\zeta_{\Gamma*}([\overline{\mathcal{M}}_1(+2, +0, +0, -2)], [\overline{\mathcal{M}}_1(+6, -2, -2, -2)]).$$

Proof of Proposition 7.1 Let (Γ, I) be an admissible graph of depth at most 1 with several stable components of level 0. Then the fiber of p over a generic point of $p(A_{g, Z, P})$ is of dimension greater than 1. That is why divisors of type 2 are not mapped to divisors while the map p restricted to D_0 is indeed of degree 1 onto its image.

Now we consider an admissible graph of depth 1 with one stable vertex of level 0. Then the graph satisfies condition $(\star\star)$ if and only if it has one vertex of level -1 .

Finally, we consider an admissible graph (Γ, I, l) of depth 1 and with no stable vertex of level 0. The projectivized stratum $\mathbb{P}A_{\Gamma, I, l} \subset \mathbb{P}\overline{\mathcal{H}}_{g, n, P}$ is empty. Indeed, Z is complete for g and P ; thus, the differential on each unstable component with a marked pole of order p is given by dz/z^p . Therefore, $A_{\Gamma, I, l}$ is a substack of the zero section of the cone $\overline{\mathcal{H}}_{g, n, P} \rightarrow \overline{\mathcal{M}}_{g, n+m}$ (see Section 2.3 for the description of the zero section). □

7.2 Classes defined by residue conditions

We recall that \mathfrak{R} is the vector space of residues, ie the subspace of \mathbb{C}^m defined by $\{(r_1, \dots, r_m) / r_1 + \dots + r_m = 0\}$. Let $R \subset \mathfrak{R}$ be vector subspace of codimension 1.

We define the following class in the rational Picard group of $\overline{\mathcal{M}}_g(Z - P)$:

$$\delta_R^{\text{res}} = p_*(\mathbb{P}\overline{A}_{g,Z,P}^R).$$

Notation 7.4 Let $1 \leq i < j \leq n + m$. We denote by $\text{SB}(Z, P)_i$ (resp. $\text{SB}(Z, P)^i$) the set of simple bicolored graphs such that the leg corresponding to i is adjacent to the vertex of level -1 (resp. to the vertex of level 0). We let $\text{SB}(Z, P)_i^j = \text{SB}(Z, P)_i \cap \text{SB}(Z, P)^j$.

Let $R \subset \mathfrak{R}$ is a vector subspace. For a simple bicolored graph, we denote by $R^0 \subset \mathfrak{R}$ the vector space defined by the linear conditions $\{r_i = 0\}$ for all $1 \leq i \leq m$ such that the leg of index $n + i$ is at level -1 . We denote by $\text{SB}(Z, P)_R$ the set of simple bicolored graphs such that the space R contains R^0 .

7.3 Classes defined by intersection

Let β be a tautological class in $\text{Pic}(\overline{\mathcal{M}}_{g,n+m}) \otimes \mathbb{Q}$. The class β determines a class in $\text{Pic}(\overline{\mathcal{M}}_g(Z - P)) \otimes \mathbb{Q}$ by taking $i^*\beta$, where i is the closed immersion of $\overline{\mathcal{M}}_g(Z - P)$ into $\overline{\mathcal{M}}_{g,n+m}$. If β is either λ_1, κ_1 or a ψ -class then we will denote by the same letter its pullback to $\text{Pic}(\overline{\mathcal{M}}_g(Z - P)) \otimes \mathbb{Q}$ if the context is clear.

The last class that we will consider is the pushforward of the ξ -class, which we denote by

$$\overline{\xi} = p_*(\xi \cdot [\mathbb{P}\overline{A}_{g,Z,P}]).$$

Theorem 6 *The following relations hold in $\text{Pic}(\overline{\mathcal{M}}_g(Z - P)) \otimes \mathbb{Q}$:*

(1) For all $1 \leq i \leq n$,

$$\overline{\xi} + (k_i + 1)\psi_1 = \sum_{(\Gamma, I) \in \text{SB}(Z, P)_i} m(I)a_{\Gamma, I}.$$

(2) For all $1 \leq i, j \leq n$,

$$(k_i + 1)\psi_i - (k_j + 1)\psi_j = \sum_{(\Gamma, I) \in \text{SB}(Z, P)_i^j} m(I)a_{\Gamma, I} - \sum_{(\Gamma, I) \in \text{SB}(Z, P)^j} m(I)a_{\Gamma, I}.$$

(3) For all $R \subset \mathfrak{R}$ vector subspace of codimension 1,

$$\overline{\xi} = \delta_R^{\text{res}} + \sum_{(\Gamma, I) \in \text{SB}(Z, P)_R} m(I)a_{\Gamma, I}.$$

(4) If $m = 0$ then

$$\lambda_1 + \kappa_Z \bar{\xi} = \frac{1}{12} \delta + \sum_{(\Gamma, I) \in \text{SB}(Z)} 2\bar{m}(I, \Gamma) a_{\Gamma, I},$$

where δ is the boundary divisor of $\bar{\mathcal{M}}_{g,n}$,

$$\kappa_Z = \frac{1}{12} \sum_{i=1}^n \frac{k_i(k_i + 2)}{k_i + 1}$$

and

$$\bar{m}(I, \Gamma) = \frac{m(I)}{12} \left(-m(I) + \sum_{i \rightarrow v^1} \frac{k_i(k_i + 2)}{k_i + 1} \right);$$

the second sum goes over all legs adjacent to the vertex of level -1 .

7.3.1 Relations (1) and (2) and double ramification cycles The second relation of Theorem 6 is a direct consequence of the first one: we write $(k_i + 1)\psi_i - (k_j + 1)\psi_j = (\bar{\xi} + (k_i + 1)\psi_i) - (\bar{\xi} + (k_j + 1)\psi_j)$. However, we choose to write relation (2) in this form for two reasons:

(1) First, because it involves only classes defined directly in the moduli space of curves.

(2) The second motivation is related to Conjectures A and B. Indeed, the classes $H_g(Z)$ and $[\bar{\mathcal{M}}_g(Z - P)]$ (see Section 1.7 for definitions) are supposed to be generalizations of double ramification cycles. In [5], the authors proved several identities between intersection of ψ -classes with double ramification cycles. One consequence of the relations proven in [5] is the existence a universal ψ -class over the double ramification cycles (independent of the choice of a marked point). For strata of differentials the following corollary gives a candidate for this universal ψ -class:

Corollary 7.5 *The class in $\text{Pic}(\bar{\mathcal{M}}_g(Z - P)) \otimes \mathbb{Q}$*

$$(k_i + 1)\psi_i - \sum_{(\Gamma, I) \in \text{SB}(Z, P)_i} m(I) a_{\Gamma, I}$$

is independent of the choice of $1 \leq i \leq n$.

Proof of relation (1) This is a direct consequence of the induction formula (see Theorem 5). We consider Z_i , the vector obtained from Z by increasing the i^{th} entry

by 1, and $R = \mathfrak{R}$ (no residue condition); then we get

$$(\xi + (k_i + 1)\psi_i) \cdot [\mathbb{P} \bar{A}_{g,Z,P}] = [\mathbb{P} \bar{A}_{g,Z_j,P}] + \sum_{(\Gamma,I) \in \text{Bic}(g,Z,P)_i} m(I)a_{\Gamma,I}.$$

We remark that $|Z_j| - |P| > 2g - 2$; thus, $[\mathbb{P} \bar{A}_{g,Z_j,P}] = 0$. Now we apply the pushforward by p to this expression. In the sum of the right-hand side, only the simple bicolored graphs will contribute and we indeed get

$$\bar{\xi} + (k_i + 1)\psi_1 = \sum_{(\Gamma,I) \in \text{SB}(Z,P)_i} m(I)a_{\Gamma,I}. \quad \square$$

7.3.2 Relation (3) To prove the third relation, we need a generalization of the induction formula. Let $R \subset \mathfrak{R}$ be a vector subspace of codimension 1. We recall that an admissible bicolored graph defines a space of residue conditions $R^0 \subset \mathfrak{R}$ (see Section 4.2 for the construction of R^0). We define $\text{Bic}(g, Z, P)_R \subset \text{Bic}(g, Z, P, \mathfrak{R})$ as the subset of bicolored graphs such that $R^0 \subset R$.

Proposition 7.6 *The following equality holds in $H^*(\mathbb{P} \bar{\mathcal{H}}_{g,n,P}; \mathbb{Q})$:*

$$[\mathbb{P} \bar{A}_{g,Z,P}^R] = \xi[\mathbb{P} \bar{A}_{g,Z,P}] - \sum_{(\Gamma,I) \in \text{Bic}(g,Z,P)_R} m(I)a_{\Gamma,I}.$$

Remark 7.7 We could have stated this proposition in larger generality (unstable disconnected base) but it will not be useful here.

Proof The proof is the same as the proof of Theorem 5. We consider the line bundle $\mathcal{O}(1) \simeq \mathcal{O}(-1)^\vee$ restricted to $\mathbb{P} \bar{A}_{g,Z,P}$ with its section

$$s: \mathcal{O}(-1) \rightarrow \mathbb{C}, \quad \alpha \mapsto \mathfrak{R}/R,$$

defined as the composition of the residue map $\mathcal{O}(-1) \rightarrow \mathfrak{R}$ and the projection $\mathfrak{R} \rightarrow \mathfrak{R}/R$. The vanishing locus of the section s is the union of $\mathbb{P} \bar{A}_{g,Z,P}^R$ and of the divisors $\mathbb{P} \bar{A}_{\Gamma,I}$ for all $(\Gamma, I) \in \text{Bic}(g, Z, P)_R$.

Now the vanishing order of s along $\mathbb{P} \bar{A}_{g,Z,P}^R$ is 1 because the residue map is a submersion. The vanishing order of s along $\mathbb{P} \bar{A}_{\Gamma,I}$ is 1 because Lemma 5.6 remains valid if we replace the section $s_{i,j}$ by the section s and the set of graphs $\text{Div}(g, Z, P, R)_{ij}$ by the set of graphs $\text{Div}(g, Z, P)_R$. □

Proof of relation (3) Relation (3) is a direct consequence of Proposition 7.6. It suffices to apply the pushforward by the forgetful map p . □

7.3.3 Relation (4) and the work of Eskin, Kontsevich and Zorich Let $g \geq 2$ and let $Z = (k_1, \dots, k_n)$ be a partition of $2g - 2$. Before proving relation (4), let us mention that Kontsevich proved that

$$\lambda_1 = -\kappa_Z \bar{\xi} + \gamma,$$

where γ is a class supported on the boundary of $\bar{\mathcal{M}}(Z)$. From this relation, he deduced an equation relating two numerical invariants of strata of differentials: the sum of the Lyapunov exponents and the Siegel–Veech constants (the complete proof of this equation was achieved in [13]). Relation (4) gives an explicit expression for the class γ .

Proof of relation (4) Let Z' be the vector equal to $(k_1, \dots, k_n, 0)$. If $\pi: \bar{\mathcal{M}}_{g,n+1} \rightarrow \bar{\mathcal{M}}_{g,Z}$ is the forgetful map of the last marked point, then we have $\bar{A}_{g,Z'} = \pi^{-1}(\bar{A}_{g,Z})$. We use the induction formula to obtain the relation

$$(\xi + \psi_{n+1})[\mathbb{P}A_{g,Z'}] = 0 + \sum_{\text{Bic}(g,Z)_{n+1}} m(I) a_{\gamma,I}.$$

We multiply this formula by ψ_{n+1} to get

$$(14) \quad \xi \psi_{n+1}[\mathbb{P}A_{g,Z'}] + \psi_{n+1}^2[\mathbb{P}A_{g,Z'}] = \sum_{\text{Bic}(g,Z')_{n+1}} m(I) \psi_{n+1} a_{\Gamma,I}.$$

Now we apply $(p_*) \circ (\pi_*)$ to this formula (we forget the last point and then the differential). We study each term separately.

Contribution of $\xi \psi_{n+1}[\mathbb{P}A_{g,Z'}]$ The classes ξ and $[\mathbb{P}A_{g,Z'}]$ are pullbacks by π ; thus,

$$\begin{aligned} p_*(\pi_*(\psi_{n+1} \xi[\mathbb{P}A_{g,Z'}])) &= p_*(\pi_*(\psi_{n+1}) \xi[\mathbb{P}A_{g,Z}]) \\ &= \kappa_0 p_*(\xi[\mathbb{P}A_{g,Z}]) \\ &= (2g - 2 + n) \bar{\xi} \end{aligned}$$

by the projection formula.

Contribution of $\psi_{n+1}^2[\mathbb{P}A_{g,Z'}]$ Still by the projection formula, we have

$$p_*(\pi_*(\psi_{n+1}^2[\mathbb{P}A_{g,Z'}])) = p_*(\pi_*(\psi_{n+1}^2)[\mathbb{P}A_{g,Z}]) = \kappa_1 = 12\lambda_1 - \delta + \sum_{i=1}^n \psi_i.$$

Now we use the first relation to write

$$\sum_{i=1}^n \psi_i = - \left(\sum_{i=1}^n \frac{1}{k_i + 1} \right) \bar{\xi} + \sum_{i=1}^n \left(\sum_{(\Gamma,I) \in \text{BS}(g,Z)_i} \frac{m(I)}{k_i + 1} a_{\Gamma,I} \right).$$

Contribution of $\sum_{\text{Bic}(g, Z')_{n+1}} m(I)\psi_{n+1}a_{\Gamma, I}$ Let (Γ, I) be a bicolored graph in $\text{Bic}(g, Z')_{n+1}$. There are two possible configurations:

- The point $n + 1$ belongs to a rational components with three special points, in which case $\psi_{n+1}a_{\Gamma, I} = 0$.
- The point $n + 1$ is carried by a general vertex of level -1 which is not contracted after the forgetful map.

In the second case, we denote by (Γ', I') the twisted graph obtained after forgetting the marked point. We get

$$\pi_*(\psi_{n+1}a_{\Gamma, I}) = (2g_{\Gamma', I', 1} - 2 + n_{\Gamma', I', 1})a_{\Gamma', I'},$$

where $g_{\Gamma, 1}$ and $n_{\Gamma, 1}$ denote the genus and valency of the vertex of level -1 . Thus,

$$(p_* \circ \pi_*) \sum_{\text{Bic}(g, Z')_{n+1}} m(I)\psi_{n+1}a_{\Gamma, I} = \sum_{(\Gamma, I) \in \text{BS}(g, Z)} m(I)(2g_{\Gamma', I', 1} - 2 + n_{\Gamma', I', 1})a_{\Gamma, I}.$$

We obtain relation (4) by replacing all the terms in (14) by their expressions in terms of simple bicolored graphs. □

References

- [1] **E Arbarello, M Cornalba, P A Griffiths**, *Geometry of algebraic curves, II*, Grundle. Math. Wissen. 268, Springer (2011) MR
- [2] **M Bainbridge, D Chen, Q Gendron, S Grushevsky, M Möller**, *Compactification of strata of abelian differentials*, Duke Math. J. 167 (2018) 2347–2416 MR
- [3] **L Battistella, N Nabijou**, *A Lefschetz-type theorem for quasimap invariants via relative quasimaps*, preprint (2017) arXiv
- [4] **C Boissy**, *Connected components of the strata of the moduli space of meromorphic differentials*, Comment. Math. Helv. 90 (2015) 255–286 MR
- [5] **A Buryak, S Shadrin, L Spitz, D Zvonkine**, *Integrals of ψ -classes over double ramification cycles*, Amer. J. Math. 137 (2015) 699–737 MR
- [6] **L Caporaso**, *A compactification of the universal Picard variety over the moduli space of stable curves*, J. Amer. Math. Soc. 7 (1994) 589–660 MR
- [7] **H-L Chang, J Li, W-P Li**, *Witten’s top Chern class via cosection localization*, Invent. Math. 200 (2015) 1015–1063 MR
- [8] **D Chen**, *Square-tiled surfaces and rigid curves on moduli spaces*, Adv. Math. 228 (2011) 1135–1162 MR

- [9] **D Chen**, *Strata of abelian differentials and the Teichmüller dynamics*, J. Mod. Dyn. 7 (2013) 135–152 MR
- [10] **D Chen, Q Chen**, *Spin and hyperelliptic structures of log twisted differentials*, Selecta Math. 25 (online publication February 2019) MR
- [11] **A Chiodo**, *A construction of Witten’s top Chern class in K -theory*, from “Gromov–Witten theory of spin curves and orbifolds” (T J Jarvis, T Kimura, A Vaintrob, editors), Contemp. Math. 403, Amer. Math. Soc., Providence, RI (2006) 21–29 MR
- [12] **T Ekedahl, S Lando, M Shapiro, A Vainshtein**, *Hurwitz numbers and intersections on moduli spaces of curves*, Invent. Math. 146 (2001) 297–327 MR
- [13] **A Eskin, M Kontsevich, A Zorich**, *Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow*, Publ. Math. Inst. Hautes Études Sci. 120 (2014) 207–333 MR
- [14] **C Faber, R Pandharipande**, *Hodge integrals and Gromov–Witten theory*, Invent. Math. 139 (2000) 173–199 MR
- [15] **G Farkas, R Pandharipande**, *The moduli space of twisted canonical divisors*, J. Inst. Math. Jussieu 17 (2018) 615–672 MR
- [16] **W Fulton**, *Intersection theory*, 2nd edition, Ergeb. Math. Grenzgeb. 2, Springer (1998) MR
- [17] **A Gathmann**, *Absolute and relative Gromov–Witten invariants of very ample hypersurfaces*, Duke Math. J. 115 (2002) 171–203 MR
- [18] **Q Gendron**, *The Deligne–Mumford and the incidence variety compactifications of the strata of $\Omega\mathcal{M}_g$* , Ann. Inst. Fourier (Grenoble) 68 (2018) 1169–1240 MR
- [19] **Q Gendron, G Tahar**, *Différentielles à singularités prescrites*, preprint (2017) arXiv
- [20] **D Gieseker**, *A degeneration of the moduli space of stable bundles*, J. Differential Geom. 19 (1984) 173–206 MR
- [21] **T Graber, R Pandharipande**, *Constructions of nontautological classes on moduli spaces of curves*, Michigan Math. J. 51 (2003) 93–109 MR
- [22] **J Guéré**, *A generalization of the double ramification cycle via log-geometry*, preprint (2016) arXiv
- [23] **F Janda, R Pandharipande, A Pixton, D Zvonkine**, *Double ramification cycles on the moduli spaces of curves*, Publ. Math. Inst. Hautes Études Sci. 125 (2017) 221–266 MR
- [24] **M Kazarian, S Lando, D Zvonkine**, *Universal cohomological expressions for singularities in families of genus 0 stable maps*, Int. Math. Res. Not. 2018 (2018) 6817–6843 MR
- [25] **M Kontsevich, A Zorich**, *Connected components of the moduli spaces of Abelian differentials with prescribed singularities*, Invent. Math. 153 (2003) 631–678 MR

- [26] **D Korotkin, P Zograf**, *Tau function and moduli of differentials*, Math. Res. Lett. 18 (2011) 447–458 MR
- [27] **S Mullane**, *Divisorial strata of abelian differentials*, Int. Math. Res. Not. 2017 (2017) 1717–1748 MR
- [28] **D Mumford**, *Towards an enumerative geometry of the moduli space of curves*, from “Arithmetic and geometry, II” (M Artin, J Tate, editors), Progr. Math. 36, Birkhäuser, Boston, MA (1983) 271–328 MR
- [29] **A Okounkov, R Pandharipande**, *Gromov–Witten theory, Hurwitz theory, and completed cycles*, Ann. of Math. 163 (2006) 517–560 MR
- [30] **R Pandharipande, A Pixton, D Zvonkine**, *Relations on $\overline{\mathcal{M}}_{g,n}$ via 3–spin structures*, J. Amer. Math. Soc. 28 (2015) 279–309 MR
- [31] **R Pandharipande, A Pixton, D Zvonkine**, *Tautological relations via r –spin structures*, preprint (2016) arXiv
- [32] **A Polishchuk**, *Moduli spaces of curves with effective r –spin structures*, from “Gromov–Witten theory of spin curves and orbifolds” (T J Jarvis, T Kimura, A Vaintrob, editors), Contemp. Math. 403, Amer. Math. Soc., Providence, RI (2006) 1–20 MR
- [33] **A Polishchuk, A Vaintrob**, *Algebraic construction of Witten’s top Chern class*, from “Advances in algebraic geometry motivated by physics” (E Previato, editor), Contemp. Math. 276, Amer. Math. Soc., Providence, RI (2001) 229–249 MR
- [34] **A Sauvaget**, *Volumes and Siegel–Veech constants of $\mathcal{H}(2G - 2)$ and Hodge integrals*, Geom. Funct. Anal. 28 (2018) 1756–1779 MR
- [35] **D Zvonkine**, *Strata in the projectivization of the hodge bundle*, personal communication (2012)

Université Pierre et Marie Curie
Paris, France

adrien.sauvaget@inj-prg.fr

Proposed: Jim Bryan

Seconded: Dan Abramovich, Anna Wienhard

Received: 28 February 2017

Revised: 7 August 2018

