



HAL
open science

Domain-specific physical activity and sedentary behavior during pregnancy and postpartum depression risk in the French EDEN and ELFE cohorts

Judith van Der Waerden, Aurelie Nakamura, Laura Pryor, Marie-Aline Charles, Fabienne El-Khoury, Patricia Dargent-Molina

► To cite this version:

Judith van Der Waerden, Aurelie Nakamura, Laura Pryor, Marie-Aline Charles, Fabienne El-Khoury, et al.. Domain-specific physical activity and sedentary behavior during pregnancy and postpartum depression risk in the French EDEN and ELFE cohorts. *Preventive Medicine*, 2019, 121, pp.33-39. 10.1016/j.ypmed.2019.02.012 . hal-02178980

HAL Id: hal-02178980

<https://hal.sorbonne-universite.fr/hal-02178980>

Submitted on 10 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Domain-specific physical activity and sedentary behavior during pregnancy and**
2 **postpartum depression risk in the French EDEN and ELFE cohorts**

3
4 Judith van der Waerden^{a*}, Aurelie Nakamura^a, Laura Pryor^{ab}, Marie-Aline Charles^{cde}, Fabienne El-
5 Khoury^a, Patricia Dargent-Molina^{cd} on behalf of The EDEN Mother–Child Cohort Study Group¹

6 (a) INSERM, Sorbonne Université, Institut Pierre Louis d'épidémiologie et de Santé Publique,
7 IPLESP UMR-S1136, F75012 Paris, France

8 (b) Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

9 (c) INSERM, UMR1153 Epidemiology and Biostatistics Sorbonne Paris Cité Center (CRESS),
10 Developmental Origins of Health and disease (ORCHAD) Team, Villejuif, F-94807 France;

11 (d) Paris Descartes University France

12 (e) INED-INSERM-EFS joint unit Elfe

13
14 ¹ The EDEN Mother–Child Cohort Study Group includes I. Annesi-Maesano, J.Y. Bernard, J. Botton,
15 M.A. Charles, P. Dargent-Molina, B. de Lauzon-Guillain, P. Ducimetière, M. de Agostini, B. Foliguet,
16 A. Forhan, X. Fritel, A. Germa, V. Goua, R. Hankard, B. Heude, M. Kaminski, B. Larroque†, N.
17 Lelong, J. Lepeule, G. Magnin, L. Marchand, C. Nabet, F. Pierre, R. Slama, M.J. Saurel-Cubizolles,
18 M. Schweitzer, O. Thiebaugeorges.

19
20 * **Corresponding author**

21 Judith van der Waerden

22 Pierre Louis Institute of Epidemiology and Public Health,

23 INSERM & Sorbonne Université

24 27 Rue de Chaligny

25 75571 Paris Cedex 12, France

26 Tel: +33 (0) 1 85 56 02 42

27 Mail : judith.van-der-waerden@inserm.fr

28
29 **Keywords:** domain-specific physical activity, sedentary behavior, pregnancy, postpartum depression

30 **Word count main text/abstract:** 4044/248

1 **Abstract**

2 The potential beneficial effects of physical activity during pregnancy on postpartum depressive
3 symptoms (PPD) remain inconclusive. Using data from two prospective French birth cohorts, we
4 aimed to examine the relationship between domain-specific physical activity (including leisure-time
5 sedentary behavior) in pregnancy and the occurrence of postpartum depressive symptoms. Participants
6 of the ELFE cohort (N=15538) completed the Pregnancy Physical Activity Questionnaire (PPAQ),
7 which assesses the following physical activity/sedentary behavior domains: household/caregiving,
8 occupational, sports/exercise, transportation and leisure-time sedentary behavior during the third
9 pregnancy trimester. In the EDEN cohort (n = 1745) women completed the Baecke Questionnaire
10 (BQ) measuring occupational, sports/exercise, and leisure-time/locomotive activity during the first
11 trimester of pregnancy. Depressive symptoms in the first postpartum year were measured using the
12 Edinburgh Postnatal Depression Scale in both cohorts. Associations of physical activity/sedentary
13 behavior with PPD symptoms were determined by logistic regression analysis, with adjustment on
14 potential confounding factors. In the adjusted models, higher levels of household/ caregiving activities
15 (OR=1.10 (95% CI 1.01-1.19)) and leisure-time sedentary behavior (OR= 1.16 (95% CI 1.06-1.23)), in
16 the third pregnancy trimester were associated with an increased odds of PPD. No significant
17 associations were found for physical activity domains during the first pregnancy trimester. Higher
18 levels of household and caregiving activities and leisure time sedentary behavior in the last trimester
19 of pregnancy appear to increase the likelihood of postpartum depression. Purpose and context of
20 physical activity should be taken into account when encouraging physical activity as a strategy to help
21 prevent postpartum mental health problems from pregnancy onwards.

22

23

24

25

26

27

28

29

30

1 **Introduction**

2 Postpartum depression (PPD) is an important public health issue, with 2.6 to 35.0 % of women
3 experiencing major or minor depression sometime in the first year after giving birth (1, 2). PPD
4 negatively affects women's functioning, personal relationships, and children's social and cognitive
5 developmental outcomes (3). Engaging in regular exercise and physical activity has been demonstrated
6 as a successful treatment modality for major and minor depression in the general population (4), and
7 the potential benefits of exercise on perinatal depression are beginning to emerge as well. Several
8 systematic reviews and meta-analyses, found evidence that exercise may be effective in reducing
9 depression during both pregnancy (5) and the postpartum (6,7). Outcomes of a recent review, although
10 based on a small number of studies, suggest that physical activity during pregnancy may likewise
11 serve to prevent subsequent PPD (8).

12 Physical activity (PA) may include a variety of activities other than recreational ones, such as
13 occupational, transportation, care-giving and household activities (9). Leisure-time physical activity in
14 particular may be most beneficial in terms of mental health promotion and the prevention of mental ill-
15 health compared to other domains (10). The role of domain-specific PA is equally important when
16 studying its effects in reducing the risk for postnatal depression. Previous research has either focused
17 on future mothers' total or leisure-time physical activities, ignoring other domains, in particular
18 household and child-care activities, which comprise a substantial portion of women's PA (11).
19 Further, studies show that most pregnant women do not practice any form of exercise and many of
20 them decrease previously established levels of activity both in intensity and duration across domains
21 from the first to the third trimester (9, 12). Therefore, to determine their potential impact on PPD, it is
22 important to look at physical activity domains beyond total or leisure- time activities. To our
23 knowledge, the study by Demissie et al (13) is the only one to investigate the role of physical activity
24 domains during pregnancy in the development of depressive symptoms among postpartum women. In
25 a prospectively followed sample of 529 women, they found no significant associations between total
26 or domain-specific physical activity at either 17–22 weeks or 27–30 weeks' gestation and postpartum
27 depressive symptoms. Further studies with larger samples of women are needed to confirm the
28 influence of physical activity across different domains on postpartum depressive symptoms.

29 Likewise, emerging evidence indicates a relationship between sedentary behavior and
30 depression in both the general and pregnant populations (14, 15). Sedentary behavior is any waking
31 behavior characterized by an energy expenditure ≤ 1.5 metabolic equivalents (METs), while in a
32 sitting, reclining or lying posture (16). While it has been reported that pregnant women spent more
33 than 50% of their time (range 57% to 78%) in sedentary activities (17), research exploring sedentary
34 behaviors as a potential risk factor for postpartum depression is limited.

1 Taking these knowledge gaps into account, our main objective was to examine the relationship
2 between total and domain-specific physical activity in pregnancy and postpartum depressive
3 symptoms. We hypothesized that the purpose and context of physical activity might be more important
4 than the overall level of physical activity for reducing the presence of postnatal depressive symptoms
5 with, for instance, higher levels of leisure-time physical activity being associated with a decreased risk
6 of PPD, but higher levels of household/care-giving being associated with an increased risk of
7 postpartum depression. Our secondary objective was to explore the potential effect of leisure-time
8 sedentary behavior during pregnancy on postpartum depressive symptoms.

9 The results of this study will add to the limited literature on the role of domain-specific physical
10 activity and sedentary behavior in pregnancy and their associations with PPD. From a public health
11 perspective, pregnancy is a period wherein women may be more apt to incorporate changes into their
12 routine and offers a window of opportunity for prevention (18). Therefore, women may more readily
13 incorporate changes in PA habits during the pregnancy period (19). Our results may help to inform the
14 development of potentially feasible strategies for preventing depression in new mothers, as well as
15 refining public health recommendations concerning appropriate physical activity engagement for
16 mental health benefits.

17

18 **Methods**

19 ***Study design and sample***

20 The ELFE (*Etude Longitudinale Française depuis l'Enfance*) study is a multidisciplinary, nationally
21 representative birth cohort, which included 18,312 children born in 349 maternity units in France in
22 2011. The protocol, design and recruitment procedures of the ELFE survey have been previously
23 described (20). Participating mothers and children were recruited in maternity wards and have been
24 followed up since then. Exclusion criteria were stillbirth, birth <33 weeks of gestation, multiple births,
25 plans to move out of mainland France in the following 3 years. Mothers had to be capable of giving
26 informed consent in one of the study languages (French, English, Arabic, and Turkish). Maternity data
27 were collected using standardized interviews conducted by trained interviewers and through self-
28 completed questionnaires. Information about obstetrical characteristics was collected from maternal
29 recall and from medical records. At two months postpartum, data were collected via phone interviews.
30 The ELFE study received approval from bodies overseeing ethical aspects of data collection in France
31 (*Comission nationale informatique et liberté-CNIL* and *Comité Consultatif sur le Traitement des*
32 *Informations pour la Recherche en Santé – CCTIRS*).

33 The EDEN (*Etude sur les déterminants pré et postnataux précoces du Développement et de la santé de*
34 *l'ENfant*) mother-child cohort assesses the pre- and postnatal nutritional, social, and environmental
35 determinants of infant and child development and health (21). Pregnant women (n=2002) were

1 recruited before 24 weeks of gestation from two maternity wards (Poitiers and Nancy University
2 hospitals) between September 2003 and January 2006. Exclusion criteria were multiple pregnancies, a
3 known history of diabetes, inability to speak and read French or plans to move out of the study region
4 in the following 3 years. During pregnancy and after birth, socio-demographic and biomedical data on
5 the mother and child were gathered from a) medical records, b) face-to-face interviews and c) self-
6 completed questionnaires. Written consent was obtained from the mother for herself at inclusion and
7 for her newborn child after delivery. The EDEN study was approved by the *Comité Consultatif de*
8 *Protection des Personnes pour la Recherche Biomédicale* (CPP, Ethics Committee, Kremlin Bicêtre
9 Hospital) and by the *CNIL*.

10

11 *Symptoms of postpartum depression*

12 Depressive symptoms during the first year after the child's birth were assessed using the Edinburgh
13 Postnatal Depression Scale (EPDS), a 10-item questionnaire designed to detect postnatal depression
14 (score ranging 0-30) (22). In this study we used the cut-off ≥ 12 , which has been recommended for the
15 French translation and has demonstrated its validity for research purposes (23). EPDS scores were
16 available at 2 months postpartum for the ELFE cohort and at 4, 8 and 12 months postpartum for the
17 EDEN cohort.

18

19 *Physical activity measures*

20 In the ELFE cohort, 15, 575 women completed the Pregnancy Physical Activity Questionnaire
21 (PPAQ), a self-administered questionnaire specifically designed to assess physical activity levels
22 among pregnant women and validated in this population (24, 25). Respondents reported the time spent
23 participating in 32 activities during the last three months of pregnancy characterizing
24 household/caregiving, occupational settings, sports/exercise, transportation, and "inactivity". This last
25 domain includes questions regarding the following activities: watching TV or a video, sitting and
26 using a computer or writing (while not at work) and sitting and reading, talking or on the phone (while
27 not at work). As these items are typical sedentary behaviors, we will refer to this domain as leisure-
28 time sedentary activity. Self-reported time spent in each activity is multiplied by its intensity
29 (pregnancy-specific estimates for walking and light- to moderate intensity household tasks, and
30 Ainsworth compendium-based Metabolic Equivalent of Task (MET) values (26) for the remainder of
31 the PPAQ activities) to arrive at a measure of average weekly energy expenditure (MET-hr/wk)
32 attributable to each activity and domain or as total activity (sum of all domain scores).

33 In the EDEN cohort, women completed an adapted French version of the Baecke
34 Questionnaire (BQ), a validated measure of habitual physical activity for adults (27, 28). This
35 questionnaire was administered at the 24-28 week pregnancy visit and refers to the frequency of
36 physical activity performed during the first trimester of pregnancy. BQ includes a total of 16 questions
37 scored on a five-point Likert scale and classified into three domains: Occupational, Sports, and

1 Leisure-time activity. The Occupational and Sports scales include some comparative and self-
2 analytical questions that have previously been reported as difficult to answer (29) and have been
3 omitted from the French version. Scoring for the three domains follows the system described by
4 Baecke et al (27) with scores ranging from 1 (low) to 5 (high). The three domain scores can be
5 summed to create an overall score of physical activity (range 3-15). Additionally, women reported
6 whether they had interrupted their regular sport practice since the beginning of their pregnancy.

8 *Covariates*

9 Our analyses were adjusted for several socio demographic, psychosocial and pregnancy-related
10 characteristics identified in the literature and significantly associated ($p < .10$) with both physical
11 activity and postnatal depressive symptoms in our samples. *Sociodemographic* characteristics included
12 maternal age at delivery, nationality (French vs. other), highest educational level, family situation
13 (married/cohabitating with partner vs. separated), employment in pregnancy (no vs. yes), average
14 household income in Euros per month (ordered categorically: 1: <800, 2: 800-1500, 3: 1500-2300, 4:
15 2300-3000, 5: 3000-3800, 6: >3800), financial difficulties (no vs. yes) and study center. *Psychosocial*
16 *characteristics* included history of mental health problems (no vs. yes), psychological distress during
17 pregnancy (no vs. yes), maternal smoking and alcohol use during pregnancy, social support (no vs.
18 yes). *Health and pregnancy related characteristics* included presence of chronic health conditions,
19 maternal pre-pregnancy body mass index (BMI) (kg/m²), average gestational weight gain (kg), parity,
20 at least one pregnancy complication (derived from medical chart, no vs. yes), hospitalization during
21 pregnancy (no vs. yes). Finally, we included prenatal care utilization (n prenatal visits/scans) and
22 attendance to birth preparation classes (no vs. at least once).

24 *Statistical analyses*

25 Logistic regression modeling was utilized to calculate odds ratios (OR) and 95% confidence intervals
26 (CIs) for associations between total physical activity and domain-specific scores on the one hand, and
27 postpartum depressive symptoms on the other hand. Unadjusted and adjusted logistic regression
28 models were computed with the adjusted model controlling for previously listed covariates. Outcomes
29 were analyzed separately for the EDEN and ELFE cohorts. To be able to interpret the ORs of the
30 ELFE study in a clinically meaningful way, they are based on an 1.5 MET hr/wk increase in physical
31 activity, which corresponds, for instance, to 30 minutes of moderate intensity activity per week. Rates
32 of missing data on maternal, child and family characteristics ranged from 0.2% to 35.9%
33 (psychological distress in pregnancy), and were accounted for using multiple imputations models with
34 fully conditional specification. Excluding individuals with missing data from our analyses did not
35 significantly alter our results. Analyses were conducted with SPSS version 20 and SAS 9.4.

36

1 **Results**

2 Data on postpartum depression were available for 15,538 and 1,745 women in the ELFE and EDEN
3 cohorts respectively. In the ELFE study, factors associated with attrition from study baseline included
4 parental low occupational grade and unemployment, young maternal age and multiparity. For EDEN,
5 attrition was highest in young mothers, those with low educational level, low income and financial
6 difficulties, of non-French origin, who did not live with the father of their child, as well as those with
7 prenatal depression. Sociodemographic and pregnancy related characteristics of both samples are
8 summarized in **Table 1**. Women were on average 30 years old at time of delivery and mainly of
9 French origin (91.7% for ELFE and 98.1% for EDEN). The majority of women was married or
10 cohabiting (94.5% and 92.1%), had on average 13 years of education and was employed at the time of
11 enrolment (77.8% and 77.2%). About 11% reported having experienced mental health problems
12 before pregnancy and 13% reported psychological distress during pregnancy.

13 -- TABLE 1—

14

15 Prevalence of depressive symptoms was 12.3% at 2 months postpartum in the ELFE cohort,
16 and ranged between 10.6% and 12.60 in the year following the birth for women in the EDEN cohort.
17 During the first pregnancy trimester 22.9% of EDEN women practiced at least one sport/exercise
18 activity. However, by the second trimester 29.8% of expectant mothers reported interrupting their pre-
19 pregnancy sports practice. ELFE cohort women were spending most of their total physical activity in
20 the third trimester on household/caregiving (46.4%) and occupational activities (14.6%) and were
21 spending the smallest proportion on sports/exercise (3.6%). Moreover, a considerable proportion
22 (23.6%) was dedicated to sedentary behaviors.

23

24 In the ELFE cohort, univariate analyses showed significant associations between postpartum
25 depression at 2 months and women's Total physical activity (OR=1.20 (95% CI 1.11-1.26)) during the
26 last 3 months of pregnancy (Table 2). These associations remained significant in the multivariate
27 model (OR=1.12 (95% CI 1.03-1.19)). In the EDEN cohort, the physical activity sum score was
28 inversely associated with odds for depressive symptoms in the first postpartum year (OR^{4 months} = 0.79
29 (95% CI 0.67-0.94); OR^{8 months} = 0.79 (95% CI 0.63-0.99); OR^{12 months} = 0.78 (95% CI 0.64-0.94)) in
30 unadjusted analyses, but these became non-significant after adjustment.

31

32 When examining effects of specific activity domains in the ELFE cohort, Household/
33 caregiving activities (OR=1.23 (95% CI 1.16-1.31)) and Transportation (OR=1.16 (95% CI 1.08-
34 1.21)), as well as Leisure-time sedentary behavior (OR=1.16 95% CI 1.07-1.22)) were significantly
35 associated with postpartum depression in the univariate analyses. In the multivariate model,
36 associations remained significant for Household/ caregiving activities (OR=1.10 (95% CI 1.01-1.19))
37 and Leisure-time sedentary behavior (OR= 1.16 95% CI 1.06-1.23)).

1 In the EDEN cohort, occupational physical activity during the first trimester was inversely
2 associated with odds for depressive symptoms in the first postpartum year while no associations were
3 found for Leisure time/locomotive and Sport activities in the unadjusted logistic regression models.
4 These effects all became statistically non-significant after taking into account covariates (**table 2**).

5 -- TABLE 2--

6
7 In sensitivity analyses, we re-ran all adjusted models for domain-specific measures of physical
8 activity, adding a covariate for all physical activity outside of the specific domain of interest to
9 account for potential confounding. The models did not substantially change when other domains were
10 included as covariates. Further, we tested for possible differential effects of physical activity between
11 women already reporting psychological difficulties during pregnancy and those who did not. The
12 relationship between physical activity and PPD did not change when stratifying by prenatal
13 psychological difficulties (data available upon request).

14 15 **Discussion**

16 Using data from two large prospective birth cohorts, we explored the relationship between domain-
17 specific physical activity as well as sedentary behavior in pregnancy and postpartum depressive
18 symptoms. No significant associations were found between domain-specific physical activity during
19 the first pregnancy trimester and postpartum depressive symptoms. However, higher levels of
20 Household/ caregiving activities, Leisure-time sedentary behavior and Total activities in the third
21 trimester were associated with increased likelihood of depressive symptoms at 2 months postpartum.
22 Overall, our findings suggests that physical activity is not automatically associated with greater mental
23 health and reduced mental ill-health, and that contextual factors associated with several physical
24 activity domains are probably crucial to such relationships.

25 26 *Domain specific physical activity and postpartum depression*

27 Few studies have examined the association between PA in pregnancy and postpartum depression. In
28 addition, the majority of prior studies focused on leisure-time activity and did not examine other
29 activity domains. While we found no associations between domain-specific physical activity in the
30 first trimester and depressive symptoms in the first postpartum year, women spending more time on
31 household and caregiving activities during the third pregnancy trimester have higher odds for PPD.
32 For every additional 30 minutes per week of household and caregiving activities, women have a 10%
33 increased likelihood of experiencing postpartum depression. It should be noted though that this risk
34 remains modest and that our large sample partly contributed to finding significant associations.
35 Importantly, and consistent with previous cross-sectional studies (30, 31), household/caregiving
36 activities constituted the largest source of physical activity energy expenditure in our population. It has
37 been suggested that household activities, in comparison to leisure-time activities, are not usually

1 performed for enjoyment or by choice and could be stressful or burdensome, thereby contributing to
2 women's depressive symptoms (10, 32). Molarius et al (33) reported that the odds of being depressed
3 or anxious increased as the more burdensome the participants rated their domestic work. This might
4 especially be true in the third trimester, when pregnancy related complaints such as tiredness, pain,
5 and a growing belly are more frequent. To tease out whether the timing of prenatal physical activity
6 might impact the risk of postnatal depression, it may be important to assess levels of PA over the
7 entire length of pregnancy or even from pre-pregnancy onwards.

8
9 A further key finding from this study is that women who engage in more sedentary behaviors
10 at the end of their pregnancy have an increased likelihood of postpartum depression. This is of
11 particular importance given the fact that pregnant women spend at least 70% of time awake in
12 sedentary behaviors (34) and that sedentary time increases even in those that maintain recommended
13 physical activity levels during pregnancy (30). We are aware of only one other study that examined the
14 association between sedentary behavior in pregnancy and postpartum depression risk. Guida et al (35)
15 reported that sedentary behavior during pregnancy increases the odds of PPD symptoms by
16 approximately 34%, compared to 10% in our sample. However, they defined antenatal sedentary
17 behavior as engaging in less than one day of physical activity per week, while the inactivity score on
18 the PPAQ questionnaire corresponds more closely to the definition for sedentary behaviors (16).
19 Psychosocial mechanisms have been suggested as possible mediators in the association between
20 sedentary behaviors and depression with passive sedentary activities encouraging social isolation (14).
21 As research investigating the association between sedentary behavior and postnatal depressive
22 symptoms remains limited, further studies are needed to confirm this relationship.

23
24 Previous studies indicate that leisure-time physical activities pursued before or after birth reduce the
25 risk for postpartum depression (36, 37). They possibly results in changes in physical self-worth and
26 self-esteem due to mastering new tasks, having a greater sense of personal control, or distraction from
27 negative or more stressful aspects of everyday life (11). Additionally, recreational exercise may be
28 performed in a group, which could be a source of support for pregnant women by reducing their
29 symptoms through social mechanisms. Yet, our results did not indicate a protective effect of leisure-
30 time/sport activities in either the first or the third trimester on the occurrence of postpartum
31 depression. This could be attributed to the fact that activities in the sports/exercise domains
32 contributed relatively little to the total activity score, especially during the last trimester of pregnancy.
33 While 22.9% of the EDEN participants practiced at least one sport at the beginning of their pregnancy,
34 a considerable part discontinued their habitual sports practice subsequently. Some women may
35 discontinue physical activity because of physical or emotional difficulties that may predict depression
36 risk. As discontinuation of sport practice was not associated with previous or current mental health
37 problems for the EDEN participants, future research is needed to confirm these hypotheses.

1 *Methodological considerations*

2 Previously, Demissie et al (13) failed to establish any correlation between domain-specific activity
3 during pregnancy and risk for postpartum depression. As they evaluated levels of moderate to
4 vigorous physical activity, few women in their sample reached these activity levels resulting in low
5 variation in the total activity score. By taking into account all possible PA intensities in our analyses,
6 we were able to evaluate more globally habitual activity levels in pregnant women. Also, the
7 proportion of physical activity across domains reported by Demissie differed from our study, with a
8 smaller percentage of women engaging in household or caregiving activities. While reporting on
9 physical activity in the first trimester gives a reasonable representation of women's habitual level of
10 physical activity, decreasing levels of overall and recreational activity during pregnancy have been
11 reported in previous studies (38, 39). Unfortunately, we did not have information on the evolution of
12 physical activity scores on the BQ questionnaire throughout pregnancy. Conversely, it is very likely
13 that participants of the ELFE cohort who were physically active in their third trimester were equally so
14 during their first trimester.

15
16 *Strengths and limitations*

17 Our study has several strengths, including its prospective design and the use of two birth cohorts,
18 making it one of very few studies of this magnitude to study physical activity in pregnancy and
19 subsequent depression. We used validated physical activity measures assessing the type, frequency,
20 and duration of different domains of activity and studied their association with subsequent depression,
21 which has rarely been done in a population of perinatal women. Finally, we controlled for a number of
22 important confounding factors, including prenatal depression. However, we need to acknowledge
23 some limitations. First, both cohorts suffered from selective attrition, which might have impacted the
24 generalizability of our findings to more disadvantaged populations. Secondly, maternal depressive
25 symptoms were ascertained using women's self-reports, rather than clinical diagnoses. However, the
26 EPDS was designed with the purpose of identifying women who are depressed after childbirth and is
27 widely used as a screening tool for PPD, with satisfactory sensitivity, specificity and positive
28 predictive value (40). Third, physical activity measurement relied on self-report tools which might be
29 subject to recall bias. Actual measures of physical activity or energy expenditure, such as activity
30 monitoring using an accelerometer, heart rate monitor or double-labeled water could give more
31 objective outcomes (41). However, these measures do not make it possible to characterize different
32 domains of physical activity, which might be more pertinent for mental health effects than their
33 intensity (42). Also, the Baecke Questionnaire was not specifically designed for use with pregnant
34 women and did not include household and caregiving activities or inactivity which are particularly
35 relevant for pregnant populations. Furthermore, the leisure-time activity score of the BQ is a mixture
36 of sport activities, active locomotive activities and a sedentary behavior. This makes the score on this
37 domain difficult to interpret compared to the distinct domains described by the PPAQ questionnaire.

1 which is one of the few validated questionnaires designed to assess physical activity among pregnant
2 women. Finally, despite its prospective design and the control for a large number of potential
3 confounders, we cannot entirely rule out the possibility of residual confounding, in particular due to
4 non-measured postnatal characteristics (for instance maternal sleep quality). However, we did control
5 for prior history of mental health problems and psychological distress during pregnancy, making it less
6 plausible that our results are due to reverse causality.

7

8 **Conclusions**

9 Our results add to the very limited number of studies examining the relationship between physical
10 activity domains during pregnancy and levels of depressive symptoms postnatally. Our study does not
11 support findings from previous work that physical activity during pregnancy may help ameliorate
12 mood disorders in postpartum mothers once confounding factors have been taken into account.
13 Additionally, higher levels of household and caregiving activities and in particular sedentary behavior
14 appear to increase the risk of postpartum depression. This indicates that it is important for studies on
15 physical activity and mental health to look beyond indicators of total energy expenditure and also
16 consider more nuanced aspects of the context in which PA is performed (43). Nevertheless, remaining
17 active during pregnancy provides benefits for maternal health and quality of life and may also have a
18 positive effect on fetal growth and fetal adaptation (44). Moreover, physical activity promotion efforts
19 targeting this period may be linked to reduced risk of antenatal depression (5), which in turn is an
20 important predictor for postnatal depression (1). While current recommendations for exercise during
21 pregnancy support moderate intensity physical activity for a minimum of 15–30 minutes at least 3–5
22 times weekly (45), pregnant women are generally inclined to reduce their physical activity levels.
23 Health care providers may need to be careful when recommending increasing physical activity as its
24 purpose and context appears to be important in promoting mental wellbeing. Further, targeting
25 reductions in sedentary behavior may be a propitious strategy for better mental health in this
26 population. This may seem more attainable than participating in physical activity toward the end of
27 pregnancy when a mother is likely to be most uncomfortable. Theory-based, face-to-face, telephone,
28 and print-based lifestyle interventions effectively increase exercise levels among sedentary adults and
29 could potentially be effective for pregnant and postpartum women (46). Additionally, modifications in
30 sedentary behavior may be more easily maintained and could represent an important strategy to help
31 manage postpartum mental health problems among women.

32

33

34

35

36

37

1 Literature

- 2 1. Banti S, Mauri M, Oppo A, Borri C, Rambelli C, Ramacciotti D, et al. From the third month
3 of pregnancy to 1 year postpartum. Prevalence, incidence, recurrence, and new onset of depression.
4 Results from the Perinatal Depression–Research and Screening Unit study. *Compr Psychiatry*.
5 2011;52(4):343-51.
- 6 2. Norhayati MN, Hazlina NH, Asrenee AR, Emilin WM. Magnitude and risk factors for
7 postpartum symptoms: a literature review. *J Affect Dis*. 2015;175:34-52.
- 8 3. Letourneau NL, Dennis CL, Benzies K, Duffett-Leger L, Stewart M, Tryphonopoulos PD, et
9 al. Postpartum depression is a family affair: addressing the impact on mothers, fathers, and children.
10 Issues Ment Health Nurs. 2012; 33(7):445-57.
- 11 4. Cooney GM, Dwan K, Greig CA, Lawlor DA, Rimer J, Waugh FR, et al. Exercise for
12 depression. The Cochrane database of systematic reviews. 2013(9):CD004366.
- 13 5. Daley AJ, Foster L, Long G, Palmer C, Robinson O, Walmsley H, et al. The effectiveness of
14 exercise for the prevention and treatment of antenatal depression: systematic review with meta-
15 analysis. *BJOG*. 2015;122(1):57-62.
- 16 6. Pritchett RV, Daley AJ, Jolly K. Does aerobic exercise reduce postpartum depressive
17 symptoms? a systematic review and meta-analysis. *Br J Gen Pract*. 2017;67(663):e684.
- 18 7. McCurdy AP, Boulé NG, Sivak A, Davenport MH. Effects of exercise on mild-to-moderate
19 depressive symptoms in the postpartum period: a meta-analysis. *Obstet Gyn*. 2017;129(6):1087-97.
- 20 8. Teychenne M, York R. Physical activity, sedentary behavior, and postnatal depressive
21 symptoms: a review. *Am J Prev Med*. 2013;45(2):217-27.
- 22 9. Borodulin KM, Evenson KR, Herring AH. Physical activity patterns during pregnancy through
23 postpartum. *BMC Women's Health*. 2009;9:32.
- 24 10. White RL, Babic MJ, Parker PD, Lubans DR, Astell-Burt T, Lonsdale C. Domain-Specific
25 Physical Activity and Mental Health: A Meta-analysis. *Am J Prev Med*. 2017;52(5):653-66.
- 26 11. Kull M, Ainsaar M, Kiive E, Raudsepp L. Relationship between low depressiveness and
27 domain specific physical activity in women. *Health Care Women Int*. 2012;33(5):457-72.
- 28 12. Gaston A, Cramp A. Exercise during pregnancy: a review of patterns and determinants. *J Sci*
29 *Med Sport*. 2011;14(4):299-305.
- 30 13. Demissie Z, Siega-Riz AM, Evenson KR, Herring AH, Dole N, Gaynes BN. Physical activity
31 during pregnancy and postpartum depressive symptoms. *Midwifery*. 2013;29(2):139-47.
- 32 14. Padmapriya N, Bernard JY, Liang S, Loy SL, Shen Z, Kwek K, et al. Association of physical
33 activity and sedentary behavior with depression and anxiety symptoms during pregnancy in a
34 multiethnic cohort of Asian women. *Arch Womens Ment Health*. 2016;19(6):1119-28.
- 35 15. de Wit L, van Straten A, Lamers F, Cuijpers P, Penninx B. Are sedentary television watching
36 and computer use behaviors associated with anxiety and depressive disorders? *Psychiatry Res*.
37 2011;186(2-3):239-43.
- 38 16. Tremblay MS, Aubert S., Barnes JD, Saunders TJ, Carson V, Latimer-Cheung AE et al,
39 Sedentary Behavior Research Network (SBRN) - Terminology Consensus Project process and
40 outcome. *Int J Behav Nutr Phys Act*. 2017;14(1), 75.
- 41 17. Fazzi C, Saunders DH, Linton K, Norman JE, Reynolds RM. Sedentary behaviours during
42 pregnancy: a systematic review. *IJBNPA*. 2017;14(1):32.
- 43 18. Edvardsson K, Ivarsson A, Eurenus E, Garvare R, Nystrom ME, Small R, et al. Giving
44 offspring a healthy start: parents' experiences of health promotion and lifestyle change during
45 pregnancy and early parenthood. *BMC Public Health*. 2011;11:936.
- 46 19. Cioffi J, Schmied V, Dahlen H, Mills A, Thornton C, Duff M, et al. Physical activity in
47 pregnancy: women's perceptions, practices, and influencing factors. *J Midwifery Womens Health*.
48 2010;55(5):455-61.
- 49 20. Vandentorren S, Bois C, Pirus C, Sarter H, Salines G, Leridon H. Rationales, design and
50 recruitment for the Elfe longitudinal study. *BMC Pediatrics*. 2009;9:58.
- 51 21. Heude B, Forhan A, Slama R, Douhaud L, Bedel S, Saurel-Cubizolles MJ, et al. Cohort
52 Profile: The EDEN mother-child cohort on the prenatal and early postnatal determinants of child
53 health and development. *Int J Epidemiol*. 2016;45:353-63.

- 1 22 Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-
2 item Edinburgh Postnatal Depression Scale. *Br J Psychiatry*. 1987;150:782-6.
- 3 23. Guedeney N, Fermanian J. Validation study of the French version of the Edinburgh Postnatal
4 Depression Scale (EPDS): new results about use and psychometric properties. *Eur Psychiatry*.
5 1998;13(2):83-9.
- 6 24. Chasan-Taber L, Schmidt MD, Roberts DE, Hosmer D, Markenson G, Freedson PS.
7 Development and validation of a Pregnancy Physical Activity Questionnaire. *Med Sci Sports Exerc*.
8 2004;36(10):1750-60.
- 9 25. Chandonnet N, Saey D, Almeras N, Marc I. French Pregnancy Physical Activity
10 Questionnaire compared with an accelerometer cut point to classify physical activity among pregnant
11 obese women. *PloS one*. 2012;7(6):e38818.
- 12 26. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Jr., Tudor-Locke C, et al.
13 2011 Compendium of Physical Activities: a second update of codes and MET values. *Med Sci Sports
14 Exerc*. 2011;43(8):1575-81.
- 15 27. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual
16 physical activity in epidemiological studies. *Am J Clin Nutr*. 1982;36(5):936-42.
- 17 28. Bigard AX, Duforez F, Portero P, Gezenec CY. Détermination de l'activité physique par
18 questionnaire: validation du questionnaire de Baecke. *Sci Sport*. 1992;7:215-21.
- 19 29. Pols, M. A., Peeters, P. H., Kemper, H. C., & Collette, H. J. (1996). Repeatability and relative
20 validity of two physical activity questionnaires in elderly women. *Med Sci Sports Exerc*, 28(8), 1020-
21 1025.
- 22 30. Huberty JL, Buman MP, Leiferman JA, Bushar J, Adams MA. Trajectories of objectively-
23 measured physical activity and sedentary time over the course of pregnancy in women self-identified
24 as inactive. *Prev Med Rep*. 2016;3:353-60.
- 25 31. Nascimento SL, Surita FG, Godoy AC, Kasawara KT, Morais SS. Physical Activity Patterns
26 and Factors Related to Exercise during Pregnancy: A Cross Sectional Study. *PloS one*.
27 2015;10(6):e0128953.
- 28 32. Asztalos M, Wijndaele K, De Bourdeaudhuij I, Philippaerts R, Matton L, Duvigneaud N, et al.
29 Specific associations between types of physical activity and components of mental health. *J Sci Med
30 Sport*. 2009;12(4):468-74.
- 31 33. Molarius, A., Berglund, K., Eriksson, C., Eriksson, H. G., Linden-Bostrom, M., Nordstrom, et
32 al. Mental health symptoms in relation to socio-economic conditions and lifestyle factors--a population
33 based study in Sweden. *BMC Public Health*. 2009; 9, 302.
- 34 34. Di Fabio DR, Blomme CK, Smith KM, Welk GJ, Campbell CG. Adherence to physical
35 activity guidelines in mid-pregnancy does not reduce sedentary time: an observational study. *IJBNPA*.
36 2015;12:27.
- 37 35. Guida J, Sundaram S, Leiferman J. Antenatal physical activity: Investigating the effects on
38 postpartum depression. *Health*. 2012;4:1276-86.
- 39 36. Daley AJ, Jolly K, MacArthur C. The effectiveness of exercise in the management of post-
40 natal depression: systematic review and meta-analysis. *Fam Pract*. 2009;26(2):154-62.
- 41 37. Strom M, Mortensen EL, Halldorson TI, Osterdal ML, Olsen SF. Leisure-time physical
42 activity in pregnancy and risk of postpartum depression: a prospective study in a large national birth
43 cohort. *J Clin Psychiatry*. 2009;70(12):1707-14.
- 44 38. Borodulin KM, Evenson KR, Wen F, Herring AH, Benson AM. Physical activity patterns
45 during pregnancy. *Med Sci Sports Exerc*. 2008;40(11):1901-8.
- 46 39. Haakstad LA, Voldner N, Henriksen T, Bo K. Why do pregnant women stop exercising in the
47 third trimester? *Acta Obstet Gynecol Scand*. 2009;88(11):1267-75.
- 48 40. Gibson, J., McKenzie-McHarg, K., Shakespeare, J., Price, J., & Gray, R. (2009). A systematic
49 review of studies validating the Edinburgh Postnatal Depression Scale in antepartum and postpartum
50 women. *Acta Psychiatr Scand*, 119(5), 350-364.
- 51 41. Aparicio-Ugarriza R, Mielgo-Ayuso J, Benito PJ, Pedrero-Chamizo R, Ara I, Gonzalez-Gross M.
52 Physical activity assessment in the general population; instrumental methods and new technologies.
53 *Nutricion Hospitalaria*. 2015;31 Suppl 3:219-26.

- 1 42. Evenson KR, Chasan-Taber L, Symons Downs D, Pearce EE. Review of self-reported
2 physical activity assessments for pregnancy: summary of the evidence for validity and reliability.
3 *Paediatr Perinat Epidemiol*. 2012;26(5):479-94.
- 4 43. Silva KS, Garcia LM, Rabacow FM, de Rezende LF, de Sa TH. Physical activity as part of daily
5 living: Moving beyond quantitative recommendations. *Prev Med*. 2017;96:160-2.
- 6 44. Nascimento SL, Surita FG, Cecatti JG. Physical exercise during pregnancy: a systematic
7 review. *Curr Opin Obstet Gynecol*. 2012;24(6):387-94.
- 8 45. Evenson KR, Barakat R, Brown WJ, Dargent-Molina P, Haruna M, Mikkelsen EM, et al.
9 Guidelines for Physical Activity during Pregnancy: Comparisons From Around the World. *Am J*
10 *Lifestyle Med*. 2014;8(2):102-21.
- 11 46. Currie S, Sinclair M, Murphy MH, Madden E, Dunwoody L, Liddle D. Reducing the decline
12 in physical activity during pregnancy: a systematic review of behaviour change interventions. *PLoS*
13 *one*. 2013;8(6):e66385.

14 **Disclosure of interest**

15 The authors declare that they have no conflicts of interest concerning this article.

16

17 **Funding**

18 The Elfe survey is a joint project between the French Institute for Demographic Studies (INED) and
19 the National Institute of Health and Medical Research (INSERM), in partnership with the French
20 blood transfusion service (Etablissement français du sang, EFS), Santé publique France, the National
21 Institute for Statistics and Economic Studies (INSEE), the Direction générale de la santé (DGS, part of
22 the Ministry of Health and Social Affairs), the Direction générale de la prévention des risques (DGPR,
23 Ministry for the Environment), the Direction de la recherche, des études, de l'évaluation et des
24 statistiques (DREES, Ministry of Health and Social Affairs), the Département des études, de la
25 prospective et des statistiques (DEPS, Ministry of Culture), and the Caisse nationale des allocations
26 familiales (CNAF), with the support of the Ministry of Higher Education and Research and the Institut
27 national de la jeunesse et de l'éducation populaire (INJEP). Via the RECONAI platform, it receives a
28 government grant managed by the National Research Agency under the "Investissements d'avenir"
29 programme (ANR-11-EQPX-0038).

30 The EDEN study was funded by Fondation pour la Recherche Médicale (FRM); French
31 Ministry of Research: IFR Program; INSERM Human Nutrition National Research Program and
32 Diabetes National Research Program (through a collaboration with the French Association of Diabetic
33 Patients (AFD)); French Ministry of Health, French Agency for Environment Security (AFSSET);
34 French National Institute for Population Health Surveillance (InVS); Paris-Sud University; French
35 National Institute for Health Education (INPES); Nestlé; Mutuelle Générale de l'Education Nationale
36 (MGEN); French speaking association for the study of diabetes and metabolism (ALFEDIAM);
37 National Agency for Research (ANR nonthematic program) and National Institute for Research in
38 Public Health (IRESP: TGIR Cohorte Santé 2008 programme).

39 The funding sources were not involved in the study design or in the collection, analysis and
40 interpretation of data.

Table 1 Characteristics of ELFE and EDEN cohort study participants (N (%) or mean (sd)), France 2003-2011				
	ELFE (n=15538)		EDEN (n=1745)	
	N (%)	mean (SD)	N (%)	mean (SD)
Sociodemographic characteristics				
Maternal age at delivery (years)		30.64 (5.06)		29.48 (4.89)
Nationality (Other than French)	1467 (8.27)		33 (1.89)	
Highest educational level (years)		13.33 (2.29)		13.58 (2.67)
Family situation (Separated)	978 (5.50)		151 (7.93)	
Employment in pregnancy (no)	3875 (22.21)		436 (22.8)	
Household income (1-6)		4.36 (1.34)		3.71 (1.31)
Financial difficulties (Yes)	8888 (49.13)		382 (20.02)	
Psychosocial characteristics				
History of mental health problems (yes)	1280 (10.79)		209 (10.96)	
Psychological distress during pregnancy (yes)	2222 (12.56)		264 (13.84)	
Substance use (smoking/alcohol) during pregnancy (yes)	4526 (25.36)		867 (49.68)	
Chronic health conditions (yes)	2391 (13.56)		293 (15.36)	
Social support (no)	1401 (8.93)		15 (0.85)	
Pregnancy related characteristics				
Parity				
0	8116 (45.94)		854 (44.80)	
1	6153 (34.78)		689 (36.11)	
2+	3407 (19.28)		264 (19.09)	
Pre-pregnancy body mass index (BMI) (kg/m ²)		23.48 (4.81)		23.27 (4.63)
Average gestational weight gain (kg)		13.17 (5.61)		13.41 (4.84)
Delivery type (C-section)	3173 (20.4)		301 (17.24)	
At least one pregnancy complication (yes)	5351 (30.49)		1102 (63.15)	
Hospitalization during pregnancy (yes)	2828 (18.20)		458 (26.24)	
Number of prenatal visits		8.60 (3.09)		8.67 (2.33)
Number of Echographies		4.70 (2.34)		4.13 (1.50)
Any birth preparation classes (no)	9765 (55.38)		859 (49.22)	
Physical activity during pregnancy * :				
Scores from Baecke Questionnaire (1st trimester of pregnancy) (range 1-5)				
Occupational				2.40 (1.30-3.40)
Leisure-time				2.25 (2.00-2.75)
Sports/exercise				1.00 (1.00-1.00)
Sum score of the above 3 scores; range 3-15				5.85 (4.40-6.90)
Scores from Pregnancy Physical Activity Questionnaire (3rd trimester of pregnancy) (MET-h.wk⁻¹)				
Household/caregiving		69.47 (41.1-110.25)		
Occupational		19.00 (0.00-49.46)		
Transportation		9.50 (3.60-19.00)		
Leisure-time sedentary behavior		35.70 (19.95-55.65)		
Sports/exercise		3.38 (1.16-8.45)		
Total (sum of the above 5 scores)		158.46 (116.97-211.32)		

Interruption of sport practice since the beginning of pregnancy (yes)	--		568 (<u>29.78</u>)	
Postpartum depression (EPDS) (<i>cut-off score ≥12</i>)				
2 months	1902 (<u>12.27</u>)		--	
4 months	--		240 (<u>12.60</u>)	
8 months	--		220 (<u>11.56</u>)	
12 months	--		202 (<u>10.60</u>)	
* Physical activity scores are presented as Median (IQR- interquartile range) due to their non-normal distribution				

Table 2 Univariate and multivariate logistic regression analysis between physical activity in pregnancy and postpartum depressive symptoms, France 2003-2011								
Physical activity during pregnancy	ELFE (n= 15,538)		EDEN (n=1,745)					
	PPD 2 months		PPD 4 months		PPD 8months		PPD 12 months	
	Unadjusted OR (95% CI)	Adjusted ¹ OR (95% CI)	Unadjusted OR (95% CI)	Adjusted ¹ OR (95% CI)	Unadjusted OR (95% CI)	Adjusted ¹ OR (95% CI)	Unadjusted OR (95% CI)	Adjusted ¹ OR (95% CI)
PPAQ scores (3 rd trimester of pregnancy) per 1 MET-h.wk increase								
Household/caregiving	<u>1.23 (1.16-1.31)</u>	<u>1.10 (1.01-1.19)</u>						
Occupational	<u>0.92 (0.82-0.99)</u>	<u>0.99 (0.91-1.08)</u>						
Transportation	<u>1.16 (1.08-1.21)</u>	<u>1.08 (1.00-1.15)</u>						
Leisure-time sedentary behavior	<u>1.16 (1.07-1.22)</u>	<u>1.16 (1.06-1.23)</u>						
Sports/exercise	<u>0.98 (0.90-1.06)</u>	<u>1.02 (0.93-1.09)</u>						
Total	<u>1.20 (1.11-1.26)</u>	<u>1.12 (1.03-1.19)</u>						
Baecke scores (1 st trimester of pregnancy) per 1 point increase								
Occupational			0.81 (0.66-1.00)	0.93 (0.69-1.24)	0.83 (0.67-1.04)	1.05 (0.81-1.36)	0.81 (0.69-0.96)	1.05 (0.84-1.31)
Leisure-time			0.91 (0.78-1.07)	1.01 (0.86-1.19)	0.89 (0.75-1.08)	1.02 (0.84-1.23)	0.81 (0.59-1.12)	0.90 (0.64-1.28)
Sports/exercise			0.94 (0.81-1.09)	0.99 (0.85-1.16)	0.93 (0.81-1.08)	0.98 (0.84- 1.15)	0.91 (0.75- 1.10)	0.95 (0.78-1.17)
Total			0.79 (0.67-0.94)	0.94 (0.79-1.12)	0.79 (0.63-0.99)	0.99 (0.77-1.28)	0.78 (0.64-0.94)	0.98 (0.76-1.28)

¹ Adjusted for: age at delivery, nationality, maternal educational level, family situation, average household income, financial difficulties, study center, employment in pregnancy, history of mental health problems, psychological distress in pregnancy, smoking and alcohol use during pregnancy, social support, chronic health problems, pre pregnancy BMI, averages gestational weight gain, parity, pregnancy complications; hospitalization during pregnancy, prenatal care utilization

PPD: Postpartum Depression; PPAQ: Pregnancy Physical Activity Questionnaire