

Fragmentations with self-similar branching speeds Jean-Jil Duchamps

▶ To cite this version:

Jean-Jil Duchamps. Fragmentations with self-similar branching speeds. 2019. hal-02179793v1

HAL Id: hal-02179793 https://hal.sorbonne-universite.fr/hal-02179793v1

Preprint submitted on 11 Jul 2019 (v1), last revised 26 Apr 2021 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fragmentations with self-similar branching speeds

Jean-Jil Duchamps

Sorbonne Université

July 11, 2019

Abstract

We consider fragmentation processes with values in the space of marked partitions of \mathbb{N} , i.e. partitions where each block is decorated with a nonnegative real number. Assuming that the marks on distinct blocks evolve as independent positive self-similar Markov processes and determine the speed at which their blocks fragment, we get a natural generalization of the self-similar fragmentations of [3]. Our main result is the characterization of these generalized fragmentation processes: a Lévy-Khinchin representation is obtained, using techniques from positive self-similar Markov processes and from classical fragmentation processes. We then give sufficient conditions for their absorption in finite time to a frozen state, and for the genealogical tree of the process to have finite total length.

Contents

1	Introduction	2
	1.1 Self-similar fragmentations	2
	1.2 Partitions with marks	4
2	Extended self-similar fragmentations	6
	2.1 Definitions, first properties	6
	2.2 Stopping lines, changing the index of self-similarity	10
3	Main results	11
	3.1 Decomposition of ESSF processes	11
	3.2 Absorption in finite time	14
Α	Proofs	16
	A.1 Proof of Proposition 2	16
	A.2 Proof of Proposition 10	16
	A.3 Proof of Proposition 11	19
	A.4 Proof of Theorem 13	20
	A.5 Proof of Proposition 16	30
References 3		

Keywords and phrases. self-similar; branching process; exchangeable; fragmentation; partition; random tree; Lévy process.

MSC 2010 Classification. 60G09,60G18,60J80,60G51,60J25.

1 Introduction

A fragmentation process is a system of particles evolving in time in a Markovian way, where each particle is assigned a mass and may dislocate at random times, distributing its mass among newly created particles. It is usually assumed that particles evolve independently of one another, in a way depending only on their mass. Self-similar fragmentations are processes where the speed of fragmentation of a particle is accelerated proportionally to a function of its mass – which then must be a power function, characterized by an exponent $\alpha \in \mathbb{R}$. These processes are said to be homogeneous when $\alpha = 0$. Homogeneous and self-similar fragmentations have been characterized in the early 2000s (see [2, 3], or [5] for a general introduction), and their connections to random trees have been developed in e.g. [1] or [14, 15].

These studies have been made under a conservative assumption, which prevents the total mass in the system from increasing. This assumption allows for instance the representation of fragmentation processes in terms of exchangeable partition-valued processes, which are convenient objects allowing one to naturally recover *discrete* genealogical structures in fragmentation processes.

The primary goal of this article is to extend the self-similar assumption while staying in a conservative setting. To this aim, we assume that particles are described by a pair mass-mark which evolves jointly in a Markovian way, such that a) the total mass does not increase, and b) it is now the *mark* – which may a priori fluctuate in any way – of a particle which determines the speed at which it fragments. The conservative assumption allows us to model this idea with Markov processes taking values in marked partitions of the integers, with very little restriction concerning marks. Consequently, if one ignores the masses of particles, our processes essentially give constructions for quite general non-conservative fragmentations. Related and inspiring works include self-similar branching Markov chains [17], the recent so-called branching Lévy processes of [7], as well as many recent developments which have been published on self-similar growth-fragmentation processes (see e.g. [10, 13, 21]), introduced by Bertoin [6], which allow masses of particles to fluctuate as a positive Markov process.

The article is organized as follows. In the remainder of the introduction, we recall some definitions and basic results of usual self-similar fragmentations, and define the space of marked partitions in which our processes live. In Section 2 we define our extended self-similar fragmentation (ESSF) processes, and point out their basic properties. We characterize ESSF processes with a type of Lévy-Khinchin representation in Section 3, and then give sufficient conditions for a process to almost surely a) reach an absorbing state in finite time b) have a genealogy where the sum of lengths of all branches is finite. Because most proofs are somewhat technical, we defer them to Appendix A to ease the exposition.

1.1 Self-similar fragmentations

To study processes with values in the space of partitions of \mathbb{N} , let us recall some classical notation and definitions. First define $[n] := \{1, 2, ..., n\}$ for $n \in \mathbb{N}$ and $[\infty] := \mathbb{N} := \{1, 2, ...\}$. Now for $n \in \mathbb{N} \cup \{\infty\}$, we denote by \mathcal{P}_n the space of partitions of [n]. We often see a partition $\pi \in \mathcal{P}_n$ as the equivalence relation \sim^{π} it represents on [n]. We will denote by $\mathbf{0}_n$ (resp. $\mathbf{1}_n$) the partition of [n] into singletons (resp. the partition with a single block $\{[n]\}$). We will often omit the subscript nand write only $\mathbf{0}$ or $\mathbf{1}$ when the context is clear.

For $n < m \leq \infty$ and $\pi \in \mathcal{P}_m$, we denote by $\pi_{|[n]}$ its restriction to the set $[n] \subset [m]$. \mathcal{P}_{∞} may be understood as the projective limit of the sets $(\mathcal{P}_n, n \in \mathbb{N})$, and as such, a natural metric which

makes this space compact may be defined on it by

$$d(\pi, \pi') = \sup\{n \in \mathbb{N}, \ \pi_{|[n]} = \pi'_{|[n]}\}^{-1},\$$

where by convention $(\sup \mathbb{N})^{-1} = 0$. We will consider the action of permutations of \mathbb{N} on \mathcal{P}_{∞} , and more generally we can define, for any $1 \leq n \leq m \leq \infty$, any *injection* $\sigma : [n] \to [m]$ and any $\pi \in \mathcal{P}_m$, the partition $\pi^{\sigma} \in \mathcal{P}_n$ defined by:

$$i \sim^{\pi^{\sigma}} j \iff \sigma(i) \sim^{\pi} \sigma(j), \quad i, j \in [n].$$

Note that in this paper, a permutation $\sigma : \mathbb{N} \to \mathbb{N}$ is a bijection with *finite* support $\{n \in \mathbb{N}, \sigma(n) \neq n\}$. We usually label the blocks of a partition $\pi = \{\pi_1, \pi_2, \ldots\}$ in the unique way such that the sequence $(\min \pi_k, k \ge 1)$ is increasing. This way, π_1 is necessarily the block containing 1, π_2 is the block containing the lowest integer not in the same block as 1, etc. By convention, if π has a finite number of blocks, say *K*, we define $\pi_{K+l} = \emptyset$ for all $l \ge 1$. It will be useful to define a fragmentation operator Frag : $\mathcal{P}_{\infty} \times (\mathcal{P}_{\infty})^{\mathbb{N}} \to \mathcal{P}_{\infty}$ by

Frag
$$(\pi, \pi^{(\cdot)}) = \{\pi_k \cap \pi_l^{(k)}, k, l \ge 1\},\$$

where (π_k) are the ordered blocks of π and $(\pi_l^{(k)})$ the ordered blocks of $\pi^{(k)}$. In words, blocks of the new partition are formed from the restriction of the *k*-th partition of the sequence $\pi^{(\cdot)}$ to π_k , for each $k \ge 1$.

Now let us recall the definition of partition-valued fragmentation processes (see e.g. [5]). For this definition, we restrict ourselves to the space of partitions that have asymptotic frequencies, i.e. $\pi \in \mathcal{P}_{\infty}$ such that for all $k \geq 1$,

$$|\pi_k| := \lim_{n \to \infty} \frac{\#\pi_k \cap [n]}{n}$$
 exists.

In this case, we write $|\pi|^{\downarrow}$ for the nonincreasing reordering of the sequence $(|\pi_1|, |\pi_2|, ...)$. Let us write \mathcal{P}'_{∞} for the space of partitions of \mathbb{N} with asymptotic frequencies.

Definition 1. A self-similar fragmentation process is a càdlàg Markov process ($\Pi(t), t \ge 0$) with values in \mathcal{P}'_{∞} , such that almost surely for all $k \in \mathbb{N}$, the map $t \mapsto |\Pi_k(t)|$ is right-continuous and for which the following properties hold.

(i) *Exchangeability*: for all $\pi \in \mathcal{P}'_{\infty}$, for all $\sigma : \mathbb{N} \to \mathbb{N}$ permutation,

$$(\Pi(t)^{\sigma}, t \ge 0)$$
 under $\mathbb{P}_{\pi} \stackrel{(a)}{=} (\Pi(t), t \ge 0)$ under $\mathbb{P}_{\pi^{\sigma}}$,

where \mathbb{P}_{π} denotes the distribution of the process started from π .

(ii) Self-similar branching: there exists $\alpha \in \mathbb{R}$ such that if (Ω, \mathbb{P}) is a probability space where $(\Pi^{(\cdot)}(t), t \ge 0)$ is a sequence of independent copies of the process started from 1, then for any $\pi \in \mathcal{P}'_{\infty}$, we have

$$(\Pi(t), t \ge 0) \text{ under } \mathbb{P}_{\pi} \stackrel{(d)}{=} (\operatorname{Frag}(\pi, \widetilde{\Pi}^{(\cdot)}(t)), t \ge 0) \text{ under } \mathbb{P},$$
(1)

where $\widetilde{\Pi}^{(\cdot)}$ is the sequence of time-changed processes defined by

$$\widetilde{\Pi}^{(k)}(t) = \Pi^{(k)}(|\pi_k|^{\alpha} t), \qquad k \ge 1, t \ge 0.$$

Note that a fragmentation with self-similarity index $\alpha = 0$ is called homogeneous. It is well-known (we refer to [5, Section 1 to 3] for a detailed account on the theory of partition-valued fragmentations) that self-similar fragmentations can be characterized in terms of their self-similarity index α ,

a so-called erosion coefficient $c \ge 0$ and a dislocation measure ν on the (metric and compact when equipped with the uniform distance) space

$$\mathscr{S}^{\downarrow} := \{ \mathbf{s} = (s_1, s_2, \ldots) \in [0, 1]^{\mathbb{N}} \text{ where } s_1 \ge s_2 \ge \ldots \ge 0 \text{ and } \sum_k s_k \le 1 \},\$$

satisfying

$$\int_{\mathscr{S}^{\downarrow}} (1-s_1) \, \nu(\mathrm{d}\mathbf{s}) < \infty.$$

In words, *c* is the rate at which each singleton detaches from "macroscopic" blocks and *v* is a measure giving the rates of "sudden dislocations", i.e. a block with asymptotic frequency *x* fragments at rate v(ds) into (possibly infinitely many) blocks with frequencies given by $xs = (xs_1, xs_2, ...) -$ these dislocations of blocks are usually represented by a so-called paintbox process, which we will define in the context of marked partitions in the next section. The self-similarity index α of a fragmentation encodes, through property (1), the speed at which blocks fragment, depending on their size. For instance, if α is negative, then there is a random time *T* which is finite almost surely at which $\Pi(T)$ is the partition into singletons, whereas it is never the case when $\alpha \ge 0$ and $v(s_1 = 0) = 0$. Note that $\alpha = 0$ means that there is no time change – in that case the sequence $\Pi^{(\cdot)}$ in (1) is simply $\Pi^{(\cdot)}$ – the process is then said to be homogeneous.

Our goal is to generalize these objects and define processes $(\Pi(t), \mathbf{V}(t), t \ge 0)$, where Π is partitionvalued and $\mathbf{V}(t) = (V_n(t), n \ge 1)$ is a random map $\mathbb{N} \to [0, \infty)$ playing the role of $(|\pi_k|^{\alpha}, k \ge 1)$, i.e. dictating the speed of fragmentation of different blocks of Π . To define this we need first to introduce the formalism of marked partitions and processes in this space.

1.2 Partitions with marks

Let us consider partitions where each block is decorated with a mark. For convenience, we consider that the space of marks is the space $[0, \infty]$ where 0 is identified with ∞ . Topologically it is a circle so we will denote it by S^1 , but throughout the paper elements of S^1 will be identified with their unique representative in $[0, \infty)$, and this enables us to consider for instance the maps

$$m_x: v \mapsto xv$$
 and $p_\alpha: v \mapsto v^\alpha$

as well-defined and continuous, where *x* is in S^1 or $[0, \infty)$ and $\alpha \in \mathbb{R} \setminus \{0\}$. Note that for a technical reason, we choose to use throughout the article the convention $0^0 = 0$, so that $0^\alpha = 0$ for any $\alpha \in \mathbb{R}$, and $v^0 = \mathbb{1}_{v\neq 0}$ for any $v \in S^1$. For convenience and with a slight abuse we will often identify an element of $[0, \infty)$ with the corresponding element of S^1 .

For $n \in \mathbb{N} \cup \{\infty\}$, we consider the space of marked partitions defined by

$$\mathcal{M}_n := \left\{ x = (\pi, \mathbf{v}) \in \mathcal{P}_n \times (S^1)^{[n]}, \forall i, j \in [n], i \sim^{\pi} j \implies v_i = v_j \right\}.$$

It is a closed subset of $\mathcal{P}_n \times (S^1)^{[n]}$, which, endowed with the product topology, is compact metrizable, therefore Polish. Note that by definition, if $(\pi, \mathbf{v}) \in \mathcal{M}_n$ where $\pi = \mathbf{1}$ is the partition into a single block, then \mathbf{v} is of the form (v, v, \ldots) for a unique $v \in S^1$. For this reason we will use the abuse of notation $(\mathbf{1}, v)$ to denote this element. We see $x = (\pi, \mathbf{v})$ as the partition π where each block is given a mark. Therefore, we will sometimes say *B* is a block of *x* with mark *v* if $B \in \pi$ and $v_i = v$ for some (hence all) $i \in B$. Similarly, we will use the notation $i \sim^x j$ if *i* and *j* are in the same block of π .

Note that for $n < m \le \infty$ and $x = (\pi, \mathbf{v}) \in \mathcal{M}_m$, we can naturally consider the restrictions $x_{|[n]} = (\pi, \mathbf{v})_{|[n]} := (\pi_{|[n]}, (v_1, v_2, \dots, v_n)) \in \mathcal{M}_n$, which are clearly continuous maps.

Similarly, we can extend the action of injections $\sigma : [n] \to [m]$ to our context and define for $x = (\pi, \mathbf{v}) \in \mathcal{M}_m$,

$$x^{\sigma} = (\pi, \mathbf{v})^{\sigma} = (\pi^{\sigma}, \mathbf{v}^{\sigma}) := (\pi^{\sigma}, (v_{\sigma(i)}, i \in [n])) \in \mathcal{M}_n$$

We say that a random variable *X* with values in \mathcal{M}_{∞} is exchangeable if for all $\sigma : \mathbb{N} \to \mathbb{N}$ permutations,

$$X^{\sigma} \stackrel{(d)}{=} X.$$

Finally we can also extend the fragmentation operator Frag to marked partitions by setting

$$\operatorname{Frag}((\pi, \mathbf{v}), (\pi^{(\cdot)}, \mathbf{v}^{(\cdot)})) := (\operatorname{Frag}(\pi, \pi^{(\cdot)}), \widetilde{\mathbf{v}}),$$

where, for $i \ge 1$, \tilde{v}_i is defined by $v_i v_i^{(k_i)}$, where k_i is the label of the block containing i – so that i is in the k_i -th block of π .

We say that a marked partition $x \in \mathcal{M}_{\infty}$ is *non-degenerate* if every finite block has mark 0, and we denote the space of non-degenerate marked partitions by

$$\mathcal{M}_{\infty}^{\star} := \left\{ x = (\pi, \mathbf{v}) \in \mathcal{M}_{\infty}, \forall i \ge 1, i \text{ in a finite block of } \pi \implies v_i = 0 \right\}$$

In particular for singleton blocks, $\{i\} \in \pi$ implies $v_i = 0$. Note that this space is still Polish [see e.g. 22, Theorem 2.2.1] as a G_{δ} -subset – a countable intersection of open sets – of \mathcal{M}_{∞} . Indeed, letting for all $i \in \mathbb{N}$, $N_i : \mathcal{M}_{\infty} \to \mathbb{N} \cup \{\infty\}$ be the map associating (π, \mathbf{v}) with the cardinality of the block of π containing *i*, then, taking *d* to be any metric on S^1 compatible with its topology, we can write

$$\mathcal{M}_{\infty}^{\star} = \bigcap_{i \ge 1} \{ N_i < \infty \implies v_i = 0 \}$$
$$= \bigcap_{i,j,k \ge 1} \left(\{ N_i \ge j \} \cup \{ d(v_i, 0) < 1/k \} \right),$$

which is a countable intersection of open subsets of \mathcal{M}_{∞} . Note that if *n* is finite, one cannot define an analogous property of non-degeneracy for marked partitions in \mathcal{M}_n .

Now let us define paintbox processes for exchangeable marked partitions. Consider the space $([0, 1] \times [0, \infty), \leq)$ equipped with the lexicographic order, that is if $z = (s, v) \in [0, 1] \times [0, \infty)$ and $z' = (s', v') \in [0, 1] \times [0, \infty)$, then

$$z \leq z' \iff s < s' \text{ or } (s = s' \text{ and } v \leq v').$$

Let us define

$$\mathscr{Z}_{0}^{\downarrow} := \{ \mathbf{z} = (z_{1}, z_{2}, \ldots) \in ([0, 1] \times [0, \infty))^{\mathbb{N}}, z_{1} \ge z_{2} \ge \ldots, \text{ and } \sum_{k} s_{k} \le 1 \},\$$

and note that, endowed with the product topology, it is a Polish space. Indeed, it can be written

$$\mathscr{Z}_0^{\downarrow} = \Big\{ \sum_k s_k \le 1 \Big\} \cap \bigcap_{i \ge 1} \Big(\{ s_i > s_{i+1} \} \cup \{ s_i = s_{i+1} \text{ and } v_i \ge v_{i+1} \} \Big),$$

which is a countable intersection of closed and open subsets of $([0, 1] \times [0, \infty))^{\mathbb{N}}$. This space being Polish, closed sets are G_{δ} , and so $\mathscr{Z}_{0}^{\downarrow}$ is Polish. Because this will be consistent with our previous definition of $\mathcal{M}_{\infty}^{\star}$, we want to ignore the possible indices $k \geq 1$ such that $s_{k} = 0$. Therefore, we will rather use the space

$$\begin{aligned} \mathscr{Z}^{\downarrow} &:= \left\{ \mathbf{z} \in \mathscr{Z}_{0}^{\downarrow}, \ \forall k \ge 1, \ s_{k} = 0 \implies v_{k} = 0 \right\} \\ &= \bigcap_{k,l \ge 1} \left\{ \mathbf{z} \in \mathscr{Z}_{0}^{\downarrow}, \ s_{k} > 0 \text{ or } v_{k} < 1/l \right\}, \end{aligned}$$

which is still Polish.

Similarly as in the usual case, we say that $x = (\pi, \mathbf{v}) \in \mathcal{M}_{\infty}$ has asymptotic frequencies if π has asymptotic frequencies. In that case, we define $|x|^{\downarrow} \in \mathscr{Z}^{\downarrow}$ as the nonincreasing reordering (with respect to the lexicographic order \leq on $[0, 1] \times [0, \infty)$) of the sequence of pairs

 $((|B_i|, v_i), B_i \text{ is the } i\text{-th infinite block of } x \text{ such that } |B_i| > 0 \text{ and with mark } v_i).$

Note that we consider only blocks *B* satisfying |B| > 0 in the previous display since in general the set $\{(|B_i|, v_i), B_i \text{ is the } i\text{-th block of } x, \text{ with mark } v_i\}$ may be impossible to enumerate in nonincreasing order.

Now let us introduce a paintbox construction for marked partitions. Consider $\mathbf{z} = (\mathbf{s}, \mathbf{v}) \in \mathscr{Z}^{\downarrow}$, and let $(U_n, n \ge 1)$ be an i.i.d. sequence of [0, 1]-uniform random variables. Define $X = (\Pi, \mathbf{V})$ as the $\mathcal{M}_{\infty}^{\star}$ -valued random variable given by the following relation:

$$i \sim^{II} j \iff i = j \text{ or } \exists n \ge 1, \ t_{n-1} \le U_i, U_j < t_n,$$
$$V_i := \begin{cases} v_n & \text{if } t_{n-1} \le U_i < t_n, \text{ for } n \ge 1, \\ 0 & \text{if } \sum_k s_k \le U_i, \end{cases}$$

where $t_n := \sum_{k=1}^n s_k$, with $t_0 := 0$ by convention. It is easily checked that the random variable X is exchangeable. Also, recall the definition of asymptotic frequencies for a marked partition, and note that the law of large numbers implies $|X|^{\downarrow} = \mathbf{z}$ almost surely. We denote by $\varrho_{\mathbf{z}}$ the distribution of X. We will also make use of the distribution of $X_{\lfloor [n]}$ for $n \in \mathbb{N}$, which we denote by $\varrho_{\mathbf{z}}^n$. Note that for any $v \in S^1$, if $\mathbf{z} = (\mathbf{s}, \mathbf{v}) \in \mathscr{Z}^{\downarrow}$ is the unique element such that $s_1 = 1$ and $v_1 = v$, then $\varrho_{\mathbf{z}} = \delta_{(1,v)}$. For this reason, we will again abuse notation and let $(\mathbf{1}, v) \in \mathscr{Z}^{\downarrow}$ denote this element, so that $\varrho_{(1,v)} = \delta_{(1,v)}$.

It is well-known since the work of Kingman [16, Theorem 2] that the law of an exchangeable partition can be expressed as a mixture of paintbox processes. Using the same arguments, one obtains the following result for marked partitions.

Proposition 2. Let X be an exchangeable random variable with values in $\mathcal{M}_{\infty}^{\star}$. Then there exists a unique probability measure v on \mathscr{Z}^{\downarrow} such that

$$\mathbb{P}(X \in \cdot) = \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}(\cdot) \, \nu(\mathrm{d}\mathbf{z}).$$
(2)

П

Proof. See Appendix A.1.

This setting of marked partitions being in place, we can now define our objects of study.

2 Extended self-similar fragmentations

2.1 Definitions, first properties

Let us now define self-similar fragmentation processes with values in \mathcal{M}_{∞} . For this, let us introduce a family of *self-similar fragmentation* operators (ssFrag_{α}, $\alpha \in \mathbb{R}$), defined as follows. For $n \in \mathbb{N} \cup \{\infty\}$, consider a marked partition $x = (\pi, \mathbf{v}) \in \mathcal{M}_n$ and a sequence $\bar{x}^{(\cdot)}$ of càdlàg maps $\bar{x}^{(k)}$: $[0, \infty) \to \mathcal{M}_n$, satisfying $\bar{x}^{(k)}(0) = (\mathbf{1}, \mathbf{1})$. Writing for all $k \geq 1$ and $t \geq 0$, $\bar{x}^{(k)}(t) = (\bar{\pi}^{(k)}(t), \bar{\mathbf{v}}^{(k)}(t))$, we define

ssFrag_{$$\alpha$$} $(x, \bar{x}^{(\cdot)}) := (\hat{\pi}, \hat{\mathbf{v}})$

as the map $[0, \infty) \rightarrow \mathcal{M}_n$ such that

$$\begin{split} \hat{v}_i(t) &= v_i \bar{v}^{(k_i)}(v_i^{\alpha} t) \\ i \sim^{\hat{\pi}(t)} j \iff i \sim^{\pi} j \text{ and } i \sim j \text{ in } \bar{\pi}^{(k_i)}(v_i^{\alpha} t), \end{split}$$

where k_i is defined as the label of the block of π containing i (i.e. such that i is in the k_i -th block of π). Note that thanks to this definition, if a block B of x has mark 0, then the process ssFrag_{α} $(x, \bar{x}^{(\cdot)})$ is frozen at block B, in the sense that for all $t \ge 0$, B is a block of $\hat{\pi}(t)$, and every $j \in B$ will have $\hat{v}_j(t) = 0$. Also, the assumptions on the maps $\bar{x}^{(k)}$ imply that ssFrag_{α} $(x, \bar{x}^{(\cdot)})$ is càdlàg and satisfies ssFrag_{α} $(x, \bar{x}^{(\cdot)})(0) = x$.

Remark 3.

(i) Consider here a convergent sequence $x_n = (\pi_n, \mathbf{v}_n) \to x = (\pi, \mathbf{v}) \in \mathcal{M}_{\infty}$, and assume that $v_{n,i} = 0$ for all $n \ge 1$ whenever $v_i = 0$ for some *i*. If additionally we have for some $t \ge 0$, for all $i \ge 1$ such that $v_i > 0$, and for all $k \ge 1$,

$$\bar{x}^{(k)}(v_{n,i}^{\alpha}t) \xrightarrow[n \to \infty]{} \bar{x}^{(k)}(v_{i}^{\alpha}t),$$

then it is a straightforward consequence of the definition that

$$\mathrm{ssFrag}_{\alpha}(x_n, \bar{x}^{(\cdot)})(t) \xrightarrow[n \to \infty]{} \mathrm{ssFrag}_{\alpha}(x, \bar{x}^{(\cdot)})(t).$$

(ii) Note that one could define $ssFrag_{\alpha}$ in terms of Frag because we have the equality

ssFrag_{$$\alpha$$} $(x, \bar{x}^{(\cdot)})(t) = \text{Frag}(x, \bar{x}^{(\cdot)}(w^{\alpha}_{(\cdot)}t)),$

where $w_{(\cdot)}$ is the vector defined by $w_{(k)} = v_i$, for any *i* in the *k*-th block of π .

We can now define the following generalization of self-similar fragmentations.

Definition 4. Let $X(t) = (\Pi(t), \mathbf{V}(t), t \ge 0)$ be a stochastic process with values in \mathcal{M}_{∞} . We say that X is an *extended self-similar fragmentation* (ESSF) process if it is a *stochastically continuous strong Markov* process with càdlàg sample paths, for which the following properties hold:

(i) *Exchangeability*: for all permutations $\sigma : \mathbb{N} \to \mathbb{N}$, for all $x \in \mathcal{M}_{\infty}$,

$$(X(t)^{\sigma}, t \ge 0)$$
 under $\mathbb{P}_{x} \stackrel{(d)}{=} (X(t), t \ge 0)$ under $\mathbb{P}_{x^{\sigma}}$,

where \mathbb{P}_x denotes the distribution of the Markov process started from *x*.

(ii) Self-similar branching: there exists $\alpha \in \mathbb{R}$ such that for all $x \in \mathcal{M}_{\infty}$,

X under
$$\mathbb{P}_x \stackrel{(d)}{=} \operatorname{ssFrag}_{\alpha}(x, X^{(\cdot)}),$$

where $X^{(.)}$ is an i.i.d. sequence of copies of the process started from (1, 1). As usual, we call α the index of self-similarity, and we will say for conciseness that X is an α -ESSF. For the special case $\alpha = 0$, we will sometimes say the process X is *homogeneous*.

An ESSF process X will be called *non-degenerate* if for all $x \in \mathcal{M}_{\infty}^{\star}$, the process has sample paths in $\mathcal{M}_{\infty}^{\star}$, \mathbb{P}_{x} -almost surely.

Remark 5.

(i) Consider $X = (\Pi, \mathbf{V})$ an α -ESSF and $\gamma \in \mathbb{R} \setminus \{0\}$. Let us define $Y := (\Pi, \mathbf{V}^{\gamma})$, where $\mathbf{V}^{\gamma}(t)$ is simply the vector $(V_i(t)^{\gamma}, i \ge 1)$. Then it is easily checked that Y is an ESSF again, with index of self-similarity α/γ . Therefore, if $\alpha \ne 0$ and $\beta \ne 0$, taking $\gamma = \alpha/\beta$, one can transform any α -ESSF into a β -ESSF, but note that one cannot get a homogeneous process with this

transformation. As a result, there are really two classes of ESSF processes to consider: the α -ESSF with $\alpha \neq 0$, which are a simple transformation away from being 1-ESSF processes, and the so-called homogeneous 0-ESSF processes.

(ii) Note that this definition extends the classical case of Definition 1. Indeed, if Π is a usual α -self-similar fragmentation process started from 1, then by definition, almost surely for all $t \ge 0$ and $i \in \mathbb{N}$, $\Pi(t)$ has asymptotic frequencies and one can define $V_i(t) := |B|$ if B is the block containing i in $\Pi(t)$. Now consider an independent sequence $X^{(\cdot)}$ of copies of (Π, \mathbf{V}) , and define for any $x \in \mathcal{M}_{\infty}$,

$$X_x = \mathrm{ssFrag}_{\alpha}(x, X^{(\cdot)}).$$

Then X_x is the distribution of an α -ESSF started from x, which extends the usual self-similar fragmentation Π – consider x = (1, 1) to obtain the original process. Note also that in this case X is non-degenerate, because finite blocks have asymptotic frequency equal to 0.

As a first remark about ESSF processes, let us show a projective Markov property. It is very analogous to [8, Lemma 3.2] and [12, Proposition 2], but we need another statement in the present context.

Lemma 6. Let X be an ESSF process. Then for any $n \in \mathbb{N}$, the process $(X(t)_{|[n]}, t \ge 0)$ is Markovian in \mathcal{M}_n . More precisely, there exists a transition kernel $(p_t^n, t \ge 0)$ on \mathcal{M}_n such that for any initial state $x \in \mathcal{M}_{\infty}$,

$$\mathbb{P}_{x}(X(t)_{|[n]} \in \cdot) = p_{t}^{n}(x_{|[n]}, \cdot).$$

Proof. We only need to prove that

$$\mathbb{P}_{x}(X(t)_{|[n]} \in \cdot) = \mathbb{P}_{x'}(X(t)_{|[n]} \in \cdot)$$

for any two initial states $x, x' \in \mathcal{M}_{\infty}$ such that $x'_{[n]} = x_{[n]}$.

Consider a probability space (Ω, \mathbb{P}) such that $X^{(\cdot)}$ is a sequence of i.i.d. copies of the ESSF process started from (1, 1), and let $\alpha \in \mathbb{R}$ be the self-similarity index of *X*. By the branching property, we have

$$\mathbb{P}_{x}(X(t)_{|[n]} \in \cdot) = \mathbb{P}(\operatorname{ssFrag}_{\alpha}(x, X^{(\cdot)})(t)_{|[n]} \in \cdot)$$

and $\mathbb{P}_{x'}(X(t)_{|[n]} \in \cdot) = \mathbb{P}(\operatorname{ssFrag}_{\alpha}(x', X^{(\cdot)})(t)_{|[n]} \in \cdot).$

It remains to notice that by definition, ssFrag_{α}(x, $X^{(\cdot)}$)(t)_{$|[n]} is in fact a functional which depends only on <math>x_{|[n]}$ and $X^{(\cdot)}$. Therefore, because $x'_{[n]} = x_{|[n]}$, we have</sub>

$$\mathrm{ssFrag}_{\alpha}(x, X^{(\cdot)})(t)_{|[n]} = \mathrm{ssFrag}_{\alpha}(x', X^{(\cdot)})(t)_{|[n]}$$

everywhere on Ω , which implies by the preceding display that

$$\mathbb{P}_{x}(X(t)_{|[n]} \in \cdot) = \mathbb{P}_{x'}(X(t)_{|[n]} \in \cdot),$$

concluding the proof.

The previous lemma shows that given an ESSF process *X*, one can define its law started from any $x_0 \in \mathcal{M}_n$, for any $n \in \mathbb{N}$, as the law of the restriction $X_{|[n]}$ of the initial process started from any $x \in \mathcal{M}_\infty$ such that $x_{|[n]} = x_0$.

As a result, the restriction $X_{|[1]}$ of an ESSF process $X = (\Pi, \mathbf{V})$ to $\mathcal{M}_1 = \mathcal{P}_1 \times S^1$ is a Markov process. Since the space \mathcal{P}_1 is a singleton, the lemma implies that the real-valued process $V_1 = (V_1(t), t \ge 0)$ is a Markov process in S^1 and note that by exchangeability, the process V_i has the same marginal distribution for all $i \ge 1$. Further, Definition 4 implies that it is an a.s. càdlàg strong Markov process satisfying a self-similar property; more precisely, for $v \ge 0$ let P_v denote the distribution of V_1

started at *v* on the Skorokhod space of càdlàg maps $[0, \infty) \rightarrow S^1$, and let *V* denote the canonical process on that space. Then

$$(V(t), t \ge 0)$$
 under $P_v \stackrel{(d)}{=} (vV(v^{\alpha}t), t \ge 0)$ under P_1 ,

where α is the self-similarity index of *X*. In other words, *V* is a positive self-similar Markov process (pssMp). Note that in the literature, the index of self-similarity of a pssMp refers in general to $-\alpha$ [19] or $-1/\alpha$ when $\alpha \neq 0$, e.g. in [18] where Lamperti calls this the *order* of the process rather than the index. Here we use the convention found in the self-similar fragmentation literature, e.g. [2, 3, 17]. Let us summarize in a proposition some properties of *V* that can be deduced from the well-developed theory of self-similar Markov processes. First, if $X = (\Pi, \mathbf{V})$ is an ESSF process, for each $i \ge 1$ define $\zeta_i := \inf\{t \ge 0, V_i(t) = 0\}$, and for $t \in [0, \zeta_i]$,

$$\varphi_i(t) := \int_0^t V_i(t)^\alpha \, \mathrm{d}s.$$

Note that φ_i is continuous and increasing. We define its right-continuous inverse $\tau_i(t)$, for $t \in [0, \infty)$, by

$$\tau_i(t) := \begin{cases} \varphi_i^{-1}(t) & \text{if } t < \varphi_i(\zeta_i), \\ \infty & \text{if } t \ge \varphi_i(\zeta_i). \end{cases}$$

We need a convention for infinite times, so we let $V_i(\infty) \equiv 0$, so that $V_i(\tau_i(t))$ is always defined. Also, note that the definition of the Frag operator implies that a.s., Π has nonincreasing sample paths for the *finer-than* partial order – π is finer than π' if the blocks of π' can be written as unions of blocks of π . Since a.s. for all $n \ge 1$, $\Pi(t)_{|[n]}$ is nonincreasing in a finite set, it is eventually constant. This implies that $\Pi(t)$ converges a.s. when $t \to \infty$, and we may denote its limit by $\Pi(\infty)$. Let us now state the proposition.

Proposition 7. Let $X = (\Pi, \mathbf{V})$ be an α -ESSF process and $i \ge 1$, and define

$$\begin{aligned} \zeta_i &:= \inf\{t \ge 0, \ V_i(t) = 0\},\\ \varphi_i(t) &:= \int_0^t V_i(t)^\alpha \, \mathrm{d} s, \qquad t \in [0, \ \zeta_i]\\ \tau_i(t) &:= \varphi_i^{-1}(t), \qquad t \ge 0 \end{aligned}$$

Then the following properties hold.

- Either $\zeta_i < \infty \mathbb{P}_{(1,1)}$ -a.s., or $\zeta_i = \infty \mathbb{P}_{(1,1)}$ -a.s.
- Either $\varphi_i(\zeta_i) < \infty \mathbb{P}_{(1,1)}$ -a.s., or $\varphi_i(\zeta_i) = \infty \mathbb{P}_{(1,1)}$ -a.s.
- In the case $\zeta_i < \infty$, either V_i reaches 0 continuously $\mathbb{P}_{(1,1)}$ -a.s., or V_i eventually jumps to 0 $\mathbb{P}_{(1,1)}$ -a.s.
- $\varphi_i(\zeta_i) = \infty$ iff V_i reaches 0 continuously.
- The process $\xi_i := \log(V_i \circ \tau_i), -i.e.$ defined by

$$\xi_i(t) = \log V_i(\tau_i(t)), \qquad 0 \le t < \varphi_i(\zeta_i),$$

is a (killed in the case $\varphi_i(\zeta_i) < \infty$) Lévy process called the inverse Lamperti transform of V_i .

Proof. These are classical results on pssMp, we refer to [18] for a proof.

This proposition tells us that it is natural to consider the time-changed processes $V_i \circ \tau_i$ for $n \ge 1$, which behave as exponentials of Lévy processes. However, there is no unique time-change that could make the whole process *X* behave nicely. Instead, we have to rely on stopping lines, which are tools generalizing stopping times in the context of branching Markov processes (see e.g. [9] for their use in branching Brownian motion, or [3, 5] in the context of fragmentations).

2.2 Stopping lines, changing the index of self-similarity

First let us define some filtrations associated with an ESSF process $X = (\Pi, \mathbf{V})$. To this aim, let us endow the power set $2^{\mathbb{N}} := \{A \subset \mathbb{N}\}$ with the topology generated by the metric $d(A, B) := (\sup\{n \in \mathbb{N}, A \cap [n] = B \cap [n]\})^{-1}$, which makes $2^{\mathbb{N}}$ a compact space. Now for $i \in \mathbb{N}$, let us define the block process $(B_i(t), t \ge 0)$ as the $2^{\mathbb{N}}$ -valued càdlàg process such that for all $t \ge 0$, $B_i(t)$ is the block of $\Pi(t)$ containing *i*, that is:

$$B_i(t) = \{ j \in \mathbb{N}, \ i \sim^{\Pi(t)} j \}.$$

Now we can define a sequence of natural filtrations associated to X by

$$\mathcal{G}_i = (\mathcal{G}_i(t), t \ge 0)$$
 with $\mathcal{G}_i(t) = \sigma(B_i(s), V_i(s), s \in [0, t]), \quad i \ge 1, t \ge 0.$

Definition 8. Let $X = (\Pi, \mathbf{V})$ be an ESSF process. A sequence $L = (L_i, i \ge 1)$ of random variables with values in $[0, \infty]$ is called a stopping line if

- (i) for all $i \ge 1$, L_i is a \mathcal{G}_i -stopping time.
- (ii) for $i, j \ge 1$, if $i \sim^{\prod(L_i)} j$, then $L_i = L_j$.

Since (ii) entails that $i \sim^{\Pi(L_i)} j$ is an equivalence relation, its equivalence classes form a well-defined partition of \mathbb{N} which we denote by $\Pi(L)$ with a slight abuse of notation. Also, denoting $\mathbf{V}(L)$ as the vector ($V_i(L_i), i \geq 1$), it is clear that $X(L) := (\Pi(L), \mathbf{V}(L))$ is a well-defined (random) element of \mathcal{M}_{∞} .

Remark 9. A fixed time $t \ge 0$ can be seen as a stopping line (an *L* for which $L_i \equiv t$ for all $i \ge 1$), and it is easily checked that for a stopping line *L*, one can define L + t and $L \land t$ by

$$(L+t)_i = L_i + t$$
 and $(L \wedge t)_i = L_i \wedge t$,

which are again stopping lines. Thus for a stopping line *L* we will be able to consider the processes $X(L + \cdot) := (X(L + t), t \ge 0)$ and $X(L \land \cdot) := (X(L \land t), t \ge 0)$. Since it will be useful, we define the following σ -algebra:

$$\mathcal{G}_L := \sigma(X(L \wedge t), \ t \ge 0).$$

We can now state the Markov property for stopping lines, which is analogous to what can be found in [5, Lemma 3.14].

Proposition 10 (Stopping line Markov property). Let X be an α -ESSF, and L be a stopping line. Then conditional on \mathcal{G}_L , the following equality in distribution holds:

$$X(L+\cdot) \stackrel{(d)}{=} \operatorname{ssFrag}_{\alpha} (X(L), X^{(\cdot)}), \tag{3}$$

where $X^{(\cdot)}$ is an independent, i.i.d. sequence of copies of the process started from (1, 1).

Proof. See Appendix A.2.

The next step in the analysis of ESSF processes is to bring the index of self-similarity to 0. This will be done via the random time changes ($\tau_i(t), i \ge 1, t \ge 0$) defined above by

$$au_i(t) := \varphi_i^{-1}(t), \quad \text{where} \quad \varphi_i(u) := \int_0^u V_i(s)^\alpha \, \mathrm{d} s, \quad u \ge 0$$

These time changes enable us to turn an α -ESSF into a homogeneous ESSF. The following proposition makes this claim more precise.

Proposition 11. Let $X = (\Pi, \mathbf{V})$ be an α -ESSF, with $\alpha \in \mathbb{R}$. Let $\beta \in \mathbb{R}$ and define the random times

$$\tau_i^{\beta}(t) = \left(\int_0^t V_i(s)^{\beta} \, \mathrm{d}s\right)^{-1}(t), \qquad i \ge 1, \ t \ge 0,$$

Then for each $t \ge 0$, $\tau^{\beta}(t)$ is a stopping line, and the process $X \circ \tau^{\beta} := (X(\tau^{\beta}(t)), t \ge 0)$ is an $(\alpha - \beta)$ -ESSF. Furthermore, if X is non-degenerate, then $X \circ \tau^{\beta}$ is also non-degenerate.

Proof. See Appendix A.3.

By bringing the index of self-similarity to 0 we can transform any ESSF into a homogeneous process. Let us now study further those 0-ESSF.

3 Main results

3.1 Decomposition of ESSF processes

Let us consider here a homogeneous 0-ESSF process $X = (\Pi, \mathbf{V})$, started from $(\mathbf{1}, \mathbf{1})$. We know by Lemma 6 that it satisfies a projective Markov property, i.e. for all $n \in \mathbb{N}$, $X_{|[n]}$ defines a Markov process with values in \mathcal{P}_n . Let $n \in \mathbb{N}$ be fixed, and define the stopping time

$$T_n := \inf\{t \ge 0, \ \Pi(t)|_{[n]} \neq \mathbf{1}_n \text{ or } V_1(t) = 0\},\$$

as well as the killed process

$$\tilde{\xi}_n := (\log V_1(t), \ 0 \le t < T_n).$$

Note that homogeneity implies that the pair $(\tilde{\xi}_n - \log v, T_n)$ has the same distribution under every $\mathbb{P}_{(1,v)}$ for all $v \in S^1 \setminus \{0\}$. Therefore for $t \ge 0$, conditional on $\{T_n > t\}$, the Markov property applied at time *t* shows that $(\tilde{\xi}_n(t + \cdot) - \tilde{\xi}_n(t), T_n - t)$ has the same distribution as $(\tilde{\xi}_n, T_n)$ under $\mathbb{P}_{(1,1)}$. This shows that the killed process $\tilde{\xi}_n$ is distributed as

$$\tilde{\xi}_n \stackrel{(d)}{=} (\xi_n(t), \ 0 \le t < T_n),$$

(1)

where ξ_n is a Lévy process and T_n is an independent exponential random variable. Note that this implies that if $T_n < \infty$, then $V_1(T_n -) = \exp(\xi_n(T_n)) > 0$. Now for $n \in \mathbb{N}$ such that $T_n < \infty$ almost surely, consider D_n , the dislocation (or freezing) at time T_n , defined by

$$D_n := (\Pi(T_n), \mathbf{V}(T_n)/V_1(T_n-))_{|[n]} \in \mathcal{M}_n,$$

where the division $\mathbf{V}(T_n)/V_1(T_n-)$ is to be understood coordinate-wise. Equivalently, D_n is the unique random marked partition such that

$$X(T_n)_{|[n]} = \operatorname{Frag}(X(T_n-)_{|[n]}, D_n)$$

with a slight abuse of notation in this case since $X(T_n-)_{[n]}$ has only one block (D_n is not a sequence but additional terms are useless to define a fragmentation of a single block).

Note that this implies that D_n has the same distribution under every $\mathbb{P}_{(1,v)}$ for all $v \in S^1 \setminus \{0\}$. Thus for any bounded measurable maps $g : \mathbb{R} \to \mathbb{R}$, $h : \mathcal{M}_n \to \mathbb{R}$ and $t \ge 0$, applying the Markov property at time $t \ge 0$, one gets

$$\mathbb{E}_{(1,1)}[g(\xi_n(t))\mathbb{1}_{T_n>t}h(D_n)] = \mathbb{E}_{(1,1)}[g(\xi_n(t))\mathbb{1}_{T_n>t}]\mathbb{E}_{(1,1)}h(D_n),$$

which shows that the killed Lévy process (ξ_n, T_n) and the marked partition D_n are independent. Let us define \mathcal{D}_n as the law of D_n , and notice also that exchangeability of $X_{|[n]}$ implies that \mathcal{D}_n is an exchangeable probability measure on \mathcal{M}_n .

Since (ξ_n, T_n) is a killed Lévy process, one can define uniquely $d_n \in \mathbb{R}$, $\beta_n \ge 0$, $J_n \ge 0$ and λ_n a measure on $\mathbb{R} \setminus \{0\}$ satisfying $\int 1 \wedge y^2 \lambda_n(dy) < \infty$, such that

• the process ξ_n is a Lévy process with characteristic exponent

$$\psi_n(\theta) := \log \mathbb{E}[e^{i\theta\xi_n(1)}] = id_n\theta - \frac{\beta_n}{2}\theta^2 + \int_{\mathbb{R}} \left(e^{i\theta y} - 1 - i\theta y\mathbb{1}_{|y| \le 1}\right) \lambda_n(\mathrm{d}y),$$

• ξ_n is killed at rate is $J_n = 1/\mathbb{E}T_n$, which may be 0 if $T_n = \infty$ almost surely.

Remark 12. Note that knowing $(\psi_n, J_n, \mathcal{D}_n)$ for $n \in \mathbb{N}$ is enough to reconstruct the process *X*. Indeed, starting from $(\mathbf{1}, v)$, the process $X_{|[n]}$ up to time T_n has distribution equal to that of

$$Y_n := ((\mathbf{1}_n, v e^{\xi_n(t)}), \ 0 \le t < T_n)$$

and at time T_n jumps to $(\Pi, v e^{\xi_n(T_n)} \mathbf{V})$, where (Π, \mathbf{V}) is independently drawn according to \mathcal{D}_n .

By the branching property, one only needs to iterate this construction at each jump time, independently for each marked block, to get the whole process $X_{|[n]}$. By Kolmogorov's extension theorem – since $(X_{|[m]})_{|[n]} = X_{|[n]}$ for each $n \le m$ – these distributions characterize the distribution of X.

Let us now state our main result which identifies the form that those characteristics can take.

Theorem 13. Let X be a non-degenerate 0-ESSF and for each n, write $(\psi_n, J_n, \mathcal{D}_n)$ for the characteristics describing the law of $X_{|[n]}$. Then there is a unique quadruple (c, d, β , Λ), where c, $\beta \ge 0$, $d \in \mathbb{R}$, and Λ is a measure on $\mathscr{Z}^{\downarrow} \setminus \{(1, 1)\}$, which satisfies necessarily

$$\int_{\mathscr{Z}^{\downarrow}} \left(1 - s_1 \mathbb{1}_{v_1 > 0} + (\log v_1)^2 \wedge 1 \right) \Lambda(\mathrm{d}\mathbf{z}) < \infty, \tag{4}$$

such that for all $n \in \mathbb{N}$,

(i)
$$\psi_n(\theta) = id\theta - \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{j \ge 1 \\ v_j > 0}} s_j^n \left(e^{i\theta \log v_j} - 1 \right) - i\theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} \Lambda(\mathrm{d}\mathbf{z}).$$

(*ii*)
$$J_n = nc + \int_{\mathscr{Z}^{\downarrow}} \left(1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n\right) \Lambda(\mathrm{d}\mathbf{z}).$$

(iii) if $J_n > 0$, $\mathcal{D}_n = \frac{1}{J_n} \Big(\sum_{i=1}^n c \delta_{\mathfrak{e}_i^n} + \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}^n \Big(\cdot \cap \{ \pi \neq \mathbf{1}_n \text{ or } (\pi, \mathbf{v}) = (\mathbf{1}_n, 0) \} \Big) \Lambda(\mathrm{d}\mathbf{z}) \Big)$, where $\varrho_{\mathbf{z}}^n$ is the paintbox process defined in Section 1.2, and $\delta_{\mathfrak{e}_i^n}$ denotes the Dirac point measure on \mathfrak{e}_i^n , the marked partition defined as

$$\mathbf{e}_{i}^{n} := \left(\left\{ [n] \setminus \{i\}, \{i\} \right\}, (1, \ldots, 1, \underbrace{0}_{i-th \ index}, 1, \ldots, 1) \right).$$

Conversely if c, $\beta \ge 0$, $d \in \mathbb{R}$ and Λ is a measure on $\mathscr{Z}^{\downarrow} \setminus \{(1, 1)\}$, satisfying (4), then there exists a 0-ESSF with characteristics as above.

Proof. See Appendix A.4.

Remark 14. It is an immediate consequence of the theorem that the process describing the block of *X* containing 1 can be constructed in the following Poissonian way. Consider \mathcal{N} a Poisson point process on $[0, \infty) \times \mathcal{M}_{\infty}^{\star}$ with intensity

$$\mathrm{d} t \otimes \Big(\sum_{i=1}^{\infty} c \delta_{\mathfrak{e}_i^n} + \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}^n \big(\cdot\big) \Lambda(\mathrm{d} \mathbf{z})\Big),$$

and define

$$\mathcal{N}' := \{ (t, \log v_1), (t, x) \in \mathcal{N} \text{ with } x = (\pi, \mathbf{v}) \text{ and } v_1 \notin \{0, 1\} \},\$$

which has intensity $dt \otimes \lambda_1$, where λ_1 is defined by

$$\int_{\mathbb{R}} f \, \mathrm{d}\lambda_1 = \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{j \ge 1 \\ v_j \notin \{0,1\}}} s_j f(\log v_j) \, \Lambda(\mathrm{d}\mathbf{z}),$$

and is the Lévy measure of the process ξ_1 . It is clear that one can build a Lévy process $(\xi_1(t), t \ge 0)$ having characteristic exponent ψ_1 given by (*i*) in the theorem and whose point process of jumps is exactly \mathcal{N}' .

Define $(B(t), t \ge 0)$ as the 2^N-valued process given by

$$B(t) = \bigcap_{\substack{0 \le s < t \\ (s,x) \in \mathcal{N}}} A(x)$$

where $A(x) \subset \mathbb{N}$ denotes the block of *x* containing 1. Also, for any $n \in \mathbb{N}$, define

 $\widetilde{T}_n := \inf \left\{ t \ge 0, \ (t, x) \in \mathcal{N} \text{ with } x = (\pi, \mathbf{v}) \text{ such that } \pi_{|[n]} \neq \mathbf{1} \text{ or } v_1 = 0 \right\} \sim \operatorname{Exp}(J_n).$

Now $(B(t), e^{\xi_1(t)}, 0 \le t < \tilde{T}_1)$ is distributed as the marked block containing 1 in *X* and by construction, we also get the following equality in distribution

$$\left(X(T_n)_{\mid [n]}, n \in \mathbb{N}\right) \stackrel{(d)}{=} \left(\left(\pi_n, e^{\xi_1(\widetilde{T}_n-)}\mathbf{v}_n\right)_{\mid [n]}, n \in \mathbb{N}\right),$$

where $x_n = (\pi_n, \mathbf{v}_n)$ is the element of $\mathcal{M}_{\infty}^{\star}$ such that $(\widetilde{T}_n, x_n) \in \mathcal{N}$.

Combining Theorem 13 with Proposition 11, we get the following characterization of all ESSF processes.

Corollary 15. Let X be a non-degenerate α -ESSF. Then there exists a unique quadruple (c, d, β , Λ) as in Theorem 13 such that if ($\tau^{\alpha}(t), t \ge 0$) are the stopping lines as defined in Proposition 11, then $X \circ \tau^{\alpha}$ is a homogeneous ESSF with characteristics (c, d, β , Λ).

Let us point out that condition (4) is surprisingly nonrestrictive. There are no integrability assumptions concerning the marks of the smallest blocks (with labels greater than 1). Consequently, the point measure $\sum_k \delta_{\tilde{V}_k(t)}$, where $\tilde{V}_k(t)$ denotes the mark of the *k*-th block in X(t), might assign infinite mass to any interval $(a, b) \subset [0, \infty)$ for any t > 0. Indeed it suffices for instance that $\Lambda(d\mathbf{z})$ be of the form

$$\int_{\mathscr{Z}^{\downarrow}} \prod_{i \ge 1} F_i(s_i, v_i) \Lambda(d\mathbf{z}) = \int_{(0,1) \times S^1} F_1(s_1, v_1) \mathbb{E} \prod_{i \ge 2} F_i\left(\frac{1-s_1}{2^{i-1}}, Z_i\right) \nu(dz_1),$$

where *v* is a measure on $(0, 1) \times S^1$ with infinite mass and satisfying

$$\int_{(0,1)\times S^1} \left(1 - s_1 \mathbb{1}_{v_1 > 0} + (\log v_1)^2 \wedge 1\right) \nu(\mathrm{d} z_1) < \infty,$$

and Z_2, Z_3, \ldots are i.i.d. Exp(1) random variables. On the other hand, if one assumes an integrability condition such as

$$\int_{\mathscr{Z}^{\downarrow}} \left(\sum_{i \ge 1} v_i^{\theta} \right) - 1 - \theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} \Lambda(\mathrm{d}\mathbf{z}) < \infty$$

for some $\theta \in \mathbb{R}$, then one observes a process of point measures $(\sum_k \delta_{\tilde{V}_k(t)}, t \ge 0)$ that is nice in the sense that for all $t \ge 0$, $\mathbb{E} \sum_k \tilde{V}_k(t)^{\theta} < \infty$. This is the object of the next section.

3.2 Absorption in finite time

Consider here a non-degenerate α -ESSF with characteristics (c, d, β , Λ), started from (1, 1). We are interested in the case where the pssMp V_1 reaches 0 in finite time and, if T_i denotes the hitting time of 0 by the process V_i , we aim at giving a sufficient condition for which

$$\sup_{i\in\mathbb{N}}T_i<\infty\qquad\text{a.s}$$

If this holds, then at this random time the process is frozen in a state $X(\infty) = (\Pi(\infty), \mathbf{V}(\infty))$ where $\mathbf{V}(\infty) = \mathbf{0} = (0, 0, ...)$, and we say the process X is *absorbed in finite time*. We first put aside a trivial case and assume that

$$c > 0$$
 or $\int_{\mathscr{Z}^{\downarrow}} (1 - s_1) \Lambda(\mathrm{d}\mathbf{z}) > 0$,

since otherwise Π would be almost surely constant equal to the coarsest partition {N}. Recall that a classical self-similar fragmentation which is absorbed in finite time (this is always true when $\alpha < 0$ [4, Proposition 2]) is always totally fragmented in the limit, in the sense that $\Pi(\infty)$ is the partition of N into singletons. Clearly in our case, because of the possible freezing of blocks at dislocation events, $\Pi(\infty)$ is almost surely totally fragmented if and only if

$$\forall i \ge 1, \ s_i > 0 \implies v_i > 0 \qquad \Lambda \text{-a.e. on } \mathscr{Z}^{\downarrow}$$

A stronger property than absorption in finite time is the following: we say X has finite total length if

$$\int_0^\infty \# X(t) \, \mathrm{d}t < \infty \quad \text{a.s.},$$

where #x denotes the number of blocks with positive mark in the marked partition x. One can interpret this quantity as the total length of the tree describing the genealogy of blocks in the fragmentation, hence the name. Note that this implies that for a fixed time $t \ge 0$, #X(t) is almost surely finite, which is well-known [4, Proposition 2] in the classical self-similar fragmentation case for $\alpha < -1$.

In this section our aim is to provide sufficient conditions for ESSF processes to be absorbed in finite time and to have finite total length. The following result extends the classical setting, and makes use of natural martingales appearing in the homogeneous case. In order to be able to state it, we need a couple of additional definitions. For a marked partition $x = (\pi, \mathbf{v}) \in \mathcal{M}_n$ with $n \in \mathbb{N} \cup \{\infty\}$, and $\theta \in \mathbb{R}$, let us write

$$S_{\theta}(x) := \sum_{k \in \mathbb{N}} \tilde{v}_k^{\theta}, \tag{5}$$

where \tilde{v}_k denotes the mark associated with the *k*-th block of *x*. Let us also introduce $\kappa : \mathbb{R} \to (-\infty, \infty]$ defined by

$$\kappa(\theta) := d\theta + \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^{\downarrow}} \left(\sum_{i\geq 1} v_i^{\theta}\right) - 1 - \theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} \Lambda(\mathrm{d}\mathbf{z}).$$
(6)

Note that the integral in the last display is well-defined with values in $(-\infty, \infty]$, since

$$\begin{aligned} 1 + \theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} - \sum_{i \ge 1} v_i^{\theta} \le \left(1 + \theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} - v_1^{\theta} \right)_+ \\ \le C \left((\log v_1^2) \land 1 \right) \end{aligned}$$

where *C* is a positive constant which depends on θ , so the negative part of the integrand in the definition of κ is Λ -integrable.

Proposition 16. Let X be a non-degenerate 0-ESSF with characteristics (c, d, β, Λ) started from (1, 1). For all $\theta \in \mathbb{R}$ and $t \ge 0$,

$$\mathbb{E}S_{\theta}(X(t)) = \mathrm{e}^{t\kappa(\theta)},$$

with S_{θ} and $\kappa(\theta)$ respectively defined as in (5) and (6), and where these quantities may be infinite. If there is $\theta \in \mathbb{R}$ such that $\kappa(\theta) < \infty$, then the process

$$\left(e^{-t\kappa(\theta)} S_{\theta}(X(t)), t \geq 0 \right)$$

is a martingale. If there is $\theta \neq 0$ such that $\kappa(\theta) < 0$, then for any $\alpha \in \mathbb{R}$:

- if $-\alpha/\theta > 0$, the α -ESSF with characteristics (c, d, β, Λ) is absorbed in finite time.
- *if* $-\alpha/\theta \ge 1$, the α -ESSF with characteristics (c, d, β , Λ) has finite total length.

Proof. See Appendix A.5.

Remark 17. For a classical self-similar fragmentation with erosion coefficient $c \ge 0$ and dislocation measure v, we have

$$\kappa(\theta) = -c\theta + \int_{\mathscr{S}^{\downarrow}} \left(\sum_{i\geq 1} s_i^{\theta} - 1\right) \nu(\mathrm{d}\mathbf{s}).$$

Since $\sum_i s_i \le 1$ *v*-a.e., for all $\theta > 1$ we have $\kappa(\theta) < 0$, so we recover absorption in finite time for any $\alpha < 0$ and finite total length for any $\alpha < -1$.

Remark 18. Let us also mention that one can model branching Brownian motion in our setting. Indeed, consider a homogeneous ESSF where the logarithm of marks follow drifted Brownian motion and blocks dislocate at rate one into two blocks (say both with asymptotic frequency equal to half of the mother block) carrying the same mark. More precisely, take a 0-ESSF with characteristics $c = 0, d \in \mathbb{R}, \beta = 1$ and with $\Lambda(d\mathbf{z})$ a Dirac measure on $((\frac{1}{2}, 1), (\frac{1}{2}, 1), 0, \ldots)$.

Then the point process recording the positions of the logarithm of marks

$$\sum_{k \in \mathbb{N}} \delta_{\log \tilde{V}_k(t)}, \qquad t \ge 0$$

is a classical binary branching Brownian motion with drift d. One gets a cumulant function

$$\kappa(\theta) = d\theta + \frac{\theta^2}{2} + 1.$$

This polynomial in θ takes negative values if and only if $d^2 - 2 > 0$, and we essentially recover the well-known fact that if $d > \sqrt{2}$, the lowest particle of a branching Brownian with drift *d* goes to $+\infty$.

A Proofs

A.1 Proof of Proposition 2

Let us write as usual $X = (\Pi, \mathbf{V})$. First, note that Π is an exchangeable partition with values in $\mathcal{M}_{\infty}^{\star}$, therefore it has asymptotic frequencies – so $|X|^{\downarrow}$ exists almost surely – and the finite blocks of Π (if any) are necessarily singletons. For the uniqueness part of the proposition, notice that any ν satisfying (2) must be equal to $\mathbb{P}(|X|^{\downarrow} \in \cdot)$.

For the existence, let $(U_k, k \ge 1)$ be an i.i.d. sequence of uniform random variables on [0, 1], independent of *X*. For every $i \in \mathbb{N}$, let $Z_i = (U_k, V_i) \in [0, 1] \times S^1$, where *k* is the label of the block containing *i*. Then the sequence $(Z_i, i \ge 1)$ is exchangeable, with values in a Polish space. Therefore by de Finetti's theorem, there is a random probability measure $\theta \in \mathcal{M}_1([0, 1] \times S^1)$ such that conditional on θ , the sequence $(Z_i, i \ge 1)$ is i.i.d. with distribution θ . Let

$$(a_k, k \ge 1) = (u_k, v_k, k \ge 1)$$

denote the collection of atoms of θ . Note that $i \sim^X j$ iff $Z_i = Z_j$, and therefore the law of large numbers ensures us that the blocks of X correspond to those atoms, i.e. for each $k \ge 1$, there is a block B of X with an asymptotic frequency $|B| = \theta(a_k)$ and a mark equal to v_k . Conversely any block which is not reduced to a singleton must be formed in this way. Furthermore, note that singleton blocks have mark 0 because of the assumption that $X \in \mathcal{M}_{\infty}^{\star}$, so the knowledge of the atoms $(a_k, k \ge 1)$ and their mass is sufficient to reconstruct the sequence $(Z_i, i \ge 1)$, and therefore the marked partition X.

Now define for all $k \in \mathbb{N}$, $z_k := (\theta(a_k), v_k)$. Up to a reordering, we can assume that $\mathbf{z} = (z_k, k \ge 1)$ is in \mathscr{Z}^{\downarrow} (if the sequence of atoms is finite, we concatenate to \mathbf{z} infinitely many (0, 0) terms). The previous discussion means that conditional on θ , the asymptotic frequencies of X are exactly

$$|X|^{\downarrow} = \mathbf{z} \in \mathscr{Z}^{\downarrow},$$

and conditional on \mathbf{z} , the marked partition X is drawn according to $\rho_{\mathbf{z}}$. Note that the map $\theta \mapsto \mathbf{z}$ is measurable. Indeed, by standard point processes arguments [see e.g. 11, Lemma 9.1.XIII], there exists a measurable enumeration $(a_k, k \ge 1)$ of the atoms of θ , and it is elementary that the nonincreasing reordering of this sequence is measurable. Therefore, defining $v = \mathbb{P}(|X|^{\downarrow} \in \cdot)$, which is the push-forward of the distribution of θ by the map $\theta \mapsto \mathbf{z}$, we see that it satisfies (2).

A.2 Proof of Proposition 10

In this section, it will be helpful to consider the restriction of an ESSF process *X* to a more general (and possibly random) subset $A \subset \mathbb{N}$, considered as a random variable living on the compact space $2^{\mathbb{N}}$. We first consider a fixed – non random – $A \subset \mathbb{N}$ with cardinality $\#A \in \mathbb{N} \cup \{\infty\}$, and define a canonical enumeration of *A* by

$$\sigma_A : \begin{cases} [\#A] & \to \mathbb{N} \\ i & \mapsto \min\{n \in \mathbb{N}, \ \#(A \cap [n]) = i\}, \end{cases}$$

such that $A = \{\sigma_A(1), \sigma_A(2), \ldots\}$, with $\sigma_A(1) < \sigma_A(2) < \ldots$. Now recall the definition of the action of injections on \mathcal{M}_{∞} . For any $x \in \mathcal{M}_{\infty}$, one can see $x^{\sigma_A} \in \mathcal{M}_{\#A}$ as the restriction of x to the set A.

As an inverse operation, for any $x' \in \mathcal{M}_{\#A}$, $x'' \in \mathcal{M}_{\#A^c}$, where $A^c := \mathbb{N} \setminus A$, we can define

$$x' \stackrel{A}{\oplus} x'' \in \mathcal{M}_{\infty}$$

as the pair (π, \mathbf{v}) such that

$$\forall i, j \in \mathbb{N}, \qquad i \sim^{\pi} j \iff \begin{cases} i, j \in A \text{ and } \sigma_A^{-1}(i) \sim^{\pi'} \sigma_A^{-1}(j) \\ \text{or} \quad i, j \in A^{\text{c}} \text{ and } \sigma_{A^{\text{c}}}^{-1}(i) \sim^{\pi''} \sigma_{A^{\text{c}}}^{-1}(j) \end{cases}$$

$$\text{and } v_i = \begin{cases} v'_{\sigma_A^{-1}(i)} & \text{if } i \in A \\ v''_{\sigma_A^{-1}(i)} & \text{if } i \in A^{\text{c}}. \end{cases}$$

Similarly, for processes $X' = (X'(t), t \ge 0)$ and X'', we write for conciseness

$$X' \stackrel{A}{\oplus} X'' := \left(X'(t) \stackrel{A}{\oplus} X''(t), \ t \ge 0 \right)$$

For $x = (\pi, \mathbf{v}) \in \mathcal{M}_{\infty}$ and $A \subset \mathbb{N}$, we will say that A is x-compatible if it is a union of a family of blocks of π – i.e. if A is such that $i \in A$, $j \notin A \implies i \not\sim^{\pi} j$. These definitions enable us to reformulate the branching property as follows.

Lemma 19. Let X be an ESSF process, $x = (\pi, \mathbf{v}) \in \mathcal{M}_{\infty}$, and $A \subset \mathbb{N}$ an x-compatible set. Defining $X' := X^{\sigma_A}$ and $X'' := X^{\sigma_{A^c}}$, then under \mathbb{P}_x , X' and X'' are two independent copies of the process X, respectively started at x^{σ_A} and $x^{\sigma_{A^c}}$, and

$$X = X' \stackrel{A}{\oplus} X''.$$

Proof. This is an immediate consequence of the branching property (*ii*) of Definition 4 and of the definition of the ssFrag operator.

Let us now tackle the proof of the Markov property for stopping lines (3). We write as usual $X = (\Pi, \mathbf{V})$. We first assume that there exist $0 \le t_1 < t_2 < \ldots < t_k \le \infty$ such that for all $i \in \mathbb{N}$, L_i takes values in the finite set $\{t_1, \ldots, t_k\}$. We prove the Markov property for such stopping lines by induction on k. For k = 1 and $t_1 < \infty$, this amounts to the simple Markov property, so (3) holds by definition. If $t_1 = \infty$, then for all $t \ge 0$, for all $i \ge 1$, $L_i + t \equiv \infty$. By convention $\mathbf{V}(\infty)$ is the null vector, and by definition ssFrag $(X(L), X^{(\cdot)})$ is the process which is a.s. constant equal to $X(\infty) = (\Pi(\infty), \mathbf{V}(\infty))$, so (3) holds again.

Now assume that k > 1, and that the stopping line Markov property has been proven for all stopping lines taking at most k - 1 distinct values. By Remark 9, $L \wedge t_{k-1}$ is a stopping line taking at most k - 1 distinct values. Therefore, one can apply the induction hypothesis, which says that conditional on $\mathcal{G}_{L \wedge t_{k-1}}$, the process $X(L \wedge t_{k-1} + \cdot)$ has the distribution of a copy of X started from $X(L \wedge t_{k-1})$. Now we define the random set

$$A := \{i \in \mathbb{N}, L_i = t_k\},\$$

which is $\mathcal{G}_{L \wedge t_{k-1}}$ -measurable. Indeed let us show that $\{i \in A\} \in \mathcal{G}_{L \wedge t_{k-1}}$. Since $\{L_i = t_k\} = \{L_i > t_{k-1}\} \in \mathcal{G}_i(t_{k-1})$, one can write the indicator of this event as $\mathbb{1}_{\{L_i > t_{k-1}\}} = F(B_i(s), V_i(s), s \in [0, t_{k-1}])$ for a measurable functional F – recall that $\mathcal{G}_i(t_{k-1}) = \sigma(B_i(s), V_i(s), s \in [0, t_{k-1}])$, where $B_i(s)$ is the block of $\Pi(s)$ containing i. Now on the event $\{L_i \ge t_{k-1}\}$, we have

$$F(B_i(s), V_i(s), s \in [0, t_{k-1}]) = F(B_i(s \wedge L_i \wedge t_{k-1}), V_i(s \wedge L_i \wedge t_{k-1}), s \in [0, t_{k-1}]),$$

therefore we can write

$$\{i \in A\} = \{L_i \ge t_{k-1}\} \cap \{F(B_i(s \land L_i \land t_{k-1}), V_i(s \land L_i \land t_{k-1}), s \in [0, t_{k-1}]) = 1\} \in \mathcal{G}_{L \land t_{k-1}}, s \in [0, t_{k-1}]\}$$

so finally *A* is $\mathcal{G}_{L \wedge t_{k-1}}$ -measurable. Now notice that because *L* is a stopping line, *A* is compatible with $\Pi(L \wedge t_{k-1})$ in the sense that *A* is necessarily a union of blocks of $\Pi(L \wedge t_{k-1})$. Therefore, it is immediate by definition that

$$X(L \wedge t_{k-1} + \cdot) = X' \stackrel{A}{\oplus} X'',$$

with

$$X' = X(L \wedge t_{k-1} + \cdot)^{\sigma_A}$$

and $X'' = X(L \wedge t_{k-1} + \cdot)^{\sigma_{A^c}}$.

Now by Lemma 19, conditional on $\mathcal{G}_{L \wedge t_{k-1}}$, X' and X'' are two independent copies of X started respectively from $X(L \wedge t_{k-1})^{\sigma_A}$ and $X(L \wedge t_{k-1})^{\sigma_{A^c}}$.

Also, notice that by definition of the random set A, we have the equality

$$X(L+\cdot) = X'(t_k - t_{k-1} + \cdot) \stackrel{A}{\oplus} X'', \tag{7}$$

and for the same reason, the following equality between σ -algebras holds:

$$\mathcal{G}_L = \mathcal{G}_{L \wedge t_{k-1}} \lor \sigma (X'(s), s \in [0, t_k - t_{k-1}]).$$

Clearly X' and X'' are still independent conditional on \mathcal{G}_L , and the distribution of $X'(t_k - t_{k-1} + \cdot)$ conditional on \mathcal{G}_L is by the Markov property at time $t_k - t_{k-1}$ the law of X started from $X'(t_k - t_{k-1}) = X(L)^{\sigma_A}$. Finally, using again Lemma 19, conditional on \mathcal{G}_L , the process

$$X'(t_k - t_{k-1} + \cdot) \stackrel{A}{\oplus} X''$$

has simply the distribution of a copy of X started at X(L). So by (7) the Markov property for stopping lines holds for L, and so by induction it holds for all stopping lines taking at most a finite number of values.

Now fix a general stopping line *L*, a time $t \ge 0$, and let us assume that our probability space contains an independent sequence $X^{(\cdot)}$ of i.i.d. copies of the process started from (1, 1). To conclude, it is enough to prove that conditional on \mathcal{G}_L ,

$$X(L+t) \stackrel{(d)}{=} \mathrm{ssFrag}_{\alpha}(X(L), X^{(\cdot)})(t), \tag{8}$$

because then for any $0 \le t_1 < t_2 < ... t_k$ one can apply successively (8) to the stopping lines $L + t_i$, $1 \le i \le k$, which implies that (3) holds for any finite dimensional distributions. Therefore it remains only to prove

$$\mathbb{E}\left[F(X(L+t))Z\right] = \mathbb{E}\left[F\left(\operatorname{ssFrag}_{\alpha}\left(X(L), X^{(\cdot)}\right)(t)\right)Z\right]$$

for any fixed continuous bounded map $F : \mathcal{M}_{\infty}^{\star} \to \mathbb{R}$ and Z a \mathcal{G}_L -measurable bounded random variable. To show this, consider the sequence of stopping lines $(L^n, n \ge 1)$ defined by

$$L_i^n = 2^{-n} \lceil 2^n L_i \rceil \mathbb{1}_{L_i \le n} + \infty \mathbb{1}_{L_i > n}, \quad i \ge 1,$$

where $\lceil \cdot \rceil$ denotes the usual ceiling function. This is a classical transformation for stopping times, and it is easily checked that L^n is a stopping line for all $n \ge 1$. Furthermore, right-continuity of the process implies that $X(L^n + t)$ converges a.s. to X(L + t) as n tends to ∞ . Therefore

$$\mathbb{E}\left[F(X(L^n+t))Z\right] \xrightarrow[n\to\infty]{} \mathbb{E}\left[F(X(L+t))Z\right],$$

Now because L^n only takes values in a finite set for all *n*, we can apply (3), so

$$\mathbb{E}\left[F(X(L^{n}+t))Z\right] = \mathbb{E}\left[F\left(\operatorname{ssFrag}_{\alpha}\left(X(L^{n}), X^{(\cdot)}\right)(t)\right)Z\right].$$
(9)

This holds because Z is \mathcal{G}_{L^n} -measurable since $\mathcal{G}_L \subset \mathcal{G}_{L^n}$ for all $n \ge 1$. For the convergence of the right-hand side, recall that $X(L^n + t) \rightarrow X(L + t)$ and note also that if $V_i(L_i) = 0$ for $i \ge 1$, then

since L_i is a stopping time, by the strong Markov property $V_i(L_i + t)$ is also zero for all $t \ge 0$, and in particular $V_i(L_i^n) = 0$ for all $n \ge 1$. Now by definition an ESSF process is stochastically continuous, so in particular for any $i \ge 1$, on the event $\{V_i(L_i) > 0\}$, we have

$$X^{(j)}(V_i(L_i)^{\alpha}t) \xrightarrow[n \to \infty]{} X^{(j)}(V_i(L_i)^{\alpha}t) \qquad \forall j \ge 1 \ a.s.$$

We can now invoke the continuity property of the operator $ssFrag_{\alpha}$ pointed out in Remark 3 and deduce

$$\operatorname{ssFrag}_{\alpha}(X(L^n), X^{(\cdot)})(t) \xrightarrow[n \to \infty]{} \operatorname{ssFrag}_{\alpha}(X(L), X^{(\cdot)})(t) \quad a.s$$

Taking limits in (9) yields the equality needed to end the proof.

A.3 Proof of Proposition 11

Let $X = (\Pi, \mathbf{V})$ be an α -ESSF process, $t \ge 0$, and recall the definition of $\tau^{\beta}(t)$ as in the proposition, i.e.

$$\tau_i^{\beta}(t) = \left(\int_0^{\cdot} V_i(s)^{\beta} \,\mathrm{d}s\right)^{-1}(t), \qquad i \ge 1,$$

where the inverse is to be understood as the right-continuous inverse. For conciseness and because β is fixed, let us write simply τ instead of τ^{β} throughout the proof. First, let us see that $\tau(t)$ is a stopping line. Fix $i \in \mathbb{N}$, and note that for $T \ge 0$,

$$\{\tau_i(t) \leq T\} = \left\{\int_0^T V_i(s)^\beta \, \mathrm{d}s \geq t\right\} \in \mathcal{G}_i(T),$$

so $\tau_i(t)$ is a \mathcal{G}_i -stopping time. Furthermore, conditional on $\tau_i(t) = T$, for any $i, j \in \mathbb{N}$, if $i \sim j$ in $\Pi(T)$, then almost surely for all $0 \leq s \leq T$, we have $i \sim j$ in $\Pi(s)$ so $V_i(s) = V_j(s)$, and necessarily $\tau_j(t) = \tau_i(t) = T$.

Therefore $\tau(t)$ is indeed a stopping line, so the process $X \circ \tau = (X(\tau(t), t \ge 0))$ is well defined. We claim that its sample paths are càdlàg in \mathcal{M}_{∞} . Indeed, by definition, for each $i \in \mathbb{N}$, τ_i is a non-decreasing right-continuous map. Now almost surely the following holds: X has càdlàg sample paths, so for each $i \in \mathbb{N}$, and $t \in [0, \infty)$,

$$X(\tau_i(s)) \xrightarrow[s \downarrow t]{} X(\tau_i(t))$$
 and $X(\tau_i(s)) \xrightarrow[s \uparrow t]{} X(\tau_i(t)-)$ if $t > 0$.

Now note that for each stopping line *L* and integer $n \in \mathbb{N}$, by definition $X(L)_{|[n]}$ is a (deterministic) continuous functional of the variables $(X(L_1), \ldots, X(L_n))$. Applying this to $L = \tau(s)$ and letting $s \to t$, it follows that almost surely

$$X(\tau(s))_{|[n]} \xrightarrow{s \downarrow t} X(\tau(t))_{|[n]}$$
 and $X(\tau(s))_{|[n]}$ converges in \mathcal{M}_n when $s \uparrow t$, in the case $t > 0$.

The integer *n* being generic, this shows that $X \circ \tau$ is an almost surely càdlàg process.

Let $t \ge 0$ be fixed. Since $\tau(t)$ is a stopping line, we can apply Proposition 10, and assume that the process $X(\tau(t) + \cdot)$ is given by

ssFrag_{$$\alpha$$} ($X(\tau(t)), X^{(\cdot)}$),

where $X^{(.)}$ is an independent sequence of i.i.d. copies of the process started from (1, 1). For each $k \in \mathbb{N}$, let $(\tau^{(k)}(s), s \ge 0)$ denote the stopping lines corresponding to $X^{(k)}$, i.e.

$$\tau_i^{(k)}(s) = \left(\int_0^{\cdot} V_i^{(k)}(u)^{\beta} \, \mathrm{d}u\right)^{-1}(s), \qquad i \ge 1, \ s \ge 0$$

Our aim is to show that

$$X(\tau(t+\cdot)) = \mathrm{ssFrag}_{\alpha-\beta}(X(\tau(t)), X^{(\cdot)} \circ \tau^{(\cdot)}).$$
(10)

Now let us fix $i \in \mathbb{N}$, and work conditional on $\mathcal{G}_{\tau(t)}$. On the event $\{V_i(\tau_i(t)) = 0\}$, then by definition of the operators ssFrag_{α}, the block containing *i* is constant in time and has mark 0 in both processes in (10), so there is equality for index *i*. Now we condition on $V_i(\tau_i(t)) = v$ with v > 0. Note that $\{V_i(\tau_i(t)) > 0\} \subset \{\tau_i(t) < \infty\}$, so in that case we have $\tau_i(t) < \infty$ almost surely, so there is the equality

$$\int_0^{\tau_i(t)} V_i(s)^\beta \,\mathrm{d}s = t.$$

Therefore we can write, for $s \ge 0$,

$$\tau_i(t+s) = \left(\int_0^{\cdot} V_i(u)^{\beta} \, \mathrm{d}u\right)^{-1} (t+s)$$

= $\tau_i(t) + \left(\int_0^{\cdot} V_i(\tau_i(t)+u)^{\beta} \, \mathrm{d}u\right)^{-1} (s).$

Now for all $u \ge 0$, because $X(\tau(t) + \cdot) = \text{ssFrag}_{\alpha}(X(\tau(t)), X^{(\cdot)})$, we have $V_i(\tau_i(t) + u) = vV_i^{(k)}(v^{\alpha}u)$, for *k* such that *i* is in the *k*-th block of $\Pi(\tau(t))$. This implies

$$\begin{aligned} \tau_i(t+s) - \tau_i(t) &= \left(\int_0^{\cdot} V_i(\tau_i(t)+u)^{\beta} \,\mathrm{d}u\right)^{-1}(s) \\ &= \left(\int_0^{\cdot} v^{\beta} V_i^{(k)}(v^{\alpha}u)^{\beta} \,\mathrm{d}u\right)^{-1}(s) \\ &= \left(v^{\beta-\alpha} \int_0^{v^{\alpha}} V_i^{(k)}(u)^{\beta} \,\mathrm{d}u\right)^{-1}(s) \\ &= v^{-\alpha} \left(\int_0^{\cdot} V_i^{(k)}(u)^{\beta} \,\mathrm{d}u\right)^{-1}(v^{\alpha-\beta}s) \\ &= V_i(\tau_i(t))^{-\alpha} \tau_i^{(k)}(V_i(\tau_i(t))^{\alpha-\beta}s). \end{aligned}$$

Now, defining $L_i(s)$ as the quantity given by the preceding display, the definition of the operators $ssFrag_{\alpha}$ yields for all $s \ge 0$,

$$\begin{aligned} X(\tau(t+s)) &= \mathrm{ssFrag}_{\alpha} \big(X(\tau(t)), X^{(\cdot)} \big) (L_i(s)) \\ &= \mathrm{ssFrag}_{\alpha-\beta} \big(X(\tau(t)), X^{(\cdot)} \circ \tau^{(\cdot)}) \big) (s). \end{aligned}$$

showing that $X \circ \tau$ is an $(\alpha - \beta)$ -ESSF.

Finally, note that if *X* is non-degenerate, then for all $x \in \mathcal{M}_{\infty}^{\star}$, \mathbb{P}_{x} -almost surely for any time $t \ge 0$, any block of X(t) is either infinite or has mark 0. So \mathbb{P}_{x} -almost surely, for all possible stopping lines *L*, the blocks of X(L) satisfy the same condition, i.e. $X(L) \in \mathcal{M}_{\infty}^{\star}$. Therefore, \mathbb{P}_{x} -almost surely $X \circ \tau$ has sample paths with values in $\mathcal{M}_{\infty}^{\star}$.

A.4 Proof of Theorem 13

Note that the whole process (X(t), $t \ge 0$) defines a coupling of all ξ_n for $n \ge 1$. By definition, one has

$$\xi_{n+1}(t) = \xi_n(t), \quad \forall t \le T_{n+1},$$

and at time T_{n+1} , either ξ_n is killed on the event $\{T_n = T_{n+1}\}$, or, conditional on $\{T_n > T_{n+1}\}$, the process ξ_n jumps, independently of the past, according to the probability

$$\eta_{n+1}(\cdot) := \mathcal{D}_{n+1}(\log v_1 \in \cdot \mid \pi_{\mid [n]} = \mathbf{1}_n \text{ and } v_1 \neq 0),$$

and goes on independently of the past, its remaining part $(\xi_n(T_{n+1}+t) - \xi_n(T_{n+1}), 0 \le t \le T_n - T_{n+1})$ being independent from ξ_{n+1} and equal in distribution to ξ_n by the strong Markov property. Let us first compute the probability p_n that $T_{n+1} = T_n$, which is by construction

$$p_n = \mathcal{D}_{n+1}(\pi_{||n|} \neq \mathbf{1}_n \text{ or } v_1 = 0).$$
(11)

From the previous description, one can write

$$T_n = T_{n+1} + ZT'_n,$$

where $Z = \mathbb{1}_{\{T_n \neq T_{n+1}\}} \sim \text{Be}(1 - p_n)$ is a Bernoulli random variable with parameter $(1 - p_n)$, and T'_n is a random variable equal in distribution to T_n , and independent from T_{n+1} and Z. Then, T_{n+1} , Z and T'_n are independent because Z is simply a function of the marked partition $D_{n+1} = (\Pi(T_{n+1}), \mathbf{V}(T_{n+1})/V_1(T_{n+1}-))_{|[n+1]}$ which is independent from ξ_{n+1} , and T'_n is independent of (ξ_{n+1}, D_{n+1}) because of the strong Markov property and the fact that $\alpha = 0$. Taking expectations yields

$$\frac{1}{J_n} = \frac{1}{J_{n+1}} + \frac{1 - p_n}{J_n},$$

which implies that $p_n = J_n/J_{n+1}$.

Now let us rebuild the coupling between ξ_n and ξ_{n+1} to show that their respective Lévy measures λ_n and λ_{n+1} satisfy

$$\lambda_n = \lambda_{n+1} + (J_{n+1} - J_n)\tilde{\eta}_{n+1},\tag{12}$$

where

$$\widetilde{\eta}_{n+1} := \eta_{n+1} \big(\cdot \cap \mathbb{R} \setminus \{0\} \big) = \mathcal{D}_{n+1} \big(\{ \log v_1 \in \cdot \} \cap \{v_1 \neq 1\} \mid \pi_{\mid [n]} = \mathbf{1}_n \text{ and } v_1 \neq 0 \big).$$
(13)

Consider the process ξ_{n+1} a Lévy process with characteristic exponent ψ_{n+1} , and let $T_n \sim \text{Exp}(J_n)$ independent. Now independently define $T' \sim \text{Exp}(J_{n+1} - J_n)$, and let $T_{n+1} := T_n \wedge T'$. We see that $T_{n+1} \sim \text{Exp}(J_{n+1})$, that it is independent of ξ_{n+1} and of the event $\{T_n = T_{n+1}\}$, which has probability $J_n/J_{n+1} = p_n$. Now conditional on (T_n, T_{n+1}) , let us define

$$D_{n+1} = \begin{cases} D' & \text{if } T_n = T_{n+1}, & \text{where } D' \sim \mathcal{D}_{n+1}(\cdot \mid \pi_{\mid [n]} \neq \mathbf{1}_n \text{ or } v_1 = 0) \\ D'' & \text{if } T_n \neq T_{n+1}, & \text{where } D'' \sim \mathcal{D}_{n+1}(\cdot \mid \pi_{\mid [n]} = \mathbf{1}_n \text{ and } v_1 \neq 0), \end{cases}$$

where D' and D'' are mutually independent and independent of everything else. Note that D_{n+1} is independent of T_{n+1} and of ξ_{n+1} , and because of (11), D_{n+1} has indeed distribution \mathcal{D}_{n+1} . Let us define $J = \log v_1$, where v_1 is the mark associated with the integer 1 in the marked partition D_{n+1} , and ξ_n a Lévy process with characteristic exponent ψ_n . Now putting everything together, define $\tilde{\xi}_{n+1}$ as the killed Lévy process ($\xi_{n+1}(t)$, $0 \le t < T_{n+1}$), and define $\tilde{\xi}_n$ as

$$\tilde{\xi}_n(t) = \begin{cases} \xi_{n+1}(t) & \text{if } t < T_{n+1} \\ \xi_{n+1}(T_{n+1}) + J + \xi_n(t - T_{n+1}) & \text{if } T_{n+1} \le t < T_n. \end{cases}$$

By construction, the joint distribution of $(\tilde{\xi}_n, \tilde{\xi}_{n+1})$ is equal to the one we get from the original process *X*, and it should now be clear that the point process of jumps of ξ_n is equal in distribution to the point process of jumps of ξ_{n+1} with additional jumps distributed as $J = \log v_1$, arising at rate $(J_{n+1} - J_n)$. Note that by construction, *J* has distribution η_{n+1} , so finally we have proven (12). The

fact that $(\lambda_n, n \ge 1)$ is a nonincreasing sequence of σ -finite measures ensures the existence of a limiting measure λ_{∞} on $\mathbb{R} \setminus \{0\}$ such that for all $n \in \mathbb{N}$,

$$\lambda_n = \lambda_\infty + \sum_{k>n} (J_k - J_{k-1}) \widetilde{\eta}_k.$$
(14)

Recall that we wrote the characteristic exponent of ξ_n in the following way:

$$\psi_n(\theta) = id_n\theta - \frac{\beta_n}{2}\theta^2 + \int_{\mathbb{R}} \left(e^{i\theta y} - 1 - i\theta y \mathbb{1}_{|y| \le 1} \right) \lambda_n(\mathrm{d}y).$$

From the previous discussion, one can construct a coupling between the two Lévy processes such that $(\xi_n(t) - \xi_{n+1}(t), t \ge 0)$ is simply a compound Poisson process with jump measure $(J_{n+1} - J_n)\tilde{\eta}_{n+1}$, so it is clear that necessarily $\beta_n = \beta_{n+1}$, and

$$d_n = d_{n+1} + (J_{n+1} - J_n) \int_{|y| \le 1} y \, \widetilde{\eta}_{n+1}(\mathrm{d}y) = d_{n+1} + \int_{|y| \le 1} y \, (\lambda_n - \lambda_{n+1})(\mathrm{d}y).$$

To summarize, letting $\beta := \beta_1$, the following holds for all $n \in \mathbb{N}$

$$\beta_n = \beta$$
 and $d_n = d_1 - \int_{|y| \le 1} y (\lambda_1 - \lambda_n) (dy),$ (15)

where $(\lambda_1 - \lambda_n)$ denotes the positive measure given by

$$(\lambda_1 - \lambda_n) = \sum_{k=2}^n (J_k - J_{k-1}) \widetilde{\eta}_k$$

Let us now examine the consistency properties of the measures \mathcal{D}_n . From this point on, for the sake of clarity, we decompose the proof in a series of steps.

Step 1. We prove the existence and uniqueness of a measure \mathcal{D} on \mathcal{M}_{∞} satisfying

$$\mathcal{D}(\pi = \mathbf{1} \text{ and } v_1 \neq 0) = 0 \tag{16}$$

and such that for all $n \in \mathbb{N}$,

$$\mathcal{D}(\{x_{|[n]} \in \cdot\} \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\}) = J_n \mathcal{D}_n,$$
(17)

then we show that this measure is exchangeable.

First, note that for the previous construction to be consistent, the random variable $D'_{[n]}$ must have distribution \mathcal{D}_n . Indeed, on the event $\{T_{n+1} < T_n\}$, the strong Markov property at time T_{n+1} implies that the process $X_{[n]}$ jumps according to \mathcal{D}_n , independently of the past, so on the complement this must hold as well, so

$$\mathcal{D}_{n+1}(x_{|[n]} \in \cdot \mid \pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0) = \mathcal{D}_n,$$

which can be rewritten

$$J_{n+1}\mathcal{D}_{n+1}(\{x_{|[n]} \in \cdot\} \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\}) = J_n \mathcal{D}_n.$$
(18)

Now for all integers $n \leq m$, let us define a measure on \mathcal{M}_m by

$$\mu_n^m := J_m \mathcal{D}_m \big(\cdot \cap \{ \pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = \mathbf{0} \} \big), \quad m \ge n.$$

Let us prove that for all integers $n \le k \le m$, we have

$$\mu_n^m(x_{|[k]} \in \cdot) = \mu_n^k. \tag{19}$$

Note that there is nothing to prove in the case k = m. Now suppose this is proven for fixed $n \le k \le m$. Then,

$$\mu_n^{m+1}(x_{|[k]} \in \cdot) = J_{m+1}\mathcal{D}_{m+1}(\{x_{|[k]} \in \cdot\} \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\})$$

= $J_{m+1}\mathcal{D}_{m+1}(\{(x_{|[m]})_{|[k]} \in \cdot\} \cap \{(\pi_{|[m]})_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\})$
 $\cap \{\pi_{|[m]} \neq \mathbf{1}_m \text{ or } v_1 = 0\})$
= $J_m\mathcal{D}_m(\{x_{|[k]} \in \cdot\} \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\})$
= $\mu_n^m(x_{|[k]} \in \cdot) = \mu_n^k,$

where we have used (18) and the fact that $\{\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\} \subset \{\pi_{|[m]} \neq \mathbf{1}_m \text{ or } v_1 = 0\}$. By induction on *m* this proves (19) for any integers $n \leq k \leq m$. Note that in particular, taking k = n, we see that the total mass of μ_n^m is equal to that of μ_n^n , which is J_n . In summary, for any $n \in \mathbb{N}$, the sequence $(\mathcal{M}_m, \mu_n^m/J_n, m \geq n)$ defines a inverse system of compact probability spaces, and by the Kolmogorov extension theorem, there exists a unique measure μ_n (with total mass J_n) on the inverse limit $\lim_{k \to m} \mathcal{M}_m = \mathcal{M}_\infty$ such that for each $m \geq n$, $\mu_n(x_{|[m]} \in \cdot) = \mu_n^m$. Now notice that by definition, for any integers $n_1 \leq n_2 \leq m$, we have

$$\mu_{n_2}^m$$
 (· $\cap \{\pi_{|[n_1]} \neq \mathbf{1}_{n_1} \text{ or } v_1 = 0\}$) = $\mu_{n_1}^m$,

which implies by construction

$$\mu_{n_2}(\cdot \cap \{\pi_{|[n_1]} \neq \mathbf{1}_{n_1} \text{ or } v_1 = 0\}) = \mu_{n_1}$$

This means that the sequence of measures $(\mu_n, n \ge 1)$ on \mathcal{M}_{∞} is increasing, and one can define the limit as \mathcal{D} . This measure then satisfies by construction

$$\mathcal{D}(\{x_{\mid [n]} \in \cdot\} \cap \{\pi_{\mid [n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\}) = \mu_n(x_{\mid [n]} \in \cdot)$$
$$= \mu_n^n$$
$$= J_n \mathcal{D}_n,$$

which is indeed (17).

Secondly, note that since for any $n \in \mathbb{N}$, clearly $\mu_n(\pi = 1 \text{ and } v_1 \neq 0) = 0$, where μ_n are the measures defined above, so in the limit (16) holds. Let us now show uniqueness. If a measure \mathcal{D}' on \mathcal{M}_{∞} satisfies (17) and (16) then

$$\mathcal{D}'(\cdot \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\}) = \mu_n,$$

and letting $n \to \infty$, $\mathcal{D}' = \mathcal{D}$, which proves uniqueness.

Finally, \mathcal{D} is exchangeable. Indeed if $\sigma : \mathbb{N} \to \mathbb{N}$ is a permutation, let $m \in \mathbb{N}$ such that $\sigma(k) = k$ for all $k \ge m$. Now for all $n \ge m$, using the exchangeability of the probability measures ($\mathcal{D}_k, k \ge 1$), we get

$$\mathcal{D}((x^{\sigma})_{|[n]} \in \cdot) = \lim_{k \to \infty} \mathcal{D}\left(\{(x^{\sigma})_{|[n]} \in \cdot\} \cap \{\pi_{|[k]} \neq \mathbf{1}_k \text{ or } v_1 = 0\}\right)$$
$$= \lim_{k \to \infty} J_k \mathcal{D}_k((x^{\sigma})_{|[n]} \in \cdot)$$
$$= \lim_{k \to \infty} J_k \mathcal{D}_k(x_{|[n]} \in \cdot)$$
$$= \lim_{k \to \infty} \mathcal{D}(\{x_{|[n]} \in \cdot\} \cap \{\pi_{|[k]} \neq \mathbf{1}_k \text{ or } v_1 = 0\})$$
$$= \mathcal{D}(x_{|[n]} \in \cdot).$$

As this is true for all $n \ge m$, necessarily $\mathcal{D}(x^{\sigma} \in \cdot) = \mathcal{D}$, i.e. \mathcal{D} is exchangeable.

Step 2. We prove that $\mathcal{D}(\mathcal{M}_{\infty} \setminus \mathcal{M}_{\infty}^{\star}) = 0$ by using that *X* is non-degenerate. For this, we need to show first that the process $(B_1(t), t \ge 0)$ of the block of *X* containing 1 is equal in distribution to a process $(B(t), t \ge 0)$ constructed from a Poisson point process of intensity $dt \otimes \mathcal{D}$.

More precisely, define \mathcal{N} a Poisson point process on $[0, \infty) \times \mathcal{M}_{\infty}$ with intensity $dt \otimes (\mathcal{D} + \widetilde{\lambda}_{\infty})$, where $\widetilde{\lambda}_{\infty}$ is the push-forward of λ_{∞} by the map $y \in \mathbb{R} \mapsto (\mathbf{1}, e^y) \in \mathcal{M}_{\infty}^{\star}$. Let us define

$$\mathcal{N}' := \{ (t, \log v_1), (t, x) \in \mathcal{N} \text{ with } x = (\pi, \mathbf{v}) \text{ and } v_1 \notin \{0, 1\} \},\$$

which is then a Poisson point process on $[0, \infty) \times \mathbb{R}$ with intensity

$$\mathrm{d} t \otimes \Big(\mathcal{D}\big(\{ \log v_1 \in \cdot \} \cap \{ v_1 \notin \{ 0, 1 \} \} \big) + \lambda_{\infty} \Big).$$

However, note that using (16), (17) and finally (13), we get

$$\mathcal{D}(\{\log v_1 \in \cdot\} \cap \{v_1 \notin \{0, 1\}\})$$

= $\sum_{n \in \mathbb{N}} \mathcal{D}(\{\log v_1 \in \cdot\} \cap \{\pi_{|[n]} = \mathbf{1}_n \text{ and } v_1 \notin \{0, 1\}\} \cap \{\pi_{|[n+1]} \neq \mathbf{1}_{n+1}\})$
= $\sum_{n \in \mathbb{N}} J_{n+1} \mathcal{D}_{n+1}(\{\log v_1 \in \cdot\} \cap \{\pi_{|[n]} = \mathbf{1}_n \text{ and } v_1 \notin \{0, 1\}\})$
= $\sum_{n \in \mathbb{N}} (J_{n+1} - J_n) \widetilde{\eta}_{n+1},$

and by (14), we find $\mathcal{D}(\{\log v_1 \in \cdot\} \cap \{v_1 \notin \{0, 1\}\}) + \lambda_{\infty} = \lambda_1$. Therefore, it is possible to define a Lévy process ξ with characteristic exponent ψ_1 , such that the point process of its jumps is precisely \mathcal{N}' . Let us also define a process $B = (B(t), t \ge 0)$ with càdlàg sample paths with values in $2^{\mathbb{N}}$ the subsets of \mathbb{N} , such that B has the distribution of $(B_1(t), t \ge 0)$ the block containing 1 in X. First define T as the first time $t \in [0, \infty)$ such that there is an atom $(t, (\pi, \mathbf{v})) \in \mathcal{N}$ with $v_1 = 0$. If there is none, then let $T = \infty$. Then, for each $n \in \mathbb{N}$, let $(t_1, x_1), (t_2, x_2), \ldots$ be the whole sequence (finite or infinite) of atoms of \mathcal{N} with $t_1 < t_2 < \ldots \leq T$ such that for each $i, x_i = (\pi_i, \mathbf{v}_i)$ with $(\pi_i)_{|[n]} \neq \mathbf{1}_n$ and $(v_i)_1 > 0$ (or such that $(v_i)_1 = 0$ for the possible last atom, at time T). We can define $\widetilde{B}_0^n = [n]$, and inductively for each $i \ge 1$

$$\widetilde{B}_i^n := \widetilde{B}_{i-1}^n \cap (A_i \cap [n]),$$

where A_i is the block of π_i containing 1. Now define, for $t \in [0, \infty)$,

$$B^n(t) = \overline{B}^n_i \text{ if } t \in [t_i, t_{i+1}).$$

where we let $t_0 := 0$, and in the case $T < \infty$, i.e. if the sequence of atoms is finite, say with length $k \in \mathbb{N}$, we let $t_{k+1} := \infty$. It is readily checked that this construction is consistent in the sense that for each $t \ge 0$ there is a single $B(t) \in 2^{\mathbb{N}}$ such that $B^n(t) = B(t) \cap [n]$. Let us show that this process $(B(t), \xi(t), t \ge 0)$ has the same distribution as the marked block containing 1 in X, i.e. $(B_1(t), \log V_1(t), t \ge 0)$. For fixed $n \in \mathbb{N}$ and $x \in \mathcal{M}_n$, recall that

$$X_{|[n]}$$
 under $\mathbb{P}_x \stackrel{(a)}{=} \operatorname{ssFrag}_0(x, X^{(\cdot)}),$

where $X^{(\cdot)}$ is an independent i.i.d. sequence of copies of X started from (1, 1). Using the same notation, for any $A \subset [n]$ with $1 \in A$ and v > 0, the law of the process $(B_1(t) \cap [n], \log V_1(t), t \ge 0)$ started from $(A, \log v)$ can be deduced from that of $X^{(1)} = (\Pi^{(1)}, \mathbf{V}^{(1)})$. More precisely, $\log V_1(t)$ behaves as a Lévy process with characteristic exponent ψ_n started from $\log v$, until an independent time $T_n \sim \operatorname{Exp}(J_n)$ when $\Pi^{(1)}_{|[n]}$ first jumps. At that time, $D_n^{(1)} = (\Pi^{(1)}(T_n), \mathbf{V}^{(1)}(T_n)/V_1(T_n-))_{|[n]}$ is independently drawn according to \mathcal{D}_n and then writing $D_n^{(1)} = (\pi, \mathbf{v})$,

$$\log V_1(T_n) = \log V_1(T_n) + \log v_1$$
 and $B_1(T_n) = B_1(T_n) \cap A_1 = A \cap A_1$

where A_1 is the block of π containing 1. Note that there is a non-zero probability that $B_1(T_n) = B_1(T_n-)$ (even with $v_1 > 0$) when $A \neq [n]$. Now in our construction, if (t_1, x) is the first atom of N such that $\pi \neq \mathbf{1}_n$ or $v_1 = 0$, where $x = (\pi, \mathbf{v})$, then t_1 is exponentially distributed with parameter $\mathcal{D}(\pi_{||n|} \neq \mathbf{1}_n$ or $v_1 = 0) = J_n$, and x is independent of t_1 , distributed as

$$\frac{1}{J_n}\mathcal{D}(\cdot \cap \{\pi \neq \mathbf{1}_n \text{ or } v_1 = 0\}),$$

so that $x_{|[n]}$ has distribution \mathcal{D}_n . It remains only to show that $(\xi(s), 0 \le s < t_1)$ is distributed as a Lévy process with characteristic exponent ψ_n (killed at t_1). The point process of its jumps is

$$\mathcal{N}' \cap [0, t_1) \times \mathbb{R},$$

which conditional on t_1 has intensity

$$\mathrm{d} t \otimes \Big(\mathcal{D}\big(\{ \log v_1 \in \cdot \} \cap \{ \pi_{|[n]} = \mathbf{1}_n \text{ and } v_1 \notin \{0, 1\} \} \big) + \lambda_{\infty} \Big).$$

Let us show that this intensity is equal to $dt \otimes \lambda_n$. Note that

$$\mathcal{D}(\{\log v_1 \in \cdot\} \cap \{\pi_{|[n]} = \mathbf{1}_n \text{ and } v_1 \notin \{0, 1\}\})$$

$$= \sum_{m \ge n} \mathcal{D}(\{\log v_1 \in \cdot\} \cap \{\pi_{|[m]} = \mathbf{1}_m \text{ and } v_1 \notin \{0, 1\}\} \cap \{\pi_{|[m+1]} \neq \mathbf{1}_{m+1}\})$$

$$= \sum_{m \ge n} J_{m+1} \mathcal{D}_{m+1}(\{\log v_1 \in \cdot\} \cap \{\pi_{|[m]} = \mathbf{1}_m \text{ and } v_1 \notin \{0, 1\}\})$$

$$= \sum_{m \ge n} (J_{m+1} - J_m) \widetilde{\eta}_{m+1},$$

so (14) shows that

$$\lambda_n = \lambda_\infty + \mathcal{D}\big(\{\log v_1 \in \cdot\} \cap \{\pi_{|[n]} = \mathbf{1}_n \text{ and } v_1 \notin \{0, 1\}\}\big)$$
(20)

therefore the Lévy measure of ($\xi(s)$, $0 \le s < t_1$) is indeed λ_n . In the end, we have shown that

$$(B(t), \xi(t), t \ge 0) \stackrel{(d)}{=} (B_1(t), \log V_1(t), t \ge 0).$$

From this construction, we see that for each atom $(t, x) \in N$, the process *B* jumps, with

$$B(t) = B(t-) \cap A,$$

where *A* is the block of *x* containing 1. Let us show that this implies $\mathcal{D}(\mathcal{M}_{\infty} \setminus \mathcal{M}_{\infty}^{\star}) = 0$. Assuming the opposite, there is a non-zero probability that there is an atom $(t, x) \in \mathcal{N}$ with t < T such that *x* contains a finite block with mark not equal to zero. By exchangeability of \mathcal{D} , and from the description of the jumps of *B*, there is a non-zero probability that there is a jump $B(t) = B(t-) \cap A$ where *A* is finite and t < T. This contradicts the assumption of non-degeneracy of *X*, as then we would have $X(t) \in \mathcal{M}_{\infty} \setminus \mathcal{M}_{\infty}^{\star}$.

From now on, we view \mathcal{D} as an exchangeable measure on $\mathcal{M}^{\star}_{\infty}$, satisfying (16) and the σ -finiteness assumption

$$\forall n \in \mathbb{N}, \quad \mathcal{D}(\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0) = J_n < \infty.$$

It remains essentially to study $\mathcal D$ in order to express it as a mixture of paintbox processes.

Step 3. Let us decompose

 $\mathcal{D} = \mathcal{D}\big(\cdot \cap \{\pi = 1\}\big) + \mathcal{D}\big(\cdot \cap \{\pi \neq 1 \text{ and } |\pi|^{\downarrow} = 1\}\big) + \mathcal{D}\big(\cdot \cap \{\pi \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}\big)$

and show that there exist $c \ge 0$ a constant and Λ' a σ -finite measure on \mathscr{Z}^{\downarrow} such that

(a) $\mathcal{D}(\cdot \cap \{\pi = 1\}) = \mathcal{D}(\pi = 1)\delta_{(1,0)},$

(b) $\mathcal{D}(\cdot \cap \{\pi \neq 1 \text{ and } |\pi|^{\downarrow} = 1\}) = c \sum_{n \in \mathbb{N}} \delta_{e_n}$, where

$$\mathbf{e}_n := \left(\left\{ \{n\}, \mathbb{N} \setminus \{n\} \right\}, \ (1, \ldots, 1, \underbrace{0}_{n-\text{th index}}, 1, \ldots) \right) \in \mathcal{M}_{\infty}^{\star},$$

(c)
$$\mathcal{D}(\cdot \cap \{\pi \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}) = \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}(\cdot) \Lambda'(\mathrm{d}\mathbf{z}),$$

We use similar arguments as in [5, Theorem 3.1], as we have already done in the context of nested fragmentations [12, Proposition 19]. First note that by (16), \mathcal{D} -a.e. on the event { $\pi = 1$ } we have x = (1, 0), in other words (a) holds. Note also that $\mathcal{D}(\pi = 1) \leq \mathcal{D}(v_1 = 0) = J_1 < \infty$.

Let us now study the measure $\mathcal{D}(\cdot \cap \{\pi \neq 1\})$. Note that $\mathcal{D}(\{\pi \in \cdot\} \cap \{\pi \neq 1\})$ is an exchangeable measure on \mathcal{P}_{∞} satisfying for all $n \geq 1$,

$$\mathcal{D}(\pi_{|[n]} \neq \mathbf{1}_n) < \infty.$$

A consequence of [5, Theorem 3.1] is that π has asymptotic frequencies \mathcal{D} -a.e. – recall that $|\pi|^{\downarrow} \in \mathscr{S}^{\downarrow} \subset [0, 1]^{\mathbb{N}}$ denotes the nonincreasing reordering of asymptotic frequencies of blocks of π – and one can write

$$\mathcal{D}(\{\pi \in \cdot\} \cap \{\pi \neq \mathbf{1} \text{ and } |\pi|^{\downarrow} = (1, 0, 0, \ldots)\}) = c \sum_{n \in \mathbb{N}} \delta_{\{\{n\}, \mathbb{N} \setminus \{n\}\}}$$

where $c \ge 0$. For conciseness – and again with some abuse of notation – we will from now on let $\mathbf{1} := (1, 0, 0...) \in \mathscr{S}^{\downarrow}$. Now let us examine the distribution of $x = (\pi, \mathbf{v})$ on the event $\{\pi = \{\{n\}, \mathbb{N} \setminus \{n\}\}\}$. Since $x \in \mathcal{M}^{\star}_{\infty} \mathcal{D}$ -a.e., the singleton block must have mark 0, while the other block may have a positive mark. Let η be the distribution of this mark on S^1 , that is

$$\eta := \mathcal{D}(\{v_1 \in \cdot\} \cap \{\pi = \{\{n\}, \mathbb{N} \setminus \{n\}\}\}),$$

which is a measure of total mass *c*, for any fixed n > 1 (by exchangeability, η does not depend on the value of *n*). First, note that $\eta(\{0\}) = 0$. Indeed, since the events

$$\left\{\pi = \left\{\{n\}, \mathbb{N} \setminus \{n\}\right\}\right\}, \qquad n > 1$$

are disjoint, the following holds.

$$\sum_{n>1} \eta(\{0\}) = \sum_{n>1} \mathcal{D}(\left\{v_1 = 0 \text{ and } \pi = \{\{n\}, \mathbb{N} \setminus \{n\}\}\right\})$$
$$= \mathcal{D}\left(\bigcup_{n>1} \left\{v_1 = 0 \text{ and } \pi = \{\{n\}, \mathbb{N} \setminus \{n\}\}\right\}\right)$$
$$\leq \mathcal{D}(v_1 = 0)$$
$$= J_1 < \infty,$$

which implies necessarily $\eta(\{0\}) = 0$. Now recall that $\mathcal{D}(\{\log v_1 \in \cdot\} \cap \{v_1 \notin \{0, 1\}\})$ is a Lévy measure, so for all $\varepsilon > 0$,

$$\sum_{n>1} \eta(|\log v_1| > \varepsilon) = \sum_{n>1} \mathcal{D}(\{|\log v_1| > \varepsilon \text{ and } \pi = \{\{n\}, \mathbb{N} \setminus \{n\}\}\})$$
$$= \mathcal{D}(\bigcup_{n>1} \{|\log v_1| > \varepsilon \text{ and } \pi = \{\{n\}, \mathbb{N} \setminus \{n\}\}\})$$
$$\leq \mathcal{D}(|\log v_1| > \varepsilon \text{ and } v_1 \neq 0) < \infty,$$

which implies necessarily $\eta(|\log v_1| > \varepsilon) = 0$. Letting $\varepsilon \to 0$, we have $\eta(|\log v_1| > 0) = 0$, so in the end $\eta = c\delta_1$, and (b) follows.

Let us now decompose the measure $\mathcal{D}(\cdot \cap \{|\pi|^{\downarrow} \neq 1\})$. Recall that by construction,

$$\mathcal{D}(\cdot \cap \{\pi_{|[n]} \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}) \leq \mathcal{D}(\cdot \cap \{\pi_{|[n]} \neq 1 \text{ or } v_1 = 0\}) = \mu_n,$$

which is a finite measure with total mass J_n . Now let us introduce the injection $\theta_n : \mathbb{N} \to \mathbb{N}, k \mapsto n + k$, and consider

$$\overleftarrow{\mu}_n := \mathcal{D}(\{x^{\theta_n} \in \cdot\} \cap \{\pi_{|[n]} \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}),$$

which can be seen as the distribution of the marked partition restricted to $\{n + 1, n + 2, ...\}$, on the event $\{\pi_{|[n]} \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}$. It is readily checked that this measure is exchangeable on $\mathcal{M}_{\infty}^{\star}$ and since it is finite, by Proposition 2,

$$\overleftarrow{\mu}_n = \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}(\cdot) \Lambda_n(\mathrm{d}\mathbf{z}),$$

with $\Lambda_n = \overleftarrow{\mu}_n(|x|^{\downarrow} \in \cdot)$ a finite measure on \mathscr{Z}^{\downarrow} . Asymptotic frequencies are such that $|x|^{\downarrow} = |x^{\theta_n}|^{\downarrow}$ for all $x \in \mathcal{M}^{\star}_{\infty}$, therefore Λ_n can also be written

$$\Lambda_n = \mathcal{D}(\{|x|^{\downarrow} \in \cdot\} \cap \{\pi_{|[n]} \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}),$$

and taking nondecreasing limits one can define

$$\Lambda' := \mathcal{D}\big(\{|x|^{\downarrow} \in \cdot\} \cap \{\pi \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}\big).$$

To show (c), fix $k, n \in \mathbb{N}$ and consider the permutation $\tau : \mathbb{N} \to \mathbb{N}$ given by

$$\tau(i) = \begin{cases} i+k & \text{if } i \le n \\ i-n & \text{if } n < i \le n+k \\ i & \text{otherwise.} \end{cases}$$

Now notice that

$$\mathcal{D}\big(\{x_{|[k]} \in \cdot\} \cap \{\pi \neq 1 \text{ and } |\pi|^{\downarrow} \neq 1\}\big) = \lim_{n \to \infty} \mathcal{D}\big(\{x_{|[k]} \in \cdot\} \cap \{(\pi^{\theta_k})_{|[n]} \neq 1_n \text{ and } |\pi|^{\downarrow} \neq 1\}\big),$$

which can be written, using the exchangeability of \mathcal{D} ,

$$\mathcal{D}\left(\{x_{|[k]} \in \cdot\} \cap \{(\pi^{\theta_k})_{|[n]} \neq \mathbf{1}_n \text{ and } |\pi|^{\downarrow} \neq \mathbf{1}\}\right)$$

= $\mathcal{D}\left(\{(x^{\tau \circ \theta_n})_{|[k]} \in \cdot\} \cap \{(\pi^{\tau})_{|[n]} \neq \mathbf{1}_n \text{ and } |\pi^{\tau}|^{\downarrow} \neq \mathbf{1}\}\right)$
= $\mathcal{D}\left(\{(x^{\theta_n})_{|[k]} \in \cdot\} \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ and } |\pi|^{\downarrow} \neq \mathbf{1}\}\right)$
= $\int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}^k(\cdot) \Lambda_n(d\mathbf{z}),$

where ρ_z^k is the paintbox process restricted to *k* elements defined in Section 1.2. Taking limits and because *k* is generic, we have indeed (c).

Step 4. It remains to define the measure Λ correctly and we will be able to complete the proof of Theorem 13. Recall the definition of $\tilde{\lambda}_{\infty}$ as the push-forward of λ_{∞} by the map $y \in \mathbb{R} \mapsto (1, e^y) \in \mathcal{M}_{\infty}^{\star}$, and note that

$$\widetilde{\lambda}_{\infty} = \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}(\cdot) \, \widehat{\lambda}_{\infty}(\mathrm{d}\mathbf{z}),$$

where $\widehat{\lambda}_{\infty}$ is the push-forward of λ_{∞} by the map $y \in \mathbb{R} \mapsto (1, e^y) \in \mathscr{Z}^{\downarrow}$. In the end, let us define

$$\Lambda = \Lambda' + \mathcal{D}(\pi = \mathbf{1})\delta_{(\mathbf{1},0)} + \widehat{\lambda}_{\infty}.$$

Putting everything together, we have

$$\mathcal{D} + \widetilde{\lambda}_{\infty} = c \sum_{n \in \mathbb{N}} \delta_{\mathfrak{e}_n} + \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}(\cdot) \Lambda(d\mathbf{z}).$$

We can almost complete the proof. First, fix $n \in \mathbb{N}$ and recall that by definition $J_n = \mathcal{D}(\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0)$. Since by definition $\widetilde{\lambda}_{\infty}(\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0) = 0$ for all n, point (ii) of Theorem 13 is proven:

$$J_n = nc + \int_{\mathscr{Z}^{\downarrow}} \left(1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n \right) \Lambda(\mathrm{d}\mathbf{z}).$$

This implies that $(1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n)$ is Λ -integrable for any $n \ge 1$. Furthermore, note that for any $\mathbf{z} = (\mathbf{s}, \mathbf{v}) \in \mathscr{Z}^{\downarrow}$,

$$1 - s_1 \le 1 - s_1 \left(\sum_{\substack{i \ge 1 \\ v_i > 0}} s_i\right) \le 1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^2, \quad \text{and} \quad s_1 \mathbb{1}_{v_1 = 0} \le 1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i,$$

therefore summing the two expressions yields

$$\int_{\mathscr{Z}^{\downarrow}} 1 - s_1 \mathbb{1}_{v_1 > 0} \Lambda(\mathrm{d}\mathbf{z}) < \infty.$$
(21)

We keep this in mind for later use and go back to the expression of the measures \mathcal{D}_n . If $J_n > 0$, then $\mathcal{D}_n = \mathcal{D}(\{x_{|[n]} \in \cdot\} \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ or } v_1 = 0\})/J_n$, so point (iii) of Theorem 13 is proven:

$$\mathcal{D}_n = \frac{1}{J_n} \Big(\sum_{i=1}^n c \delta_{\mathbf{e}_i^n} + \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}^n \big(\cdot \cap \{ \pi \neq \mathbf{1}_n \text{ or } v_1 = 0 \} \big) \Lambda(\mathrm{d}\mathbf{z}) \Big),$$

where $e_i^n \in \mathcal{M}_n$ is defined as

$$e_i^n := \left(\{ [n] \setminus \{i\}, \{i\} \}, (1, \dots, 1, \underbrace{0}_{i-\text{th index}}, 1, \dots, 1) \right).$$

It remains for the first part of the theorem to express ψ_n correctly and to show the integrability condition (4). Recall from (20) that $\lambda_n = \lambda_\infty + \mathcal{D}(\{\log v_1 \in \cdot\} \cap \{\pi_{|[n]} = \mathbf{1}_n \text{ and } v_1 \notin \{0, 1\}\})$, which shows that

$$\psi_n(\theta) = id_n\theta - \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{j \ge 1 \\ v_j > 0}} s_j^n \left(e^{i\theta \log v_j} - 1 - i\theta \log v_j \mathbb{1}_{|\log v_j| \le 1} \right) \Lambda(\mathrm{d}\mathbf{z}),$$

but, from (15), we have

$$d_n = d_1 - \int_{|y| \le 1} y (\lambda_1 - \lambda_n) (dy)$$

= $d_1 - \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{j \ge 1 \\ v_j > 0}} s_j (1 - s_j^{n-1}) \log v_j \mathbb{1}_{|\log v_j| \le 1} \Lambda(d\mathbf{z})$

where we used $(\lambda_1 - \lambda_n) = \mathcal{D}(\{\log v_1 \in \cdot\} \cap \{\pi_{|[n]} \neq \mathbf{1}_n \text{ and } v_1 \neq 0\})$ which is again deduced from (20). Putting the last two displays together, we get

$$\psi_n(\theta) = id_1\theta - \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{j \ge 1 \\ v_j > 0}} \left(s_j^n \left(e^{i\theta \log v_j} - 1 \right) - i\theta s_j \log v_j \mathbb{1}_{|\log v_j| \le 1} \right) \Lambda(\mathrm{d}\mathbf{z}).$$

In order to simplify this notation, note that

$$\left|\sum_{\substack{j\geq 1\\v_j>0}} s_j \log v_j \mathbb{1}_{|\log v_j|\leq 1} - \log v_1 \mathbb{1}_{|\log v_1|\leq 1}\right| \leq (1 - s_1 \mathbb{1}_{v_1>0}) |\log v_1| \mathbb{1}_{|\log v_1|\leq 1} + \sum_{\substack{j\geq 2\\v_j>0}} s_j \leq 2(1 - s_1 \mathbb{1}_{v_1>0}),$$

which we proved to be Λ -integrable in (21). Therefore, we can finally define

$$d := d_1 + \int_{\mathscr{Z}^{\downarrow}} \log v_1 \mathbb{1}_{|\log v_1| \le 1} - \sum_{\substack{j \ge 1 \\ v_j > 0}} s_j \log v_j \mathbb{1}_{|\log v_j| \le 1} \Lambda(\mathrm{d}\mathbf{z})$$

in order to get point (i) of Theorem 13, that is

$$\psi_n(\theta) = id\theta - \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{j \ge 1 \\ v_j > 0}} s_j^n \left(e^{i\theta \log v_j} - 1 \right) - i\theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} \Lambda(d\mathbf{z}).$$

Now let us show (4). From (21), it remains only to check that $(\log v_1)^2 \wedge 1$ is Λ -integrable. Since λ_1 must be a Lévy measure, we have

$$\begin{split} \int_{\mathbb{R}} (y^2 \wedge 1) \,\lambda_1(\mathrm{d} y) &= \int_{\mathscr{Z}^{\downarrow}} \int_{\mathbb{R}} (y^2 \wedge 1) \,\varrho_{\mathbf{z}}^1 \big\{ \log v_1 \in \mathrm{d} y \big\} \cap \{v_1 \neq 0\} \big) \,\Lambda(\mathrm{d} \mathbf{z}) \\ &= \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{i \geq 1 \\ v_i > 0}} s_i \big((\log v_i)^2 \wedge 1 \big) \,\Lambda(\mathrm{d} \mathbf{z}) < \infty. \end{split}$$

Now note that for all $\mathbf{z} \in \mathscr{Z}^{\downarrow}$,

$$(\log v_1)^2 \wedge 1 \le s_1 \mathbb{1}_{v_1 > 0} \left((\log v_1)^2 \wedge 1 \right) + (1 - s_1 \mathbb{1}_{v_1 > 0})$$

$$\le \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n \left((\log v_i)^2 \wedge 1 \right) + (1 - s_1 \mathbb{1}_{v_1 > 0}),$$

which is Λ -integrable. This proves (4), and ends the proof of the main result of Theorem 13.

For the converse part, let (c, d, β, Λ) be a given quadruple, where $c, \beta \ge 0, d \in \mathbb{R}$, and Λ is a measure on $\mathscr{Z}^{\downarrow} \setminus \{(1, 1)\}$ satisfying (4). Then $(\psi_n, J_n, \mathcal{D}_n)$ for all $n \in \mathbb{N}$ are well-defined as in the theorem if one checks that

$$\int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n \left((\log v_i)^2 \wedge 1 \right) \Lambda(\mathrm{d}\mathbf{z}) < \infty$$
and
$$\int_{\mathscr{Z}^{\downarrow}} \left(1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n \right) \Lambda(\mathrm{d}\mathbf{z}) < \infty.$$
(22)

To that aim, note that for all $\mathbf{z} \in \mathscr{Z}^{\downarrow}$,

$$\begin{split} \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n \big((\log v_i)^2 \wedge 1 \big) + \Big(1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n \Big) &\leq s_1^n \big((\log v_1)^2 \wedge 1 \big) + \sum_{\substack{i \ge 2 \\ v_i > 0}} s_i^n + \Big(1 - \sum_{\substack{i \ge 1 \\ v_i > 0}} s_i^n \Big) \\ &\leq \big((\log v_1)^2 \wedge 1 \big) + (1 - s_1^n \mathbb{1}_{v_1 > 0}) \\ &\leq \big((\log v_1)^2 \wedge 1 \big) + n(1 - s_1 \mathbb{1}_{v_1 > 0}), \end{split}$$

which is Λ -integrable by (4), so (22) is proven. Now the construction of a 0-ESSF *X* with characteristics as above is done via Remark 12.

A.5 Proof of Proposition 16

Consider a non-degenerate homogeneous ESSF $X = (\Pi, \mathbf{V})$ with characteristics (c, d, β, Λ) . We make use of a natural genealogy appearing in our construction: jointly for all $n \in \mathbb{N}$, we define processes $(F_n(t), t \ge 0)$ taking values in the subsets of \mathbb{N} , starting at $F_n(0) = [n]$, whose role is to "follow" integers along a discrete genealogy of blocks. We will make this statement more precise, but first let us explain the idea. We will build the F_n deterministically from a sample path of X, such that for each time $t \ge 0$,

$$[n] \subset F_n(t) \subset F_{n+1}(t),$$

and so by defining

$$S_{\theta}^{(n)}(t) = \sum_{B \text{ block of } X(t)|_{F_n(t)}} \tilde{V}_B(t)^{\theta},$$
(23)

where $\tilde{V}_B(t)$ denotes the mark of a block *B* of *X*(*t*), it is clear that

$$S_{\theta}^{(n)}(t) \leq S_{\theta}^{(n+1)}(t) \xrightarrow[n \to \infty]{} S_{\theta}(X(t)) = \sum_{B \text{ block of } X(t)} \tilde{V}_{B}(t)^{\theta}.$$

The way to define the sets F_n is the following. Recall that for any $n \in \mathbb{N}$, T_n denotes the first time t when [n] is no longer part of a unique block with positive mark in X(t). Therefore let $F_n(t) = [n]$ for any $0 \le t < T_n$. Since X is homogeneous, T_n is an exponential random variable with parameter J_n , and conditional on T_n , the mark $(V_1(t), 0 \le t < T_n)$ behaves as the exponential of a Lévy process ξ_n with characteristic exponent

$$\psi_n: \theta \mapsto id\theta - \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^\downarrow} \sum_{\substack{j \ge 1 \\ v_i > 0}} s_j^n (e^{i\theta \log v_j} - 1) - i\theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} \Lambda(\mathrm{d}\mathbf{z}).$$

Then at time T_n , $(\Pi(T_n), \mathbf{V}(T_n)/V_1(T_n-))_{|[n]}$ is independent of the past and has distribution \mathcal{D}_n . Let us recall that \mathcal{D}_n is expressed in terms of Λ by

$$\mathcal{D}_n = \frac{1}{J_n} \Big(\sum_{i=1}^n c \delta_{\mathfrak{e}_i^n} + \int_{\mathscr{Z}^{\downarrow}} \varrho_{\mathbf{z}}^n \big(\cdot \cap \{ \pi \neq \mathbf{1}_n \text{ or } (\pi, \mathbf{v}) = (\mathbf{1}_n, \mathbf{0}) \} \Big) \Lambda(\mathrm{d}\mathbf{z}) \Big).$$

At this time, for each block *B* among the newly created blocks of $X(T_n)$, if $B \cap [n] \neq \emptyset$ and if *B* has positive mark, let $F_B \subset B$ be the subset consisting of exactly the first *n* integers that are part of block *B* (necessarily *B* contains infinitely many integers so the first *n* ones exist). Now we define

$$F_n(T_n) := F_n(T_n-) \cup \bigcup_B F_B,$$

where the union is taken over all newly created blocks of $X(T_n)$ with positive mark and nonempty intersection with $F_n(T_n-)$. After this first step, $X(T_n)_{|F_n(T_n)}$ consists of a random but finite (bounded by *n*) number of blocks, those with positive marks containing exactly *n* integers. The construction is recursive: if at time *t* the marked partition $X(t)_{|F_n(t)}$ contains *K* blocks of size *n*, then after an exponential time *T* with parameter KJ_n , one of them dislocates exactly as in the first step and *n* integers are selected for each newly created block in this dislocation. At the time of dislocation $F_n(t + T)$ is modified accordingly, and between time *t* and t + T, the branching property ensures us that each block has a mark behaving independently as e^{ξ_n} , where ξ_n is a Lévy process with characteristic exponent ψ_n . This recursion defines the process for all $t \ge 0$, and the construction is designed so that if $S_{\rho}^{(n)}$ is defined by (23), then

$$(S_{\theta}^{(n)}(t), t \ge 0) \stackrel{(d)}{=} \Big(\sum_{i} e^{\theta \xi_{n}^{i}(t)}, t \ge 0\Big),$$

where $(\xi_n^i(t), i \ge 1, t \ge 0)$ is a system of branching particles started from a unique particle at position 0, which can be described by:

- particles move independently as Lévy processes equal to ξ_n in distribution.
- a particle branches at rate J_n into a random set of K particles at positions $y + (y_1, \ldots, y_K)$, where y is the position of the mother particle at the time of branching and (y_1, \ldots, y_K) is a vector independent of the past and with distribution given by

$$\mathbb{E}f(y_1,\ldots,y_K)=\int_{\mathcal{M}_n}f(\log v_1,\ldots,\log v_K)\mathcal{D}_n(\mathrm{d} x),$$

where in the right-hand side integrand, the vector (v_1, \ldots, v_K) denotes the non-zero marks of *x*.

At this point we need the following lemma, which results from standard branching processes arguments. I could not find a reference which proves this result entirely in this form, so a short, straightforward proof is given below.

Lemma 20. We have

$$\mathbb{E}S_{\theta}^{(n)}(t) = e^{t\kappa^{(n)}(\theta)}, \quad with \quad \kappa^{(n)}(\theta) = A^{(n)}(\theta) + J_n B^{(n)}(\theta), \tag{24}$$

where $A^{(n)}$ corresponds to the movement of particles, with

$$A^{(n)}(\theta) = d\theta + \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^{\downarrow}} \sum_{\substack{i \ge 1\\v_i > 0}} s_i^n \left(v_i^{\theta} - 1\right) - \theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} \Lambda(\mathrm{d}\mathbf{z})$$
(25)

and $B^{(n)}$ corresponds to the branching, with

$$B^{(n)}(\theta) = \int_{\mathcal{M}_n} \left(S_{\theta}(x) - 1 \right) \mathcal{D}_n(\mathrm{d}x) = \frac{1}{J_n} \int_{\mathscr{Z}^{\downarrow}} \int_{\{\pi \neq \mathbf{1}_n \text{ or } \nu_1 = 0\}} \left(S_{\theta}(x) - 1 \right) \varrho_{\mathbf{z}}^n(\mathrm{d}x) \Lambda(\mathrm{d}\mathbf{z}).$$

Proof. It is standard in the theory of Lévy processes (see e.g. [20, Theorem 25.17]) that $A^{(n)}(\theta) < \infty$ if and only if $\mathbb{E}e^{\theta\xi_n(t)} < \infty$ for all $t \ge 0$, and in that case $\mathbb{E}e^{\theta\xi_n(t)} = e^{tA^{(n)}(\theta)}$. Now fix 0 < s < t and consider the event

 $A_s^t := \{$ the initial particle branches at time *s* and no other branching occurs before time *t* $\}$.

Then conditional on A_s^t , the branching construction yields

$$\mathbb{E}\left[\sum_{i} e^{\theta \xi_{n}(t)} \left| A_{s}^{t} \right] = \mathbb{E}\left[\int_{\mathcal{M}_{n}} \sum_{i} e^{\theta(\xi_{n}(s) + \log v_{i} + \widetilde{\xi}_{n}^{(i)}(t-s))} \mathcal{D}_{n}(dx) \left| A_{s}^{t} \right] \right]$$
$$= \left(\int_{\mathcal{M}_{n}} \sum_{i} v_{i}^{\theta} \mathcal{D}_{n}(dx)\right) \mathbb{E}\left[e^{\theta(\xi_{n}(s) + \widetilde{\xi}_{n}^{(1)}(t-s))}\right]$$
$$= \left(\int_{\mathcal{M}_{n}} S_{\theta}(x) \mathcal{D}_{n}(dx)\right) \mathbb{E}e^{\theta \xi_{n}(t)}$$
$$= \left(B^{(n)}(\theta) + 1\right) e^{tA^{(n)}(\theta)},$$

where ξ_n and the $\tilde{\xi}_n^{(i)}$, $i \ge 1$ are i.i.d. Lévy processes started from 0. This quantity does not depend on *s*, so one may write, if A^t is the event of a single branching occurring before time *t*,

$$\mathbb{E}\Big[\sum_{i} e^{\theta \xi_n^i(t)} \mid A^t\Big] = (B^{(n)}(\theta) + 1)e^{tA^{(n)}(\theta)}.$$

and in particular,

$$\mathbb{E}S_{\theta}^{(n)}(t) = \mathbb{E}\left[\sum_{i} e^{\theta \xi_{n}^{i}(t)}\right] \ge \mathbb{P}(A^{t}) \left(B^{(n)}(\theta) + 1\right) e^{tA^{(n)}(\theta)}$$

which shows that if $A^{(n)}(\theta) = \infty$ or $B^{(n)}(\theta) = \infty$, then $\mathbb{E}S^{(n)}_{\theta}(t) = \infty$. Now let us assume that both quantities are finite, and prove (24). First note that the argument above readily extends to

$$\mathbb{E}\Big[\sum_{i} e^{\theta \xi_{n}^{i}(t)} \mid A^{t,k}\Big] = (B^{(n)}(\theta) + 1)^{k} \mathbb{E} e^{\theta \xi_{n}(t)}.$$

where $A^{t,k}$ is the event of exactly k particles branching before time t. We now bound from above the probability of $A^{t,k}$. Let $t_0 := 0 < t_1 < t_2 < ...$ denote the branching times in our particle system. Note that as particles produce at most n offspring, the time between consecutive branching times $t_j - t_{j-1}$ is greater than an exponential random variable with parameter nJ_nj . Therefore we can compare the process counting branching times in our process and a Yule process with birth rate nJ_n , which yields

$$\mathbb{P}(A^{t,k}) \le \mathbb{P}(t_k < t) \le (1 - \mathrm{e}^{-tnJ_n})^k.$$

Now if t^* is small enough so that $(B^{(n)}(\theta) + 1)(1 - e^{-tnJ_n}) < 1$ for all $0 \le t \le t^*$, then we have

$$\forall 0 \le t \le t^*, \qquad \mathbb{E}S_{\theta}^{(n)}(t) = \sum_{k \ge 0} \mathbb{P}(A^{t,k}) \big(B^{(n)}(\theta) + 1 \big)^k \mathbb{E}e^{\theta \xi_n(t)} < \infty,$$

so the map $f : t \mapsto \mathbb{E}S_{\theta}^{(n)}(t)$ takes finite values before time t^* . Now note that for any times $t, s \ge 0$ the branching property applied at time t yields $\mathbb{E}[S_{\theta}(t + s) | S_{\theta}(t)] = S_{\theta}(t)f(s)$, and so taking expectations, f(t + s) = f(t)f(s). Since f takes finite value for $0 \le t \le t^*$, this shows that f(t) is finite for all $t \ge 0$. Now let us compute f(t) by applying the branching property at the first branching time t_1 :

$$\begin{split} f(t) &= \mathbb{P}(t_1 > t) \mathbb{E} e^{\theta \xi_n(t)} + \int_0^t \int_{\mathcal{M}_n} \sum_i \mathbb{E} e^{\theta(\xi_n(s) + \log v_i)} f(t-s) \mathcal{D}_n(\mathrm{d}x) \, \mathbb{P}(t_1 \in \mathrm{d}s) \\ &= e^{-J_n t} e^{tA^{(n)}(\theta)} + \int_0^t J_n e^{-J_n s} e^{sA^{(n)}(\theta)} \big(B^{(n)}(\theta) + 1 \big) f(t-s) \, \mathrm{d}s, \end{split}$$

and it is easily checked that the only solution of this equation is indeed

$$\mathbb{E}S_{\theta}^{(n)}(t) = f(t) = e^{t(A^{(n)}(\theta) + J_n B^{(n)}(\theta))} = e^{t\kappa^{(n)}(\theta)}$$

so (24) is proved.

Now note that for any $\mathbf{z} \in \mathscr{Z}^{\downarrow}$, one can write

$$\sum_{\substack{i\geq 1\\v_i>0}} s_i^n (v_i^\theta - 1) = \int_{\{\pi=\mathbf{1}_n \text{ and } v_1>0\}} \left(S_\theta(x) - 1\right) \varrho_{\mathbf{z}}^n(\mathrm{d}x),$$

so plugging this into the expression (25) for $A_{\theta}^{(n)}$ and putting everything together, we have

$$\kappa^{(n)}(\theta) = d\theta + \frac{\beta}{2}\theta^2 + \int_{\mathscr{Z}^{\downarrow}} \int_{\mathcal{M}_n} \left(S_{\theta}(x) - 1 \right) \varrho_{\mathbf{z}}^n(\mathrm{d}x) - \theta \log v_1 \mathbb{1}_{|\log v_1| \le 1} \Lambda(\mathrm{d}\mathbf{z}).$$

Now it is a consequence of the law of large numbers that for all $z \in \mathscr{Z}^{\downarrow}$, the following convergence holds (and is nondecreasing)

$$\int_{\mathcal{M}_n} \left(S_{\theta}(x) - 1 \right) \varrho_{\mathbf{z}}^n(\mathrm{d}x) \underset{n \to \infty}{\longrightarrow} \sum_{i \ge 1} v_i^{\theta} - 1,$$

and in the end, Lemma 20 and monotone convergence yield

$$\mathbb{E}S_{\theta}(X(t)) = \mathrm{e}^{t\kappa(\theta)}.$$

Now if $\kappa(\theta)$ is finite, it is a simple consequence of the Markov property of the process *X* that $(e^{-t\kappa(\theta)}S_{\theta}(X(t)), t \ge 0)$ is a martingale. Since it is nonnegative it converges almost surely as $t \to \infty$ so it is almost surely bounded by a random variable which we denote by $C = C_{\theta} > 0$. Now assume furthermore that $\kappa(\theta) < 0$ for some $\theta \neq 0$. Notice that almost surely for all $i \ge 1$ and $t \ge 0$,

$$V_i(t)^{\theta} \leq S_{\theta}(X(t)) \leq C \mathrm{e}^{t\kappa(\theta)},$$

so for any $\alpha \in \mathbb{R}$ such that $-\alpha/\theta > 0$,

$$V_i(t)^{-\alpha} \leq \left(C \mathrm{e}^{t\kappa(\theta)}\right)^{-\alpha/\theta},$$

and so almost surely,

$$\sup_{i\geq 1}\int_0^\infty V_i(t)^{-\alpha}\,\mathrm{d}t \le \frac{\theta C^{-\alpha/\theta}}{\alpha\,\kappa(\theta)} < \infty. \tag{26}$$

Now recall the stopping lines

$$\tau_i^{-\alpha}(t) = \left(\int_0^{\cdot} V_i(s)^{-\alpha} \, \mathrm{d}s\right)^{-1}(t), \qquad i \ge 1, \ t \ge 0,$$

which we used to change the self-similarity index. Proposition 11 tells us that the time-changed process $X \circ \tau^{-\alpha}$ is an α -ESSF process with characteristics (c, d, β , Λ), and note that for each $i \ge 1$, the integral

$$\zeta_i = \int_0^\infty V_i(s)^{-\alpha} \, \mathrm{d}s$$

is the hitting time of 0 by the pssMp $V_i \circ \tau_i^{-\alpha}$. Clearly (26) shows that $X \circ \tau^{-\alpha}$ reaches absorption before time $\frac{\theta C^{-\alpha/\theta}}{\alpha \kappa(\theta)}$.

It remains to show the finite total length property in the case $-\alpha/\theta \ge 1$. Recall that

$$S_{\theta}(X(t)) = \sum_{k \ge 1} \tilde{V}_k(t)^{\theta} \le C \mathrm{e}^{t\kappa(\theta)}.$$

It is elementary (because for any summable sequence u, $||u||_p \le ||u||_1$ for any $p \ge 1$) that for any α such that $-\alpha/\theta \ge 1$ this implies

$$S_{-\alpha}(X(t)) = \sum_{k\geq 1} \tilde{V}_k(t)^{-\alpha} \leq \left(C \mathrm{e}^{t\kappa(\theta)}\right)^{-\alpha/\theta}.$$

We claim the time change is such that

$$\int_0^\infty \# X \circ \tau^{-\alpha}(t) \, \mathrm{d}t = \int_0^\infty S_{-\alpha}(X(t)) \, \mathrm{d}t \le \frac{\theta C^{-\alpha/\theta}}{\alpha \kappa(\theta)} < \infty \quad \text{a.s}$$

To make this claim entirely justified, let us define for all $x \in \mathcal{M}_{\infty}^{\star}$ the (finite of infinite) set $I(x) = \{i_1, i_2, \ldots\}$ where i_k is the first integer contained in the *k*-th block with positive mark of *x*. Notice that by definition for any $i \ge 1$, for all $t \le \zeta_i$, $d\tau_i(t) = V_i^{\alpha}(\tau_i^{-\alpha}(t))dt$, therefore

$$\int_0^\infty \#X \circ \tau^{-\alpha}(t) dt = \int_0^\infty \sum_{i \in I(X \circ \tau^{-\alpha}(t))} 1 dt$$
$$= \sum_{i \ge 1} \int_0^\infty \mathbb{1}_{i \in I(X(\tau_i^{-\alpha}(t)))} \frac{V_i^\alpha(\tau_i^{-\alpha}(t)) dt}{V_i^\alpha(\tau_i^{-\alpha}(t))}$$
$$= \sum_{i \ge 1} \int_0^\infty \mathbb{1}_{i \in I(X(t))} V_i^{-\alpha}(t) dt$$
$$= \int_0^\infty S_{-\alpha}(X(t)) dt$$

and the proof is complete.

Acknowledgements. I thank the *Center for Interdisciplinary Research in Biology* (Collège de France) for funding, and my advisor Amaury Lambert for a careful reading of this manuscript and many helpful comments.

References

- D. Aldous and J. Pitman. The standard additive coalescent. *The Annals of Probability*, 26(4): 1703–1726, Oct. 1998. doi:10.1214/aop/1022855879.
- J. Berestycki. Ranked Fragmentations. ESAIM: Probability and Statistics, 6:157–175, 2002/ed. doi:10.1051/ps:2002009.
- [3] J. Bertoin. Self-similar fragmentations. Annales de l'Institut Henri Poincaré (B) Probability and Statistics, 38(3):319–340, 2002. doi:10.1016/S0246-0203(00)01073-6.
- [4] J. Bertoin. The asymptotic behavior of fragmentation processes. Journal of the European Mathematical Society, 5(4):395–416, Nov. 2003. doi:10.1007/s10097-003-0055-3.
- [5] J. Bertoin. Random Fragmentation and Coagulation Processes. Cambridge University Press, 2006. ISBN 978-0-521-86728-3. doi:10.1017/CBO9780511617768.
- [6] J. Bertoin. Markovian growth-fragmentation processes. *Bernoulli*, 23(2):1082–1101, May 2017. doi:10.3150/15-BEJ770.
- [7] J. Bertoin and B. Mallein. Infinitely ramified point measures and branching Lévy processes. *The Annals of Probability (to appear)*, 2019+. arXiv:1703.08078.
- [8] A. Blancas, J.-J. Duchamps, A. Lambert, and A. Siri-Jégousse. Trees within trees: Simple nested coalescents. *Electronic Journal of Probability*, 23(0), 2018. doi:10.1214/18-EJP219.
- [9] B. Chauvin. Product Martingales and Stopping Lines for Branching Brownian Motion. *The Annals of Probability*, 19(3):1195–1205, July 1991. doi:10.1214/aop/1176990340.
- [10] B. Dadoun. Asymptotics of self-similar growth-fragmentation processes. *Electronic Journal of Probability*, 22, 2017. doi:10.1214/17-EJP45.
- [11] D. J. Daley and D. Vere-Jones. An Introduction to the Theory of Point Processes, volume II of Probability and Its Applications. Springer New York, New York, NY, 2008. ISBN 978-0-387-21337-8 978-0-387-49835-5. doi:10.1007/978-0-387-49835-5.
- [12] J.-J. Duchamps. Trees within trees II: Nested fragmentations. Ann. Inst. H. Poincaré Probab. Statist. (to appear), 2019+. arXiv:1807.05951.
- F. G. Ged. Profile of a self-similar growth-fragmentation. *Electronic Journal of Probability*, 24, 2019. doi:10.1214/18-EJP253.
- [14] B. Haas and G. Miermont. The Genealogy of Self-similar Fragmentations with Negative Index as a Continuum Random Tree. *Electronic Journal of Probability*, 9:57–97, 2004. doi:10.1214/EJP.v9-187.
- [15] B. Haas, G. Miermont, J. Pitman, and M. Winkel. Continuum tree asymptotics of discrete fragmentations and applications to phylogenetic models. *The Annals of Probability*, 36(5): 1790–1837, Sept. 2008. doi:10.1214/07-AOP377.
- [16] J. Kingman. The coalescent. Stochastic processes and their applications, 13(3):235–248, 1982. doi:10.1016/0304-4149(82)90011-4.

- [17] N. Krell. Self-Similar Branching Markov Chains. In C. Donati-Martin, M. Émery, A. Rouault, and C. Stricker, editors, *Séminaire de Probabilités XLII*, volume 1979, pages 261–280. Springer Berlin Heidelberg, 2009. doi:10.1007/978-3-642-01763-6 10.
- [18] J. Lamperti. Semi-stable Markov processes. I. Zeitschrift f
 ür Wahrscheinlichkeitstheorie und Verwandte Gebiete, 22(3):205–225, 1972. doi:10.1007/BF00536091.
- [19] J. C. Pardo and V. Rivero. Self-similar Markov processes. Boletín de la Sociedad Matemática Mexicana: Tercera Serie, 19(2):201-235, 2013. url:https://www.cimat.mx/~jcpardo/ survey.pdf.
- [20] K. Sato. Lévy Processes and Infinitely Divisible Distributions. Number 68 in Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, U.K.; New York, 1999. ISBN 978-0-521-55302-5.
- [21] Q. Shi. Growth-fragmentation processes and bifurcators. *Electronic Journal of Probability*, 22, 2017. doi:10.1214/17-EJP26.
- [22] S. M. Srivastava. A Course on Borel Sets, volume 180 of Graduate Texts in Mathematics. Springer Berlin Heidelberg, 1998. doi:10.1007/978-3-642-85473-6.