



**HAL**  
open science

## Transplanting gravid lizards to high elevation alters maternal and embryonic oxygen physiology, but not reproductive success or hatchling phenotype

Laura Kouyoumdjian, Eric J. Gangloff, Jérémie Souchet, Gerardo A Cordero, Andréaz Dupoué, Fabien Aubret

### ► To cite this version:

Laura Kouyoumdjian, Eric J. Gangloff, Jérémie Souchet, Gerardo A Cordero, Andréaz Dupoué, et al.. Transplanting gravid lizards to high elevation alters maternal and embryonic oxygen physiology, but not reproductive success or hatchling phenotype. *Journal of Experimental Biology*, 2019, 222 (14), pp.jeb.206839. 10.1242/jeb.206839 . hal-02186233

**HAL Id: hal-02186233**

**<https://hal.sorbonne-universite.fr/hal-02186233>**

Submitted on 3 Aug 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## RESEARCH ARTICLE

# Transplanting gravid lizards to high elevation alters maternal and embryonic oxygen physiology, but not reproductive success or hatchling phenotype

Laura Kouyoumdjian<sup>1,\*</sup>, Eric J. Gangloff<sup>1,\*‡</sup>, Jérémie Souchet<sup>1</sup>, Gerardo A. Cordero<sup>2</sup>, Andréaz Dupoué<sup>1,3</sup> and Fabien Aubret<sup>1,4</sup>

## ABSTRACT


Increased global temperatures have opened previously inhospitable habitats, such as at higher elevations. However, the reduction of oxygen partial pressure with increase in elevation represents an important physiological constraint that may limit colonization of such habitats, even if the thermal niche is appropriate. To test the mechanisms underlying the response to ecologically relevant levels of hypoxia, we performed a translocation experiment with the common wall lizard (*Podarcis muralis*), a widespread European lizard amenable to establishing populations outside its natural range. We investigated the impacts of hypoxia on the oxygen physiology and reproductive output of gravid common wall lizards and the subsequent development and morphology of their offspring. Lowland females transplanted to high elevations increased their haematocrit and haemoglobin concentration within days and maintained routine metabolism compared with lizards kept at native elevations. However, transplanted lizards suffered from increased reactive oxygen metabolite production near the oviposition date, suggesting a cost of reproduction at high elevation. Transplanted females and females native to different elevations did not differ in reproductive output (clutch size, egg mass, relative clutch mass or embryonic stage at oviposition) or in post-oviposition body condition. Developing embryos reduced heart rates and prolonged incubation times at high elevations within the native range and at extreme high elevations beyond the current range, but this reduced oxygen availability did not affect metabolic rate, hatching success or hatchling size. These results suggest that this opportunistic colonizer is capable of successfully responding to novel environmental constraints in these important life-history stages.

**KEY WORDS:** Climate change, Development, Embryo, Hypoxia, Metabolic rate, Physiological plasticity, *Podarcis muralis*, Reactive oxygen metabolites, Reproductive output

<sup>1</sup>Station d'Ecologie Théorique et Expérimentale du CNRS-UMR 5321, Moulis, France. <sup>2</sup>Fachbereich Geowissenschaften, Eberhard Karls Universität Tübingen, Tübingen, Germany. <sup>3</sup>EES Paris, Sorbonne Université, CNRS, UMR 7618, Tours 44-45, 4 place Jussieu, 75005 Paris, France. <sup>4</sup>School of Molecular and Life Sciences, Curtin University, 6102 WA, Australia.

\*These authors contributed equally to this work

‡Author for correspondence (eric.gangloff@sete.cnrs.fr)

 L.K., 0000-0003-3201-7862; E.J.G., 0000-0002-9074-8009; J.S., 0000-0001-9296-1332; G.A.C., 0000-0002-9137-1741; A.D., 0000-0002-2501-464X; F.A., 0000-0002-7783-8659

## INTRODUCTION

In mountainous regions, the impacts of climate change are particularly pronounced owing to shifts in the altitudinal climate envelope (Chen et al., 2011; Dirnböck et al., 2011; Nogués-Bravo et al., 2008; Walther et al., 2002). Such shifts may invoke biological changes through the processes of adaptation, plasticity and migration; or may result in extirpation (Parmesan, 2006; Sinervo et al., 2010). Just as low-lying valleys provided refuge for a number of organisms during periods of glaciation, it is possible that high-elevation areas will play a similar role in times of global warming (Hewitt, 1999; Tzedakis, 2004). Indeed, the migration of some populations to higher elevations is currently observed in mountainous landscapes (Bani et al., 2019; Bassler et al., 2013; Chen et al., 2011; Freeman et al., 2018; Hickling et al., 2006; Pottier, 2012; Pauchard et al., 2015) and the preservation of such habitats has been proposed specifically as a refuge for lizards (Sinervo et al., 2018). As a result, it is important to identify factors that both permit and limit colonization of higher-elevation habitats. Terrestrial ectothermic vertebrates, such as non-avian reptiles, are sensitive to variation in environmental temperatures and exhibit substantial metabolic plasticity (Huey, 1982). As such, they provide excellent models to understand the impacts of climate change across geographic scales (Diele-Viegas and Rocha, 2018; Huey et al., 2018; Le Galliard et al., 2012; Walther et al., 2002). However, their flexible physiologies and complex interactions among pathways when responding to many environmental factors make broad predictions difficult as the effects of changing habitats may vary dramatically across taxa (Levy et al., 2017; Moore et al., 2018; Pontes-da-Silva et al., 2018; Sinervo et al., 2010; Weatherhead et al., 2012).

One potential constraint to upslope migration is the reduced oxygen partial pressure at high elevations, which causes high-altitude hypoxia (Bouverot, 1985; Powell and Hopkins, 2010). This phenomenon imposes important physiological challenges on a variety of organisms (reviewed in Storz et al., 2010). Importantly, oxygen availability interacts with temperature in a context-dependent manner to influence thermal performance curves and thermal limits in terrestrial ectotherms (Gangloff and Telemeco, 2018; Jackson, 2007). Acute and chronic effects of hypoxia have been fairly well studied in some mammals (humans in particular), birds and some reptiles (Monge and Leon-Velarde, 1991; Weathers and McGrath, 1972), and the responses can be diverse (Storz et al., 2010). Acute effects usually include hyperventilation and tachycardia, as for example in humans brought to high altitude (Saito et al., 1988). Chronic effects may involve changes to the cardiorespiratory systems, such as increased size of the lungs, changes to cardiac morphology, and increased blood pressure (Cunningham et al., 1974; He et al., 2013; Hillyard, 1980; Powell and Hopkins, 2010). In lizards specifically, chronic hypoxia may also result in hematological and muscle composition changes

(González-Morales et al., 2015; Lu et al., 2015; Weathers and McGrath, 1972). Such physiological shifts in oxygen-carrying capacity, however, may be insufficient to maintain performance traits or compensate for reduced oxygen in the long term. Hypoxia exposure during early development, such as embryos *in ovo*, can also dramatically affect physiology and development. Most common responses include mechanisms to facilitate oxygen diffusion and transport, such as increased vascularization of chorioallontoric membranes, increased haematocrit and cardiac hypertrophy (Corona and Warburton, 2000; Crossley and Altimiras, 2005; Jochmans-Lemoine and Joseph, 2018; Kam, 1993; Nechaeva, 2011; Warburton et al., 1995). Under certain conditions, oxygen restrictions can also result in depressed metabolism, decreased growth and reduced survivorship to hatching (Iungman and Piña, 2013; Warburton et al., 1995). These early developmental effects can persist into later life phenotypes (Sun et al., 2014; Wearing et al., 2017).

Previous work demonstrates the impacts of hypoxia on adult male and embryo common wall lizards (*Podarcis muralis*) transplanted to extreme high elevation beyond the current range. Transplanted adult males increased haematocrit and haemoglobin within 3 weeks, but were unable to sustain this response for longer. Despite these changes in oxygen-carrying parameters, lizards transplanted to high elevation suffered reduced running performance and a reduction in body condition compared to lizards kept at lowland native elevation (Gangloff et al., 2019). Embryos of this species developing in high-altitude hypoxia, however, demonstrate a potential to sustain growth and development at least until hatching. High-altitude hypoxia affected embryo physiology, resulting in reduced metabolism, cardiac hypertrophy, and hyperventilation. These physiological and morphological adjustments seemed to buffer embryos from negative impacts of hypoxia, resulting in similar hatching success and hatchling body size compared with embryos developing at native lowland elevation (Cordero et al., 2017a). While studies such as these provide important data on the effects of high-altitude hypoxia across life stages, relatively little is known about the effects of high-altitude hypoxia exposure on maternal energetics and reproductive allocation in natural populations. If reduced oxygen availability imposes a physiological constraint, we might expect that gestating females will be less able to provide adequate resources or environments for developing embryos. This is particularly important in the context of responses to climate change, given the potential for transgenerational plasticity (e.g. maternal effects) to buffer developing offspring from the impacts of novel environmental conditions (Sinervo et al., 2018; Warner, 2014). These effects have the potential to promote offspring success in future environments, such as when maternal exposure to warmer temperatures increases offspring survival in future warm environments (Shama et al., 2014; Sun et al., 2018).

With this study, we quantify physiology and reproduction at different levels of oxygen availability on native and transplanted common wall lizards (*P. muralis*). Such data are needed to assess the potential importance of the effects of high-altitude hypoxia on reproduction and development, which, in turn, will determine the colonization potential of this species with a wide and expanding range (Pottier, 2012; Speybroeck et al., 2016). Our goals are to quantify how reproducing females physiologically respond to changes in oxygen availability; how these responses can influence reproductive output and capacity; how reduced oxygen availability affects embryo development; and how such responses might differ between lizards native to different elevations. We first measured aspects of maternal physiology and reproduction from replicate low- and high-elevation populations at native elevations and after

translocation to high elevations. Then, we measured embryo development and physiology at native, high (within the natural range) and extreme high (beyond current range limit) elevations, to test five primary hypotheses.

#### **Energetic limitations imposed by hypoxia will force a trade-off resulting in reduced relative reproductive investment**

Because reduced oxygen availability affects the shape of performance curves (Gangloff and Telemeco, 2018), we expected that short-term (i.e. weeks) exposure to hypoxic conditions might reduce efficiency of physiological functions involved in energy processing for both gravid females and developing embryos. If lizards are unable to compensate, we expected that a maternal allocation trade-off would reduce egg mass of lizards transplanted to high elevation, resulting in a smaller relative clutch mass (RCM), a metric of reproductive investment. Because clutch size is likely determined before our experimental manipulations and females are unlikely to resorb eggs (Blackburn, 1998), we did not expect to see an effect of transplant on clutch size. If female lizards at high elevation are able to maintain the same level of reproductive investment (egg size and RCM), we expected to observe prolonged egg retention and thus laying of eggs at a later developmental stage, as a consequence of reduced oxygen availability on gravid mothers (see Mathies and Andrews, 1995). Alternatively, if oxygen limitation imposes a constraint on reproduction, mothers may sustain investment into offspring at a physiological cost to themselves, for instance resulting in reduced post-oviposition body condition.

#### **Hypoxia and reproductive stage will interact to affect reactive oxygen metabolite production**

Reactive oxygen metabolites (ROMs) are produced as by-products of oxidative metabolism and can damage a variety of subcellular molecules. ROMs have been implicated in senescence and as a by-product of reproduction in reptiles (Costantini, 2016; Dowling and Simmons, 2009; Robert et al., 2007; Stahlschmidt et al., 2013; Stier et al., 2017; Webb et al., 2018). ROM production generally increases as a result of cellular hypoxia (reviewed in Guzy and Schumacker, 2006; Harrison et al., 2015; Solaini et al., 2010). Exploring the interaction of reproduction and hypoxia on ROM production is essential to both understand short-term physiological changes and potential long-term consequences of reproduction at altitude. We expected that the reactive oxygen metabolite profile of lizards would shift across reproduction and that transplanted lizards would suffer from relatively increased levels compared with lizards kept at native elevation.

#### **Developing embryos will exhibit changes in physiology in response to hypoxia to meet metabolic demands important to development and differentiation**

Reptilian embryos *in ovo* often experience naturally occurring hypoxia due to flooding events or limited gas exchange in subterranean nests (reviewed in Packard and Packard, 1987; Ackerman and Lott, 2004; Booth, 1998). Given the resilience to these conditions exhibited by a variety of taxa, we predicted that embryos would adjust physiology and development, such as through changes in heart rate and incubation time. With these responses, embryos can sustain differentiation, growth, and organogenesis, critical processes that must be fuelled by aerobic metabolism and are likely buffered from moderate hypoxia (Crossley et al., 2017; Iungman, and Piña, 2013; Warburton et al., 1995). Physiological or morphological responses will facilitate the maintenance of oxygen consumption ( $\dot{V}_{O_2}$ , an integrated index of

general metabolism), although late-period increases in demand might present challenges at high elevation. Furthermore, we predicted that these adjustments would allow embryos to maintain important fitness-related traits such as size and mass at birth; if energetic constraints are present this would be demonstrated by reduced body size of embryos developing at extreme high elevation.

### Effects of hypoxia on post-oviposition development will be reduced in embryos that are exposed to hypoxia during gestation

Exposure to hypoxia earlier in development will provide both more time for embryos to respond and a potential window for anticipatory maternal effects. We expected that the change in heart rate and incubation time predicted above would be less pronounced in embryos from females that were transplanted to high altitudes compared with embryos translocated after oviposition. Furthermore, maternal effects in response to hypoxia may provide a mechanism to prime embryos for subsequent development in low-oxygen environments, thus reducing the potentially negative impacts on development in embryos transplanted to high or extreme high elevation.

### Lizards from high-elevation populations will demonstrate local adaptation in traits related to oxygen capacity

We predicted that the responses to hypoxia of transplanted females would differ from that of lizards native to high elevation, while embryos from high-elevation populations would show reduced effects on development at extreme high elevations compared to embryos from low-elevation populations. As has been found in previous studies of transplanted lizards (He et al., 2013; Gangloff et al., 2019; Weathers and McGrath, 1972) we predicted physiological plasticity in parameters related to oxygen-carrying capacity in blood chemistry (haematocrit and haemoglobin concentration) in reproductive females from low elevations transplanted to high elevations. Importantly, we did not expect the physiological profile of lowland lizards transplanted to high elevation to match those of lizards from high-elevation populations, as often the short-term plastic response differs from that of locally adapted populations (He et al., 2013; Jochmans-Lemoine et al., 2015; Reyes et al., 2018; Storz et al., 2010; Velotta et al., 2018). We predicted that such shifts in oxygen capacity would allow lizards to maintain resting metabolic rates but perhaps bear other physiological or reproductive consequences given the increased energetic demands of reproduction (Angilletta and Sears, 2000; Foucart et al., 2014). Furthermore, we predicted that embryos from high-elevation populations would respond less dramatically to development at extreme high elevations compared with lizard embryos native to low elevations.

### Research objectives

Taken together, these observations allow us to evaluate how hypoxia may constrain physiology in reproducing and developing lizards and how these responses may differ between populations native to different elevations. Given the dependence of successful colonization on recruitment (Aubret, 2013; Warner, 2014; While et al., 2015), testing these effects in reproductive females and their developing offspring is critical to predict the dynamics of colonization in common wall lizards in mountainous environments affected by climate change.

## MATERIALS AND METHODS


### Source populations and husbandry

The common wall lizard [Lacertidae: *Podarcis muralis* (Laurenti 1768)] is widespread across southern Europe in a variety of habitats

(Arnold et al., 1988; Speybroeck et al., 2016). Its geographic distribution is restricted by the thermal environment because embryos are incapable of completing development at cold temperatures (Strijbosch et al., 1980; Van Damme et al., 1991; While et al., 2015). However, wall lizards have recently been observed at higher areas of mountainous regions in the south of France, extending up to 2600 m above sea level (ASL) (Pottier, 2012). At low elevations, females produce two or three clutches per season with an average of 6 eggs per clutch (Le Hénanff, 2011). We monitored populations in the Pyrénées Ariégeoises (France) from the beginning of the active season and sampled adult female lizards during their first reproductive event of the season in April and May 2018 at low (382–472 m ASL) and high (1402–1795 m ASL) elevations. A total of 47 gravid females were sampled from three low-elevation and two high-elevation populations (see Table S1 for sampling details). Individuals were caught by using the lasso method (Blomberg and Shine, 1996; Fitzgerald, 2012) and marked using a cautery pen (Vervust and Van Damme, 2009). On the day of capture, we measured mass to the nearest 0.01 g with a precision balance (mean±s.d.: 5.73±1.06 g) and snout–vent length (SVL) with a digital caliper to the nearest 0.01 mm (mean±s.d.: 61.97±4.08 mm). Neither  $M_b$  nor SVL at capture differed between lizards from low- and high-elevation populations (Wilcoxon rank-sum test,  $M_b$ :  $W=206$ ,  $P=0.28$ ; SVL:  $W=176$ ,  $P=0.082$ ). On the day of capture, we transported animals to facilities either at the Station d'Ecologie Théorique et Expérimentale du CNRS à Moulis (low-elevation treatment: 42°57'26.8"N, 1°05'08.3"E; 436 m ASL;  $P_{O_2}$  ~20.1 kPa) or in the commune of La Mongie (high-elevation treatment: 42°54'34.8"N, 0°10'53.6"E; 1735 m ASL;  $P_{O_2}$  ~17.4 kPa; see Fig. 1 for experimental design). We maintained females individually in identical plastic enclosures (38×26×23 cm) containing a thin layer of substrate, a water bowl, a plastic shelter/basking platform (15×5×3.5 cm) and a nest box (18×14×10 cm) filled with wet sand. Enclosures were misted with water 3–4 times a week. Every 2 days, five live mealworms (*Tenebrio* sp. larvae) were distributed to each female and standing water was supplied *ad libitum*. Light was provided by UV lamps for 12 h per day and the enclosures were warmed by heat lamps (42 W) for 5 h per day at 1 h intervals, providing a gradient of ~25–40°C. Additionally, we monitored temperatures in the enclosures with iButton thermal data loggers (Model DS1921G-F5, Maxim Integrated, San Jose, CA, USA) placed in the cool end of two of the enclosures in each treatment. Temperature regimes were nearly identical across labs, with a diel range from 15°C at night to 25°C during the day. Furthermore, we rotated the enclosures on their shelves every several days to ensure no position effects.

### Maternal metabolic rate

We measured resting metabolic rate after 2 days in captivity and approximately every week thereafter until 1 week after egg-laying (measures were skipped when lizards were actively nesting; mean time between measures±s.d.: 8.5±2.9 days) Lizards were offered their first meal after their initial measurement (2 days after capture) and food was withheld for 2 days before each subsequent measure. Each individual was placed in a custom-made 250 ml opaque metabolic chamber, which was then placed in an incubator set at 32°C for 1 h before measurement, a temperature intermediate between thermal preference (32.6°C) and field body temperatures (30.6°C) for gravid females in this area (Trochet et al., 2018; F.A., unpublished data). This acclimation time allows body temperature to equilibrate and reduces effects of handling stress (Braña, 1993; Tosini and Avery, 1996). We used pull-mode respirometry (Foxbox-C Field  $O_2$  and  $CO_2$  Analysis System, Sable Systems, Inc., Las Vegas, NV, USA) to measure gas



**Fig. 1. Experimental design.** Gravid lizards were collected from three low-elevation (382–472 m ASL) and two high elevation (1402–1795 m ASL) populations in the Pyrénées of southern France (see Table S1 for details). We maintained all lizards in common-garden conditions in labs at low elevation (436 m ASL;  $PO_2 \sim 20.1$  kPa) or high elevation (1735 m ASL;  $PO_2 \sim 17.4$  kPa), where they completed gestation and laid eggs. Egg clutches were then split and incubated at low, high or extreme high elevation (2877 m ASL;  $PO_2 \sim 15.3$  kPa). Illustration credit: Bea Angelica Andersson.

exchange [oxygen consumption ( $\dot{V}_{O_2}$ ) and carbon dioxide production ( $\dot{V}_{CO_2}$ ), corrected for barometric pressure] continuously for 30 min. Air was pumped at a rate of  $500 \text{ ml min}^{-1}$  through the metabolic chamber, dried of water vapour with Drierite, then measured for both  $CO_2$  and  $O_2$  content. Data were analysed with ExpeData software (v.1.7.30, Sable Systems, Inc.) to calculate  $\dot{V}_{CO_2}$  and  $\dot{V}_{O_2}$ . We extracted the average of the lowest values of gas exchange over a 10 min interval for our analysis of metabolic rate, which allows the elimination of elevated metabolic rates during the measurement period (presumably because of activity). We recorded the body temperature of each lizard immediately following measurement using a cloacal thermometer (Miller & Weber T-6000, Ridgewood, NY, USA). All metabolic rate measures were conducted during the normal activity period for these diurnal lizards (07:30–20:00 h), and so represent resting metabolic rate (Andrews and Pough, 1985).

#### Blood sampling and hematological measures

The day after metabolic rate measurements, blood samples were collected by placing a heparinized glass capillary in the retro-orbital sinus of lizards (MacLean et al., 1973; Meylan et al., 2003). We collected 15–25  $\mu\text{l}$  of blood in less than 3 min (bleed time mean  $\pm$  s.d.:  $1.66 \pm 0.62$  min), which was stored on ice until processing. Haematocrit values represent the volume of packed red blood cells relative to the total volume as measured with a caliper after spinning 10–15  $\mu\text{l}$  of whole blood for 5 min at 5000  $g$ . Haemoglobin concentration was measured with the colorimetric cyanmethemoglobin method using 5  $\mu\text{l}$  whole blood, following the manufacturer's instructions (Drabkin's reagent, Sigma-Aldrich, St Louis, MO, USA; cat. no. D5941). Samples were run in duplicate on five plates, with a pooled sample run repeatedly on each plate to provide estimates of intra- and interplate variation of 6.5% and 5.7%,

respectively. Additionally, we quantified the amount of hyperoxides in blood plasma with a reactive oxygen metabolite (d-ROM) test kit (MC001, Diacron International, Italy) as a proxy for overall plasma oxidative status (Costantini, 2016). After centrifugation of whole blood at 3000  $g$  for 5 min, we removed plasma by pipette and stored plasma at  $-80^\circ\text{C}$  until the time of assay. We followed the manufacturer's instructions modified for use with a 4  $\mu\text{l}$  lizard plasma sample (A.D., unpublished data). We calculated the amount of ROMs corrected for control blanks and report values in equivalents of  $\text{mg H}_2\text{O}_2 \text{ dl}^{-1}$ . Samples were run on three plates, with a pooled sample run repeatedly on each plate to provide estimates of intra- and interplate variation of 7.0% and 8.6%, respectively. In some cases, plasma volume limitations and other logistical constraints precluded all measurements on each sample, so the number of observations varies slightly for each measure (see Table 1).

#### Reproductive measures

The nest boxes were checked every 24–48 h and eggs were immediately removed upon discovery. Eggs were individually weighed with a digital scale and the female was immediately re-weighed to obtain a post-oviposition mass. We calculated the relative clutch mass (RCM) as the ratio of total clutch mass to female post-oviposition mass. One egg from each clutch was randomly selected for dissection and staging. This egg was kept in moist vermiculite at room temperature for 1–4 days before we removed approximately one-third of the yolk with a syringe needle to facilitate embryo isolation and fixed the remaining embryo and tissue in 10% buffered formalin. Fixed eggs were sequentially transferred to 40 and 70% ethanol and then washed in a Tween 20/phosphate-buffered saline solution (PBS-T), following the embryo dissection protocol of Cordero and Janzen (2014). Incisions were

**Table 1. Results of linear mixed model analyses of physiological measures in gravid female *Podarcis muralis* lizards**

Source of variation	Hct (N=186)	[Hb] (N=187)	Resting $\dot{V}_{O_2}$ (N=175)	ROMs (N=155)
Treatment	Low-high>high-high>low-low	Low-high>high-high>low-low		High-high>low-high>low-low
	<i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>			
Relative oviposition date linear	Negative <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>	Negative <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>	1.37 (2, 9.68) 0.30	3.49 (2, 23.8) 0.047*
Relative oviposition date quadratic	See Fig. 2A <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>	–	–	–
Treatment×relative oviposition date linear	–	–	–	See Fig. 2D 4.48 (2, 144.7) 0.013*
log <sub>10</sub> (Mass)	–	–	Positive <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>	–
Pre-oviposition	Pre>post <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>	–	Post>pre <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>	Post>pre <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>
Body temperature	–	–	Positive <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>	–
Size-corrected clutch mass	–	–	–	Negative <i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) Pr> <i>F</i>

See Materials and Methods for statistical details. Significant factors are indicated by \* $P < 0.05$  and \*\* $P < 0.001$  along with directionality. *N* indicates the number of observations; d.f.<sub>n</sub> is the numerator degrees of freedom; d.f.<sub>d</sub> is the denominator degrees of freedom. Hct, haematocrit; [Hb], haemoglobin concentration;  $\dot{V}_{O_2}$ , oxygen consumption rate; ROMs, reactive oxygen metabolite concentration.

made on extreme longitudinal poles of the egg and egg contents were gently removed and placed in PBS-T. The embryo was then located, transferred to 70% ethanol, and placed on a dissecting microscope for observation and staging, following the criteria of Dufaure and Hubert (1961). Since stage at oviposition is generally invariable within a clutch (Mathies and Andrews, 1995), we can infer the stage of the randomly selected embryo to be representative of all eggs in the clutch.

We recognize that an ideal design would have included more frequent checks of nest boxes and an immediate fixing of the randomly selected embryo. However, this was impossible due to logistical constraints. We combated bias by checking nest-boxes in both treatments at the same intervals and through an analysis of the potential effect of the number of possible days the eggs were incubating in the nest boxes before being discovered. The number of potential days in the nest box did not affect mass of eggs at oviposition ( $F_{1,46,1} = 2.99$ ,  $P = 0.09$ ) nor did the number of days before fixation affect embryo stage at oviposition ( $F_{1,37} = 0.57$ ,  $P = 0.46$ ).

### Egg incubation and treatments

The remaining eggs of each clutch ( $N = 209$ ) were individually incubated in plastic cups containing moist vermiculite (1:5 water to vermiculite by volume; Cordero et al., 2017a; While et al., 2015) and sealed with plastic film to retain moisture. Incubators contained an open tray of water to maintain high humidity. We verified the total cup mass weekly and added water to maintain this ratio throughout incubation. Eggs were incubated at a constant 28°C, which provides optimal hatching success in this species (Van Damme et al., 1991). Within the first week after oviposition (mean ± s.d.: 1.8 ± 2.2 days), eggs were dispatched to the three different incubation treatments (low elevation, high elevation and extreme high elevation). Within each maternal treatment, mass of eggs allocated to different incubation treatments did not differ (all  $P > 0.15$ ). Low- and high-elevation treatments matched the locations

of the maternal treatments (see above) while the extreme high-elevation treatment was conducted at the laboratory of Pic du Midi de Bigorre (42°56'11.0"N, 0°08'32.9"E; 2877 m ASL;  $P_{O_2}$ : ~15.3 kPa; see Fig. 1 for complete experimental design and sample sizes). We rotated egg position in incubators twice a week to ensure no position effects.

### Embryo heart and metabolic rates

Beginning 1 week post-oviposition, we measured the heart rate of all embryos repeatedly throughout incubation (up to four times per embryo, mean ± s.d. interval between measures: 12.8 ± 4.2 days), spacing our measurements to ensure an equal sampling across the incubation period and amongst treatment groups. We measured the heart rate with a Buddy digital egg monitor (MK2; Avitronics, Truro, UK) placed inside the incubator at 28°C. Eggs were quickly moved from their incubation cup to the instrument and we used a stopwatch to measure the time from egg removal to when a consistent heart rate was observed (mean ± s.d. time: 121.6 ± 73.7 s). If a reliable measurement could not be made in under 5 min, we replaced the egg in the incubator and re-measured later in the day to reduce any effects of handling or temperature change (Cordero et al., 2017a; Hulbert et al., 2017).

In a subset of eggs, we also measured metabolic rate on the same day as we measured heart rate. Each egg was placed in a small cup with moist vermiculite and inside a 250 ml metabolic chamber. We replaced the chamber in the incubator at 28°C and used closed-system respirometry to measure gas exchange. We flushed the chamber for 15 min at a flow rate of 300 ml min<sup>-1</sup> then closed valves to seal the chamber for 90 min. We then opened the valves to re-establish air flow, dried air of water vapour with Drierite, and measured O<sub>2</sub> and CO<sub>2</sub> as above. We used ExpeData software to calculate  $\dot{V}_{O_2}$  and  $\dot{V}_{CO_2}$  by integrating the change in instantaneous gas concentrations over the period the chamber was sealed (Lighton, 2008).

### Hatchling morphology

Eggs were checked for hatching every 48 h. After emergence from the egg, we measured SVL using digital calipers (mean $\pm$ s.d.: 25.1 $\pm$ 1.2 mm) and mass using an electronic balance (mean $\pm$ s.d.: 0.363 $\pm$ 0.053 g). Of the 209 eggs incubated, 184 hatched (88.0% hatching success rate).

Field sampling and experimental protocols were conducted under permit provided by the Direction régionale de l'environnement, de l'aménagement et du logement (DREAL) Midi-Pyrénées (Arrêté Préfectoral no. 2017-s-02 30 March 2017), under current ethical committee approval (APAFIS#16359-201808011445465 v4), and in accordance with Directive 2010/63/EU on protection of animals used for scientific purposes.

### Statistics

For our measures of the reproducing females, we assessed the influence of native elevation and gestation elevation treatments, time relative to oviposition and reproductive investment on five physiological parameters (haematocrit, haemoglobin concentration,  $\dot{V}_{O_2}$ ,  $\dot{V}_{CO_2}$  and ROMs) using linear mixed models with the lme4 package (Douglas et al., 2015) in the programming language R (<https://www.r-project.org/>). We verified normal distributions of residuals in all cases and assessed the relative importance of fixed effects using type III sums of squares, correcting denominator degrees of freedom for *F*-tests (Kenward and Roger, 1997). Models included the fixed effect of experimental treatment group, which combined origin elevation and gestation elevation (low–low, low–high or high–high), relative oviposition date (both linear and quadratic to account for non-linear changes over this time period), and the interaction between treatment group and relative oviposition date (both linear and quadratic). We also included a categorical fixed effect indicating whether the measure occurred before or after oviposition (Y/N). To test for the effect of reproductive investment, we included the covariate of size-corrected clutch mass (calculated as the residuals of the total clutch mass on SVL regression). Initial models also included the covariate of body size (SVL for haematocrit, haemoglobin concentration and ROMs;  $\log_{10}$ -transformed mass for  $\dot{V}_{O_2}$  and  $\dot{V}_{CO_2}$ ). We accounted for non-independence of samples from each population and the repeated measures on individuals by including population of origin and individual as additive random effects. To simplify models, we used a backward selection procedure, removing non-significant interactions and main effects (all  $P > 0.10$ ) except for the factor of treatment, which addresses our main biological hypotheses. To meet assumptions of normal distribution of residuals,  $\dot{V}_{O_2}$ ,  $\dot{V}_{CO_2}$  and ROMs data were  $\log_{10}$ -transformed before analysis.

For the reproductive parameters (stage at oviposition, clutch size, egg mass, relative clutch mass, post-oviposition body condition), we used linear mixed models with a fixed effect of treatment (as above) and size (SVL) as a covariate. We included the random effects of population of origin for all models and also maternal identity in the model of egg size to account for covariance among eggs in the same clutch. Body condition was calculated as the residual of the linear regression of  $\log_{10}$ -transformed mass on  $\log_{10}$ -transformed SVL. We reduced models following a backward-selection process as above. Because of the limited range of embryonic stages observed at oviposition, we also assessed the effect of treatment on stage at oviposition with an ordinal logistic regression using the polr function of the MASS package (Venables and Ripley, 2002), which gave results qualitatively identical to the linear mixed model analysis.

We analysed aspects of embryo physiology (heart rate and oxygen consumption throughout incubation), development


(incubation time and survival) and hatchling morphology (mass and SVL) using the same linear model structure and backward selection approach as described above. The analysis of survival utilized a generalized linear mixed model with a binomial error distribution. We categorized embryos into one of seven treatment groups based on the combination of population of origin, maternal gestation treatment, and egg incubation treatment (low–low–low, low–low–high, low–low–extreme high, low–high–high, low–high–extreme high, high–high–high or high–high–extreme high). All models included the fixed effect of treatment group. Initial models also contained the covariate of egg mass at the time of measurement (for heart rate and oxygen consumption) or egg mass at oviposition (for survival, hatchling mass and hatchling size) to account for variation in maternal energetic investment (Uller and While, 2015; Van Damme et al., 1991; Warner and Lovern, 2014), as well as the interaction of egg mass with treatment group. We also included the covariate of stage at oviposition in initial models of incubation duration to test if the developmental stage at which an egg was laid affected time to hatching. Initial models of embryo heart rate and oxygen consumption also included the linear and quadratic effects of time (days post-oviposition) to account for the expected non-linearity of these parameters over time (Burggren and Warburton, 1994; Cordero et al., 2017a; Sartori et al., 2017), as well as the interaction of treatment and the linear and quadratic effects of time. As above, models included the random effect of population of origin and maternal identity. Models of heart rate and oxygen consumption also included the random effect of individual to account for covariation of measures on the same individual. For models that demonstrated a clear effect of treatment, we then tested our specific biological hypotheses by using linear contrasts of estimated marginal means using the emmeans package (Russell, 2016, 2019), correcting for multiple comparisons with the Šidák method (see Results for details). All data figures were created with the ggplot2 package (Wickham, 2009).

## RESULTS

### Maternal physiology

Reproducing lizards from low-elevation populations increased both haematocrit and haemoglobin concentration in response to translocation to high elevation (by 23.9% and 31.8% compared with low-elevation values, respectively) and values for both parameters decreased over time in all groups. Additionally, haematocrit changed non-linearly over time and was greater before oviposition (Table 1, Fig. 2A,B).


As the results of the  $\dot{V}_{CO_2}$  and  $\dot{V}_{O_2}$  analyses are qualitatively identical, we present and discuss the results of  $\dot{V}_{O_2}$  only. Resting metabolic rate increased after oviposition and did not differ among the treatment groups (Table 1, Fig. 2C). As expected, resting metabolic rates were positively dependent on both body mass and body temperature. Reactive oxygen metabolites in blood plasma were affected by time differently among the treatment groups: ROMs decreased linearly over time for lizards from low- and high-elevation populations kept at native elevations, whereas they remained elevated for lizards from low elevation transplanted to high elevation. Despite the fact that lizards at native elevations exhibited higher levels of ROMs early in reproduction, this interaction resulted in transplanted lowland lizards exhibiting higher ROMs later in reproduction (Table 1, Fig. 2D). Additionally, ROMs after oviposition were higher than predicted by the change across oviposition date. Finally, ROMs were negatively correlated with size-corrected total clutch mass, indicating that individuals investing more in reproduction also produced less reactive oxygen metabolites (Table 1, Fig. 3).



**Fig. 2. Response of haematocrit, haemoglobin concentration, resting oxygen consumption rate ( $\dot{V}_{O_2}$ ) and reactive oxygen metabolites over time in reproducing lizards from each experimental treatment.** Haematocrit (A), haemoglobin concentration (B),  $\dot{V}_{O_2}$  (C) and reactive oxygen metabolites (ROMs; D) in reproducing lizards from different elevation treatments. Lines are linear or quadratic regressions depending on the results of mixed effect models with shaded areas representing standard error of the estimate. Significant effects shown in inset with one ( $P < 0.05$ ) or two ( $P < 0.001$ ) asterisks (see Table 1 and Materials and Methods for statistical details). Relative oviposition date 0 = date of oviposition. Low–low,  $N = 14$  individuals; low–high,  $N = 16$  individuals; high–high,  $N = 17$  individuals.

### Reproductive measures

Larger lizards had larger clutches and invested more in reproduction. After accounting for this variation, egg mass at oviposition differed slightly among the maternal origin treatment



**Fig. 3. Scatterplot showing relationship between total clutch mass (corrected for body size) and mean ROMs for all females ( $N = 47$ ).** Black line represents linear regression with standard error in grey shading. After accounting for other effects, ROMs decrease with increased relative clutch mass (see Table 1 for details).

groups, although none of the pairwise differences was significant. Lizards originating from populations at different elevations did not differ in any other reproductive parameter, nor did transplanting low-elevation lizards to high elevation affect reproduction. Female *P. muralis* laid eggs with embryos that were on average at stage 27 (range: 25–28.5) of Dufaure and Hubert (1961). Finally, we saw no effect of treatment group on post-oviposition body condition (Table 2, Fig. 4A–D).

### Embryo heart and metabolic rates

Embryonic heart rates varied amongst treatment groups in a manner that was dependent upon the interactions with both linear and quadratic age post-oviposition (Table 3, Fig. 5A). Embryos from low-elevation populations generally showed a concave response, whereby heart rates were lowest early and late in incubation. Conversely, embryos from high elevations showed a convex response, whereby heart rates were highest early and late in incubation. Using linear contrasts from the mixed models to test our specific *a priori* hypotheses demonstrates that embryos from low-elevation populations exhibited higher heart rates at their native elevation compared with embryos from high-elevation populations at native elevation, on average by 15.3 beats  $\text{min}^{-1}$  (Table 4). Furthermore, embryos from low-elevation populations reduced heart rates when incubated at high elevation (by 18.9 beats  $\text{min}^{-1}$ ) or extreme high elevation (by 9.6 beats  $\text{min}^{-1}$ ). Notably, the change was greatest in the high-elevation treatment, in which heart rates were lower than the extreme high treatment by 9.3 beats  $\text{min}^{-1}$ . Embryos from high elevation populations increased heart rates when incubated at extreme high elevation compared with native (high) elevation, on average by 9.6 beats  $\text{min}^{-1}$ .



**Table 2. Results of linear mixed model analyses of reproductive parameters in gravid female *P. muralis* lizards**

Source of variation		Stage at oviposition ( <i>N</i> =39)	Clutch size ( <i>N</i> =47)	Egg mass ( <i>N</i> =253)	Relative clutch mass ( <i>N</i> =47)	Post-oviposition body condition ( <i>N</i> =47)
Treatment	<i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) <i>P</i> > <i>F</i>	0.32 (2, 9.1) 0.73	0.37 (2, 8.4) 0.70	See Fig. 4C 4.80 (2, 7.5) 0.046*	1.32 (2, 8.3) 0.32	1.32 (2, 7.2) 0.33
Snout–vent length	<i>F</i> (d.f. <sub>n</sub> , d.f. <sub>d</sub> ) <i>P</i> > <i>F</i>	–	Positive 14.17 (1, 43.0) 0.0005**	Positive 5.58 (1, 43.2) 0.023*	Positive 5.48 (1, 43.0) 0.024*	–


See Materials and Methods for statistical details. Significant factors indicated as \**P*<0.05 or \*\**P*<0.001 along with directionality.

Embryo metabolic rate (oxygen consumption;  $\dot{V}_{O_2}$ ) increased with time post-oviposition, although this response was non-linear and dependent on a quadratic function of embryo age (Table 3; Fig. 5B). Embryos from the different treatments exhibited different patterns of oxygen consumption, with this response dependent on a linear interaction with embryo age post-oviposition. The interaction of treatment and embryo age is driven by differences in embryos incubated at extreme high elevation: embryos from high-elevation populations increased oxygen consumption across incubation, whereas embryos from low-elevation populations exhibited lower levels later in incubation. No test of our *a priori* hypotheses for oxygen consumption demonstrated a clear difference between treatment groups. Furthermore, oxygen consumption was not clearly dependent on egg mass as we would expect (Vleck and

Hoyt, 1991), most likely because of the strong effect of days post-oviposition, which is correlated with egg mass.

#### Embryo survival and hatchling morphology

We found no difference among treatment groups in probability of hatching ( $\chi^2_6=5.45$ , *P*=0.49) or an effect of oviposition egg mass on survival to hatching probability ( $\chi^2_1=2.83$ , *P*=0.09). Among embryos that survived to hatching, incubation duration was affected by incubation treatment (Table 3, Fig. 6A): embryos native to low-elevation populations took an average of 1.5 days longer to hatch at high elevation and 1.4 days longer to hatch at extreme high elevation. Egg mass at oviposition strongly and positively influenced hatchling mass and body size, although these were not affected by treatment (Table 3, Fig. 6B,C).



**Fig. 4. Reproductive parameters in females from three experimental treatments (combining origin elevation and gestation treatment).** (A) Stage at oviposition. (B) Clutch size. (C) Egg mass at oviposition. (D) Relative clutch mass. Plots are Tukey box plots showing median, interquartile range and range of raw data values. Significant differences among treatment groups shown with one asterisk (*P*<0.05; see Table 2 and Materials and Methods for statistical details). Low–low, *N*=14 individuals; low–high, *N*=16 individuals; high–high, *N*=17 individuals.

**Table 3. Results of linear mixed model analyses of embryo *P. muralis* heart rate and oxygen consumption rate during development, incubation duration and hatchling morphology**


Source of variation		Heart rate ( <i>N</i> =414)	$\dot{V}_{O_2}$ ( <i>N</i> =136)	Incubation duration ( <i>N</i> =184)	Hatchling mass ( <i>N</i> =184)	Hatchling body size (SVL) ( <i>N</i> =184)
Treatment		See Fig. 5A	See Fig. 5B	See Fig. 6A		
	<i>F</i> (d.f.n, d.f.d)	4.02 (6, 306.6)	6.14 (6, 91.3)	3.01 (6, 131.9)	1.97 (6, 144.4)	1.53 (6, 40.6)
	<i>Pr&gt;F</i>	0.0007**	<0.0001**	0.009*	0.074	0.19
Oviposition egg mass					Positive	Positive
	<i>F</i> (d.f.n, d.f.d)	–	–	1.33 (1, 71.2)	42.5 (1, 79.3)	36.0 (1, 63.1)
	<i>Pr&gt;F</i>	–	–	0.25	<0.0001**	<0.0001**
Oviposition mass×treatment						
	<i>F</i> (d.f.n, d.f.d)	–	–	2.10 (6, 136.4)	2.12 (6, 146.0)	–
	<i>Pr&gt;F</i>	–	–	0.057	0.055	–
Days post-oviposition linear			Positive			
	<i>F</i> (d.f.n, d.f.d)	1.35 (1, 300.1)	29.7 (1, 81.6)	–	–	–
	<i>Pr&gt;F</i>	0.25	<0.0001**	–	–	–
Days post-oviposition quadratic			See Fig. 5B			
	<i>F</i> (d.f.n, d.f.d)	1.41 (1, 305.0)	7.52 (1, 81.8)	–	–	–
	<i>Pr&gt;F</i>	0.24	0.008*	–	–	–
Treatment×days post-oviposition linear		See Fig. 5A	See Fig. 5B			
	<i>F</i> (d.f.n, d.f.d)	2.33 (6, 302.4)	6.55 (6, 89.9)	–	–	–
	<i>Pr&gt;F</i>	0.033*	<0.0001**	–	–	–
Treatment×days post-oviposition quadratic		See Fig. 5A				
	<i>F</i> (d.f.n, d.f.d)	2.35 (6, 307.1)	–	–	–	–
	<i>Pr&gt;F</i>	0.031*	–	–	–	–

See Materials and Methods for statistical details. Significant factors indicated as \* $P<0.05$  or \*\* $P<0.001$  along with directionality.  $\dot{V}_{O_2}$ , oxygen consumption rate; SVL, snout–vent length. *N* indicates the number of observations.

## DISCUSSION

Our results demonstrate that translocation to high elevation alters the physiology of lowland gravid lizards compared with that of lizards kept at native elevations (from both low- and high-elevation populations) but does not affect short-term reproductive investment

or output. Importantly, the concentration of ROMs did not decrease during the reproductive period in transplanted lizards, as it did in lizards kept at native elevation. This suggests that there may be a deleterious effect of gestation at high elevation for non-native lizards, even if that cost is not evident in current reproduction. Interestingly,



**Fig. 5. Heart rate and  $\dot{V}_{O_2}$  over time in lizard embryos from each experimental treatment.** (A) Heart rate and (B)  $\dot{V}_{O_2}$  quadratic regressions (lines) with shaded areas representing standard error of the estimate. Significant effects shown in inset with one ( $P<0.05$ ) or two ( $P<0.001$ ) asterisks (see Table 3 and Materials and Methods for statistical details). Embryos are from three maternal treatments (low–low, *N*=14 clutches; low–high, *N*=16 clutches; high–high, *N*=17 clutches) incubated at three elevations. See Fig. 1 for complete sample size details.

**Table 4. Results of a priori hypothesis tests**

Hypothesis test		Heart rate (N=414)	$\dot{V}_{O_2}$ (log <sub>10</sub> -transformed) (N=136)	Incubation duration (N=184)
Embryos native to low elevation versus embryos native to high elevation incubated at native elevations	Estimate (s.e.)	-15.29 (4.11)	-0.047 (0.04)	0.058 (1.00)
	<i>t</i> -Statistic (d.f.)	-3.72 (53.7)	-1.10 (6.27)	0.058 (4.4)
	Pr> <i>t</i>	0.0029*	0.89	>0.99
Embryos native to low elevation incubated at native elevation versus incubated at high elevation	Estimate (s.e.)	-18.85 (3.58)	-0.076 (0.03)	1.49 (0.42)
	<i>t</i> -Statistic (d.f.)	-5.26 (375.4)	-2.71 (56.7)	3.52 (166.6)
	Pr> <i>t</i>	<0.0001**	0.052	0.003*
Embryos native to low elevation incubated at native elevation versus incubated at extreme high elevation	Estimate (s.e.)	-9.60 (3.49)	-0.039 (0.03)	1.35 (0.47)
	<i>t</i> -Statistic (d.f.)	-2.75 (363.1)	-1.35 (33.1)	2.86 (167.3)
	Pr> <i>t</i>	0.037*	0.71	0.028*
Embryos native to low elevation incubated at high elevation versus incubated at extreme high elevation	Estimate (s.e.)	9.25 (2.92)	0.037 (0.03)	-0.14 (0.36)
	<i>t</i> -Statistic (d.f.)	3.17 (338.3)	1.25 (42.5)	-0.38 (162.3)
	Pr> <i>t</i>	0.0098*	0.77	>0.99
Embryos native to high elevation incubated at native elevation versus incubated at extreme high elevation	Estimate (s.e.)	9.55 (3.60)	-0.009 (0.03)	-0.80 (0.41)
	<i>t</i> -Statistic (d.f.)	2.65 (389.0)	-0.27 (66.6)	-1.95 (156.1)
	Pr> <i>t</i>	0.049*	>0.99	0.28
Embryos native to low elevation gestated at native elevation versus gestated at high elevation	Estimate (s.e.)	0.63 (3.41)	-0.001 (0.03)	-0.56 (0.41)
	<i>t</i> -Statistic (d.f.)	0.19 (126.1)	-0.043 (35.4)	-1.37 (62.4)
	Pr> <i>t</i>	>0.99	>0.99	0.69

Tests were constructed using linear contrasts of estimated marginal means. Estimate represents the differences in observed values between groups, taking into account covariates included in the model (see Table 3 for full model results and Materials and Methods for statistical details).


Significant factors indicated as \* $P < 0.05$  or \*\* $P < 0.001$ .  $\dot{V}_{O_2}$ , oxygen consumption rate. *N* indicates the number of observations.

reactive oxygen metabolite concentration was highest in females exhibiting the lowest relative clutch mass, contrary to assumptions of a reactive metabolite cost of reproduction. Nonetheless, at least in the short-term, there appears to be no indication that reproduction in lowland *P. muralis* is limited at higher elevations within the range of the species, assuming the environment is otherwise suitable. Furthermore, the conditions experienced by females during gestation did not affect the physiology, development or hatchling morphology of embryos post-oviposition. Thus, we did not find evidence that maternal hypoxia exposure may 'prime' embryos for the constraints of post-oviposition hypoxia. Embryos post-oviposition demonstrated a clear physiological and developmental response to hypoxia, including reduced heart rate at high elevation, altered patterns of oxygen consumption – indicating diminished late-stage metabolic rate – and increased incubation duration. These adjustments in embryo physiology and developmental timing at least partially buffer these lizards from potentially negative effects of reduced oxygen availability; development at these elevations did not affect fitness-related traits, including embryo survivorship to hatching or hatchling morphology. Taken together, these results suggest that this widespread and common species, a successful invader elsewhere (Beninde et al., 2018; Deichsel and Gist, 2001; Michaelides et al., 2015), will be likely to colonize higher-elevation habitats in future climate scenarios.

### Maternal physiology and reproduction

As predicted, gravid females from low-elevation populations transplanted to high elevation had higher haematocrit and haemoglobin concentration compared with low-elevation lizards kept at low elevation. This response is concordant with other studies showing that lizards from high-elevation populations feature higher haematocrit and haemoglobin concentration compared with lowland populations (González-Morales et al., 2015; Lu et al., 2015). Haematocrit and haemoglobin concentration was increased in lowland lizards transplanted to high elevation to match lizards native to high elevations, which does not support the prediction that lizards native to different elevations would exhibit different blood chemistry profiles. The current data do not allow us to distinguish between plasticity and local adaptation in these traits in lizards native to high elevations, but in either case, this could illustrate an adaptive response of lizards to facilitate oxygen transport at high elevation.

In parallel with these results, we did not observe differences in resting metabolic rate among the treatment groups. Even in a state of elevated energy exchange, such as occurs during reproduction (Angilletta and Sears, 2000; Foucart et al., 2014), oxygen demands are met at high elevation, possibly due to the observed plasticity in blood biochemistry. While energetic demands are increased in reproductive compared with non-reproducing females, it is



**Fig. 6. Incubation duration and hatchling morphology for lizard embryos from each experimental treatment.** Incubation duration (A), hatchling mass (B) and hatchling snout–vent length (SVL) (C). Plots are Tukey box plots showing median, interquartile range and range of raw data values. Significant differences among treatment groups shown with one asterisk ( $P < 0.05$ ; see Table 3 and Materials and Methods for statistical details). Hatchlings are from three maternal treatments (low–low,  $N = 14$  clutches; low–high,  $N = 16$  clutches; high–high,  $N = 17$  clutches) incubated at three elevations. See Fig. 1 for complete sample size details.

important to note that this increased demand occurs during vitellogenesis (Foucart et al., 2014; Van Dyke and Beaupre, 2011), before the start of this experiment. Nonetheless, the question of long-term effects of increased haemoglobin concentration and haematocrit remains because a higher concentration of red blood cells increases blood viscosity. This phenomenon could then carry an important cost for blood circulation (Dunlap, 2006; Hedrick

et al., 1986) and may not be maintained, as observed in lowland males of this species transplanted to extreme high elevation (Gangloff et al., 2019). Increasing red blood cell concentration is especially constraining during gravidity when haematocrit drops, probably to reduce viscosity, and facilitate circulation and gas transport to developing embryos (e.g. Dupoué et al., 2015).

We found little evidence of important differences among treatment groups in indicators of reproductive investment, including clutch size and relative clutch mass. After accounting for variation due to maternal size, we identified a trend whereby lizards at high elevations (either native or transplanted) produced eggs with a greater mass at oviposition. This finding is counter to our prediction that lizards at higher elevations would demonstrate reduced reproductive output. Indeed, through changes in physiological parameters, females appeared to compensate for reduced oxygen availability and maintained reproductive output across treatments. It is important to recognize that this experiment encompassed only the first reproductive event of a single year. Lowland lizards transplanted to high elevation were able to maintain reproductive output without an apparent cost in terms of body condition in this bout, but there may be a trade-off with future reproduction such as via ROM damage. Although stage at oviposition did not differ among treatment groups and was within the expected range for most oviparous lizards (reviewed in Mathies and Andrews, 1999), our findings broadly support the prediction that gestation duration is a highly stable trait whose population mean might undergo gradual intergenerational change as an adaptive evolutionary response to high-altitude conditions (e.g. temperature, Mathies and Andrews, 1995). Thus, because stage at oviposition might be highly canalized in *P. muralis*, variation in this trait may not manifest until several generations are exposed to oxygen-limited conditions, similar to the adaptive evolution of prolonged egg retention in cool environments (Telemeco et al., 2010; While et al., 2015).

We provide two important results regarding ROM production in response to both reproduction and hypoxia. First, we found that females with increased reproductive investment (size-corrected total clutch mass) had lower levels of ROMs in blood plasma. Broadly, this result does not support the common assumption that reproduction incurs an oxidative cost and thereby mediates life-history trade-offs (Dowling and Simmons, 2009; Metcalfe and Monaghan, 2013; Speakman and Garratt, 2014). Previous work in lizards concurs with our results: no such oxidative cost of reproduction was found in two other lizard species (*Niveoscincus ocellatus*, Isaksson et al., 2011; *Ctenophorus pictus*, Olsson et al., 2009, 2012). This result suggests that both ROM production and reproductive investment could be correlated via some unmeasured physiological parameter representing overall ‘individual quality’: higher-quality individuals are able to both invest more in reproduction and meet energetic demands while producing fewer potentially detrimental by-products (Wilson and Nussey, 2010). On the contrary, Webb et al. (2018) found that increased reproductive investment (as measured by follicle number) was associated with increased ROMs in the lizard *Cyclura cychlura inornata*. Importantly, Webb et al. (2018) measured this effect during vitellogenesis, thus pointing to this period as being key to understanding the relationship between reproductive investment and ROM production. A similar pattern of higher reactive oxygen metabolite production earlier in the reproductive cycle is also found in snakes (*Antaresia childreni*, Stahlschmidt et al., 2013) and birds (*Sula leucogaster*, Montoya et al., 2016). Our results agree with this pattern, whereby ROMs decreased during the gestation period, indicating that such potentially harmful by-products are greatest earlier in the reproductive process.

Our second significant finding is that both low- and high-elevation lizards kept at native elevations decreased reactive oxygen metabolite production across the reproductive period, while lowland lizards transplanted to high elevations did not. We might expect that the reactive oxygen cost of reproduction will likely be context dependent, perhaps because of resource limitations and subsequent allocation decisions (Dowling and Simmons, 2009). In our case, it appears that the reduced oxygen availability at higher elevations elicits such a constraint and prohibits females from decreasing ROM production across the reproductive period. Our results demonstrate higher reactive oxygen metabolite production later in gravidity and after oviposition in transplanted lizards. These elevated levels of ROMs have the potential to bear costs for future fitness in both the mothers and their offspring, given the potential damaging effects on proteins, membranes and DNA (Blount et al., 2016; Costantini, 2016; Speakman and Garratt, 2014). Unfortunately, plasma volume limitations precluded us from also measuring antioxidant capacity, which would provide valuable data to quantify the response to these damaging molecules. For example, antioxidant capacity may change across the reproductive period, thus mitigating the potential negative impacts of increased ROM production. Future work is needed to understand patterns of ROMs mediating physiological trade-offs at high elevations by measuring additional tissues, measuring antioxidant and repair capacity, and quantifying the long-term impacts of increased ROM levels (Speakman and Garratt, 2014).

### Embryo physiology and development

Developing *P. muralis* embryos appear to be robust to the effects of ecologically relevant hypoxia, at least until hatching and the beginning of the free-living stage. The most dramatic physiological response is the decreased embryonic heart rate in naturally occurring hypoxia at high elevation (Cordero et al., 2017a), which is similar to results seen in many oviparous species exposed to laboratory-simulated hypoxia (Crossley and Altimiras, 2005; Du et al., 2010; Nechaeva and Vladimirova, 2008; VanGolde et al., 1997). This response is coupled with a relative increase in heart size in *P. muralis* (Cordero et al., 2017a), which may facilitate the maintenance of oxygen delivery via increased stroke volume. This response is predicted under a scenario where hypervolemia increases the optimal haematocrit: oxygen delivery is maintained, despite reductions in heart rate, via increases in haematocrit and stroke volume (Birchard, 2015; Weibel et al., 1991). While work is needed to test these specific relationships among parameters important to efficient oxygen delivery under hypoxia, we found that embryos at high and extreme high elevations maintained comparable metabolic rates (as measured by oxygen consumption). This concurs with previous studies finding that incubation under hypoxia had little effect on the metabolic rates of developing alligator embryos (*Alligator mississippiensis*, Crossley et al., 2017; Warburton et al., 1995).

While oxygen consumption generally increased during incubation, the pattern was dependent on treatment. Notably, lowland embryos incubated at extreme high elevation exhibited a pattern whereby oxygen consumption decreased late in incubation. This pattern is mirrored in the pattern of heart rate variation across incubation, suggesting a late-stage metabolic suppression resulting from oxygen limitation. This metabolic constraint on low-elevation *P. muralis* embryos developing at high and extreme high elevations could also explain the increased incubation duration of these embryos. That late-stage embryos native to high elevation did not experience this metabolic suppression and did not alter incubation times at extreme high elevation suggests a role for local adaptation

in maintaining physiological and developmental robustness in hypoxia. Although late-stage embryos of sea-level chicken populations also exhibited reduced oxygen consumption when initially transplanted to high elevation, oxygen consumption rates exceeded sea-level values after >9 generations of selective breeding (based on hatching success) at 3100 m ASL (Beattie and Smith, 1975; Wangenstein et al., 1974). The fitness consequences of hypoxia-induced metabolic suppression in non-avian reptilian embryos awaits further experimentation.

While embryo physiology and development responded to high-elevation hypoxia, neither survival to hatching nor hatchling morphology were affected. This corroborates our previous work in this system in demonstrating that early-life stages are relatively resilient to hypoxia and therefore would not limit future colonization of high-elevation habitats beyond the current species range. We emphasize that even our extreme high elevation treatment represents a moderate and biologically relevant reduction in oxygen availability (approximately 72% of sea level equivalent), which explains why we did not find the clear negative effects of hypoxia as have other studies in reptiles exposed to more severe hypoxia, even for short periods of time (Andrews, 2002; Cordero et al., 2017b; Lungman and Piña, 2013; Kam, 1993). The lack of effect of our maternal gestation treatments implies that embryos pre-oviposition did not face a mismatch between oxygen supply and demand. Our data suggest a maternal ability to maintain oxygen supply to tissues, including developing embryos, by adjusting oxygen-carrying capacity (haematocrit, haemoglobin) to maintain metabolic rates. Furthermore, the early stages of embryonic development in lizards require low oxygen consumption (Thompson and Stewart, 1997), as during this time embryos only constitute ~10% of their hatchling body mass (Andrews, 2004). A fruitful direction for future experiments is to transplant reproductive females earlier in the reproductive process (i.e. before vitellogenesis) to test for trade-offs in reproductive investment between offspring size and number and also for earlier maternal exposure to affect offspring development, either via adaptive maternal effects (e.g. hormonal mechanisms) or because of energetic constraints imposed by hypoxia.

We report this apparent resilience of embryos to hypoxia with the important caveat that oxygen availability interacts with temperature to exacerbate the performance limitations and negative effects of hypoxia at higher temperatures (Gangloff and Telemeco, 2018). This may be most notable in the late embryonic stages, in which oxygen uptake depends on diffusion across the chorioallantoic membrane and behavioural responses to the environment are highly restricted (Cordero et al., 2018; Telemeco et al., 2016; but see Shine and Du, 2018). Work in other reptile taxa has demonstrated a clear pattern whereby the detrimental effects of hypoxia are more severe at higher temperatures (Flewelling and Parker, 2015; Lungman and Piña, 2013; Liang et al., 2015; Smith et al., 2015). The interaction of temperature and hypoxia likely also explains why we found longer incubation periods for low-elevation embryos incubated at higher elevations, while our previous work did not find such an effect. In the current study, we incubated eggs at 28°C, whereas previously we used 24°C (Cordero et al., 2017a). The impact of hypoxia on prolonging development times was more pronounced at higher temperatures, which is probably due to limitations of the embryo to meet the corresponding increased oxygen demand, such as found previously in caimans (*Caiman latirostris*; Lungman and Piña, 2013). While the effects of reduced oxygen availability and high temperatures interact to impact physiology, it is also possible that shared response pathways offer protection against these simultaneous stressors (e.g. Teague et al., 2017). Exploring the inter-relation of these response

pathways, as well as how responses may change across life-history stages, remains an important avenue for research.

Our results concur with other recent work demonstrating the apparent flexibility of reptilian embryos in response to novel environments (e.g. Tiatragul et al., 2017). This is not surprising, given that reptile nests are frequently exposed to low-oxygen conditions in natural settings (Ackerman and Lott, 2004; Booth, 1998; Packard and Packard, 1987). Thus, evolution may have promoted physiological mechanisms to deal with hypoxia in embryos that can then facilitate development under situations of reduced oxygen partial pressure, as found at high elevations. Physiological plasticity to adjust to hypoxia could thus be stage-specific, explaining the apparent resilience of *P. muralis* embryos but not adults (Cordero et al., 2017a; Gangloff et al., 2019). The long-term effects of hypoxia exposure during development on phenotype and fitness at later life-history stages remain important research questions. For example, development in hypoxia has long-term effects on post-hatching cardiovascular phenotypes in snapping turtles (*Chelydra serpentina*; Wearing et al., 2016, 2017), decreases swimming performance in snakes (*Natrix maura*; F.A. and J.S., unpublished results), and impacts cognitive ability in lizards (*Eremias argus*; Sun et al., 2014). The adaptive significance of embryonic plasticity in response to hypoxia can only be evaluated by quantifying the long-term effects on individual fitness (Mitchell et al., 2018).

## Conclusion

This study addresses two important dimensions of reproduction, with implications for understanding both life histories and range expansion in mountainous regions in common wall lizards: ROM production and the effects of hypoxia. Females transplanted to high elevation matched the reproductive output of lizards native to both low-elevation and high-elevation populations, and embryos are physiologically resilient to development at high elevations, even above the current range limits. This suggests that these life stages will not limit initial colonization of high-elevation habitats. However, population establishment will depend on the long-term costs associated with life in reduced oxygen availability on adults (such as damage from increased ROM production) and the consequences of reduced performance (Gangloff et al., 2019). Colonization of high-altitude areas is important for conservation efforts as it will likely put high-elevation specialist lizards endemic to small mountaintop ranges, such as *Iberolacerta* spp., at further risk via competition for resources and direct interactions (Žagar et al., 2015, 2017). With our work in this system, we hope to quantify the relative potential of physiological plasticity and local adaptation across life history stages to facilitate range expansion in a species likely to benefit under climate change scenarios.

## Acknowledgements

We gratefully acknowledge logistical support from Y. Ardrif, S. Bonzom, M. Bousquet, H. Le Chevalier, E. Darnet, A. Dolfi, M. Huet, M. Istoczek, A. Jossierand, L. Rabardelle, and the staff of the Observatoire Midi-Pyrénées. A. Chaine provided insightful manuscript feedback and B. Joos at Sable Systems provided valuable technical advice, as usual.

## Competing interests

The authors declare no competing or financial interests.

## Author contributions

Conceptualization: L.K., E.J.G., J.S., G.A.C., F.A.; Methodology: E.J.G., A.D.; Investigation: L.K., E.J.G., J.S., G.A.C., A.D., F.A.; Resources: F.A.; Data curation: L.K., E.J.G.; Writing - original draft: L.K., E.J.G.; Writing - review & editing: L.K., E.J.G., J.S., G.A.C., A.D., F.A.; Supervision: E.J.G., F.A.; Project administration: F.A.; Funding acquisition: E.J.G., F.A.

## Funding

This work was supported by the Laboratoire d'Excellence (LabEx) TULIP (ANR-10-LABX-41), the Interreg POCTEFA ECTOPYR (no. EFA031/15), and the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement (752299). G.A.C. was supported by a Company of Biologists Travelling Fellowship.

## Data availability

Data are archived in Mendeley at DOI: 10.17632/94xsf6wv6c.1.

## Supplementary information

Supplementary information available online at <http://jeb.biologists.org/lookup/doi/10.1242/jeb.206839.supplemental>

## References

- Ackerman, R. A. and Lott, D. B. (2004). Thermal, hydric and respiratory climate of nests. In *Reptilian Incubation: Environment, Evolution and Behavior* (ed. D. C. Deeming), pp. 15-44. Nottingham, UK: Nottingham University Press.
- Andrews, R. M. (2002). Low oxygen: a constraint on the evolution of viviparity in reptiles. *Physiol. Biochem. Zool.* **75**, 145-154. doi:10.1086/339388
- Andrews, R. M. (2004). Patterns of embryonic development. In *Reptilian Incubation: Environment, Evolution and Behavior* (ed. D. C. Deeming), pp. 75-102. Nottingham, UK: Nottingham University Press.
- Andrews, R. M. and Pough, F. H. (1985). Metabolism of squamate reptiles: allometric and ecological relationships. *Physiol. Zool.* **58**, 214-231. doi:10.1086/physzool.58.2.30158569
- Angilletta, M. J. and Sears, M. W. (2000). The metabolic cost of reproduction in an oviparous lizard. *Funct. Ecol.* **14**, 39-45. doi:10.1046/j.1365-2435.2000.00387.x
- Arnold, E. N., Burton, J. A., Ovenden, D. and Coppo, G. (1988). *Tous les reptiles et amphibiens d'Europe en couleurs*. Paris: Bordas Paris.
- Aubret, F. (2013). Role and evolution of adaptive plasticity during the colonization of novel environments. In *Phenotypic Plasticity: Molecular Mechanisms, Evolutionary Significance and Impact on Speciation* (ed. J. B. Valentino and P. C. Harrelson), pp. 1-34. New York: Novinka.
- Bani, L., Luppi, M., Rocchia, E., Dondina, O. and Orioli, V. (2019). Winners and losers: how the elevational range of breeding birds on Alps has varied over the past four decades due to climate and habitat changes. *Ecol. Evol.* **9**, 1289-1305. doi:10.1002/ece3.4838
- Bassler, C., Hotborn, T., Brandl, R. and Müller, J. (2013). Insects overshoot in the expected upslope shift caused by climate warming. *PLoS ONE* **8**, e0065842. doi:10.1371/journal.pone.0065842
- Beattie, J. and Smith, A. H. (1975). Metabolic adaptation of the chick embryo to chronic hypoxia. *Am. J. Physiol.* **228**, 1346-1350. doi:10.1152/ajplegacy.1975.228.5.1346
- Beninde, J., Feldmeier, S., Veith, M. and Hochkirch, A. (2018). Admixture of hybrid swarms of native and introduced lizards in cities is determined by the cityscape structure and invasion history. *Proc. R. Soc. B* **285**, 20180143. doi:10.1098/rspb.2018.0143
- Birchard, G. F. (2015). Optimal hematocrit: theory, regulation and implications. *Am. Zool.* **37**, 65-72. doi:10.1093/icb/37.1.65
- Blackburn, D. G. (1998). Resorption of oviductal eggs and embryos in squamate reptiles. *Herpetol. J.* **8**, 65-71.
- Blomberg, S. and Shine, R. (1996). Reptiles. In *Ecological Census Techniques* (ed. W. J. Sutherland), pp. 218-226. Cambridge: Cambridge University Press.
- Blount, J. D., Vitikainen, E. I. K., Stott, I. and Cant, M. A. (2016). Oxidative shielding and the cost of reproduction. *Biol. Rev. Camb. Philos. Soc.* **91**, 483-497. doi:10.1111/brv.12179
- Booth, D. T. (1998). Nest temperature and respiratory gases during natural incubation in the broad-shelled river turtle, *Chelodina expansa* (Testudinata: Chelidae). *Au. J. Zool.* **46**, 183-191. doi:10.1071/ZO98003
- Bouverot, P. (1985). *Adaptation to Altitude-Hypoxia in Vertebrates*. Berlin: Springer-Verlag Berlin Heidelberg.
- Braña, F. (1993). Shifts in body temperature and escape behaviour of female *Podarcis muralis* during pregnancy. *Oikos* **66**, 216. doi:10.2307/3544807
- Burggren, W. W. and Warburton, S. J. (1994). Patterns of form and function in developing hearts: Contributions from non-mammalian vertebrates. *Cardioscience* **5**, 183-191.
- Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B. and Thomas, C. D. (2011). Rapid range shifts of species associated with high levels of climate warming. *Science* **333**, 1024-1026. doi:10.1126/science.1206432
- Cordero, G. A. and Janzen, F. J. (2014). An enhanced developmental staging table for the painted turtle *Chrysemys picta* (Testudininae: Emydidae). *J. Morphol.* **275**, 442-455. doi:10.1002/jmor.20226
- Cordero, G. A., Andersson, B. A., Souchet, J., Micheli, G., Noble, D. W. A., Gangloff, E. J., Uller, T. and Aubret, F. (2017a). Physiological plasticity in lizard embryos exposed to high-altitude hypoxia. *J. Exp. Zool. A Ecol. Integr. Physiol.* **327**, 423-432. doi:10.1002/jez.2115

- Cordero, G. A., Karnatz, M. L., Svendsen, J. C. and Gangloff, E. J. (2017b). Effects of low-oxygen conditions on embryo growth in the painted turtle, *Chrysemys picta*. *Integr Zool.* **12**, 148-156. doi:10.1111/1749-4877.12206
- Cordero, G. A., Telemeco, R. S. and Gangloff, E. J. (2018). Reptile embryos are not capable of behavioral thermoregulation in the egg. *Evol. Dev.* **20**, 40-47. doi:10.1111/ede.12244
- Corona, T. B. and Warburton, S. J. (2000). Regional hypoxia elicits regional changes in chorioallantoic membrane vascular density in alligator but not chicken embryos. *Comp. Biochem. Physiol.* **125A**, 51-56. doi:10.1016/S1095-6433(99)00159-2
- Costantini, D. (2016). Oxidative stress ecology and the d-ROMs test: facts, misfacts and an appraisal of a decade's work. *Behav. Ecol. Sociobiol.* **70**, 809-820. doi:10.1007/s00265-016-2091-5
- Crossley, D. A. and Altimiras, J. (2005). Cardiovascular development in embryos of the american alligator *Alligator mississippiensis*: effects of chronic and acute hypoxia. *J. Exp. Biol.* **208**, 31-39. doi:10.1242/jeb.01355
- Crossley, D. A., Ling, R., Nelson, D., Gillium, T., Conner, J., Hapgood, J., Elsey, R. M. and Eme, J. (2017). Metabolic responses to chronic hypoxic incubation in embryonic American alligators (*Alligator mississippiensis*). *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **203**, 77-82. doi:10.1016/j.cbpa.2016.08.017
- Cunningham, E. L., Brody, J. S. and Jain, B. P. (1974). Lung growth induced by hypoxia. *J. Appl. Physiol.* **37**, 362-366. doi:10.1152/jappl.1974.37.3.362
- Deichsel, G. and Gist, D. H. (2001). On the origin of the common wall lizards *Podarcis muralis* (Reptilia: Lacertidae) in Cincinnati, Ohio. *Herpetol. Rev.* **32**, 230-232.
- Diele-Viegas, L. M. and Rocha, C. F. D. (2018). Unraveling the influences of climate change in *Lepidosauria* (Reptilia). *J. Therm. Biol.* **78**, 401-414. doi:10.1016/j.jtherbio.2018.11.005
- Dirnböck, T., Essl, F. and Rabitsch, W. (2011). Disproportional risk for habitat loss of high-altitude endemic species under climate change. *Global Change Biol.* **17**, 990-996. doi:10.1111/j.1365-2486.2010.02266.x
- Douglas, B., Maechler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed-effects models using lme4. *J. Stat. Soft.* **67**, 1-48. doi:10.18637/jss.v067.i01
- Dowling, D. K. and Simmons, L. W. (2009). Reactive oxygen species as universal constraints in life-history evolution. *Proc. R. Soc. B* **276**, 1737-1745. doi:10.1098/rspb.2008.1791
- Du, W.-G., Thompson, M. B. and Shine, R. (2010). Facultative cardiac responses to regional hypoxia in lizard embryos. *Comp. Biochem. Physiol. A* **156**, 491-494. doi:10.1016/j.cbpa.2010.04.005
- Dufaure, J. P. and Hubert, J. (1961). Table de développement du lézard vivipare-*Lacerta (Zootoca) vivipara* Jacquin. *Archives d'anatomie microscopique et de morphologie expérimentale* **50**, 309.
- Dunlap, K. D. (2006). Ontogeny and scaling of hematocrit and blood viscosity in Western Fence Lizards, *Sceloporus occidentalis*. *Copeia* **2006**, 535-538.
- Dupoué, A., Brischox, F., Angelier, F., DeNardo, D. F., Wright, C. D. and Lourdais, O. (2015). Intergenerational trade-off for water may induce a mother-offspring conflict in favour of embryos in a viviparous snake. *Funct. Ecol.* **29**, 414-422. doi:10.1111/1365-2435.12349
- Fitzgerald, L. A. (2012). Finding and capturing reptiles. In *Reptile Biodiversity: Standard Methods for Inventory and Monitoring* (ed. R. W. McDiarmid, M. S. Foster, C. Guyer, N. Chernoff and J. W. Gibbons), pp. 77-88. Berkeley: University of California Press.
- Flewelling, S. and Parker, S. L. (2015). Effects of temperature and oxygen on growth and differentiation of embryos of the ground skink, *Scincella lateralis*. *J. Exp. Zool. A* **323**, 445-455. doi:10.1002/jez.1940
- Foucart, T., Lourdais, O., DeNardo, D. F. and Heulin, B. (2014). Influence of reproductive mode on metabolic costs of reproduction: insight from the bimodal lizard *Zootoca vivipara*. *J. Exp. Biol.* **217**, 4049-4056. doi:10.1242/jeb.104315
- Freeman, B. G., Scholer, M. N., Ruiz-Gutierrez, V. and Fitzpatrick, J. W. (2018). Climate change causes upslope shifts and mountaintop extirpations in a tropical bird community. *Proc Nat Acad Sci. USA* **115**, 11982-11987. doi:10.1073/pnas.1804224115
- Gangloff, E. J. and Telemeco, R. S. (2018). High temperature, oxygen, and performance: Insights from reptiles and amphibians. *Integr Comp Biol.* **58**, 9-24. doi:10.1093/icb/icy005
- Gangloff, E. J., Sorlin, M., Cordero, G. A., Souchet, J. and Aubret, F. (2019). Lizards at the peak: Physiological plasticity does not maintain performance in lizards transplanted to high altitude. *Physiol. Biochem. Zool.* **92**, 189-200. doi:10.1086/701793
- González-Morales, J. C., Quintana, E., Díaz-Albiter, H., Guevara-Fiore, P. and Fajardo, V. (2015). Is erythrocyte size a strategy to avoid hypoxia in Wiegmann's Torquate Lizards (*Sceloporus torquatus*)? Field evidence. *Can. J. Zool.* **93**, 377-382. doi:10.1139/cjz-2014-0265
- Guzy, R. D. and Schumacker, P. T. (2006). Oxygen sensing by mitochondria at complex III: the paradox of increased reactive oxygen species during hypoxia. *Exp. Physiol.* **91**, 807-819. doi:10.1113/expphysiol.2006.033506
- Harrison, J. F., Shingleton, A. W. and Callier, V. (2015). Stunted by developing in hypoxia: linking comparative and model organism studies. *Physiol. Biochem. Zool.* **88**, 455-470. doi:10.1086/682216
- He, J., Xiu, M., Tang, X., Yue, F., Wang, N., Yang, S. and Chen, Q. (2013). The different mechanisms of hypoxic acclimatization and adaptation in lizard *Phrynocephalus vlangalii* living on Qinghai-Tibet Plateau. *J. Exp. Zool. A Ecol. Integr. Physiol.* **319**, 117-123. doi:10.1002/jez.1776
- Hedrick, M. S., Duffield, D. A. and Cornell, L. H. (1986). Blood viscosity and optimal hematocrit in a deep-diving mammal, the northern elephant seal (*Mirounga angustirostris*). *Can. J. Zool.* **64**, 2081-2085. doi:10.1139/z86-317
- Hewitt, G. M. (1999). Post-glacial re-colonization of European biota. *Biol. J. Linn. Soc.* **68**, 87-112. doi:10.1111/j.1095-8312.1999.tb01160.x
- Hickling, R., Roy, D. B., Hill, J. K., Fox, R. and Thomas, C. D. (2006). The distributions of a wide range of taxonomic groups are expanding polewards. *Glob. Change Biol.* **12**, 450-455. doi:10.1111/j.1365-2486.2006.01116.x
- Hillyard, S. D. (1980). Respiratory and cardiovascular adaptations of amphibians and reptiles to altitude. In *Environmental Physiology: Aging, Heat and Altitude* (ed. S. M. Horvath and M. K. Yousef), pp. 363-377. New York: Elsevier.
- Huey, R. B. (1982). Temperature physiology, and the ecology of reptiles. In *Biology of the Reptilia, Physiology C: Physiological Ecology*, Vol. 12. (ed. C. Gans and F. H. Pough), pp. 25-91. New York: Academic Press.
- Huey, R. B., Buckley, L. B. and Du, W. (2018). Biological buffers and the impacts of climate change. *Integr. Zool.* **13**, 349-354. doi:10.1111/1749-4877.12321
- Hulbert, A. C., Mitchell, T. S., Hall, J. M., Guiffre, C. M., Douglass, D. C. and Warner, D. A. (2017). The effects of incubation temperature and experimental design on heart rates of lizards embryos. *J. Exp. Zool. A* **327**, 466-476. doi:10.1002/jez.2135
- Isaksson, C., While, G. M., Olsson, M., Komdeur, J. and Wapstra, E. (2011). Oxidative stress physiology in relation to life history traits of a free-living vertebrate: the spotted snow skink, *Niveoscincus ocellatus*. *Integr. Zool.* **6**, 140-149. doi:10.1111/j.1749-4877.2011.00237.x
- Jungman, J. L. and Piña, C. I. (2013). Hypoxia and temperature: does hypoxia affect caiman embryo differentiation rate or rate of growth only? *J. Therm. Biol.* **38**, 407-418. doi:10.1016/j.jtherbio.2013.05.003
- Jackson, D. C. (2007). Temperature and hypoxia in ectothermic tetrapods. *J. Therm. Biol.* **32**, 125-133. doi:10.1016/j.jtherbio.2007.01.007
- Jochmans-Lemoine, A. and Joseph, V. (2018). Case study: Developmental physiology at high altitude. In *Development and Environment* (ed. W. Burggren and B. Dubansky), pp. 435-457. Switzerland: Springer.
- Jochmans-Lemoine, A., Villalpando, G., Gonzales, M., Valverde, I., Soria, R. and Joseph, V. (2015). Divergent physiological responses in laboratory rats and mice raised at high altitude. *J. Exp. Biol.* **218**, 1035-1043. doi:10.1242/jeb.112862
- Kam, Y.-C. (1993). Physiological effects of hypoxia on metabolism and growth of turtle embryos. *Respir Physiol.* **92**, 127-138. doi:10.1016/0034-5687(93)90033-7
- Kenward, M. G. and Roger, J. H. (1997). Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics* **53**, 983-997. doi:10.2307/2533558
- Le Galliard, J. F., Massot, M., Baron, J. P. and Clobert, J. (2012). Ecological effects of climate change on European reptiles. In *Wildlife conservation in a changing climate* (ed. J.F. Brodie, E.S. Post and D.F. Doak), pp. 179-203. Chicago: University of Chicago Press.
- Le Héaniff, M. (2011). Stratégie reproductrice d'une espèce de lézard à pontes multiples (*Podarcis muralis*) dans un environnement contraignant. Sciences pour l'Environnement Gay Lussac. 212p. *Master's Thesis*: Université de Poitiers.
- Levy, O., Borchert, J. D., Rusch, T. W., Buckley, L. B. and Angilletta, M. J. (2017). Diminishing returns limit energetic costs of climate change. *Ecology* **98**, 1217-1228. doi:10.1002/ecy.1803
- Liang, L., Sun, B.-J., Ma, L. and Du, W.-G. (2015). Oxygen-dependent heat tolerance and developmental plasticity in turtle embryos. *J. Comp. Physiol. B* **185**, 257-263. doi:10.1007/s00360-014-0874-4
- Lighton, J. R. B. (2008). *Measuring Metabolic Rates: A Manual for Scientists*. USA: Oxford University Press.
- Lu, S., Xin, Y., Tang, X., Yue, F., Wang, H., Bai, Y., Niu, Y. and Chen, Q. (2015). Differences in hematological traits between high- and low-altitude lizards (Genus *Phrynocephalus*). *PLoS ONE* **10**, e0125751. doi:10.1371/journal.pone.0125751
- MacLean, G. S., Lee, A. K. and Wilson, K. J. (1973). A simple method of obtaining blood from lizards. *Copeia* **1973**, 338-339. doi:10.2307/1442973
- Mathies, T. and Andrews, R. M. (1995). Thermal and reproductive biology of high and low elevation population of the lizard *Sceloporus scalaris*: Implications for the evolution of viviparity. *Oecologia* **104**, 101-111. doi:10.1007/BF00365568
- Mathies, T. and Andrews, R. M. (1999). Determinants of embryonic stage at oviposition in the lizard *Urosaurus ornatus*. *Physiol. Biochem. Zool.* **72**, 645-655. doi:10.1086/316707
- Metcalfe, N. B. and Monaghan, P. (2013). Does reproduction cause oxidative stress? An open question. *Trends Ecol. Evol.* **28**, 347-350. doi:10.1016/j.tree.2013.01.015
- Meylan, S., Dufty, A. M. and Clobert, J. (2003). The effect of transdermal corticosterone application on plasma corticosterone levels in pregnant *Lacerta vivipara*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **134**, 497-503. doi:10.1016/S1095-6433(02)00343-4
- Michaelides, S. N., While, G. M., Zajac, N. and Uller, T. (2015). Widespread primary, but geographically restricted secondary, human introductions of wall lizards, *Podarcis muralis*. *Mol. Ecol.* **24**, 2702-2714. doi:10.1111/mec.13206

- Mitchell, T. S., Janzen, F. J. and Warner, D. A. (2018). Quantifying the effects of embryonic phenotypic plasticity on adult phenotypes in reptiles: a review of current knowledge and major gaps. *J. Exp. Zool. A Ecol. Integr. Physiol.* **329**, 203-214. doi:10.1002/jez.2187
- Monge, C. and Leon-Velarde, F. (1991). Physiological adaptation to high altitude: oxygen transport in mammals and birds. *Physiol. Rev.* **71**, 1135-1172. doi:10.1152/physrev.1991.71.4.1135
- Montoya, B., Valverde, M., Rojas, E. and Torres, R. (2016). Oxidative stress during courtship affects male and female reproductive effort differentially in a wild bird with biparental care. *J. Exp. Biol.* **219**, 3915-3926. doi:10.1242/jeb.141325
- Moore, D., Stow, A. and Kearney, M. R. (2018). Under the weather?-The direct effects of climate warming on a threatened desert lizard are mediated by their activity phase and burrow system. *J. Anim. Ecol.* **87**, 660-671. doi:10.1111/1365-2656.12812
- Nechaeva, M. V. (2011). Physiological responses to acute changes in temperature and oxygenation in bird and reptile embryos. *Respir. Physiol. Neurobiol.* **178**, 108-117. doi:10.1016/j.resp.2011.04.003
- Nechaeva, M. and Vladimirova, I. (2008). Effect of hypoxia on amnion rhythmic contractions and heart rate in the embryonic European pond turtle *Emys orbicularis* (Reptilia: Emydidae). *Biol. Bull.* **35**, 398. doi:10.1134/S1062359008040122
- Nogués-Bravo, D., Araújo, M. B., Lasanta, T. and Moreno, J. I. L. (2008). Climate change in Mediterranean mountains during the 21st century. *Ambio* **37**, 280-285.
- Olsson, M., Wilson, M., Uller, T., Mott, B. and Isaksson, C. (2009). Variation in levels of reactive oxygen species is explained by maternal identity, sex and body-size-corrected clutch size in a lizard. *Naturwissenschaften* **96**, 25-29. doi:10.1007/s00114-008-0444-2
- Olsson, M., Healey, M., Perrin, C., Wilson, M. and Tobler, M. (2012). Sex-specific SOD levels and DNA damage in painted dragon lizards (*Ctenophorus pictus*). *Oecologia* **170**, 917-924. doi:10.1007/s00442-012-2383-z
- Packard, G. C. and Packard, M. J. (1987). The physiological ecology of reptilian eggs and embryos. In *Biology of the Reptilia*, Vol. 16. Ecology B. Defense and Life History (ed. C. Gans and R. Huey), pp 523-606. New York: Alan R. Liss.
- Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Ann. Rev. Ecol. Evol. System* **37**, 637-669. doi:10.1146/annurev.ecolsys.37.091305.110100
- Pauchard, A., Milbau, A., Albiñ, A., Alexander, J., Burgess, T., Daehler, C., Englund, G., Essl, F., Evengård, B. and Greenwood, G. B. (2015). Non-native and native organisms moving into high elevation and high latitude ecosystems in an era of climate change: new challenges for ecology and conservation. *Biol. Inv.* **18**, 345-353. doi:10.1007/s10530-015-1025-x
- Pontes-da-Silva, E., Magnusson, W. E., Sinervo, B., Caetano, G. H., Miles, D. B., Colli, G. R., Diele-Viegas, L. M., Fenker, J., Santos, J. C. and Werneck, F. P. (2018). Extinction risks forced by climatic change and intraspecific variation in the thermal physiology of a tropical lizard. *J. Therm. Biol.* **73**, 50-60. doi:10.1016/j.jtherbio.2018.01.013
- Pottier, G. (2012). Plan National d'Actions en faveur des Lézards des Pyrénées. In *Plans Nationaux d'Action pour les espèces menacées en France* (ed. Ministère de l'Écologie, du Développement durable et de l'Énergie). Bagnères de Bigorre: Nature Midi-Pyrénées.
- Powell, F. L. and Hopkins, S. R. (2010). Vertebrate life at high altitude. In *Respiratory Physiology of Vertebrates* (ed. G. E. Nilsson), pp. 265-299. New York: Cambridge University Press.
- Reyes, R. V., Díaz, M., Ebersperger, G., Herrera, E. A., Quezada, S. A., Hernandez, I., Sanhueza, E. and Llanos, A. J. (2018). The role of nitric oxide in the cardiopulmonary response to hypoxia in highland and lowland newborn llamas. *J. Physiol.* **596**, 5907-5923. doi:10.1113/JP274340
- Robert, K. A., Brunet-Rossini, A. and Bronikowski, A. M. (2007). Testing the 'free radical theory of aging' hypothesis: physiological differences in long-lived and short-lived colubrid snakes. *Aging Cell* **6**, 395-404. doi:10.1111/j.1474-9726.2007.00287.x
- Russell, V. L. (2016). Least-squares means: the r package lsmmeans. *J. Stat. Softw.* **69**, 1-33. doi:10.18637/jss.v069.i01
- Russell, V. L. (2019). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.3.3.
- Saito, M., Mano, T., Iwase, S., Koga, K., Abe, H. and Yamazaki, Y. (1988). Responses in muscle sympathetic activity to acute hypoxia in humans. *J. Appl. Physiol.* **65**, 1548-1552. doi:10.1152/jappl.1988.65.4.1548
- Sartori, M. R., Abe, A. S., Crossley, D. A. and Taylor, E. W. (2017). Rates of oxygen uptake increase independently of changes in heart rate in late stages of development and at hatching in the green iguana, *Iguana iguana*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **205**, 28-34. doi:10.1016/j.cbpa.2016.12.020
- Shama, L. N. S., Strobel, A., Mark, F. C. and Wegner, K. M. (2014). Transgenerational plasticity in marine sticklebacks: maternal effects mediate impacts of a warming ocean. *Funct. Ecol.* **28**, 1482-1493. doi:10.1111/1365-2435.12280
- Shine, R. and Du, W.-G. (2018). How frequent and important is behavioral thermoregulation by embryonic reptiles? *J. Exp. Zool. A Ecol. Integr. Physiol.* **329**, 215-221. doi:10.1002/jez.2153
- Sinervo, B., Mendez-de-la-Cruz, F., Miles, D. B., Heulin, B., Bastiaans, E., Cruz, M. V. S., Lara-Resendiz, R., Martinez-Mendez, N., Calderon-Espinosa, M. L., Meza-Lazaro, R. N. et al. (2010). Erosion of lizard diversity by climate change and altered thermal niches. *Science* **328**, 894-899. doi:10.1126/science.1184695
- Sinervo, B., Miles, D. B., Wu, Y., Mendez de la Cruz, F. R., Kirchoff, S. Qi, Y. (2018). Climate change, thermal niches, extinction risk and maternal-effect rescue of toad-headed lizards, *Phrynocephalus*, in thermal extremes of the Arabian Peninsula to the Tibetan Plateau. *Integr. Zool.* **13**, 450-470. doi:10.1111/1749-4877.12315
- Smith, C., Telemeco, R. S., Angilletta, M. J. and VandenBrooks, J. M. (2015). Oxygen supply limits the heat tolerance of lizards embryos. *Biol. Lett.* **11**, 20150113. doi:10.1098/rsbl.2015.0113
- Solaini, G., Baracca, A., Lenaz, G. and Sgarbi, G. (2010). Hypoxia and mitochondrial oxidative metabolism. *Biochim. Biophys. Acta* **1797**, 1171-1177. doi:10.1016/j.bbabi.2010.02.011
- Speakman, J. R. and Garratt, M. (2014). Oxidative stress as a cost of reproduction: Beyond the simplistic trade-off model. *BioEssays* **36**, 93-106. doi:10.1002/bies.201300108
- Speybroeck, J., Beukema, W., Bok, B. and Van Der Voort, J. (2016). *Field Guide to the Amphibians and Reptiles of Britain and Europe*. New York: Bloomsbury Publishing.
- Stahlschmidt, Z. R., Lourda, O., Lourioux, S., Butler, M. W., Davis, J. R., Salin, K., Voituron, Y. and DeNardo, D. F. (2013). Morphological and physiological changes during reproduction and their relationships to reproductive performance in a capital breeder. *Physiol. Biochem. Zool.* **86**, 398-409. doi:10.1086/670918
- Stier, A., Dupoué, A., Picard, D., Angelier, F., Brischoux, F. and Lourda, O. (2017). Oxidative stress in a capital breeder (*Vipera aspis*) facing predation and water constraints. *J. Exp. Biol.* **220**, 1792-1796. doi:10.1242/jeb.156752
- Storz, J. F., Scott, G. R. and Cheviron, Z. A. (2010). Phenotypic plasticity and genetic adaptation to high-altitude hypoxia in vertebrates. *J. Exp. Biol.* **213**, 4125-4136. doi:10.1242/jeb.048181
- Strijbosch, H., Bonnemayer, J. and Dietvorst, P. (1980). The northernmost population of *Podarcis muralis* (Lacertilia, Lacertidae). *Amphib-Reptilia* **1**, 161-172. doi:10.1163/156853880X00150
- Sun, B.-J., Wang, T.-T., Pike, D. A., Liang, L. and Du, W.-G. (2014). Embryonic oxygen enhances learning ability in hatchling lizards. *Front. Zool.* **11**, 21. doi:10.1186/1742-9994-11-21
- Sun, B.-J., Wang, Y., Lu, H.-L. and Du, W.-G. (2018). Anticipatory parental effects in a subtropical lizard in response to experimental warming. *Front. Zool.* **15**, 51. doi:10.1186/s12983-018-0296-3
- Teague, C., Youngblood, J. P., Ragan, K., Angilletta, M. J. and VanderBrooks, J. M. (2017). A positive genetic correlation between hypoxia tolerance and heat tolerance supports a controversial theory of heat stress. *Biol. Lett.* **13**, 20170309. doi:10.1098/rsbl.2017.0309
- Telemeco, R. S., Radder, R. S., Baird, T. A. and Shine, R. (2010). Thermal effects on reptile reproduction: adaptation and phenotypic plasticity in a montane lizard. *Biol. J. Linn. Soc.* **100**, 642-655. doi:10.1111/j.1095-8312.2010.01439.x
- Telemeco, R. S., Gangloff, E. J., Cordero, G. A., Mitchell, T. S., Bodensteiner, B. L., Holden, K. G., Mitchell, S. M., Polich, R. L. and Janzen, F. J. (2016). Reptile embryos lack the opportunity to thermoregulate by moving within the egg. *Am. Nat.* **188**, E13-E27. doi:10.1086/686628
- Thompson, M. B. and Stewart, J. R. (1997). Embryonic metabolism and growth in lizards of the genus *Eumeces*. *Comp. Biochem. Physiol. A Physiol.* **118**, 647-654. doi:10.1016/S0300-9629(97)00081-9
- Tiatragul, S., Kurniawan, A., Kolbe, J. J. and Warner, D. A. (2017). Embryos of non-native anoles are robust to urban thermal environments. *J. Therm. Biol.* **65**, 119-124. doi:10.1016/j.jtherbio.2017.02.021
- Tosini, G. and Avery, R. A. (1996). Pregnancy decreases set point temperatures for behavioural thermoregulation in the wall lizard *Podarcis muralis*. *Herpetol. J.* **6**, 94-96.
- Trochet, A., Dupoué, A., Souchet, J., Bertrand, R., Deluen, M., Murarasu, S., Calvez, O., Martinez-Silvestre, A., Verdaguer-Foz, I., Darnet, E. et al. (2018). Variation of preferred body temperatures along an altitudinal gradient: A multi-species study. *J. Therm. Biol.* **77**, 38-44. doi:10.1016/j.jtherbio.2018.08.002
- Tzedakis, P. C. (2004). The Balkans as prime glacial refugial territory of European temperate trees. In *Balkan Biodiversity: Pattern and Process in the European Hotspot* (ed. H. I. Griffiths, B. Kryštufek and J. M. Reed), pp. 49-68. Dordrecht: Springer Netherlands.
- Uller, T. and While, G. M. (2015). The evolutionary ecology of reproductive investment in lizards. In *Reproductive Biology and Phylogeny of Lizards and Tuatara* (ed. J. L. Rheubert, D. S. Siegel and S. E. Trauth), pp. 425-447. Boca Raton: CRC Press.
- Van Damme, R., Bauwens, D. and Verheyen, R. F. (1991). The thermal dependence of feeding behaviour, food consumption and gut-passage time in the lizard *Lacerta vivipara* Jacquin. *Funct. Ecol.* **5**, 507. doi:10.2307/2389633
- Van Dyke, J. U. and Beaupre, S. J. (2011). Bioenergetic components of reproductive effort in viviparous snakes: costs of vitellogenesis exceed costs of pregnancy. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **160**, 504-515. doi:10.1016/j.cbpa.2011.08.011



- VanGolde, J., Mulder, T. and Blanco, C. E.** (1997). Changes in mean chorioallantoic artery blood flow and heart rate produced by hypoxia in the developing chick embryo. *Pediat. Res.* **42**, 293-298. doi:10.1203/00006450-199709000-00008
- Velotta, J. P., Ivy, C. M., Wolf, C. J., Scott, G. R. and Cheviron, Z. A.** (2018). Maladaptive phenotypic plasticity in cardiac muscle growth is suppressed in high-altitude deer mice. *Evolution* **72**, 2712-2727. doi:10.1111/evo.13626
- Venables, W. N. and Ripley, B. D.** (2002). *Modern Applied Statistics with S*, 4th edn. New York: Springer.
- Vervust, B. and Van Damme, R.** (2009). Marking lizards by heat branding. *Herpetol. Rev.* **40**, 173-174.
- Vleck, C. M. and Hoyt, D. F.** (1991). Metabolism and energetics of reptilian and avian embryos. In *Egg Incubation: its Effects on Embryonic Development in Birds and Reptiles* (ed. D. C. Deeming and M. W. J. Ferguson), pp. 285-306. Cambridge: Cambridge University Press.
- Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., Fromentin, J.-M., Hoegh-Guldberg, O. and Bairlein, F.** (2002). Ecological responses to recent climate change. *Nature* **416**, 389. doi:10.1038/416389a
- Wangensteen, O. D., Rahn, H., Burton, R. R. and Smith, A. H.** (1974). Respiratory gas exchange of high altitude adapted chick embryos. *Respir. Physiol.* **21**, 60-70. doi:10.1016/0034-5687(74)90007-3
- Warburton, S. J., Hastings, D. and Wang, T.** (1995). Responses to chronic hypoxia in embryonic alligators. *J. Exp. Zool.* **273**, 44-50. doi:10.1002/jez.1402730106
- Warner, D. A.** (2014). Fitness consequences of maternal and embryonic responses to environmental variation: using reptiles as models for studies of developmental plasticity. *Integr. Comp. Biol.* **54**, 757-773. doi:10.1093/icb/ictu099
- Warner, D. A. and Lovern, M. B.** (2014). The maternal environment affects offspring viability via an indirect effect of yolk investment on offspring size. *Physiol. Biochem. Zool.* **87**, 276-287. doi:10.1086/674454
- Wearing, O. H., Eme, J., Rhen, T. and Crossley, D. A.** (2016). Phenotypic plasticity in the common snapping turtle (*Chelydra serpentina*): long-term physiological effects of chronic hypoxia during embryonic development. *Am. J. Regul. Integr. Comp. Physiol.* **310**, R176-R184. doi:10.1152/ajpregu.00293.2015
- Wearing, O. H., Conner, J., Nelson, D., Crossley, J. and Crossley, D. A.** (2017). Embryonic hypoxia programmes postprandial cardiovascular function in adult common snapping turtles (*Chelydra serpentina*). *J. Exp. Biol.* **220**, 2589-2597. doi:10.1242/jeb.160549
- Weatherhead, P. J., Sperry, J. H., Carfagno, G. L. F. and Blouin-Demers, G.** (2012). Latitudinal variation in thermal ecology of North American ratsnakes and its implications for the effect of climate warming on snakes. *J. Therm. Biol.* **37**, 273-281. doi:10.1016/j.jtherbio.2011.03.008
- Weathers, W. W. and McGrath, J. J.** (1972). Acclimation to simulated altitude in the lizard *Dipsosaurus dorsalis*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **42**, 263-268. doi:10.1016/0300-9629(72)90107-7
- Webb, A. C., Iverson, J. B., Knapp, C. R., DeNardo, D. F. and French, S. S.** (2018). Energetic investment associated with vitellogenesis induces an oxidative cost of reproduction. *J. Anim. Ecol.* **88**, 461-472. doi:10.1111/1365-2656.12936
- Weibel, E. R., Taylor, C. R. and Hoppeler, H.** (1991). The concept of symmorphosis: a testable hypothesis of structure-function relationship. *Proc. Natl. Acad. Sci. USA* **88**, 10357-10361. doi:10.1073/pnas.88.22.10357
- White, G. M., Williamson, J., Prescott, G., Horvathova, T., Fresnillo, B., Beeton, N. J., Halliwell, B., Michaelides, S. and Uller, T.** (2015). Adaptive responses to cool climate promotes persistence of a non-native lizard. *Proc. R. Soc. B* **282**, 20142638. doi:10.1098/rspb.2014.2638
- Wickham, H.** (2009). *ggplot2: Elegant Graphics for Data Analysis* (ed. H. Wickham). New York: Springer.
- Wilson, A. J. and Nussey, D. H.** (2010). What is individual quality? An evolutionary perspective. *Trends Ecol. Evol.* **25**, 207-214. doi:10.1016/j.tree.2009.10.002
- Žagar, A., Carretero, M. A., Osojnik, N., Sillero, N. and Vrezec, A.** (2015). A place in the sun: interspecific interference affects thermoregulation in coexisting lizards. *Behav. Ecol. Sociobiol.* **69**, 1127-1137. doi:10.1007/s00265-015-1927-8
- Žagar, A., Carretero, M. A., Vrezec, A., Drašler, K., Kaliontzopoulou, A. and Portugal, S.** (2017). Towards a functional understanding of species coexistence: ecomorphological variation in relation to whole-organism performance in two sympatric lizards. *Funct. Ecol.* **31**, 1780-1791. doi:10.1111/1365-2435.12878