



**HAL**  
open science

## In vitro axenic germination and cultivation of mixotrophic Pyroloideae (Ericaceae) and their post-germination ontogenetic development

Tomáš Figura, Edita Tylová, Jan Šoch, Marc-André Selosse, Jan Ponert

### ► To cite this version:

Tomáš Figura, Edita Tylová, Jan Šoch, Marc-André Selosse, Jan Ponert. In vitro axenic germination and cultivation of mixotrophic Pyroloideae (Ericaceae) and their post-germination ontogenetic development. *Annals of Botany*, 2019, 123 (4), pp.625-639. 10.1093/aob/mcy195 . hal-02187965

**HAL Id: hal-02187965**

**<https://hal.sorbonne-universite.fr/hal-02187965>**

Submitted on 18 Jul 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 ***In vitro* axenic germination and cultivation of mixotrophic Pyroloideae (Ericaceae) and**  
2 **their post-germination ontogenetic development**

3

4 Tomáš FIGURA<sup>1,2\*</sup>, Edita TYLOVÁ<sup>1</sup>, Jan ŠOCH<sup>1</sup>, Marc-André SELOSSE<sup>2,3</sup>, Jan  
5 PONERT<sup>1,4</sup>

6

7 1 Department of Experimental Plant Biology, Faculty of Science, Charles University, Viničná  
8 5, 128 44, Prague, Czech Republic

9 2 Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire  
10 naturelle, CNRS, Sorbonne Université, EPHE, CP 39, 57 rue Cuvier, 75005 Paris, France

11 3 Department of Plant Taxonomy and Nature Conservation, University of Gdańsk, Wita  
12 Stwosza 59, 80-308 Gdańsk, Poland

13 4 Prague Botanical Garden, Trojská 800/196, 171 00, Prague, Czech Republic

14

15 \*corresponding author: E-mail: figura.tomas@gmail.com

16

17 RUNNING TITLE: Ontogenetic development of Pyroloideae revealed by *in vitro* culture

18

19 6134 words, 7 figures, 3 suppl. tables and 9 suppl. figures

20 **Abstract**

21 *Background and Aims.* Pyroloids, forest subshrubs of the Ericaceae family, are an important  
22 model for their mixotrophic nutrition, which mixes carbon from photosynthesis and from their  
23 mycorrhizal fungi. They have medical uses but are difficult to cultivate *ex situ*, in particular  
24 their dust seeds contain undifferentiated, few-celled embryos, whose germination is normally

25 fully supported by fungal partners. Their germination and early ontogenesis thus remain  
26 elusive.

27 *Methods.* Optimised *in vitro* cultivation system of five representatives from the subfamily  
28 Pyroloideae, to study the strength of seed dormancy and the effect of different media and  
29 conditions (incl. light, gibberellins and soluble saccharides) on germination. The obtained  
30 plants were analysed for morphological, anatomical and histochemical development.

31 *Key Results.* Thanks to this novel cultivation method, which breaks dormancy and achieved  
32 up to 100% germination, leafy shoots were obtained *in vitro* for representatives of all pyroloid  
33 genera (*Moneses*, *Orthilia*, *Pyrola* and *Chimaphila*). In all cases, the first post-germination  
34 stage is an undifferentiated structure, from which a root meristem later emerges, well before  
35 formation of an adventive shoot.

36 *Conclusions.* This cultivation method can be used for further research or for *ex situ*  
37 conservation of pyroloid species. After strong seed dormancy is broken, the tiny globular  
38 embryo of pyroloids germinates into an intermediary zone, which is anatomically and  
39 functionally convergent with the protocorm of other plants with dust seeds such as orchids.  
40 Like the orchid protocorm, this intermediary zone produces a single meristem: however,  
41 unlike orchids, which produce a shoot meristem, pyroloids first generate a root meristem.

42

## 43 **Keywords**

44 *Pyrola*, *Monotropa*, Ericaceae, seed germination, *in vitro* culture, *Chimaphila*, protocorm,  
45 orchid, convergent evolution, seed dormancy, mixotrophy, *Moneses*

## 46 **Introduction**

47 Mycorrhiza is worldwide symbiosis of most plants (Smith and Read, 2008). Usually,  
48 plants provide photosynthates in exchange for fungal mineral nutrients (van der Heijden *et al.*,  
49 2015). However, mycorrhizal exchange may be less reciprocal, and perhaps exploitative, in  
50 mycoheterotrophic plants that derive carbon resources from their mycorrhizal fungi (Leake,  
51 1994). Mycoheterotrophic plants evolved independently in various plant lineages (Merckx,  
52 2013), and beyond full mycoheterotrophs, which are achlorophyllous, some green species  
53 display a mixed strategy, obtaining carbon from both mycorrhizal fungi and their own  
54 photosynthesis, so-called mixotrophy (Selosse and Roy, 2009). There are two non-exclusive  
55 mixotrophic conditions (Merckx, 2013): some species are mixotrophic at adulthood, while  
56 other species germinate in a mycoheterotrophic way, before switching to autotrophy or  
57 mixotrophy during ontogeny (such as orchids; Dearnaley *et al.*, 2016; Selosse *et al.*, 2016,  
58 Těšitel *et al.* 2018).

59 Little is known about the biology of mycoheterotrophic and mixotrophic plants,  
60 perhaps due to the difficulty of studying them (Merckx, 2013). The only cultivable group of  
61 mixotrophic plants is the orchid family, although many of them have so far escaped  
62 cultivation (Rasmussen, 1995). For this reason, orchids are the most explored mixotrophic  
63 models, but unfortunately it is largely unknown which results obtained for them can be  
64 generalised to other mixotrophic plants. Developing cultivation protocols for other  
65 mycoheterotrophic or mixotrophic plants is therefore urgently needed to understand general  
66 mechanisms connected with mycoheterotrophy.

67 Within Ericaceae, the subfamily Pyroloideae (hereafter pyroloids) comprises initial  
68 mycoheterotrophs that develop, depending on the species, into auto- or mixotrophic adults  
69 (Tedersoo *et al.*, 2007; Matsuda *et al.*, 2012; Hynson *et al.*, 2013a), with the exception of one

70 completely mycoheterotrophic species (Hynson and Bruns, 2009). This similarity with  
71 orchids is an evolutionary convergence and it is therefore interesting to compare adaptations  
72 of both groups to mycoheterotrophy. Pyroloids encompass about 40 subshrub species divided  
73 into four genera, mostly distributed in northern temperate and boreal ecosystems (Takahashi,  
74 1993; Liu *et al.*, 2010), and are of medicinal interest, mostly in Asia (e.g. Ma *et al.*, 2014;  
75 Wang *et al.*, 2014).

76 The seeds of pyroloids consist of oval, central living tissue surrounded by a coat of  
77 dead cells called a testa (Fürth, 1920; Christoph, 1921; Lück, 1941; Pyykkö, 1968; Takahashi,  
78 1993). Despite the fact that the oval living part seems to be anatomically homogeneous, it is  
79 composed of a one-layered endosperm and triploid nutritive tissue of 40 to 50 cells  
80 surrounding a smaller embryo inside (Hofmeister, 1858; Fürth, 1920; Christoph, 1921;  
81 Pyykkö, 1968). Similar seed structure was observed in the related, mycoheterotrophic genus  
82 *Monotropa*, where the embryo consists of only two to three cells (Olson, 1993) to 5-9 cells  
83 (Goebel, 1887) in addition to the endosperm, although minute seeds evolved independently in  
84 this Ericaceae genus (Lallemand *et al.*, 2016). The embryo of pyroloids is larger, with 8 or 16  
85 cells (Goebel 1887) to 30 cells (Christoph 1921). Similarly to *Monotropa*, the living part of  
86 the seed contains limited reserves of lipids and proteins (Fürth, 1920; Christoph, 1921; Lück,  
87 1940). Other mycotrophic plants exhibit a similar seed structure, but, for example, orchids  
88 lack endosperm and the only living part of the seed is undifferentiated globular embryo  
89 (Arditti and Ghani, 2000).

90 Post-germination development of such seeds in pyroloids with undifferentiated  
91 embryo has attracted attention for a long time. First, researchers searched for seedlings in  
92 nature (Irmish, 1855; Velenovský, 1892, 1905; Fürth, 1920), albeit with limited success. They  
93 found only older seedlings, which revealed that a small root-like structure grows into an  
94 extensively branched structure which is formed before the first shoot emerges (Irmish, 1855;

95 Velenovský, 1892; Fürth, 1920). By contrast, orchids, which also start mycoheterotrophic  
96 development from an undifferentiated embryo, first form a specific structure called a  
97 protocorm, from which shoots and roots later develop (Rasmussen, 1995; Dearnaley *et al.*,  
98 2016). This raised three questions: Are the structures observed in germinating pyroloids true  
99 roots? How does this root-like structure develop from a tiny undifferentiated embryo? What is  
100 its relationship to a protocorm?

101         Many attempts have been made to answer these questions, but with only ambiguous  
102 results to the date. Velenovský (1892, 1905) believed this belowground structure is "neither a  
103 root nor a stem" and called it "prokaulom," while Christoph (1921) suggested that this  
104 structure is a root. Later, Lihnell (1942) and Copeland (1947) did the same anatomical  
105 analyses and found a typical root anatomical structure, where the radial vascular bundle is  
106 diarch, rarely triarch (i.e. has two or rarely three xylem strands). The middle layer of the root  
107 called the cortex is made of three to four (or rarely more) cell layers from which the innermost  
108 forms the endodermis (Lihnell, 1942; Copeland, 1947). The outermost root layer, the  
109 epidermis, is one-layered and consists of isodiametric cells. Lateral branches grow from the  
110 outer layer of the stele (Lihnell, 1942).

111         The method of burying seed packets at natural sites yields pyroloid seedlings easily,  
112 but the anatomical structure of such seedlings has not been studied in detail. These root-like  
113 structures, called "root-like structures" (Hashimoto *et al.*, 2012; Johansson and Eriksson,  
114 2013), live heterotrophically below ground for months, if not years (Lihnell, 1942; Hynson *et*  
115 *al.*, 2013a; Johansson *et al.*, 2017).

116         Other researchers tried to germinate pyroloid seeds *in vitro* axenically (Christoph,  
117 1921; Lück, 1940; Lihnell, 1942) or symbiotically (Fürth, 1920; Lück, 1941; Lihnell, 1942).  
118 Germination was achieved, although the results were "mostly uneven, not very  
119 consistent"(Lihnell, 1942) and the seedlings usually stopped growing very soon after

120 germination (Christoph, 1921; Francke, 1934; Lück, 1940, 1941), sometimes being only a  
121 "few tenths of a mm long" (Christoph, 1921). Similar tiny seedlings, which ceased growth,  
122 were obtained for *Monotropa* (Francke, 1934). The best results were achieved by Lihnell  
123 (1942) and yielded a few branched root-like seedlings of *Pyrola rotundifolia*.

124         In these studies, observing different pyroloids, a polarised cone-shaped structure  
125 resembling a root with big epidermal cells grows from the tiny globular embryo (Christoph,  
126 1921; Lück, 1940, 1941; Lihnell, 1942). At this stage, the seedling breaks the testa and forms  
127 an approximately 1-mm long roll-shaped structure (Lihnell, 1942), whose central cells  
128 undergo elongation (Lück, 1941). At this point growth usually ceases (Lück, 1940; Lihnell,  
129 1942). Seedlings rarely grow further, producing a rod-shaped stage, where vessels start  
130 differentiation (Lihnell, 1942), and, soon after, a root cap. Then, the seedling becomes darker,  
131 possibly because of the accumulation of tannins (e.g. Holm, 1898; Christoph, 1921; Lück,  
132 1940; Lihnell, 1942). Lihnell (1942) showed that such older seedlings have the structure of a  
133 root and their illumination does not lead to the formation of chlorophyll. Shoots were never  
134 produced *in vitro*.

135         Growth suddenly stopped in all *in vitro* experiments (Christoph, 1921; Lück, 1940,  
136 1941; Lihnell, 1942), which raised the question of the ideal conditions for *in vitro* growth.  
137 Germination usually started four to eight months after sowing (Lück, 1940, 1941; Lihnell,  
138 1942) and it was difficult to keep the cultures moist for such a long period (Lihnell, 1942). It  
139 is also hard to say if a specific experiment was really axenic, as the authors admitted  
140 themselves (for example, Christoph, 1921, discussed whether worms were the reason for the  
141 failure of cultivation), and in symbiotic cultures it was not clear whether the fungus really  
142 formed symbiosis or not (Lihnell, 1942; indeed, sometimes "symbiotic" seedlings grew away  
143 from the fungus). Moreover, *in vitro* cultivation media often contained substances of variable  
144 composition, such as potato extract, yeast extract, malt, peptone, or even humus, soil extracts

145 (Christoph, 1921; Lück, 1941; Lihnell, 1942). Previous results from *in vitro* cultures are  
146 therefore based on a few plants only and do not indicate cultivation conditions that are ideal  
147 for more detailed observation.

148         Possible seed dormancy was also discussed. Christoph (1921) noticed that it is  
149 difficult to soak pyroloid seeds in water, indicating impermeability of the seed coat. Seeds  
150 disinfected with calcium hypochlorite solution showed the best germination after the longest  
151 bleaching time (15 and 30 minutes; Lihnell, 1942). Lihnell (1942) and Lück (1941) suggested  
152 that some "water-soluble substances" have an inhibitory effect on germination, and Harley  
153 (1959) hypothesised that some substances could be removed by soaking in solution. Although  
154 dormancy in minute seeds may seem unexpected, long bleaching of seeds enhances  
155 germination in many orchids (e.g. Burgeff, 1936; Rasmussen, 1992; Rasmussen, 1995)  
156 probably because hypochlorite solutions have high pH and strong oxidative effects on a wide  
157 range of compounds, which could break impermeable seed coats (Arditti, 1967; Rasmussen,  
158 1995; Zeng *et al.*, 2014). In some hardly germinating orchid species, pre-treatment with a  
159 weak H<sub>2</sub>SO<sub>4</sub> solution enhances germination, probably ensuring stronger degradation of seed  
160 coats (e.g. Malmgren 1993; Ponert *et al.* 2013; Malmgren *et Nyström* 2019). However, the  
161 mechanisms are unclear in orchids and strong differences exist between species. Dormancy  
162 clearly requires further study in pyroloid seeds.

163         To summarize, no reliable protocol for *in vitro* germination of pyroloids exists and  
164 their early ontogenetic development remains elusive. We therefore sought to develop an  
165 efficient protocol for *in vitro* culture. We successfully report early steps of the post-  
166 germination development from undifferentiated embryo to leafy plant, in terms of storage  
167 compounds, and morphological and anatomical development, which enabled us to clearly  
168 answer the long-standing abovementioned questions about the nature of root-like structures of

169 pyroloid seedlings, their development from a tiny undifferentiated embryo and their  
170 relationship to a protocorm.

171

## 172 **Materials and methods**

173

### 174 Plant material

175 We selected five European species as representatives of all four genera of the  
176 subfamily Pyroloideae (*Pyrola media*, *Pyrola minor*, *Orthilia secunda*, *Moneses uniflora*,  
177 *Chimaphilla umbellata*; Table S1). To compare with another subfamily of Ericaceae where  
178 minute seeds and mycoheterotrophic germination independently evolved (Freudenstein *et al.*,  
179 2016; Lallemand *et al.*, 2016), we used *Monotropa uniflora* from the Monotropeoideae (Table  
180 S1). Ripe capsules were collected, dried at room temperature, and extracted seeds were stored  
181 in the dark and dry conditions at +4°C (Table S1).

182

### 183 Cultivation media

184 To find a suitable cultivation medium for germination and growth, we tested nine  
185 media originally designed for orchid *in vitro* culture (Table S2). All media 0.7% agar (w/v,  
186 Sigma-Aldrich) and 1 – 3 % sucrose (Table S2). After pH was adjusted to 5.8 using NaOH,  
187 media were autoclaved at 144 kPa, 121°C (Tuttnauer 2540 EK-N) for 20 min and poured into  
188 5 cm plastic Petri dishes, unless otherwise indicated. Medium Knudson C with activated  
189 charcoal (Sigma-Aldrich) was used for all subsequent experiments, unless otherwise  
190 indicated. Activated charcoal is used to improve germination and growth of orchids and slow-  
191 growing tissue cultures generally, perhaps due to its ability to adsorb toxic products of plant

192 metabolism (van Waes, 1985; Thomas, 2008). To test the effect of different soluble  
193 saccharides that could mimic the carbon provided by the fungi in natural situations, sucrose  
194 was excluded or replaced with the monosaccharide 100 mM glucose or the disaccharide  
195 50 mM sucrose or trehalose. To test the effect of gibberellins, 0.01, 0.1 or 1 mg/L of GA<sub>3</sub>  
196 (Sigma-Aldrich) was added to the medium before autoclaving.

197

#### 198 Seed disinfection and sowing

199       Seeds were disinfected in 5 mL syringes and sown as a suspension in sterile deionised  
200 water as described previously (Ponert *et al.*, 2011, 2013), but the application times of  
201 disinfection solutions of H<sub>2</sub>SO<sub>4</sub> and Ca(OCl)<sub>2</sub> were modified in a fully factorial design to find  
202 proper seed treatment. All treatments of disinfection were pre-incubated in 70% ethanol for 5  
203 minutes, washed three times with deionised water (< 0.2 µm.cm<sup>-1</sup>), treated with 2% H<sub>2</sub>SO<sub>4</sub> for  
204 10 minutes or not, treated with Ca(OCl)<sub>2</sub> solution for 5, 10 or 15 minutes, and finally washed  
205 three times with sterile deionised water. Ca(OCl)<sub>2</sub> solution was prepared by dissolving 20 g of  
206 chlorinated lime (Kittfort, Czech Republic) in 100 mL of deionised water, filtering through  
207 filter paper and adding a drop of Tween 20. For all subsequent experiments, we selected the  
208 most efficient treatment: 70% ethanol for 5 minutes, washed three times with deionised water,  
209 2% H<sub>2</sub>SO<sub>4</sub> for 10 minutes, Ca(OCl)<sub>2</sub> solution for 10 minutes, washed three times with sterile  
210 deionised water. Seven Petri dishes sealed with air permeable foil (Parafilm M) were prepared  
211 for each experimental treatment.

212

#### 213 Cultivation conditions

214       Because seeds ripen late in the season, all cultures were incubated in the dark at 4°C  
215 for three months after sowing to simulate winter, and then transferred to the dark at 20°C,

216 except for experiments where the effect of light or cold stratification period was tested (see  
217 below). Cultures were observed every two weeks using a Krüss, MSZ 5400 Stereo Zoom  
218 Microscope (magnification 40x) and an Olympus Provis AX70 microscope for higher  
219 magnification. Germination rate was counted three times after end of cold stratification, at 30,  
220 60 and 90 days. The last count (3<sup>rd</sup>; after 90 days at 20°C) was used to count total germination  
221 rate, because there no further germination was observed. To count the germination rate, seeds  
222 without embryo or with obviously undeveloped embryo were excluded and well-developed  
223 seeds with broken testa were regarded as germinated seeds (Fig. S1). To estimate seedling size  
224 in selected experiments, the total length of all branches of each seedling was measured from  
225 pictures taken with a Nikon D7000 + Micro Nikkor 55/2.8 using ImageJ 1.6.0\_24 software.

226 To study the establishment of shoots in detail, we used approximately 3-5 mm long  
227 seedlings of *P. minor* pre-cultivated on Knudson C medium (Table S2) for 60 days after  
228 transfer to temperature 20°C. Each seedling (n=22) was transplanted to an individual nine cm  
229 Petri dish with BM-1 medium (Table S2). Pictures of plants were taken every 10 days. Plants  
230 with established shoots were continuously collected for anatomical analyses.

231 Plants with growing etiolated shoots were transferred to light (16 h light / 8 h dark)  
232 and cultivated for the next month to produce green leafy shoots. Plants with green shoots were  
233 deflasked, washed with water and potted in a mixture of coarse expanded perlite, fine pumice  
234 gravel, fine pine bark and loamy soil (1:1:1:2) in clay pots. Pots were sealed in polyethylene  
235 bags to keep air humidity high and were kept on a windowsill at 25°C in moderate light.

236

237 Anatomical analysis

238 Plant material (seeds, seedlings of *P. minor*) was fixed in 4% formaldehyde in  
239 phosphate buffer (0.1 M, pH 7.1). Selected samples were embedded in paraplast after

240 dehydration using an ethanol-butanol series (for details, see Soukup and Tylová, 2014).  
241 Sections (10 µm) were prepared using a Leica 2155 microtome and collected on microscopic  
242 slides coated with alum gelatine adhesive. Cryosections were prepared on a Shandon  
243 cryomicrotome. Hand sections were prepared on a Leica hand-microtome. For whole-mount  
244 preparations, samples were gradually equilibrated in 65% glycerol and mounted in NaI-based  
245 clearing solution of high refractive index (Soukup and Tylová 2014). Histochemical tests  
246 involved staining with safranin O (2-h incubation) and Fast Green FCF (2 min). Lipids were  
247 detected with Sudan Red 7B (1 h) according to Brundrett *et al.* (1991). Proteins were stained  
248 with Ponceau 2R in 2% acetic acid (10 min) and Azur II (10 s) according to Gutmann *et al.*  
249 (1996). Detection of starch involved staining with Lugol solution. Observations were made  
250 with an Olympus BX51 microscope (Olympus Corp., Tokyo, Japan) equipped with an Apogee  
251 U4000 digital camera (Apogee Imaging Systems, Inc., Roseville, CA, USA) or with a Zeiss  
252 LSM 880 confocal microscope.

253

#### 254 Endogenous starch HPLC analysis

255 To confirm the presence of starch in seedlings, we characterised the endogenous  
256 saccharide spectrum of selected pyroloids. Six-month-old (including period of cold  
257 stratification) seedlings of *Mone. uniflora*, *O. secunda* and *P. minor* (n=3 for each species)  
258 cultivated on BM-1 medium were collected in liquid nitrogen. Soluble carbohydrates were  
259 extracted following the protocol of Kubeš *et al.* (2014).

260 The pellets left after soluble saccharide extraction were carefully washed with Milli-Q  
261 ultrapure water (sonicated in 1 mL of water for 15 min, centrifuged at 14000 rpm for 15 min,  
262 supernatant removed) and used for starch analysis. Starch was enzymatically degraded by  $\alpha$ -  
263 amylase (Sigma-Aldrich) and amyloglucosidase (Sigma-Aldrich) following the protocol of

264 Steinbachova-Vojtiskova *et al.* (2006) and glucose content was quantified with the HPLC  
265 system described above, except for the use of an IEX Ca form 8 µm column in this case.

266

267 Data analysis

268 Statistical analyses were performed using statistical software R 3.2.3 (R Core Team  
269 2015). The normality of data was tested using the Shapiro-Wilk test (Shapiro and Wilk, 1965)  
270 and homogeneity of variances was tested using the Bartlett test (Bartlett, 1937). Differences  
271 between the measurements were statistically tested with ANOVA, followed by the Tukey–  
272 Kramer test (Kramer, 1956) for data with a normal distribution and the Kruskal–Wallis test  
273 (Kruskal and Wallis, 1952), followed by pairwise comparisons using Wilcoxon’s rank-sum  
274 test for data that did not have normal distribution. To compare the effects of different  
275 disinfectants on seed germination, we used two-way ANOVA, followed by a Tukey–Kramer  
276 test.

277

## 278 **Results**

279

280 Effect of seed disinfection

281 It turned out to be impossible to sow non-disinfected seeds *in vitro* because of  
282 overwhelming contamination (data not shown). The highest germination rate of all tested  
283 species was reached after H<sub>2</sub>SO<sub>4</sub> treatment and the effect of H<sub>2</sub>SO<sub>4</sub> was significant in all  
284 tested species (*P. minor*:  $F_{[1,24]}=193.96$ ;  $p=5.4 \times 10^{-13}$ ; *M. uniflora*:  $F_{[1,40]}=8.23$ ;  $p=0.0065$ ; *O.*  
285 *secunda*:  $F_{[1,23]}=24.32$ ;  $p=5.5 \times 10^{-5}$ ) except for *C. umbellata*;  $F_{[1,28]}=0.06$ ;  $p=0.8$ ; Fig 1). The  
286 effect of Ca(OCl)<sub>2</sub>; was significant in *P. minor* ( $F_{[2,24]}=57.42$ ;  $p=7.1 \times 10^{-10}$ ) and *O. secunda*

287 ( $F_{[1,23]}=3.79$ ;  $p=0.038$ ), but the optimal length of disinfection differed between these taxa.  
288 Longer incubation in  $\text{Ca}(\text{OCl})_2$  strongly promoted germination of *P. minor*, but slightly  
289 inhibited germination of *O. secunda* (Fig. 1). *Mone. uniflora* germination was also higher  
290 after longer incubation in  $\text{Ca}(\text{OCl})_2$ , especially after the pre-treatment with  $\text{H}_2\text{SO}_4$  (Fig. 1B),  
291 but the effect was not significant ( $F_{[2,40]}=2.13$ ;  $p=0.13$ ). The interaction between the effects of  
292  $\text{H}_2\text{SO}_4$  and  $\text{Ca}(\text{OCl})_2$  was significant for *P. minor* ( $F_{[2,24]}=35.97$ ;  $p=6.0 \times 10^{-8}$ ) and for *O.*  
293 *secunda* ( $F_{[1,23]}=4.05$ ;  $p=0.031$ ) only.

294

295 Effect of different media

296 Optimal media differed between species (Fig. 2A-C). Generally, the highest  
297 germination rates were observed on the media BM-1 (with or without activated charcoal), DS  
298 and MoN (Fig 2; Fig. S2). Media 1/4-2, Knudson C and MS allowed slightly lower  
299 germination generally (Fig 2; Fig. S2). Surprisingly, few seedlings of *C. umbellata*  
300 germinated on Knudson C medium ( $\chi^2_{(7)} = 8.45$ ;  $p= 0.29$ ; Fig. S3B). The best medium for  
301 germination of *P. minor* was BM-1 (with or without activated charcoal), with a germination  
302 rate of almost 100% ( $\chi^2_{(8)} = 32.59$ ;  $p= 7.29 \times 10^{-5}$ ; Fig. 2A). For *Mone. uniflora*, the highest  
303 germination rates were achieved on media BM-1 and DS ( $\chi^2_{(5)} = 7.34$ ;  $p= 0.20$ ; Fig. 2B).  
304 *Orthilia secunda* germinated on all media tested, however, after longer incubation the highest  
305 germination rate was achieved on BM-1 medium without activated charcoal ( $F_{[7,41]}=3.21$ ;  
306  $p=0.0083$ ; Fig. 2C).

307 Media also differed in their suitability for seedling growth in a more or less similar  
308 way (Fig. 2D-F). We were unable to compare the effects of the different media on *C.*  
309 *umbellata* growth because few seedlings developed on medium Knudson C. The biggest  
310 seedlings of *P. minor* developed on medium BM-1 (with or without activated charcoal) and

311 on DS medium, although the differences between treatments were not significant  
312 ( $F_{[7,16]}=0.984$ ;  $p=0.48$ ; Fig. 2D). The largest seedlings of *Mone. uniflora* developed on media  
313 DS and BM-1 without activated charcoal ( $F_{[6,15]}=21.76$ ;  $p=1.32 \times 10^{-6}$ ; Fig. 2E). The largest  
314 seedlings of *O. secunda* developed on MoN medium followed by BM-1 medium with  
315 activated charcoal ( $F_{[7,17]}=14.54$ ;  $p=1.19 \times 10^{-5}$ ; Fig. 2F).

316

### 317 Effect of light

318 The germination of both tested species (*P. minor* and *O. secunda*) was inhibited by  
319 light (16/8 h photoperiod; Fig. S4A,B). For *O. secunda*, this difference was significant (Fig.  
320 S4B;  $\chi^2_{(1)} = 5.4$ ;  $p=0.02$  in 2<sup>nd</sup> month; only one plate in the 3<sup>rd</sup> month because light cultures  
321 became contaminated). For *P. minor*, a similarly small proportion of seeds germinated after  
322 the 1<sup>st</sup> month in both treatments and further germination was only slightly lower on light  
323 (differences not significant;  $F_{[1,8]}=0.058$ ;  $p=0.82$ ; Fig. S4A). However, after another 6 months  
324 of cultivation, seedlings on light did not grow, while the seedlings in the dark grew, produced  
325 many branches and even some buds and first shoots (data not shown). Interestingly, seedlings  
326 cultivated on light displayed a reddish colour (Fig. 3B).

327

### 328 Effect of gibberellic acid and selected soluble saccharides

329 We found little effect of GA<sub>3</sub> on germination of the tested species, namely *P. minor*  
330 (Fig. S4C,D) and *O. secunda* (Fig. S4E,F). For *O. secunda*, there was no difference in  
331 germination rate between GA<sub>3</sub> concentrations on both media tested (Fig. S4E,F), except after  
332 the 1<sup>st</sup> month on Knudson C (higher germination on the highest GA<sub>3</sub> concentration;  
333  $F_{[3,15]}=7.94$ ;  $p=0.002$ ; Fig. S4F). For *P. minor*, there was a weak stimulatory effect after the 1<sup>st</sup>

334 month on BM-1 medium at the highest (1 mg/L) GA<sub>3</sub> concentration ( $\chi^2_{(3)} = 4.719$ ;  $p = 0.002$ ;  
335 Fig. S4C).

336 Tested saccharides stimulated germination of seeds of both tested species. *Pyrola*  
337 *minor* was stimulated by all saccharides, with sucrose being the best ( $\chi^2_{(3)} = 20.57$ ,  $p =$   
338  $0.0001$ , Fig. 4A), and the seeds cultivated without soluble saccharides germinated very rarely.  
339 *O. secunda* reached higher germination rates only on sucrose ( $\chi^2_{(3)} = 16.34$ ,  $p = 0.001$ , Fig.  
340 4B), while glucose and trehalose hardly enhanced germination. Seedling size was similarly  
341 affected by saccharides, with all saccharides acting on *P. minor* (Fig. S5A, Fig. S6A,  
342  $F_{[3,14]} = 27.11$ ;  $p = 4.32 \times 10^{-6}$ ) and sucrose only on *O. secunda* (Fig. S5B, Fig. S6B;  $F_{[3,16]} = 9.57$ ;  
343  $p = 0.00074$ ). After three additional months of cultivation, seedlings from all saccharide-  
344 supplemented treatments produced at least a few long roots, while those from saccharide-free  
345 cultures did not, indicating that trehalose and glucose were also partially utilised. Moreover,  
346 after cultivation for one year, the individual seedlings on media with glucose and trehalose  
347 showed significant growth (Fig. S7).

348

#### 349 Ontogenesis of seedlings

350 All species tested produced elongated seedlings of similar morphology (Fig. 3),  
351 irrespective of the medium. We used *P. minor* as a model species to analyse seedling  
352 development in detail. Like other pyroloids, *P. minor* has a minute seed with few-celled  
353 embryo surrounded by a cellular endosperm enclosed in single-layered transparent testa with  
354 pitted cell walls (Fig. 5A). Histochemical test indicates storage of lipids (Fig. S8C,E) and  
355 proteins (Fig. S8B,D) but not starch (Fig. S8A) in mature seeds. When germinated aseptically  
356 on media supplemented with soluble carbohydrate, germination starts with enlargement of the  
357 embryo at the micropylar pole (Fig. 5B) and with the formation of an apical meristematic area

358 (in seedlings < 0.5 mm long; Fig. 5F) that later passes into root meristem (Fig. 5G). The  
359 transition from embryo into growing root is fast, but gradual. Less organised tissue is  
360 produced during a short initial phase to form an “intermediary zone” (Fig. 5C-H) that exhibits  
361 a heterogeneous surface cell pattern (Fig. 5D) and differentiates vascular tissues in its central  
362 part (Fig. 5G,H). In seedlings above 0.5 mm long, typical root organisation is achieved within  
363 the growing apex (Fig. 5G). The root produced includes radial vascular bundle devoid of pith  
364 (mostly diarch), exarch xylem with protoxylem close to the pericycle, suberised endodermis,  
365 and a root cap (Fig. 5I,J, Fig. S8I,L).

366         Seedling establishment on sucrose-amended media is accompanied by gradual  
367 disappearance of lipids (Fig. S8C,F,J) and accumulation of starch (Fig. S8F,H,K). Starch  
368 grains are located in cortical cells (Fig. S8H-I,K-L). High starch content was confirmed by  
369 HPLC analyses (Table S3.). Conversely, germination on media lacking carbohydrates  
370 terminated in the very initial phase, with very limited enlargement of embryo even after 11  
371 months of cultivation (Fig. S8G).

372         The emerging root started to branch very early and produced an extensive root system  
373 (Fig. 6A), which initiates the first shoot bud much later (Fig. 6B). To observe the timing and  
374 positioning of the emergence of the first bud, we transferred seedlings individually (n=16) to  
375 new Petri dishes soon after germination on sucrose-amended BM-1 media. Seedlings  
376 produced their first buds at different times, starting 35 days after transplantation, and 7 out of  
377 16 plants did not sprout after six months. The first shoot always sprouted a few mm away  
378 from the original position of the embryo, often at the nearest root branching (8 out of 9 plants;  
379 Fig. 6C-E). Very often additional buds appeared approximately one to two months after the  
380 first one, emerging along the root axis at sites of branching (Fig. 6D). One individual  
381 displayed 7 sprouts after six months. The buds grew quickly into leafy shoots which became  
382 green when transferred to the light (Fig. 6F,G). Seedlings left in a small Petri dish with other

383 seedlings grew much slower and did not form buds till the end of the experiment. We also  
384 successfully germinated branching *Mono. uniflora* (n=3; Fig. S9A on BM-1 with activated  
385 charcoal; Fig. S3A) from which one plant produced multiple buds within a dense nest-like  
386 root cluster after one year of cultivation (Fig. S9B).

387 In summary, germination was successful to the stage of green leaves in *P. minor* (Fig.  
388 6F-G) and also in *Ch. umbellata*, *O. secunda*, *Mone. uniflora* and *P. media* (not shown). The  
389 seedlings of *P. minor* were successfully transferred to *ex-vitro* conditions, and cultivated for  
390 the next six months (not shown).

391

## 392 **Discussion**

393

394 We developed an efficient novel protocol for *in vitro* axenic sowing of pyroloids,  
395 which enable us, for the first time, to successfully germinate representatives of all pyroloid  
396 genera and a related mycoheterotroph *Monotropia uniflora*. This protocol allowed us to  
397 overcome problems with seed germination and seedling grow cessation that were previously  
398 reported. Germination is probably driven mostly by physical dormancy because gibberellins  
399 had little effect on germination, while intensive pre-treatment (chemical scarification) of  
400 seeds greatly improved germination. This new, reliable protocol allowed us to produce  
401 seedlings in great number and to describe ontogenesis clearly, from early germination to leafy  
402 plantlet. In all cases, a tiny undifferentiated embryo produces an intermediary zone, which  
403 subsequently establishes a first root. Adventitious shoots grow later from this and secondary  
404 roots (Fig. 7, on the right). Our results suggest convergence in mixotrophic plants with dust-  
405 like seeds in regressive evolution leading to a single meristem that builds mycorrhizal tissue  
406 at germination.

407

408 **Germination conditions**

409           Gibberellic acid (GA<sub>3</sub>), which greatly stimulates germination of many plant species  
410 (Shu *et al.*, 2016), had little effect on pyrolloid germination. A strikingly similar situation  
411 occurs in orchid minute seeds, where gibberellins usually have little effect on seed  
412 germination (Arditti, 1967; Rasmussen, 1995), despite occasional reports of inhibition (Van  
413 Waes and Debergh, 1986) or stimulation (Pedroza-Manrique *et al.*, 2005; Pierce and  
414 Cerabolini, 2011). Low sensitivity to gibberellins may be a general feature of dust-like seeds.  
415 The situation in orchids is sometimes explained by the absence of endosperm (Arditti and  
416 Ghani, 2000; Yeung, 2017), because the stimulation of germination by gibberellins is mostly  
417 connected to the induction of the expression of genes encoding enzymes hydrolysing the  
418 endosperm (Groot and Karssen, 1987; Groot *et al.*, 1988; Schuurink *et al.*, 1992; Leubner-  
419 Metzger *et al.*, 1996). Yet, since pyrolloid seeds have a one-layered endosperm (Hofmeister,  
420 1858; Fürth, 1920; Christoph, 1921; Pyykkö, 1968), the low sensitivity to gibberellins of dust-  
421 like seeds may not be associated with the absence of endosperm but with other characteristics  
422 yet to be clarified.

423           The strong inhibitory effect of light on germination could serve as a protection against  
424 germination on the soil surface, where mycorrhizal fungi may not be present. A similar  
425 mechanism of light inhibition is well known in many terrestrial species of orchids (Arditti,  
426 1967, 2008; Rasmussen, 1995). We also found a strong effect of disinfection on seed  
427 germination. More intensive seed disinfection stimulated germination and longer incubation  
428 in calcium hypochlorite solution after sulphuric acid treatment was often most effective.  
429 Similarly, Lihnell (1942) achieved the highest germination rate after the longest incubation in  
430 calcium hypochlorite solution, which he was unable to explain. Since these solutions also act  
431 corrosively to destroy the impermeable coats of dust-like seeds (e.g. Arditti, 1967;

432 Rasmussen, 1995; Zeng *et al.*, 2014), we propose that strong physical dormancy due to the  
433 testa explains the long pre-germination period and the low germination rates observed *in situ*  
434 (Hashimoto *et al.*, 2012; Johansson and Eriksson, 2013; Hynson *et al.*, 2013b; Johansson *et*  
435 *al.*, 2017). After proper chemical scarification, we reached 100% and 99% germination for *P.*  
436 *minor* and *O. secunda* seeds, respectively, which is much higher than observed *in situ*  
437 (Hashimoto *et al.* 2012; Hynson *et al.*, 2013b) or *in vitro* (Lück, 1940; Lihnell, 1942) in  
438 different species of the same genera. It is hard to discuss possible actions of disinfection  
439 agent's on seeds in detail, because chemical composition of pyroloid seeds is unknown. In  
440 orchids, seed coats seems to consist of lignin, lipids and other uncharacterized compounds  
441 (Barsberg *et al.*, 2013; 2018). Ethanol is a good solvent for many lipids and wax compounds  
442 and it usually makes seeds less hydrophobic and more accessible for subsequent treatment  
443 with aqueous solutions. Lignin is highly sensitive to oxidative effect of hypochlorite  
444 solutions, so we could expect disintegrating of seed coat during hypochlorite treatment. The  
445 effect of H<sub>2</sub>SO<sub>4</sub> is hard to guess, but we could expect existence of some other compound  
446 sensitive to acid hydrolysis, but not the alkaline one.

447 *Chimaphila umbellata* and *Mone. uniflora* germinated at significantly lower rates  
448 comparing with *P. minor* and *O. secunda* in our experiments. Since these two genera form a  
449 clade separate from that encompassing *Pyrola* and *Orthilia* (Freudenstein *et al.*, 2016;  
450 Lallemand *et al.*, 2016), some additional dormancy mechanism may occur in this clade, which  
451 remains to be identified. In soils, dormancy, added to photoinhibition, may increase the ability  
452 of the plant to wait for suitable fungi required for germination.

453 Our *in vitro* methods provided seedlings in great number and shed light on the early  
454 development of pyroloids, with replicates for dissection to an extent that was never reached  
455 before, allowing a clear discussion of early ontogenesis of pyroloids – assuming, of course,  
456 that our artificial, non-symbiotic conditions do not alter ontogeny, as supported below.

## 457 **Cultivation protocol**

458 Based on our results, we propose a cultivation protocol that would allow to germinate all  
459 pyroloids generally. This procedure uniquely combines different techniques used in orchid  
460 culture previously to satisfy needs of pyroloids. Seed disinfection implies: 70% ethanol for 5  
461 minutes, washed out three-times with a deionised water, then 2% H<sub>2</sub>SO<sub>4</sub> for 10 minutes,  
462 followed by Ca(OCl)<sub>2</sub> solution for 10 minutes, washed three times with sterile deionised  
463 water. The best cultivation medium is BM-1, but DS and MoN work also well. For  
464 cultivation conditions, we recommend 4°C for 3 months after sowing, and subsequently  
465 20°C, in dark. Seedlings can be transferred to the new medium when they are approximately 5  
466 mm long. When shoots will start to grow, seedlings should be transferred to the light and, at  
467 that time, it is better to put them separately into 100 ml cultivation jars.

468

## 469 **Development of pyroloid seedlings**

470 The dust seed structure with an undifferentiated globular embryo is typical for many  
471 mycoheterotrophic plants (Leake, 1994) and plants with mycoheterotrophic early  
472 development (Dearnaley *et al.*, 2016). Such dust seeds have limited reserves, the storage  
473 compounds of which are mostly unknown. In pyroloids, previous studies found storage lipids  
474 and proteins in mature seeds (Fürth 1920; Christoph 1921; Lück 1940). Similarly, ripe seeds  
475 of orchids usually contain lipids and proteins (Harrison 1977; Manning and Van Staden 1987;  
476 Rasmussen 1990; Richardson *et al.* 1992; Yam *et al.* 2002; Li *et al.* 2016), and few orchid  
477 species display starch in mature embryos (Tian and Wang 1985; Guo and Xu 1990; Yeung  
478 and Law 1992). Our histochemical tests confirmed storage lipids and proteins, but absence of  
479 starch in *P. minor* seeds, thus further amplifying the convergence with orchids.

480           During germination, lipids and storage proteins were utilised and seedlings shifted to  
481 use starch as the main storage compound, as confirmed by histochemistry and HPLC. This  
482 switch to starch was already observed in several pyroloids (Christoph, 1921; Lück, 1940;  
483 Lihnell, 1942) and is common among orchids (Manning and Van Staden, 1987; Rasmussen,  
484 1990; Richardson *et al.*, 1992). The transition from lipid and protein reserves in seeds to  
485 starch in seedlings therefore also convergently characterizes mycoheterotrophic germination.

486           Mycoheterotrophic pyroloid seedlings grow *in vitro* as branching roots for a long time  
487 before the first green shoot emerges. Such non-green seedlings grow below ground in nature  
488 very slowly, for an even longer time (Hashimoto *et al.*, 2012; Johansson and Eriksson, 2013;  
489 Hynson *et al.*, 2013b; Johansson *et al.*, 2017). Seedling growth was also promoted by more  
490 complex cultivation media (e.g. BM-1, compared to Knudson C or MS), which could indicate  
491 dependency on some other organic compounds provided in nature by fungus. Complex media  
492 are beneficial also for many mixotrophic orchids (Rasmussen, 1995; Arditti, 2008).

493           Post-germination development differs from orchids despite some parallels. Orchid  
494 germination forms a protocorm whose enlargement establishes a shoot meristem on its  
495 anterior pole (Burgeff, 1936; Leroux *et al.*, 1997; Yeung, 2017). In pyroloids, germination  
496 starts with enlargement of the embryo at the suspensor pole, which develops into a small  
497 elongated structure growing apically-the ‘cone stadium’ of Lück (1941) and Lihnell (1942).  
498 Soon after, the apical region establishes a root meristem and grows into a typical root.  
499 Although the transition is pretty fast, a short region (app. 0.5 mm) close to the former embryo  
500 exhibits a slightly different internal structure (Fig. 7), which is why we call this part the  
501 “intermediary zone” (see discussion below).

502           The first root grows and establishes lateral roots to develop an extensively branched  
503 root system as also observed *in situ* (Irmish, 1855; Velenovsky, 1892, 1905; Bobrov, 2009;  
504 Hashimoto *et al.*, 2012). For a long time, it was unclear whether these structures are true roots

505 (see Goebel, 1900; Lihnell, 1942). Velenovský (1892, 1905) suggested these are “neither root  
506 nor stem”. We clearly show a typical root organisation (incl. root cap, endogenous origin of  
507 lateral roots and radial vascular bundle; Fig. 5 H,I,J), as concluded in earlier anatomical  
508 studies (Lihnell, 1942; Copeland, 1947). The abovementioned discussions were probably  
509 caused by insufficient anatomical examination and confusion between the intermediary zone  
510 and the true roots. As in the observations of Velenovský (1905) and Lihnell (1942) and adult  
511 roots (Hynson, 2009), the roots observed had no root hairs. Our data clearly show that the first  
512 shoot sprouts from a small section between the original position of embryo and the first root  
513 branching site. Shortly after, other adventitious shoots usually emerge from other places of the  
514 root system. Such root sprouting also exists in adult pyroloids in natural conditions  
515 (Copeland, 1947; Klimešová, 2007).

516

### 517 **Is the intermediary zone a protocorm?**

518 Typical plant embryos develop from both radicle (basal pole establishing the primary  
519 root) and plumule (apical pole establishing the primary shoot) poles, while in pyroloids or  
520 orchids one embryo pole does not grow: respectively, the primary shoot or the primary root is  
521 completely missing (Fig. 7). Other mycoheterotrophic species with dust seeds that have been  
522 investigated to the date, *Afrothismia hydra* (Burmanniaceae; Imhof and Sainge, 2008) and the  
523 genus *Voyria* (Gentianaceae; Imhof, 2010; Imhof *et al.*, 2013), also germinate by a single,  
524 basal (radicle) embryo pole. This independent loss of one meristematic embryo pole in plants  
525 with dust-like seeds indicates some general selection pressure in mycoheterotrophic  
526 germinations perhaps for several non-exclusive reasons: (i) the limited resources available  
527 cannot support two growth sites (Imhof and Sainge, 2008) and /or (2) the absence of  
528 requirement for shoots and roots in mycoheterotrophy, since a single organ for interaction  
529 with the fungus is sufficient, and/or (3) the later mycorrhizal organ makes one embryo pole

530 unable to differentiate into a meristem. We therefore propose that transformation of one  
531 embryo pole into mycorrhizal tissue is a necessary evolutionary step enabling reduction of  
532 seed reserves to a minimum of the production of a true dust-like seed structure.

533           It could be argued that plants cultivated *in vitro* exhibit different development from  
534 plants in nature. However, the seedlings observed by us fully fit these observed *in situ*  
535 (Irmish, 1855; Velenovsky, 1892, 1905; Hashimoto *et al.*, 2012; Johansson and Eriksson,  
536 2013; Hynson *et al.*, 2013a). In particular, the drawings and photographs *in vitro* by Christoph  
537 (1921), Lück (1940, 1941) and Lihnell (1942) reveal similar seedlings.

538           How to classify the structure that develops between the undifferentiated globular  
539 embryo and the typical root? It starts with polarised growth at suspensor pole of the embryo  
540 and continues with cell divisions in an emerging meristematic area at its apical part. Later, it  
541 becomes thick, with extremely heterogeneous large epidermal cells and a broad central area of  
542 vascular tissues, but without a typical root structure. We therefore called it here the  
543 intermediary zone. In previous studies, it was named a “root-like structure” (Hashimoto *et al.*,  
544 2012), a “procaulom” (Velenovsky, 1892, 1905) or a “protosoma” Bobrov (2004, 2009,  
545 2014). We believe that previous studies did not explore the anatomical structure in detail (an  
546 analysis now allowed by the number of seedlings provided by our cultivation methods) and  
547 therefore did not distinguish between the intermediary zone and the true roots. Moreover,  
548 previous studies analysed seedlings that were already much larger.

549           Orchids also display an intermediary transition stage between the globular embryo and  
550 the first shoot meristem that is larger than the pyrroid intermediary zone and is called a  
551 protocorm (Rasmussen, 1995; Yeung, 2017), because it precedes formation of the typical  
552 plant cormus (i.e. the first shoot or root). Since the intermediary zone of pyrroids also  
553 precedes the cormus (here, the first root), we also suggest use of the term protocorm for the  
554 intermediary zone of pyrroids. The term protocorm was first used for post-embryonic stages

555 of clubmosses (Treub, 1890), which also form a mycoheterotrophic structure appearing before  
556 formation of the first shoot and root. This term was later transferred to orchids (Bernard,  
557 1909) and it is thus historically not orchid-limited, but really naming a convergently evolved  
558 structure. Recently, protocorm was even used in the obligately parasitic Rafflesiaceae, for an  
559 endophytic structure developing from proembryonic endophytic tissue before the formation of  
560 the flower-bearing shoot (Nikolov *et al.*, 2014; Nikolov and Davis, 2017). Interestingly,  
561 Harley (1959) also suggested the term protocorm for seedlings of *Monotropa*, which are  
562 closely related to pyroloids. The pyroloid protocorm is significantly smaller and grows on the  
563 opposite embryo pole compared to orchids, but this transitory structure precedes the typical  
564 cormus in both mycoheterotrophic developments. Investigations of the early ontogenetic  
565 development of other unrelated mycoheterotrophic plants with dust seeds (e.g Burmanniaceae,  
566 Gentianaceae, Triuridaceae; Eriksson and Kainulainen, 2011) are pending. Scarce studies  
567 have revealed that *Afrothismia hydra* (Burmanniaceae; Imhof and Sainge, 2008) and *Voyria*  
568 spp. (Gentianaceae; Imhof, 2010, 2013) produce a root as the first organ, but nothing is  
569 known about initiation of the root from the globular embryo in these plants.

570

## 571 **Conclusion**

572 We used unique combination of cultivation techniques which, for the first time  
573 allowed us to develop an efficient cultivation protocol to germinate and grow leafy plants of  
574 representatives from all pyroloid genera. Our unique methods for pyroloid cultivation may be  
575 used for conservation purposes and also for physiological investigations, since pyroloid  
576 mixotrophic nutrition at adulthood is currently of considerable interest (e.g., Lallemand *et al.*,  
577 2017). The cultures obtained allow fine analyses of germination, with the transition from lipid  
578 and protein storage in seeds to starch accumulation, based on saccharides from the  
579 environment. Seedlings form first roots, before adventitious sprouting. The first root emerges

580 from an intermediary zone (Fig. 7), which we suggest should be called a protocorm, due to its  
581 functional and developmental similarity with protocorms in other plants. Our data further  
582 support the many convergent traits in mycoheterotrophic germination and early development  
583 (especially the existence of a single meristem) in plants with dust seeds.

584

## 585 **Funding**

586 This work was supported by grant of the Czech Ministry of Education, Youth and  
587 Sports [grant no. LO1417] and by Grant Agency of Charles University [grant no. 468318].  
588 Confocal microscopy was performed in the Laboratory of Confocal and Fluorescence  
589 Microscopy co-financed by the European Regional Development Fund and the state budget of  
590 the Czech Republic [grant no. CZ.1.05/4.1.00/16.0347, CZ.2.16/3.1.00/21515].

591

## 592 **Acknowledgements**

593 We thank to Montreal Botanical Garden for sending us seeds of *C. umbellata* and  
594 *Mono. uniflora*, Yvetta Šefrnová and Marek Šírl for help with anatomical study, Radim Hédli  
595 and Jitka Klimešová for help with the old literature and Karel Fajmon and Michal Štefánek  
596 for help with sampling localities. We also thank editor Margreth Sauter and two anonymous  
597 reviewers for useful comments.

598

## 599 **Literature**

600 **Arditti J. 1967.** Factors Affecting the Germination of Orchid Seeds. *The Botanical Review*

601 **33**: 1–97.

602 **Arditti J. 2008.** *Micropropagation of orchids. Volume 1, 2nd ed.* John Willey and Sons.

603 **Arditti J, Ghani A. 2000.** Numerical and physical properties of orchid seeds and their  
604 biological implications. *New Phytologist* **145**: 367–421.

605 **Bartlett M. 1937.** Properties of sufficiency and statistical tests. *Proceedings of the Royal*  
606 *Society of London. Series A, Mathematical and Physical Sciences* **160**: 268–282.

607 **Bernard N. 1909.** L'évolution dans la symbiose. Les Orchideés et leurs champignons  
608 commensaux [In French]. *Annales des Sciences Naturelles Serie (Botanique)* **9**: 1–196.

609 **Barsberg S, Rasmussen HN, Kodahl N. 2013.** Composition of *Cypripedium calceolus*  
610 (Orchidaceae) seeds analyzed by attenuated total reflectance IR spectroscopy: in search of  
611 understanding longevity in the ground. *American journal of botany* **100**: 2066-2073.

612 **Barsberg ST, Lee YI, Rasmussen HN. 2018.** Development of C-lignin with G/S-lignin and  
613 lipids in orchid seed coats—an unexpected diversity exposed by ATR-FT-IR spectroscopy.  
614 *Seed Science Research* **28**: 41-51.

615 **Bobrov J. 2004.** On the early stages of ontogenesis of European Pyrolaceae species [In  
616 Russian]. *Botaniceskij Zhurnal* **89**: 1342–1351.

617 **Bobrov J. 2009.** Grushankovye Rossii [In Russian]. Thesis, Vyatka State University of  
618 Humanities, Kirov, Russia.

619 **Bobrov J. 2014.** Zhiznennaya forma *Moneses uniflora* (Pyroloideae, Ericaceae) [In Russian].  
620 *Vektor nauki Tol'yattinskogo gosudarstvennogo universiteta.* **29**: 21–29.

621 **Brundrett MC, Kendrick B, Peterson CA. 1991.** Efficient Lipid Staining in Plant Material  
622 with Sudan Red 7B or Fluoral Yellow 088 in Polyethylene Glycol-Glycerol. *Biotechnic &*

623 *Histochemistry* **66**: 111–116.

624 **Burgeff H. 1936.** Samenkeimung der Orchideen und Entwicklung ihrer Keimpflanzen [In  
625 German]. *Verlag von Gustav Fisher, Jena, Gemany.*

626 **Byng JW, Chase MW, Christenhusz MJM, et al. 2016.** An update of the Angiosperm  
627 Phylogeny Group classification for the orders and families of flowering plants: APG IV.  
628 *Botanical Journal of the Linnean Society* **181**: 1–20.

629 **Christoph. 1921.** Untersuchungen über mykotrophen Verhältnisse der „Ericales“ und die  
630 Keimung von Pirolaceen [In German]. *Beihefte Botanisches Centralblatt* **38**: 115–157.

631 **Copeland HF. 1947.** Observations on the structure and classification of the Pyroleae.  
632 *Madroño* **9**: 65–102.

633 **Dearnley JDW, Perotto D., Selosse M-A, 2016.** Structure and development of orchid  
634 mycorrhizas, in F. Martin (ed.) *Molecular mycorrhizal symbiosis*, p. 63-86. Springer, Berlin  
635 Heidelberg.

636 **Eriksson O, Kainulainen K. 2011.** The evolutionary ecology of dust seeds. *Perspectives in*  
637 *Plant Ecology, Evolution and Systematics* **13**: 73–87.

638 **Francke H-L. 1934.** Beiträge zur Kenntnis der Mykorrhiza von *Monotropa hypopitys* L [In  
639 German]. *Flora* **129**: 1–52.

640 **Freudenstein J V., Broe MB, Feldenkris ER. 2016.** Phylogenetic relationships at the base  
641 of ericaceae: Implications for vegetative and mycorrhizal evolution. *Taxon* **65**: 794–804.

642 **Fürth P. 1920.** Zur Biologie und Mikrochemie einiger Pirola-Arten. [In German].  
643 *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien, Mathematisch-*  
644 *Naturwissenschaftliche Klasse, Abteilung I*, **129**: 559–587.

645 **Goebel K. 1887.** *Outlines of classification and special morphology of plants.* Oxford:  
646 Clarendon Press.

647 **Goebel K. 1900.** *Organographie der Pflanzen Insbesondere der Archegoniaten und*  
648 *Samenpflanzen. Zweiter Teil. II Heft: Pteridophyten und Samenpflanzen* [In German]. Verlag  
649 von Gustav Fischer, Jena.

650 **Groot SPC, Karssen CMC. 1987.** Gibberellins regulate seed germination in tomato by  
651 endosperm weakening: a study with gibberellin-deficient mutants. *Planta* **171**: 525–531.

652 **Groot S, Kieliszewska-Rokicka B, Vermeer E, Karssen CM. 1988.** Gibberellin-induced  
653 hydrolysis of endosperm cell walls in gibberellin-deficient tomato seeds prior to radicle  
654 protrusion. *Planta* **174**: 500–504.

655 **Guo S, Xu J. 1990.** Studies on the changes of cell ultrastructure in the course of seed  
656 germination of *Bletilla striata* under fungus infection conditions. *Acta Botanica Sinica* **32**:  
657 594–598.

658 **Harley JL. 1959.** The biology of mycorrhiza. *The biology of mycorrhiza.* London: Leonard  
659 Hill.

660 **Harrison CR. 1977.** Ultrastructural and Histochemical changes during the Germination of  
661 *Cattleya aurantiaca* (Orchidaceae). *Botanical Gazette* **138**: 41–45.

662 **Hashimoto Y, Fukukawa S, Kunishi A, et al. 2012.** Mycoheterotrophic germination of  
663 *Pyrola asarifolia* dust seeds reveals convergences with germination in orchids. *New*  
664 *Phytologist* **195**: 620–630.

665 **van der Heijden MGA, Martin FM, Selosse M-A, Sanders IR. 2015.** Mycorrhizal ecology  
666 and evolution: the past, the present, and the future. *New Phytologist* **205**: 1406–1423.

667 **Hofmeister W. 1858.** Neuere Beobachtungen über Embryobildung der Phanerogamen.  
668 *Jahrbücher für Wissenschaftliche Botanik*. **1**: 82–188.

669 **Holm T. 1898.** *Pyrola aphylla*: A Morphological Study. *Botanical Gazette* **25**: 246–254.

670 **Hynson NA, Bruns TD. 2009.** Evidence of a myco-heterotroph in the plant family Ericaceae  
671 that lacks mycorrhizal specificity. *Proceedings of the Royal Society B: Biological Sciences*  
672 **276**: 4053–4059.

673 **Hynson NA, Madsen TP, Selosse M-A, et al. 2013.** The Physiological Ecology of  
674 Mycoheterotrophy In: Merckx VSFT, ed. *Mycoheterotrophy*. New York: Springer New York,  
675 297–342.

676 **Hynson NA, Weiß M, Preiss K, Gebauer G, Treseder KK. 2013.** Fungal host specificity is  
677 not a bottleneck for the germination of Pyroleae species (Ericaceae) in a Bavarian forest.  
678 *Molecular Ecology* **22**: 1473–1481.

679 **Imhof S. 2010.** Are Monocots Particularly Suited to Develop Mycoheterotrophy? In: Seberg,  
680 O, Petersen, G, Barfod, A & Davis, JI, eds. Aarhus university press: Aarhus *Diversity,*  
681 *Phylogeny, and Evolution in the Monocotyledons*: 11–23.

682 **Imhof S, Massicotte H., Melville LH, Peterson R. 2013.** Subterranean morphology and  
683 mycorrhizal structures In: Merckx VSFT, ed. *Mycoheterotrophy: The Biology of Plants Living*  
684 *on Fungi*. New York: Springer, 157–214.

685 **Imhof S, Sainge MN. 2008.** Ontogeny of the mycoheterotrophic species *Afrothismia hydra*  
686 (Burmanniaceae). *Botanical Journal of the Linnean Society* **157**: 31–36.

687 **Irmish T. 1855.** Bemerkungen über einige Pflanzen der deutschen Flora [In German]. *Flora*  
688 **13**: 625–638.

689 **Johansson VA, Bahram M, Tedersoo L, Kõljalg U, Eriksson O. 2017.** Specificity of

690 fungal associations of Pyroleae and *Monotropa hypopitys* during germination and seedling  
691 development. *Molecular Ecology* **26**: 2591–2604.

692 **Johansson VA, Eriksson O. 2013.** Recruitment limitation, germination of dust seeds, and  
693 early development of underground seedlings in six Pyroleae species. *Botany* **91**: 17–24.

694 **Klimešová J. 2007.** Root-sprouting in myco-heterotrophic plants: prepackaged symbioses or  
695 overcoming meristem limitation? *New Phytologist* **173**: 8–10.

696 **Knudson L. 1922.** Nonsymbiotic Germination of Orchid Seeds. *Botanical Gazette* **73**: 1–25.

697 **Kramer C. 1956.** Extension of multiple range tests to group means with unequal numbers of  
698 replications. *Biometrics* **12**: 307–310.

699 **Kruskal WH, Wallis WA. 1952.** Use of ranks in one-criterion variance analysis. *Journal of*  
700 *the American statistical Association.* **47**: 583–621.

701 **Kubeš M, Drážná N, Konrádová H, Lipavská H. 2014.** Robust carbohydrate dynamics  
702 based on sucrose resynthesis in developing Norway spruce somatic embryos at variable sugar  
703 supply. *In Vitro Cellular and Developmental Biology* **50**: 45–57.

704 **Lallemand F, Gaudeul M, Lambourdière J, Matsuda Y, Hashimoto Y, Selosse MA.**  
705 **2016.** The elusive predisposition to mycoheterotrophy in Ericaceae. *New Phytologist* **212**:  
706 314–319.

707 **Leake J. 1994.** Tansley Review No. 69. The biology of myco- heterotrophic (“saprophytic”)  
708 plants. *New Phytologist* **127**: 171–216.

709 **Leroux G, Barabé D, Vieth J. 1997.** Morphogenesis of the protocorm of *Cypripedium*  
710 *acaule* (Orchidaceae). *Plant systematics and evolution* **205**: 53–72.

711 **Leubner-Metzger G, Fründt C, Meins F. 1996.** Effects of gibberellins, darkness and

712 osmotica on endosperm rupture and class I  $\beta$ -1, 3-glucanase induction in tobacco seed  
713 germination. *Planta* **199**: 282–28.

714 **Li Y-Y, Chen X-M, Guo S-X, Lee Y-I. 2016.** Embryology of two mycoheterotrophic orchid  
715 species, *Gastrodia elata* and *Gastrodia nantoensis*: ovule and embryo development. *Botanical*  
716 *Studies* **57**: 1–10.

717 **Lihnell D. 1942.** Keimungsversuche mit Pyrolasamen [In German]. *Symbolae botanicae*  
718 *upsalienses* **6**: 1–37.

719 **Liu Z-W, Zhou J, Liu E-D, Peng H. 2010.** A molecular phylogeny and a new classification  
720 of *Pyrola* (Pyroleae, Ericaceae). *Taxon* **59**: 1690–1700.

721 **Lück R. 1940.** Zur Biologie der heimischen *Pirola*-arten [In German]. *Schriften der*  
722 *Physikalisch-Ökonomischen Gesellschaft zu Königsberg* **71**: 300–334.

723 **Lück R. 1941.** Zur keimung der heimischen *Pirola*-Arten [In German]. *Flora* **135**: 1–5.

724 **Ma W-D, Zou Y-P, Wang P, et al. 2014.** Chimaphilin induces apoptosis in human breast  
725 cancer MCF-7 cells through a ROS-mediated mitochondrial pathway. *Food and Chemical*  
726 *Toxicology* **70**: 1–8.

727 **Malmgren S. 1993.** Asymbiotisk fröförökning i stor skala av *Anacamptis*, *Ophrys*, *Orchis*  
728 och andra orkidéer med runda rotknölar, *Svensk Botanisk Tidskrift* **87**: 221-234.

729 **Malmgren, S., Nyström, H. 2019:** Orchid Propagation. -  
730 <http://www.lidaforsgarden.com/Orchids/engelsk.htm>. Accessed 2.9.2019

731 **Manning J, Van Staden J. 1987.** The development and mobilisation of seed reserves in  
732 some African orchids. *Australian Journal of Botany* **35**: 343–353.

733 **Matsuda Y, Shimizu S, Mori M, Ito S-I, Selosse M-A. 2012.** Seasonal and environmental  
734 changes of mycorrhizal associations and heterotrophy levels in mixotrophic *Pyrola japonica*

735 (Ericaceae) growing under different light environments. *American journal of botany* **99**:  
736 1177–1188.

737 **Merckx VSFT. 2013.** *Mycoheterotrophy: The biology of plants living on fungi*. New York:  
738 Springer.

739 **Murashige T, Skoog F. 1962.** A Revised Medium for Rapid Growth and Bio Assays with  
740 Tobacco Tissue Cultures. *Physiologia Plantarum* **15**: 473–497.

741 **Nikolov LA, Davis CC. 2017.** The big, the bad, and the beautiful: Biology of the world’s  
742 largest flowers. *Journal of Systematics and Evolution* **55**: 516–524.

743 **Nikolov LA, Staedler YM, Manickam S, et al. 2014.** Floral structure and development in  
744 Rafflesiaceae with emphasis on their exceptional gynoecea. *American journal of botany* **101**:  
745 225–243.

746 **Olson AR. 1993.** Patterns of embryo and endosperm formation in *Monotropa hypopitys*  
747 (Monotropaceae) from North America and Western Sweden. *American Journal of Botany* **80**:  
748 839–846.

749 **Pedroza-Manrique J, Fernandez-Lizarazo C, Suarez-Silva A. 2005.** Evaluation of the  
750 Effect of three growth regulators in the germination of *Comparettia falcata* seeds under *In*  
751 *vitro* conditions. *In Vitro Cellular & Developmental Biology. Plant* **41**: 838–843.

752 **Pierce S, Cerabolini BE. 2011.** Asymbiotic germination of the White Mountain Orchid  
753 (*Pseudorchis albida*) from immature seed on media enriched with complex organics or  
754 phytohormones. *Seed Science and Technology* **39**: 199–203.

755 **Ponert J, Figura T, Vosolobě S, Lipavská H, Vohník M, Jersáková J. 2013.** Asymbiotic  
756 germination of mature seeds and protocorm development of *Pseudorchis albida*  
757 (Orchidaceae) are inhibited by Nitrates even at extremely low concentrations. *Canadian*

758 *Journal of Botany* **91**: 662–670.

759 **Ponert J, Vosolsobě S, Kmecová K, Lipavská H. 2011.** European orchid cultivation – from  
760 seed to mature plant. *European journal of environmental sciences* **1**: 95–107.

761 **Pyykkö M. 1968.** Embryological and anatomical studies on Finnish species of the Pyrolaceae.  
762 *Annales Botanici Fennici* **5**: 153–165.

763 **R Core Team. 2015.** R: A Language and Environment for Statistical Computing.

764 **Rasmussen HN. 1990.** Cell differentiation and mycorrhizal infection in *Dactylorhiza majalis*  
765 (Rchb. f.) Hunt & Summerh. (Orchidaceae) during germination *in vitro*. *New Phytologist* **116**:  
766 137–147.

767 **Rasmussen HN. 1992.** Seed dormancy patterns in *Epipactis palustris* (Orchidaceae):  
768 Requirements for germination and establishment of mycorrhiza. *Physiologia Plantarum*, **86**:  
769 161-167.

770 **Rasmussen HN. 1995.** *Terrestrial orchids: from seed to mycotrophic plant*. Cambridge:  
771 Cambridge University Press.

772 **Richardson KA, Peterson RL, Currah RS. 1992.** Seed reserves and early symbiotic  
773 protocorm development of *Platanthera hyperborea* (Orchidaceae). *Canadian Journal of*  
774 *Botany* **70**: 291–300.

775 **Selosse MA, Rousset F. 2011.** The plant-fungal marketplace. *Science* **333**: 828–829.

776 **Selosse MA, Roy M. 2009.** Green plants that feed on fungi: facts and questions about  
777 mixotrophy. *Trends in Plant Science* **14**: 64–70.

778 **Selosse, M-A., Bocayuva, M. F., Kasuya, M. C. M., & Courty, P. E. 2016.** Mixotrophy in  
779 mycorrhizal plants: extracting carbon from mycorrhizal networks. *Molecular Mycorrhizal*

780 Symbiosis, 451-471.

781 **Shapiro SS, Wilk MB. 1965.** An analysis of variance test for normality (complete samples).  
782 *Biometrika* **52**: 591–611.

783 **Shu K, Liu X, Xie Q, He Z. 2016.** Two Faces of One Seed: Hormonal Regulation of  
784 Dormancy and Germination. *Molecular Plant* **9**: 34–45.

785 **Schuurink R, Sedee N, Wang M. 1992.** Dormancy of the barley grain is correlated with  
786 gibberellic acid responsiveness of the isolated aleurone layer. *Plant Physiology* **100**: 1834–  
787 1839.

788 **Smith SE, Read D. 2008.** *Mycorrhizal symbiosis*. London. Academic Press.

789 **Soukup A, Tylová E. 2014.** Essential Methods of Plant Sample Preparation for Light  
790 Microscopy In: Clifton N., ed. *Methods in molecular biology*. Humana Press, Totowa, NJ, 1–  
791 23.

792 **Steinbachová-Vojtíšková L, Tylová E, Soukup A, et al. 2006.** Influence of nutrient supply  
793 on growth, carbohydrate, and nitrogen metabolic relations in *Typha angustifolia*.  
794 *Environmental and Experimental Botany* **57**: 246–257.

795 **Takahashi H. 1993.** Seed Morphology and Its Systematic Implications in Pyroloideae  
796 (Ericaceae). *International Journal of Plant Sciences* **154**: 175–186.

797 **Tedersoo L, Pellet P, Kõljalg U, Selosse MA. 2007.** Parallel evolutionary paths to  
798 mycoheterotrophy in understorey Ericaceae and Orchidaceae: Ecological evidence for  
799 mixotrophy in Pyroloideae. *Oecologia* **151**: 206–217.

800 **Těšitel J, Těšitelová T, Minasiewicz J., Selosse M-A. 2018.** Mixotrophy in land plants : why  
801 to stay green? Trends in Plant Science, *in press*. <https://doi.org/10.1016/j.tplants.2018.05.010>

802 **Thomas, TD, 2008.** The role of activated charcoal in plant tissue culture. *Biotechnology*  
803 *advances*, 26: 618-631.

804 **Tian M, Wang F. 1985.** *In vitro* seed germination and developmental morphology of  
805 seedling in *Cymbidium ensifolium*. *Acta Botanica Sinica* **27**: 455–459.

806 **Treub M. 1890.** Études sur les Lycopodiaceès. *Annales du Jardin Botanique Buitenzorg* [In  
807 French]. *Annales du Jardin Botanique de Buitenzorg* **1**: 23–27.

808 **Velenovský J. 1892.** O biologii a morfologii rodu “ Monesis” [In Czech]. *Rozpravy*  
809 *Královskéčeské společnosti nauk, řada, matematicko-přírodovědná* **11**: 147–159.

810 **Velenovský J. 1905.** Über die Keimpflanzen der Pirolaceen [In German]. *Bulletin*  
811 *international de l’Académie des Science de Bohême*. **14**: 6–12.

812 **Van Waes, JM, 1985.** Effect of activated charcoal on *in vitro* propagation of Western  
813 European orchids. *Acta Horticulturae* 212: 131-138.

814 **Van Waes JM, Debergh PC. 1986.** *In vitro* germination of some Western European orchids.  
815 *Physiologia Plantarum* **67**: 253–261.

816 **Wang D, He F, Lv Z, Li D. 2014.** Phytochemical Composition, Antioxidant Activity and  
817 HPLC Fingerprinting Profiles of Three *Pyrola* Species from Different Regions (J Chamani,  
818 Ed.). *PLoS ONE* **9**: e96329.

819 **Yam TW, Yeung EC, Ye X-L, Zee S-Y, Arditti J. 2002.** Orchid Embryos In: Kull T, Arditti  
820 J, eds. *Orchid Biology: Reviews and Perspectives, VIII*. Dordrecht: Kluwer Academic  
821 Publisher, 287–385.

822 **Yeung EC. 2017.** A perspective on orchid seed and protocorm development. *Botanical*  
823 *Studies* **58**: 1–14.

824 **Yeung EC, Law SK. 1992.** Embryology of *Calypso bulbosa*. II. Embryo development.  
825 *Canadian Journal of Botany* **70**: 461–468.

826 **Zeng S, Zhang Y, Teixeira da Silva JA, Wu K, Zhang J, Duan J. 2014.** Seed biology and  
827 *in vitro* seed germination of *Cypripedium*. *Critical Reviews in Biotechnology* **34**: 358–371.

828 **Zimmer K, Hynson NA, Gebauer G, Allen EB, Allen MF, Read DJ. 2007.** Wide  
829 geographical and ecological distribution of nitrogen and carbon gains from fungi in pyroloids  
830 and monotropoids (Ericaceae) and in orchids. *New Phytologist* **175**: 166–175.

831

## 832 **Figure captions**

833 Fig. 1. Effect of different disinfection treatments on seed germination of (A) *Pyrola minor*,  
834 (B) *Moneses uniflora*, (C) *Chimaphila umbellata*, (D) *Orthilia secunda*. *In vitro* cultivation on  
835 medium Knudson C was evaluated after three months after the cold stratification. Median  
836 values are given (n=7). Different letters indicate significant difference according to the  
837 ANOVA followed by Tukey HSD post-hoc test for multiple comparison.

838

839 Fig. 2. Effect of different media on germination of (A) *Pyrola minor*, (B) *Moneses uniflora*,  
840 (C) *Orthilia secunda* and growth of (D) *Pyrola minor*, (E) *Moneses uniflora*, (F) *Orthilia*  
841 *secunda*. Both germination and total length of seedlings were measured 3 months after cold  
842 stratification. Different letters indicate significant difference. Results of Tukey HSD post hoc  
843 test for multiple comparisons are indicated by capital letters and results of Pairwise Wilcoxon  
844 Rank Sum test by lowercase letters.

845

846 Fig. 3. Representatives of *in vitro* grown seedlings of species used in this study. A) *Pyrola*  
847 *minor* cultivated in dark, B) *Pyrola minor* cultivated on light and C) *Moneses uniflora*, D)  
848 *Orthilia secunda*, E) *Chimaphila umbellata* and F) *Monotropa uniflora* cultivated in dark.  
849 Scale bars: 0,5 mm.

850

851 Fig. 4. Effect of selected soluble saccharides on germination of *Pyrola minor* (A) and *Orthilia*  
852 *secunda* (B). *In vitro* cultivation on Knudson C medium where sucrose was replaced by  
853 saccharide indicated. Different letters indicate significant difference. Results of Tukey HSD  
854 post hoc test for multiple comparisons indicated by capital letters and results of Pairwise  
855 Wilcoxon Rank Sum test by lowercase letters; bars indicate 1<sup>st</sup> and 3<sup>rd</sup> quantiles. Pairwise  
856 comparisons were calculated separately for each month.

857

858 Fig. 5. Germination and early development of *Pyrola minor*. (a) mature seed with globular  
859 embryo. (b) germination starts with enlargement of embryo (arrows indicate the former  
860 embryo). (c) seedling develops further into root through less organized intermediary zone  
861 with very heterogenic surface cell pattern. (d) detail of the surface cell pattern of intermediary  
862 zone. (e) seedling of 1-1.5 mm size with already established root meristem. (f) meristematic  
863 zone is established very early after germination. (g) seedling of less than 1 mm size with  
864 established meristem and differentiated vascular tissues in central part of intermediary zone  
865 close to the former embryo. (h) seedling of *ca.* 3 mm size with already established lateral root  
866 primordium (white arrow) and differentiated vascular tissues (in detail). (i) root structure is  
867 obvious at 1.5 cm from the position of former embryo in 3 cm long seedling. Abbreviations:  
868 en, endodermis; x, protoxylem vessels of radial vascular bundle; c, cortex; rh, rhizodermis. (j)  
869 root tip of 3 cm long seedling. Abbreviations: lrc, lateral root cap; p, protoderm; qc, quiescent  
870 center. a, b, e, j, whole-mounts, DIC; c, whole-mount, confocal image; d, whole-mount, UV;  
871 f-h, paraplasm sections, Safranin O + Fast Green FCF; I, hand section, UV; j, paraplasm  
872 section, Lugol + Orange G. Black arrows indicate position of former embryo; scale bars 50  
873  $\mu\text{m}$ .

874

875 Fig. 6. Transition to shoot formation in *P. minor* seedlings. (a) Nine-months old seedling with  
876 established root system prior the onset of shoot growth. (b) Nine-months old seedling with  
877 emerging shoot (arrow). (c) shoot bud emerges at first root branching site close to the position  
878 of former embryo (r, root; sh, shoot bud; iz, intermediary zone connecting the former embryo;  
879 x, xylem), section stained with Safranin O and Fast Green FCF. (d) later, additional shoot  
880 buds emerge along root axis (arrows). (e) detail of first shoot bud, whole-mount preparation.  
881 (f) fully-established *in vitro* plant. (g) adventitious roots emerging at stem node (sh, shoot).  
882 Scale bars 0.5 cm, except c, e: 200  $\mu\text{m}$ .

883

884 Fig. 7. Comparison of germination and subsequent development of orchids (left) and  
885 pyroloids (right). Embryo is highlighted in red, protocorm in blue, shoots in green and roots in  
886 brown. Abbreviations: a, adventive root; br, branching roots; e, embryo; end, endosperm; h,  
887 hilum; iz, intermediary zone; p, protocorm; ram, root meristem; sh, shoot; sam, shoot  
888 meristem; t, testa; x, xylem. Thin bar is 100 $\mu\text{m}$ , dotted bar 1mm and thick bar 0,5 cm.

889