



HAL
open science

A new chlorite geothermometer for diagenetic to low-grade metamorphic conditions

Franck Bourdelle, Teddy Parra, Christian Chopin, Olivier Beyssac

► To cite this version:

Franck Bourdelle, Teddy Parra, Christian Chopin, Olivier Beyssac. A new chlorite geothermometer for diagenetic to low-grade metamorphic conditions. *Contributions to Mineralogy and Petrology*, 2013, 165 (4), pp.723-735. 10.1007/s00410-012-0832-7 . hal-02270198

HAL Id: hal-02270198

<https://hal.sorbonne-universite.fr/hal-02270198>

Submitted on 23 Aug 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Revision 1**

2 Title: A new chlorite geothermometer for diagenetic to low-grade metamorphic conditions

3

4 Authors: Franck Bourdelle ^{a,b,c}, Teddy Parra ^a, Christian Chopin ^b, Olivier Beyssac ^c

5

6 Mailing addresses of the relevant institutions:

7

8 ^a IFP Energies Nouvelles, 1&4 avenue de Bois Préau, 92852 Rueil-Malmaison cedex, France

9 ^b Ecole normale supérieure — CNRS, Laboratoire de Géologie, 24 rue Lhomond, 75231 Paris
10 cedex 05, France

11 ^c IMPMC, UPMC-CNRS, Case courrier 115, 4 Place Jussieu, 75252 Paris cedex 05, France

12

13 Corresponding authors:

14 1. Franck Bourdelle

15 Present mailing address: IMPMC, UPMC-CNRS, Case courrier 115, 4 Place Jussieu, 75252

16 Paris cedex 05, France

17 E-mail address: franck.bourdelle@gmail.com

18 Phone number: + 33 1 44 27 25 60

19 Fax number: + 33 1 44 27 51 52

20

21 2. Teddy Parra

22 Mailing address: IFP Energies Nouvelles, 1&4 avenue de Bois Préau, 92852 Rueil-

23 Malmaison cedex, France

24 E-mail address: teddy.parra@ifpen.fr

25 Phone number: + 33 1 47 52 73 67

26

27 **Abstract**

28

29 The evolution of chlorite composition with temperature (and pressure) serves as basis to a
30 number of chlorite chemical thermometers, for which the oxidation state of iron has been
31 recognised as a recurrent issue, especially at low temperature (*T*). A new chlorite
32 geothermometer that does not require prior Fe³⁺ knowledge is formulated, calibrated on 161
33 analyses with well constrained *T* data covering a wide range of geological contexts, and tested
34 here for low-*T* chlorites (*T* < 350°C and pressures below 4 kbar). The new solid-solution
35 model used involves six end-member components (the Mg and Fe end-members of ‘Al-free

36 chlorite S', sudoite and amesite) and so accounts for all low- T chlorite compositions; ideal
37 mixing on site is assumed, with an ordered cationic distribution in tetrahedral and octahedral
38 sites.

39 Applied to chlorite analyses from three distinct low- T environments for which independent T
40 data are available (Gulf Coast, Texas; Saint Martin, Lesser Antilles; Toyoha, Hokkaido), the
41 new pure-Fe²⁺ thermometer performs at least as well as the recent models, which require an
42 estimate of Fe³⁺ content. This relief from the ferric iron issue, combined with the simple
43 formulation of the semi-empirical approach, makes the present thermometer a very practical
44 tool, well suited for, e.g., the handling of large analytical datasets – provided it is used in the
45 calibration range ($T < 350^{\circ}\text{C}$, $P < 4$ kbar).

46

47 Keywords: Chlorite, Geothermometry, Diagenesis, Low-grade metamorphism, Solid-solution.

48

49 **Introduction**

50

51 Chlorites are ubiquitous in most diagenetic and metamorphic rocks (Foster 1962;
52 Cathelineau 1988; Laird 1988; De Caritat et al. 1993; Vidal et al. 2001). Their wide
53 compositional variations, through the effect of the three main substitutions Fe²⁺ = Mg,
54 Tschermak Al^{IV}Al^{VI} = Si(Mg, Fe²⁺), and di/trioctahedral 3(Mg, Fe²⁺) = □ + 2Al^{VI} (where □
55 represents an octahedral vacancy), is known to be sensitive to the pressure (P) and
56 temperature (T) conditions of formation, to the bulk composition and to the physico-chemical
57 properties of the environment, like activities in the fluid(s) (e.g. Cathelineau and Nieva 1985;
58 Hillier and Velde 1991; Vidal and Parra 2000). This makes chlorites good indicators of rock
59 history and, for this reason, they are the basis of several empirical or thermodynamic methods
60 to estimate P - T conditions (e.g. Cathelineau 1988; Hillier and Velde 1991; Walshe 1986;
61 Vidal et al. 2001; Inoue et al. 2009).

62 Several studies showed however that an empirical approach, in which the temperature
63 is directly linked to chlorite composition, is often inappropriate (e.g. De Caritat et al. 1993;
64 Essene and Peacor 1995; Bourdelle et al. *in revision*). For example, the empirical calibrations
65 that are based on the T -dependent variation of Al^{IV} content are easily applicable, but do not
66 account for the influence of bulk-rock composition. Thus, this approach has no generality and
67 can only be applied to a restricted range of geological settings (De Caritat et al. 1993), in spite
68 of the many formulations proposed (Cathelineau and Nieva 1985; Kranidiotis and McLean
69 1987; Cathelineau 1988; Jowett 1991; Hillier and Velde 1991; Zang and Fyfe 1995; Xie et al.
70 1997). Thermodynamic methods circumvent this problem, allowing one to account for

71 mineral assemblage and/or physicochemical parameters in addition to temperature and
72 pressure (e.g. Walshe 1986; Vidal and Parra 2000; Vidal et al. 2001, 2005, 2006). However,
73 the thermodynamic approach may be also questioned, considering that the large range of
74 compositional variations in chlorites was not properly handled by thermodynamic modelling.
75 In fact, the inaccuracy of this approach is due in most cases to uncertainties in the P - T data
76 employed for the calibration, to the inaccuracy of the thermodynamic properties of chlorites
77 end-members, to the choice of the mixing model, and to the difficulty in measuring the Fe^{3+}
78 content and in accounting for it in solid-solution models (cf. Grosch et al. 2012).

79 On this basis, Inoue et al. (2009) proposed a new, better suited low- T geothermometer
80 that does not require the knowledge of end-member thermodynamic properties but includes
81 the effect of Fe^{3+} content on temperature estimation, thereby requiring independent Fe^{3+}
82 determination. However, in their comparative study based on high-spatial-resolution analysis,
83 Bourdelle et al. (*in revision*) obtained some unsatisfactory results with this thermometer for
84 low- T conditions, probably due to the difficulty in obtaining reliable values of XFe^{3+} at
85 nanometer scale. Indeed, the estimation of Fe^{3+} content needs either a Mössbauer
86 spectroscopy study (e.g. Beaufort et al. 1992), which involves grinding of the sample and loss
87 of the textural information, or *in situ* XANES analysis (e.g. Vidal et al. 2006) which remains a
88 complex procedure, or a numerical estimation using an iterative multi-equilibrium calculation
89 (Vidal et al. 2006). Thus, the presence of ferric iron is clearly an important issue.

90 The present study was undertaken as a continuation of previous works of Vidal et al.
91 (2006), Inoue et al. (2009) and Bourdelle et al. (*in revision*), with the goal of developing a
92 chlorite geothermometer for diagenetic to low-grade metamorphic conditions but avoiding the
93 problem of estimating the oxidation state of iron in chlorite. The new calibration is based on a
94 bibliographic compilation of data published during the past three decades and addressing a
95 large variety of geological environments. The validity of this geothermometer is then tested
96 by comparison of the results with those obtained with Vidal et al. (2006) and Inoue et al.
97 (2009) thermometers on three *other* sets of analytical data, for which Fe^{3+} contents are known:
98 chlorite analytical electron microscope (AEM) analyses from the Gulf Coast (Texas), where
99 Fe^{3+} content was estimated by multi-equilibrium calculation; chlorite electron-microprobe
100 (EMP) analyses from Saint Martin (Lesser Antilles), where Fe^{3+} content was estimated by
101 Mössbauer spectroscopy (Beaufort et al. 1992); and chlorite EMP analyses from Toyoha
102 (Hokkaido), where Fe^{3+} content was estimated by X-ray photoelectron spectroscopy (XPS,
103 Inoue et al. 2010).

104

105 **Sources of chlorite analyses**

106

107 *Literature data for calibration*


108

109 In order to calibrate the new geothermometer on a wide spectrum in terms of *P-T*
110 conditions, of geological environments and of analytical methods, published data concerning
111 the low-*T* chlorites and their chemical compositions were screened. The specifications
112 governing the selection of the data were as follows:

113

- 114 - A detailed quantitative analysis of chlorites by EMP or AEM should be available along
115 with temperature and pressure data estimated by independent methods, in the range of 50-
116 350°C and below 4 kbar. The temperature data were used as published when given with a
117 sufficient level of confidence (i.e. convergence of several thermometric methods or
118 discussion of the obtained data). The pressure data refer to the authors' estimates
119 whenever possible and to indications of geological context ensuring that formation
120 pressure was less than 4 kbar. In case of uncertainty or insufficient level of confidence
121 regarding the *P-T* data, the analyses were discarded. For instance, some important studies
122 such as Jowett (1991), Zang and Fyfe (1995) and Inoue et al. (2009) had to be discarded
123 because pressure estimates were not available or because temperature data were not
124 assigned to specific chlorite compositions. All relevant published data/studies are
125 summarized in Table 1.
- 126 - Only analyses of material that was not identified as detrital by the authors were retained,
127 as in Rahn et al. (1994) study, where authigenic and detrital compositions are identified.
128 All published data summarized in Table 1 are considered as referring to authigenic
129 chlorites.
- 130 - Only quartz-bearing samples were considered.
- 131 - When these first criteria were met, two additional chemical criteria were applied to
132 exclude contaminated analyses or those that cannot be expressed as a linear combination
133 of end-members. These are:

134



137

138 As a result, a total of 161 analyses and relevant formation conditions were retained for the
139 calibration (Figure 1).

140

141 *Datasets for testing the thermometers*

142

143 *Gulf Coast chlorites*

144

145 These chlorites are from sandstone core samples from nine wells of the Gulf Coast,
146 Texas. Analyses were obtained with AEM on rock ultrathin sections obtained by focused ion
147 beam (FIB) milling, in accordance with the analytical protocol described by Bourdelle et al.
148 (2012). All details regarding the location and geology methods are given by Bourdelle et al.
149 (*in revision*). In the following, four analyses of the crystal rim were used for each sample
150 because these compositions were considered to be a good representation of the last steps of
151 dissolution/recrystallisation processes and were assumed to represent the closest approach to
152 the equilibrium composition for the highest P - T conditions. The independently obtained P - T
153 formation data are bottom-hole temperature (BHT) and bottom-hole pressure (BHP) data
154 corrected following Kehle (1971) and assumed to be the maximum P - T conditions (100-
155 230°C, 0-1.2 kbar) undergone by the samples, with an estimated error range of $\pm 20^\circ\text{C}$ and
156 ± 0.2 kbar (for discussion, see Bourdelle et al. *in revision*).

157 The chemical composition of the selected Gulf Coast chlorites spans a large range
158 (Table 2), but similar to that reported by previous studies on diagenetic clays (e.g. Curtis et al.
159 1984, 1985; Velde and Medhioub 1988; Hillier and Velde 1991; Jahren and Aagaard 1989,
160 1992; Jahren 1991). If all iron is taken as Fe^{2+} , total Al contents range from 2.52 to 3.38
161 atoms per formula unit (apfu) based on a 14 oxygen anhydrous basis (which will be used
162 throughout this paper), vacancies from 0.12 to 0.49 apfu while the $\text{Fe}^{2+}/(\text{Fe}^{2+}+\text{Mg}^{2+})$ ratio
163 varies between 0.47 and 0.88 (Table 2).

164 The Fe^{3+} content was estimated using the approach of Vidal et al. (2006), which is
165 based on the convergence of four chlorite-quartz-water equilibria. This method gives the
166 minimum Fe^{3+} proportion required to reach the best convergence at specific P - T conditions; a
167 maximum Fe^{3+} proportion can be estimated when the equilibrium convergence is lost. Under
168 low- T conditions, the difference between minimum and maximum Fe^{3+} contents is small and,
169 therefore, we considered that the minimum Fe^{3+} content estimated with this approach is a fair
170 approximation of the actual value. The results show that the minimum Fe^{3+} contents are
171 heterogeneous in the temperature window of the Gulf Coast and range between 5% and 40%
172 of total iron.

173

174 *Saint Martin chlorites*

175

176 Relevant analyses have been published by Beaufort et al. (1992). The samples are
177 from volcanoclastic rocks that were hydrothermally altered by an intrusive quartz-diorite
178 pluton emplaced during the Early Oligocene (Beaufort et al. 1990, 1992). The chlorites used
179 in this study are from three different alteration zones: the epidote + chlorite + quartz
180 assemblage zone (*zone 2*), the epidote + chlorite + hematite + quartz assemblage zone (*zone*
181 *3*), and the chlorite + phengite + magnetite assemblage zone (as vein, *zone 4*).
182 Paleotemperatures were estimated by the fluid-inclusion method, and range between 220 and
183 340°C (Beaufort et al. 1992; Inoue et al. 2009).

184 Under the assumption that $Fe_{total} = Fe^{2+}$, the Al^{IV} content ranges from 0.96 to 1.21
185 apfu, vacancies from 0.05 to 0.20 apfu, and the $Fe^{2+}/(Fe^{2+}+Mg^{2+})$ ratio between 0.08 and 0.68.
186 The XFe^{3+} ratios were estimated by Mössbauer spectroscopy (Beaufort et al. 1992) and the
187 results strongly depend on the metamorphic zones: 25-30% for zone 2 chlorites, 32% for zone
188 3, and 16% for zone 4.

189

190 *Toyoha chlorites*

191

192 The seven analyses used concerning Toyoha chlorites have been published by Inoue et
193 al. (2010) and have been also separated from the literature data because they are associated
194 with precise XFe^{3+} estimations. These chlorites are from hydrothermally altered rocks in the
195 Toyoha geothermal system, southwestern Hokkaido, Japan, from the propylitic zone and the
196 ore mineralized zone. Paleotemperatures were estimated with Inoue et al. (2009) model and
197 were close both to the homogenisation temperatures of fluid inclusions and the present
198 subsurface temperatures measured through drill holes. They range between 159 and 264°C.

199 Under the assumption that $Fe_{total} = Fe^{2+}$, the Al^{IV} content ranges from 0.85 to 1.20
200 apfu, Al^{VI} from 1.08 to 1.33 apfu and vacancies from 0.25 to 0.40 apfu. The Fe^{3+} contents
201 were estimated by X-ray photoelectron spectroscopy (XPS) and the results are homogeneous
202 and range from 0.13 to 0.17 for the ore mineralized zone and from 0.20 to 0.26 for the
203 propylitic zone.

204

205 **Formulation of a new geothermometer, as an extension of previous methods**

206

207 To establish a new thermometer, it is necessary to define the cationic repartition in the
208 structure, the activity model used and appropriate solid solutions by the choice of relevant
209 end-member components. In this study, the focus is on chlorites formed at low-temperatures,
210 i.e. between 50 and 350°C, which have high Si content and octahedral vacancies due to

211 Si□R²⁺₂ exchange (e.g. Hillier and Velde 1991). Vidal et al. (2006), building on Vidal et al.
 212 (2001) and Vidal et al. (2005) studies, chose clinochlore, daphnite, Mg-sudoite, Fe-amesite
 213 and Mg-amesite as end-members. This excludes Si-rich (Si > 3 apfu) compositions (Table 3)
 214 and therefore, many of the diagenetic chlorites. Inoue et al. (2009) preferred to consider a
 215 compositional space defined by the Mg-chlorite S, Mg-amesite, daphnite and Mg-sudoite end-
 216 members (Table 3). This choice seems appropriate to cover all the compositional range of
 217 low-*T* chlorites, especially because the octahedral vacancies rarely exceed 1 apfu. However,
 218 all iron is then concentrated in one trioctahedral end-member component, daphnite.
 219 Considering these observations, the chlorite solid solution is defined in this study with six
 220 end-members: (Fe, Mg)-chlorite S, (Fe, Mg)-sudoite and (Fe, Mg)-amesite (Table 3).

221 The activity model depends on the cationic site repartition in the structure; according
 222 to the ideal structure of the trioctahedral chlorite defined by Bailey (1988), the basic layer
 223 consists in a regular alternation of talc {M1(M2)₂[(T1)₂(T2)₂]O₁₀(OH)₂} and brucite layers
 224 {(M3)₂M4(OH)₆}. Each crystallographic site entails some cationic preferences, but two
 225 assumptions remain possible: that of an ordered distribution (Vidal et al. 2005, 2006) and that
 226 of random mixing (Inoue et al. 2009). Vidal et al. (2006) assumed that Al^{IV} is restricted to T2
 227 sites, vacancies to M1, Al^{VI} fills M4 first then M1 and eventually M2-M3, while Fe-Mg fills
 228 M2-M3 sites then M1. Inoue et al. (2009) borrowed the distinction between T1 and T2 sites,
 229 with Al^{IV} exclusively on T2, but assumed a random mixing on all octahedral sites. The
 230 ordered approach seems to be closest to Bailey's (1988) observations and makes Vidal's
 231 thermometer less sensitive to Fe³⁺ content, as shown in a comparative study (Bourdelle et al.
 232 *in revision*). For this reason, an ordered model was adopted in this study to describe the
 233 compositional variations of low-*T* chlorite, as follows. To account for 'chlorite S' end-
 234 members, we unified M1 and M4 sites (Table 4). Tetrahedral Al is restricted to T2 sites,
 235 vacancies to M1-M4, Al^{VI} resulting from Tschermak exchange fills M1-M4, excess Al^{VI} is in
 236 M2-M3, Fe-Mg fills predominantly M2-M3 sites then M1-M4 (Table 4). In accordance with
 237 Vidal et al. (2005, 2006), Fe/Mg and Si/Al ratio are considered equal in the M1-M4 and M2-
 238 M3 sites, and T2 sites respectively. Regarding the special case of Fe³⁺, Vidal et al. (2005,
 239 2006) restricted it to M4 and Inoue et al. (2009) exchanged it with Al^{VI} in all M sites (with an
 240 identical Fe³⁺/Fe²⁺ ratio), whereas the new model does not consider it.

241 From the above cation site repartition, end-members ideal activities can be calculated
 242 following the mixing-on-site model of Helgeson et al. (1978) and Powell (1978):

243

$$244 \quad a_j^{ideal} = k \prod_s \prod_m (X_m)^{r_m} \quad (1)$$

245

246 where a_{ideal} is the ideal part of the activity, r_m and X_m are the number and the mole fraction of
247 cation m on the site s , and k is the normalization constant defined by:

248

$$249 \quad k = \prod_s \prod_m \left(\frac{n_s}{r_m} \right)^{r_m} \quad (2)$$

250

251 As done by Inoue et al. (2009), neglecting the non-ideal contribution of the site
252 mixing, a new semi-empirical $T = f(\log K)$ geothermometer can be formulated with our
253 specific choice of ordered solid-solution model, where K is the equilibrium constant of the
254 end-member-component reaction describing the chlorite + quartz equilibrium.

255 From a thermodynamic point of view, for any balanced chemical reaction involving j end-
256 members, the equilibrium condition can be expressed by:

257

$$258 \quad \Delta G_r^0 + R.T. \ln K = 0 \quad (3)$$

259

260 where ΔG_r^0 is the Gibbs free energy of reaction, R is the gas constant and K is the chlorite +
261 quartz equilibrium constant, which can be written as:

262

$$263 \quad K = \prod_j (a_{ideal})_j^{v_j} \quad (4)$$

264

265 with v_j as the stoichiometric reaction coefficient of phase j . The equations for the calculations
266 of end-member ideal activities are listed in Table 3, according to the cationic site mixing
267 repartition detailed in Table 4. With three end-members, the chlorite + quartz assemblage can
268 be described in the Mg system by the reaction:

269



271

272 If the effect of pressure is ignored, in accord with the $P < 4$ kbar data selection, and the
273 change in heat capacity is assumed to be naught, the $\log K$ for the reaction defined above can
274 be expressed by:

275

$$\begin{aligned}
276 \quad \log K &= \frac{-\Delta G_r^0}{2.303(R.T)} = \frac{A}{T} + B = \log \left(\frac{a_{Mg-Am}^3 \cdot a_{SiO_2}^7 \cdot a_{H_2O}^4}{a_{Mg-ChlS} \cdot a_{Mg-Sud}^3} \right) & (6) \\
&= 3 \log a_{Mg-Am} - \log a_{Mg-ChlS} - 3 \log a_{Mg-Sud} + 7 \log a_{SiO_2} + 4 \log a_{H_2O}
\end{aligned}$$

277

278 where T is temperature and A and B constants, with $A = \Delta H \div (2.303 \times R)$ and

279 $B = \Delta S \div (2.303 \times R)$, given the ΔC_p approximation.

280

281 Calibration of the thermometer

282

283 The new thermometer is calibrated from the relation between the logarithm of the
284 equilibrium constant and $1/T$, using the 161 low- P chlorite compositions ($P < 4$ kbar) and
285 their estimated formation temperature according to the literature (Table 1). For those
286 calculations, we assumed that $a_{SiO_2} = 1$, $a_{H_2O} = 1$ and $Fe_{total} = Fe^{2+}$. From the regression
287 analysis, we obtain the new linear equation:

288

$$289 \quad \log K = -\frac{9400}{T(K)} + 23.40 \quad (7)$$

290

291 A quadratic regression yields the equation

292

$$293 \quad \log K = \frac{11185729}{T(K)^2} - \frac{56598}{T(K)} + 72.3 \quad (8)$$

294

295 which provides a better fit in the range 300-350°C, but has no physical basis (cf. equation 6)
296 and so should not be used outside the range 150-350°C. The relationship between reference
297 temperatures (literature data) and temperatures calculated with the new thermometer for the
298 calibration dataset is shown in Figure 2. The standard deviation from the 1:1 line is almost
299 46°C when using the combination of equations (7) and (8) (cf. Fig 2) and 65°C when using
300 equation (7). However, the standard deviation calculation is biased by the 6 calculated
301 temperatures exceeding 450°C (Fig 2) and is variable with T : in fact, the standard deviation is
302 probably lower at low T (around $\pm 30^\circ\text{C}$) and higher at $T > 300^\circ\text{C}$ (around ± 50 -60°C). This
303 data scattering is similar to those obtained when testing the Inoue et al. (2009) and Vidal et al.
304 (2006) models on the same literature dataset (XFe^{3+} estimated with the Vidal et al. 2006
305 model). Therefore, we consider that the correlation between estimated and expected T is
306 satisfactory.

307

308 **Testing the new geothermometer**

309

310 Using equation (7) applied to the chlorite + quartz assemblage, the formation
311 temperatures of Gulf Coast, Saint Martin and Toyoha chlorites were estimated assuming all
312 iron is ferrous and 1 for H₂O activity. In fact, in all cases, the values of H₂O activity in the
313 fluid are unknown and the assumption $a_{\text{H}_2\text{O}} = 1$ seems to be reasonable in diagenetic to low-
314 grade metamorphic conditions (Inoue et al. 2009), but remains locally questionable. The
315 assumption $a_{\text{SiO}_2} = 1$ is justified in the case of quartz-bearing rocks.

316 The temperatures calculated with equation (7) are plotted (i) versus measured
317 temperatures in Figure 3 for Gulf Coast chlorites, for which the high-spatial-resolution should
318 ensure the best reliability of the analyses, (ii) versus Al^{IV} content in Figure 4 for Saint Martin
319 and Toyoha chlorites assumed to be authigenic crystals. The temperatures obtained with the
320 new thermometer range from 91 to 231°C, for an expected range of 102-232°C for the Gulf
321 Coast samples. Few analyses (BHT~180°C) give too low a temperature (Figure 3) suggesting
322 that they do not represent highest-*T* equilibrium composition. Generally, the measured and
323 calculated temperatures are very close, with less than 20°C difference. For Saint Martin
324 chlorites, the calculated temperatures range from 172 to 351°C (Figure 4) for an expected
325 220-340°C whereas for Toyoha chlorites, the temperatures range from 159 to 301°C for an
326 expected 159-264°C (a single calculated temperature is outside the expected range). In all
327 cases, the new equation appears to be highly reliable to obtain sensible temperature values,
328 despite the assumption that all iron is ferrous and regardless of the rock-type considered.

329

330 **Discussion**

331

332 *Suitability of chlorite analyses for calibration and testing*

333

334 As pointed out by Hillier and Velde (1992), De Caritat et al. (1993), Jiang et al. (1994)
335 and Essene and Peacor (1995), part of the compositional variations measured for low-*T*
336 chlorites may be ascribed to contamination by kaolinite, or to interstratification with
337 berthierine or smectite, which might also lead to erroneous structural formulae and an
338 overestimation of vacancy contents (vacancies are then considered as an analytical artifact).
339 The interstratification with smectite is assumed to be negligible here because only analyses
340 with < 1 % wt total Na₂O + CaO + K₂O were selected, both for calibration and for testing,
341 almost 90% of which actually have Na₂O + CaO + K₂O < 0.5%wt. The interstratification with

342 the 7 Å-phase berthierine, which is a common phenomenon in low- T chlorites (Xu and
343 Veblen, 1996), is difficult to distinguish chemically from a true solid-solution without high-
344 resolution structural control. In this respect, the use of a high-resolution analytical protocol
345 (with AEM) for the Gulf Coast chlorites allows for avoiding the 7 or 10 Å-layer and quartz
346 contaminations; it shows that octahedral vacancies in low- T chlorite are an actual feature and
347 that their amount varies with T . Therefore, the deficit in octahedral cations (< 6 apfu)
348 observed under all conditions (diagenetic or metamorphic) in the absence of significant
349 smectite or mica contamination is real, and should be amplified in structural formula
350 recalculation when accounting for Fe^{3+} content, which is not negligible for low- T chlorites
351 (Beaufort et al. 1992).

352 An issue is the reliability of AEM data compared to EMP. Only three AEM databases
353 were used here: Jahren and Aagaard (1992) and Lopez-Munguira et al. (2002) for calibration,
354 and Gulf Coast chlorites analyses for testing. Lopez-Munguira et al. (2002) showed that the
355 analyses obtained with both methods are highly similar. Bourdelle et al. (2012) also show the
356 suitability of a FIB-AEM protocol to obtain chlorite analyses with a level of confidence
357 similar to EMP analyses: this is the protocol used here to analyse the Gulf Coast chlorites. On
358 this basis, we consider that the reliability of the analyses used for calibration or testing is not
359 affected by the applied analytical method.

360 In a broader perspective and for practical purposes, we aimed at the largest possible
361 database for calibration, offering a wide coverage of geological parameters, like lithology,
362 temperature gradient, kinetics, fluid regimes, oxidation state, chlorite composition (including
363 XFe^{3+}), chlorite precursors, impurities and analytical contaminations (insofar as they are of
364 limited extent). This was considered the best way to limit the effect of each parameter, with
365 the aim of a general thermometer for low- T chlorites. Therefore, all the data used for
366 thermometer calibration were considered as suitable, provided they met the imposed chemical
367 criteria. As a result, the database used in this study is the largest ever gathered for low- T
368 chlorite thermometry, with 161 analyses for $T < 350^\circ\text{C}$, $P < 4\text{kbar}$, of which 94 analyses are
369 for $T < 250^\circ\text{C}$, compared to 23 analyses (for $T < 350^\circ\text{C}$) for Inoue et al. (2009; from Vidal et
370 al. 2001 appendix), with 11 analyses for $T < 250^\circ\text{C}$.

371

372 *Comparison with recent chlorite thermometers*

373

374 The results obtained on Gulf Coast, Saint Martin and Toyoha chlorites with the new
375 thermometer (using the linear equation (7) and the assumption $\text{Fe}_{\text{total}} = \text{Fe}^{2+}$) are compared in
376 Figure 5 to the two main non-empirical models (Vidal et al. 2005, 2006; Inoue et al. 2009) for

377 which Fe^{3+} content is taken into account. Note that many analyses were automatically
378 excluded in Vidal's model by their Si content higher than 3 apfu. The thermodynamic model
379 of Walshe (1986) was not considered, as it does not allow for Si contents lower than 3 apfu.

380 For the Gulf Coast chlorites, Vidal et al. (2005, 2006) and Inoue et al. (2009) models
381 give substantially similar results, with a range of formation temperatures of 84-317°C and
382 113-325°C, respectively (Figure 5a). The two sets of results are very similar (Figure 5a, inset)
383 and are very close to those obtained with equation (7). However, some analyses near ~200°C
384 branch off the 1:1 trend, yielding overestimated temperatures with Inoue and Vidal models.
385 These analyses are characterised by high Al content and low XFe^{3+} (according to Vidal
386 procedure) in contrast to the high- T analyses of the 1:1 trend, which have low Al content and
387 high XFe^{3+} (~35-40%). Important is that, regardless of these compositional variations
388 including XFe^{3+} , the present thermometer, unlike the two others, yields a single trend of
389 temperatures for these two groups, consistent with measured temperatures (cf. Figure 3).

390 For the Saint Martin chlorites, Vidal et al. (2005, 2006) and Inoue et al. (2009) models
391 yield results in the ranges 50-321°C and 172-313°C, respectively. In this case, most of the
392 temperatures obtained with Vidal's model are underestimated and differ from the two other
393 thermometers, whereas the newly proposed equation (7) gives temperatures that are very
394 similar to those obtained with Inoue's model.

395 For Toyoha chlorites, the situation is similar to the Saint Martin case, with a scatter of
396 results obtained with Vidal's thermometer (70-265°C), and results of the two other
397 thermometers close to each other and in better agreement with the expected range (159-
398 264°C).

399 For each of the three geological fields, the results are similar and in the expected
400 temperature windows for Inoue's and the newly proposed model, whereas Vidal's model
401 gives more disparate, often underevaluating results. These differences in calculated
402 temperatures can be explained by the choices of end-members (and site assignment of iron)
403 and of calibration basis (more weight of the high- T , low- Fe^{3+} analyses in Vidal and Inoue
404 databases). The excellent agreement between Inoue and Vidal's models in Figure 5a (inset) in
405 contrast to Figure 5b and c (insets) is not fortuitous as this is the sole case for which XFe^{3+}
406 input ratios are obtained with Vidal et al. (2006) iterative method. In summary, the
407 comparison of the new thermometer and Vidal and Inoue's models shows that equation (7),
408 based on an ordered distribution and the $\text{Fe}_{\text{total}} = \text{Fe}^{2+}$ assumption, performs well in the case of
409 low- T chlorites.

410

411 *Choice of a semi-empirical ordered model for low- T chlorites*

412

413 The semi-empirical approach, based on a reaction, appears as a good alternative
414 solution between a purely thermodynamic treatment, which requires knowledge of the many
415 end-member and mixing properties (as Vidal et al. 2006), and the purely empirical approach,
416 which does not consider the bulk composition (as Cathelineau 1988). As regards the cationic
417 distribution, Inoue et al. (2009) suggest that one reason for the inaccuracy in low- T domain of
418 the Vidal et al. (2001) model, rearranged by Vidal et al. (2005, 2006), is the ordered site
419 occupancy. This statement is derived from the polytype distribution (Ia, Ib, IIb) according to
420 the metamorphic grade. However, in the absence of univocal information, in particular on the
421 validity of each cationic distribution model according to polytype, we prefer to base the new
422 thermometer on a semi-ordered distribution. The results presented above show that it gives at
423 least as reliable estimates as Inoue's disordered model for the three geological fields of Gulf
424 Coast, Saint Martin and Toyoha. The cation distribution model is therefore not the main issue
425 for chlorite thermometry. On the basis of the above comparisons, it appears that the
426 calibration (T range and compositional space) and the choice of end-members determine the
427 pertinence of the various models.

428

429 *Testing the effect of non-ideality*

430

431 As done by Inoue et al. (2009), the non-ideal contribution of the site mixing was
432 neglected in the present study. This assumption was required to calibrate a semi-empirical $T =$
433 $f(K)$ geothermometer, but it needs to be evaluated, considering that sudoite and trioctahedral
434 chlorite may coexist in equilibrium in a few cases (in low- T , low- P or low- T , high- P
435 aluminous metapelites, e.g. Theye et al. 1992). For the assumption of ideal mixing to be
436 tenable, one must verify that (i) for a given chlorite composition, the contribution of the non-
437 ideality varies linearly with $1/T$, and (ii) at given T , the variation of the non-ideal contribution
438 with chlorite composition is small, so that the net effect of non-ideality is grossly linear in T
439 and so makes a linear calibration possible.

440 For the first point, the expression of the non-ideality is:

441

442
$$K = \prod_j (a_{ideal} \cdot \gamma)_j^{v_j} \text{ with } \gamma_j = \prod_s \prod_m \gamma_m^{n_s} \quad (9)$$

443

444 where, for a j component, γ is the non-ideality coefficient, n_s is the multiplicity of the site s
 445 and m are the relevant cations. The $\log/\ln K$ can be expressed as the sum of an ideal and a
 446 non-ideal part:

447

$$448 \quad R.T.\ln K = R.T.\ln\left(\prod_j a_{ideal}\right) + R.T.\ln\left(\prod_j \gamma\right) \quad (10)$$

449

450 Neglecting ternary and quaternary interactions, the non-ideal contribution may be written:

451

$$452 \quad n_s.R.T.\ln \gamma_m = \sum W_{ij}.X_i.X_j \left[\frac{Q_m}{X_m} - 1 \right] \quad (11)$$

453

454 where W_{ij} are the Margules parameters, X_m is the mole fraction of cation m and Q_m is the
 455 number of i, j subscripts that are equal to m . Clearly, from this formalism, the non-ideal
 456 contribution in the $\log/\ln K$ calculation is linear in $1/T$ for a fixed composition.

457 For the second point, Bourdelle (2011) tested the new model for various chlorite
 458 compositions and P - T conditions. Starting with an average composition and an approximation
 459 of Margules parameters defined by Vidal et al. (2001, 2005, 2006), he used equations (10)
 460 and (11) and the defined semi-ordered distribution to calculate the non-ideal contribution for
 461 several temperatures at given P , considering symmetric binary interactions. The author
 462 introduced variation in the composition and conditions, by varying the $X_{Si_{T2}}$, X_{Mg} (ratio),
 463 X_{\square} and pressure. The results of these tests are plotted in Figure 6 and show, in the range of
 464 100-350°C and neglecting the $W(M2+M3)$ parameters (following Vidal et al. 2005, 2006) as
 465 well as the pressure factor, that the variations of X_{Mg} ratio affect only very slightly the
 466 variation of $\log/\ln K$ with $1/T$. This holds especially if the tested chlorite compositions are
 467 only considered in the range of temperatures to which they refer (i.e. the Si-rich composition,
 468 $Si = 3.4$ apfu, mainly characterises very low- T chlorites, contrary to the Si-poor composition,
 469 $Si = 2.6$ apfu). The non-ideal contribution mainly affects the y-intercept of the $\ln K = f(1/T)$
 470 curve, which does not preclude a linear calibration. Thus, we conclude that the variation of
 471 the non-ideal contribution with composition (in the classical range of diagenetic chlorites) is
 472 sufficiently small for the linearity of $\log/\ln K$ in $1/T$ to be grossly preserved. Therefore, for the
 473 P - T domain and the composition range investigated here, calibration on a linear basis is
 474 justified and, actually, any effect of non-ideality is implicitly taken into account in the
 475 thermometer equation.

476

477 *Link between Fe³⁺ content, octahedral vacancies, thermometer calibration and temperature*
478 *calculation*

479

480 The main compositional features of chlorite evolution with increasing T are long
481 recognized as the Tschermak substitution with a contribution of di-trioctahedral substitution
482 at low T . This decrease of octahedral vacancies with increasing T is therefore built in any
483 calibration database, particularly when the database includes lower- T chlorites. Besides,
484 accounting for Fe³⁺ in the structural formula recalculation on a fixed oxygen basis
485 arithmetically reduces the number of cations, and implies an increase of octahedral vacancies,
486 which in turn links indirectly Fe³⁺ content and temperature in any thermometric formulation.
487 However, this effect of Fe³⁺ remains subordinate with respect to the main compositional
488 evolutions with T mentioned above (compare Figures 6a and 6b in Inoue et al. 2009).

489 Accordingly, variations of the chlorite + quartz log K remain negatively correlated with $1/T$,
490 regardless of the assumption made on the oxidation state of iron. This justifies calibrating a
491 chlorite thermometer on one or the other assumption, i.e. Fe_{total} = Fe²⁺ or real Fe³⁺ values. A
492 calibration based on the latter choice should lead to higher precision, but calibration based on
493 the former case may be a good trade-off of practicability and simplicity (Fe³⁺ content not
494 required) against precision. Admittedly, the sudoite activity is poorly evaluated, i.e. biased by
495 the non consideration of Fe³⁺ content, but this possible bias is built in the calibration. The best
496 demonstration of the validity of this trade-off is the success of the thermometer presented here
497 in its applications to various geological contexts.

498

499 **Conclusion**

500

501 Various concerns have been raised on the relevance, applicability and precision of
502 chlorite thermometry (e.g. De Caritat et al. 1993; Jiang et al. 1994; Essene and Peacor 1995).
503 However, recent developments (e.g. Vidal et al. 2001, 2005, 2006; Inoue et al. 2009) have led
504 to a reappraisal of this technique and to new opportunities in exploiting the information
505 contained in chlorite composition. Comparative studies (e.g. Inoue et al. 2009; this paper)
506 have yet pointed out that order-disorder, compositional space, non-ideality and Fe³⁺ effects
507 may be limitations and/or sources of disagreement. The approach developed here,
508 compensating these sources of variability through the use of a large database for linear
509 calibration, proves successful in its application to low- T chlorites. This success may lie in the
510 fact that the linear dependence of log K with $1/T$ either integrates these effects (were they

511 non-ideality or Fe³⁺ content), or is not significantly affected by them, the backbone of the
512 variation being the combined effects of the Tschermak and di-trioctahedral substitutions.

513 The new geothermometer allows one to account for all the low-*T* chlorite
514 compositions, especially Si-rich compositions that characterize diagenetic chlorites. The
515 comparison of the new thermometer with the recent ones that require Fe³⁺ content estimation
516 shows that it is possible to obtain reliable results ($\pm 20^\circ\text{C}$ in most cases) without any
517 measurement or assumption on the Fe³⁺ content. In this respect, the present semi-empirical
518 thermometer is a much practical tool, well suited for, e.g., the handling of large analytical
519 datasets in exploration geology. However, it should be kept in mind that the new equation
520 proposed here is only valid over its calibration range ($T < 350^\circ\text{C}$, $P < 4$ kbar) and that any use
521 out of these limits, in particular in the metamorphic realm, is not recommended – mainly
522 because of the assumptions on the heat-capacity and volume changes of the reaction, the
523 pressure effect and the water activity in the new model, which may no longer be valid. Above
524 350°C , the thermometry of Vidal et al. (2005, 2006) remains probably the most convenient
525 approach.

526

527 **Acknowledgements**

528

529 We are most grateful to the materials characterization department of IFP Energies Nouvelles-
530 Lyon, in particular to F. Moreau, and to the laboratory of CP2M-Université Aix-Marseille, for
531 technical advice. The discussions and comments of the journal editor Jochen Hoefs, of
532 Atsuyuki Inoue and two anonymous reviewers are gratefully acknowledged. Thanks are also
533 extended to K. Milliken, S. Dutton and J. Donnelly of Bureau of Economic Geology at
534 Austin, Texas. This study was financially supported by IFP Energies Nouvelles, CNRS and
535 ENS Paris.

536

537 **References**

538

539 Bailey SW (1988) Chlorites: structures and crystal chemistry. In: Bailey SW (eds) *Hydrous*
540 *Phyllosilicates (Exclusive of Micas)*, vol 19. Reviews in Mineralogy. Mineralogical
541 Society of America, Washington D.C., pp 347-403

542 Beaufort D, Patrier P, Meunier A, Ottaviani MM (1992) Chemical variations in assemblages
543 including epidote and/or chlorite in the fossil hydrothermal system of Saint Martin (Lesser
544 Antilles). *J Volcanol Geoth Res* 51:95-114

545 Beaufort D, Westercamp D, Legendre O, Meunier A (1990) The fossil hydrothermal system
546 of Saint Martin: (1) Geology and lateral distribution of alterations. *Journal of Volcanology*
547 and *Geothermal Research* 40:219-243

548 Bevins RE, Robinson D, Rowbotham G (1991) Compositional Variations in Mafic
549 Phyllosilicates from Regional Low-Grade Metabasites and Application of the Chlorite
550 Geothermometer. *J Metamorph Geol* 9 (6):711-721

551 Boles JR, Francks GS (1979) Clay diagenesis in Wilcox sandstones of Southwest Texas:
552 Implications of smectite diagenesis on sandstone cementation. *J Sediment Petrol* 49:55-70

553 Bourdelle F (2011) Thermobarométrie des phyllosilicates dans les séries naturelles:
554 Conditions de la diagenèse et du métamorphisme de bas degré. Thesis, University of Paris-
555 Sud, Orsay

556 Bourdelle F, Parra T, Beyssac O, Chopin C, Moreau F (2012) Ultrathin section preparation of
557 phyllosilicates by Focused Ion Beam milling for quantitative analysis by TEM-EDX. *Appl*
558 *Clay Sci* 59-60:121-130

559 Bourdelle F, Parra T, Chopin C, Beyssac O, Vidal O (*in revision*) Clay minerals thermometry:
560 A comparative study based on high-spatial-resolution analyses of illite and chlorite in Gulf
561 Coast sandstones (Texas, USA). *Am Min*

562 Cathelineau M (1988) Cation site occupancy in chlorites and illites as a function of
563 temperature. *Clay Miner* 23 (4):471-485

564 Cathelineau M, Nieva D (1985) A chlorite solid solution geothermometer. The Los Azufres
565 (Mexico) geothermal system. *Contrib Mineral Petrol* 91 (3):235-244

566 Curtis CD, Hughes CR, Whiteman JA, Whittle CK (1985) Compositional variation within
567 some sedimentary chlorites and some comments on their origin. *Mineral Mag* 49:375-386

568 Curtis CD, Ireland BJ, Whiteman JA, Mulvaney R, Whittle CK (1984) Authigenic chlorites:
569 Problems with chemical analysis and structural formula calculation. *Clay Miner* 19:471-
570 481

571 De Caritat P, Hutcheon I, Walshe JL (1993) Chlorite geothermometry: A review. *Clay Clay*
572 *Miner* 41 (2):219-239

573 Essene EJ, Peacor DR (1995) Clay mineral thermometry - A critical perspective. *Clay Clay*
574 *Miner* 43 (5):540-553

575 Foster MD (1962) Interpretation of the composition and a classification of the chlorites. US
576 Geological Survey Professional Paper 414-A:33

577 Grosch EG, Vidal O, Abu-Alam T, McLoughlin N (2012) P-T Constraints on the
578 metamorphic evolution of the Paleoproterozoic Kromberg type-Section, Barberton Greenstone
579 Belt, South Africa. *J Petrol* 53 (3): 513-545.

580 Helgeson HC, Delany JM, Nessbitt HW, Bird DK (1978) Summary and critique of the
581 thermodynamic properties of rock-forming minerals. *Am J Sci* 278A:1-229

582 Hillier S, Velde B (1991) Octahedral occupancy and the chemical-composition of diagenetic
583 (low-temperature) chlorites. *Clay Miner.* 26 (2):149-168

584 Hillier S, Velde B (1992) Chlorite interstratified with a 7 Å mineral: An example from
585 offshore Norway and possible implications for the interpretation of the composition of
586 diagenetic chlorites. *Clay Miner.* 27:475-486

587 Hutcheon I (1990) Clay carbonate reactions in the Venture area, Scotian Shelf, Nova Scotia,
588 Canada. The Geochemical society, Special Publication, pp 199-212

589 Inoue A, Kurokawa K, Hatta T (2010) Application of chlorite geothermometry to
590 hydrothermal alteration in Toyoha geothermal system, Southwestern Hokkaido, Japan.
591 *Resour Geol* 60 (1):52-70. doi:10.1111/j.1751-3928.2010.00114.x

592 Inoue A, Meunier A, Patrier-Mas P, Rigault C, Beaufort D, Vieillard P (2009) Application of
593 Chemical Geothermometry to Low-Temperature Trioctahedral Chlorites. *Clay Clay Miner*
594 57 (3):371-382

595 Jahren JS (1991) Evidence of Ostwald ripening related recrystallization of chlorites from
596 reservoir rocks offshore Norway. *Clay Miner* 26:169-178

597 Jahren JS, Aagaard P (1989) Compositional variations in diagenetic chlorites and illites, and
598 relationships with formation-water chemistry. *Clay Miner* 24:157-170

599 Jahren JS, Aagaard P (1992) Diagenetic Illite-Chlorite Assemblages in Arenites .1. Chemical
600 Evolution. *Clay Clay Miner* 40 (5):540-546

601 Jiang WT, Peacor DR, Buseck PR (1994) Chlorite geothermometry? - Contamination and
602 apparent octahedral vacancies. *Clay Clay Mineral* 42 (5):593-605

603 Jowett EC (1991) Fitting iron and magnesium into the hydrothermal chlorite geothermometer.
604 Paper presented at the GAC/MAC/SEG Joint annual meeting, Toronto, Canada, 27-29 May
605 1991

606 Kehle RO (1971) Geothermal survey of North America. American Association of Petroleum
607 Geologists, 31p.

608 Koroknai B, Arkai P, Horvath P, Balogh K (2008) Anatomy of a transitional brittle-ductile
609 shear zone developed in a low-*T* meta-andesite tuff: A microstructural, petrological and
610 geochronological case study from the Bukk Mts. (NE Hungary). *J Struct Geol* 30 (2):159-
611 176

612 Kranidiotis P, McLean WH (1987) Systematics of chlorite alternation at the Phelps Dodge
613 massive sulfide deposit, Matagami, Quebec. *Econ Geol* 82 (7):1898-1911

614 Laird J (1988) Chlorites: metamorphic petrology. In: Bailey SW (eds) *Hydrous Phyllosilicates*
615 (Exclusive of Micas), vol 19. The Mineralogical Society of America, Washington D.C., pp
616 405-453

617 Lopez-Munguira A, Nieto F, Morata D (2002) Chlorite composition and geothermometry: a
618 comparative HRTEM/AEM-EMPA-XRD study of Cambrian basic lavas from the Ossa
619 Morena Zone, SW Spain. *Clay Miner* 37 (2):267-281

620 Mas A, Guisseau D, Mas PP, Beaufort D, Genter A, Sanjuan B, Girard JP (2006) Clay
621 minerals related to the hydrothermal activity of the Bouillante geothermal field
622 (Guadeloupe). *J Volcanol Geoth Res* 158 (3-4):380-400

623 McDowell SD, Elders WA (1980) Authigenic layer silicate minerals in borehole Elmore 1,
624 Salton Sea geothermal field, California, USA. *Contrib Mineral Petrol* 74:293-310

625 Powell R (1978) *Equilibrium Thermodynamics in Petrology: An Introduction*. Harper & Row,
626 London

627 Rahn M, Mullis J, Erdelbrock K, Frey M (1994) Very Low-Grade Metamorphism of the
628 Taveyanne Greywacke, Glarus Alps, Switzerland. *J Metamorph Geol* 12 (5):625-641

629 Schmidt D, Schmidt ST, Mullis J, Mahlmann RF, Frey M (1997) Very low grade
630 metamorphism of the Taveyanne formation of western Switzerland. *Contrib Mineral Petrol*
631 129 (4):385-403

632 Theye T, Seidel E, Vidal O (1992) Carpholite, sudoite, and chloritoid in low-grade high-
633 pressure metapelites from Crete and the Peloponnese, Greece. *Eur J Mineral* (4): 487-507

634 Velde B, Medhioub M (1988) Approach to chemical equilibrium in diagenetic chlorites.
635 *Contrib Mineral Petrol* 98:122-127

636 Vidal O, De Andrade V, Lewin E, Munoz M, Parra T, Pascarelli S (2006) P-T-deformation-
637 Fe³⁺/Fe²⁺ mapping at the thin section scale and comparison with XANES mapping:
638 application to a garnet-bearing metapelite from the Sambagawa metamorphic belt (Japan).
639 *J Metamorph Geol* 24 (7):669-683

640 Vidal O, Parra T (2000) Exhumation paths of high-pressure metapelites obtained from local
641 equilibria for chlorite-phengite assemblages. *Geol J* 35 (3-4):139-161

642 Vidal O, Parra T, Trotet F (2001) A thermodynamic model for Fe-Mg aluminous chlorite
643 using data from phase equilibrium experiments and natural pelitic assemblages in the 100 °
644 to 600 °C, 1 to 25 kb range. *Am J Sci* 301 (6):557-592

645 Vidal O, Parra T, Vieillard P (2005) Thermodynamic properties of the Tschermak solid
646 solution in Fe-chlorite: Application to natural examples and possible role of oxidation. *Am*
647 *Mineral* 90 (2-3):347-358

648 Walshe JL (1986) A six-component chlorite solid solution model and the conditions of
649 chlorite formation in hydrothermal and geothermal systems. *Econ Geol* 81:681-703
650 Xie XG, Byerly GR, Ferrell RE (1997) Ilb trioctahedral chlorite from the Barberton
651 greenstone belt: Crystal structure and rock composition constraints with implications to
652 geothermometry. *Contrib Mineral Petrol* 126 (3):275-291
653 Xu H, Veblen DR (1996) Interstratification and other reaction microstructures in the chlorite-
654 berthierine series. *Contrib Mineral Petrol* 124:291-301
655 Zang W, Fyfe WS (1995) Chloritization of the Hydrothermally Altered Bedrock at the
656 Igarape-Bahia Gold Deposit, Carajas, Brazil. *Miner Deposita* 30 (1):30-38

657

658 **Tables**

659

660 **Table 1** Published data and methods to estimate temperature for the analyses that have been
661 used to calibrate the new thermometer

662

Study	Temperature estimation methods
Boles and Franks (1979)	<i>in situ</i> temperature measurement
McDowell and Elders (1980)	<i>in situ</i> temperature measurement
Cathelineau (1988)	<i>in situ</i> temperature measurement, data compilation
Hutcheon (1990)	<i>in situ</i> temperature measurement
Bevins et al. (1991)	chl-thermometry, mineral assemblages
Jahren and Aagaard (1992)	<i>in situ</i> temperature measurement
De Caritat et al. (1993)	data compilation, chl-thermometry, carbonates thermometry
Rahn et al. (1994)	chl-thermometry, vitrinite reflectance, fluid inclusions
Schmidt et al. (1997)	chl-thermometry, fluid inclusions, isotopy ($\Delta^{18}\text{O}_{\text{qz-calcite}}$), IC and CC index
Xie et al. (1997)	chl-polytypism, chl-thermometry
Lopez-Munguira et al. (2002)	chl-thermometry
Mas et al. (2006)	<i>in situ</i> temperature measurement
Koroknai et al. (2008)	chl-thermometry, mineral assemblages

663

664 **Table 2** AEM analyses of Gulf Coast chlorites (crystal rims analyses). Elements contents are
665 given in atom per formula unit (O = 14 apfu). All iron is considered as ferrous

666

Sample	AZ#159	AZ#159	AZ#159	AZ#159	ST#470	ST#470	ST#470	CK#2	CK#2	CK#2
	9230	9230	9230	9230	10717	10717	10717	11924	11924	11924
Analysis	chl10	chl17	chl30	chl28	chl27	chl29	chl30	chl47	chl50	chl54
BHT (°C)	102	102	102	102	121	121	121	129	129	129
BHP (bars)	300	300	300	300	590	590	590	660	660	660

Si	2.98	2.95	2.95	2.91	3.08	3.19	3.01	2.96	2.90	2.91
Ti	0.01	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.00
Al	2.76	2.76	2.75	2.86	2.55	2.60	2.77	2.90	2.98	2.86
Fe ²⁺	2.18	2.20	2.34	2.46	2.81	2.55	2.53	2.45	2.37	2.53
Mn	0.00	0.01	0.02	0.00	0.02	0.02	0.01	0.00	0.01	0.00
Mg	1.65	1.71	1.60	1.39	1.15	1.11	1.22	1.18	1.31	1.29
Ca	0.03	0.01	0.00	0.00	0.01	0.00	0.02	0.01	0.00	0.01
Na	0.00	0.00	0.01	0.03	0.00	0.03	0.00	0.02	0.00	0.00
K	0.05	0.05	0.06	0.05	0.07	0.05	0.06	0.06	0.06	0.07

667

Sample	CK#2 12196	CK#2 12196	CK#2 12196	CK#2 12196	LA#1 13559	LA#1 13559	LA#1 13559	LA#1 13559	ST#356 14501	ST#356 14501
Analysis	chl20	chl18	chl19	chl21	chl23	chl20	chl11	chl17	chl36	chl34
BHT (°C)	135	135	135	135	149	149	149	149	166	166
BHP (bars)	690	690	690	690	850	850	850	850	800	800
Si	2.96	3.05	3.02	3.02	2.88	2.86	2.83	2.78	2.84	2.71
Ti	0.00	0.02	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Al	2.85	2.83	2.91	2.81	2.91	2.92	2.95	3.07	2.85	3.00
Fe ²⁺	2.45	2.38	2.38	2.39	3.28	3.36	3.41	3.28	2.05	2.07
Mn	0.00	0.02	0.00	0.01	0.01	0.00	0.00	0.03	0.00	0.00
Mg	1.34	1.18	1.20	1.30	0.53	0.48	0.47	0.49	1.96	1.98
Ca	0.00	0.04	0.01	0.01	0.02	0.00	0.01	0.03	0.00	0.00
Na	0.00	0.00	0.00	0.00	0.03	0.02	0.00	0.02	0.03	0.00
K	0.00	0.03	0.01	0.08	0.03	0.04	0.04	0.03	0.03	0.05

668

Sample	CW#1 14277	CW#1 14277	CW#1 14277	CW#1 14277	WR#C1 17805	WR#C1 17805	WR#C1 17805	FR#1 18946	FR#1 18946	FR#1 18946
Analysis	chl13	chl14	chl32	chl31	chl13	chl18	chl16	chl26	chl30	chl28
BHT (°C)	191	191	191	191	191	191	191	204	204	204
BHP (bars)	750	750	750	750	1050	1050	1050	1150	1150	1150
Si	2.68	2.68	2.65	2.63	2.98	2.97	2.90	2.88	2.87	2.87
Ti	0.00	0.02	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.01
Al	3.38	3.22	3.24	3.21	3.00	2.91	2.95	2.64	2.62	2.60
Fe ²⁺	2.68	2.98	3.00	3.01	2.38	2.39	2.48	2.01	2.08	2.07
Mn	0.20	0.00	0.00	0.00	0.00	0.00	0.01	0.01	0.01	0.00
Mg	0.82	0.79	0.80	0.85	1.08	1.26	1.28	2.24	2.19	2.23
Ca	0.05	0.00	0.00	0.04	0.01	0.02	0.00	0.00	0.02	0.01
Na	0.00	0.00	0.07	0.00	0.03	0.00	0.01	0.00	0.00	0.00
K	0.04	0.04	0.00	0.03	0.03	0.05	0.02	0.05	0.06	0.07

669

Sample	FR#1 18946	AL#1 19110	AL#1 19110	AL#1 19110	AL#1 19110	AL#1 20711	AL#1 20711	AL#1 20711
Analysis	chl29	chl17	chl18	chl27	chl25	chl31	chl28	chl29

BHT (°C)	204	216	216	216	216	232	232	232
BHP (bars)	1150	1150	1150	1150	1150	1200	1200	1200
Si	2.84	2.66	2.64	2.66	2.61	2.94	2.88	2.85
Ti	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.00
Al	2.64	3.21	3.21	3.13	3.16	2.52	2.60	2.64
Fe ²⁺	2.15	2.47	2.53	2.43	2.44	2.74	2.95	2.93
Mn	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00
Mg	2.11	1.38	1.32	1.49	1.55	1.50	1.36	1.36
Ca	0.01	0.01	0.02	0.01	0.00	0.00	0.00	0.00
Na	0.02	0.00	0.00	0.03	0.00	0.02	0.00	0.02
K	0.07	0.01	0.07	0.04	0.05	0.06	0.07	0.06

670

671 **Table 3** Ideal activities of solid-solution end-members used in the calculation of the three
672 chlorite geothermometers

673

End-members (abbreviations)	Chemical formula* (half formula unit)	Equilibrium / Ideal activities [†]
<i>Vidal et al. (2005, 2006) model</i>		
<i>Thermodynamic formalism</i>		
Clinochlore (Clin)	(AlMg ₅)(Si ₃ Al)	2 Clin + 3 Mg-Sud = 4 Mg-Am + 7 Qtz + 4 H ₂ O = 4 (X _{Mg,M1})(X _{Mg,M2+M3}) ⁴ (X _{Si,T2})(X _{Al,T2})
Daphnite (Daph)	(AlFe ₅)(Si ₃ Al)	= 4 (X _{Fe,M1})(X _{Fe,M2+M3}) ⁴ (X _{Si,T2})(X _{Al,T2})
Mg-Amesite (Mg-Am)	(Al ₂ Mg ₄)(Si ₂ Al ₂)	= 64 (X _{□,M1})(X _{Al,M2+M3}) ² (X _{Mg,M2+M3}) ² (X _{Si,T2})(X _{Al,T2})
Fe-Amesite (Fe-Am)	(Al ₂ Fe ₄)(Si ₂ Al ₂)	= 4 (X _{□,M1})(X _{Al,M2+M3}) ² (X _{Mg,M2+M3}) ² (X _{Si,T2})(X _{Al,T2})
Mg-Sudoite (Mg-Sud)	(Al ₃ Mg ₂)(Si ₃ Al)	= (X _{Al,M1})(X _{Mg,M2+M3}) ⁴ (X _{Al,T2}) ²
<i>Inoue et al. (2009) model</i>		
$T(K) = 1 / (0.00293 - 5.13 \times 10^{-4} \times \log K + 3.904 \times 10^{-5} \times (\log K)^2)$		
Mg-Chlorite S ^a (Mg-Chl S)	(Mg ₆)(Si ₄)	Mg-Chl S + 3 Mg-Sud = 3 Mg-Am + 7 Qz + 4 H ₂ O = (X _{Mg,oct}) ⁶ (X _{Si,tet}) ²
Daphnite ^b (Daph)	(AlFe ₅)(Si ₃ Al)	= 59.720 (X _{Fe,oct}) ⁵ (X _{Al,oct})(X _{Si,tet})(X _{Al,tet})
Mg-Sudoite (Mg-Sud)	(Al ₃ Mg ₂)(Si ₃ Al)	= 1728 (X _{Mg,oct}) ² (X _{Al,oct}) ³ (X _{□,oct})(X _{Si,tet})(X _{Al,tet})
Mg-Amesite ^c (Mg-Am)	(Al ₂ Mg ₄)(Si ₂ Al ₂)	= 45.563 (X _{Mg,oct}) ⁴ (X _{Al,oct}) ² (X _{Al,tet}) ²
<i>This study model</i>		
$T(K) = 9400 / (23.40 - \log K)$		
Mg-Chlorite S (Mg-Chl S)	(Mg ₆)(Si ₄)	Mg-Chl S + 3 Mg-Sud = 3 Mg-Am + 7 Qz + 4 H ₂ O = (X _{Si,T2}) ² (X _{Mg,M2+M3}) ⁴ (X _{Mg,M1+M4}) ²
Fe-Chlorite S (Fe-Chl S)	(Fe ₆)(Si ₄)	= (X _{Si,T2}) ² (X _{Fe,M2+M3}) ⁴ (X _{Fe,M1+M4}) ²
Mg-Amesite (Mg-Am)	(Al ₂ Mg ₄)(Si ₂ Al ₂)	= (X _{Al,T2}) ² (X _{Mg,M2+M3}) ⁴ (X _{Al,M1+M4}) ²
Fe-Amesite (Fe-Am)	(Al ₂ Fe ₄)(Si ₂ Al ₂)	= (X _{Al,T2}) ² (X _{Fe,M2+M3}) ⁴ (X _{Al,M1+M4}) ²
Mg-Sudoite (Mg-Sud)	(Al ₃ Mg ₂)(Si ₃ Al)	= 256 (X _{Si,T2})(X _{Al,T2})(X _{Al,M1+M4}) (X _{□,M1+M4})(X _{Mg,M2+M3}) ² (X _{Al,M2+M3}) ²

$$\text{Fe-Sudoite (Fe-Sud)} \quad (\text{Al}_3\text{Fe}_2)(\text{Si}_3\text{Al}) \quad = 256 (X_{\text{Si},\text{T}2})(X_{\text{Al},\text{T}2})(X_{\text{Al},\text{M}1+\text{M}4}) \\ (X_{\square,\text{M}1+\text{M}4})(X_{\text{Fe},\text{M}2+\text{M}3})^2 (X_{\text{Al},\text{M}2+\text{M}3})^2$$

674 Note : $X_{j,s}$ is the mole fraction of the j cation on the s site

675 Equivalents in Inoue et al. (2009): ^a Al-free chlorite, ^b chamosite, ^c corundophilite. * Structural formula based on
676 $\text{O}_{10}(\text{OH})_8$. † Ideal activities are completed with a non-ideal contribution in the Vidal et al. (2005, 2006) model

677
678 **Table 4** Chlorite solid-solution end-members and cationic site repartition used in the new
679 geothermometer calculation

680


End-members	T1(2)	T2(2)	†	(M2+M3)(4)	†	(M1+M4)
Mg-Chlorite S	Si(2)	Si(2)		Mg(4)		Mg(2)
Fe-Chlorite S	Si(2)	Si(2)		Fe(4)		Fe(2)
Mg-Sudoite	Si(2)	SiAl		Al(2)Mg(2)		Al□
Fe-Sudoite	Si(2)	SiAl		Al(2)Fe(2)		Al□
Mg-Amesite	Si(2)	Al(2)		Mg(4)		Al(2)
Fe-Amesite	Si(2)	Al(2)		Fe(4)		Al(2)
Mg			4	Mg - Mg _{M1}	3	(Fe _{M1} + Mg _{M1}) x XMg
Fe			4	Fe - Fe _{M1}	3	(Fe _{M1} + Mg _{M1}) x XFe
Fe + Mg			2	4 - (Al ^{VI} - Al ^{IV})	2	2 - (Al _{M1+M4} + □)
Al		4 - Si	1	Al ^{VI} - Al ^{IV}	1	Al ^{IV}
□					1	(Al ^{VI} - Al ^{IV})/2

681 † indicate the sequence in which the cation assignment is made

682


683 Figures

684



685


686 **Fig. 1** P - T data for 161 chlorites analyses compiled from literature (Table 1)



688


689 **Fig. 2** Relationship between temperatures of reference (literature data) and temperatures
 690 calculated with the new thermometer (161 analyses used for thermometer calibration, listed in
 691 Table 1), considering the chemical composition of chlorites formed at $P < 4$ kbar. Black
 692 symbols: temperatures calculated with the linear equation (7) and the quadratic equation (8)
 693 when $T > 300^{\circ}\text{C}$. Grey symbols: temperatures calculated with the linear equation (7) only.
 694 The shaded zone indicates the 1:1 line $\pm 50^{\circ}\text{C}$. See text for details

695






696


697 **Fig. 3** Comparison of measured/expected temperatures ($BHT \pm 20^\circ C$) for chlorites of the Gulf
 698 Coast formation versus temperatures calculated using the new geothermometer (all Fe is
 699 ferrous)
 700



701
 702 **Fig. 4** Plot of tetrahedral Al contents versus calculated temperatures using the new
 703 geothermometer (all Fe is ferrous) for Saint Martin and Toyoha chlorites. The range of
 704 expected temperatures is indicated by the shaded area for St Martin chlorites, by the hatched
 705 area for Toyoha chlorites. The solid and dashed curves are the equations of Cathelineau
 706 (1988) and Hillier and Velde (1991) empirical thermometers, respectively
 707



709 **Fig. 5** Comparison of calculated temperatures using Inoue et al. (2009), Vidal et al. (2006)
 710 and the present chlorite geothermometers. Solid and open symbols refer to temperatures
 711 calculated by Inoue et al. (2009) and Vidal et al. (2006) models, respectively. Dashed
 712 rectangular areas correspond to the expected temperature ranges for the Gulf Coast (a), Saint
 713 Martin (b) and Toyoha (c) datasets. The shaded zone indicates the 1:1 line $\pm 50^\circ\text{C}$.
 714



715
 716 **Fig. 6** Effect of non-ideality on $\ln K$, considering symmetric binary interactions, $X_{\square} = 0.2$
 717 apfu, $W(M2-M3) = 0$ and various Si content and XMg ratio; modified after Bourdelle (2011).
 718 The solid curve corresponds to the thermometer calibration (noted $\ln K_{th}$), which is indicated
 719 as reference and is calculated according to $1/T$ (not from the fixed compositions). The
 720 variations of XMg ratio affect only very slightly the slope of the $\ln K = f(1/T)$ curve,
 721 especially for the Si-poor composition case