



HAL
open science

Arachnoid webs: preoperative differential diagnosis and surgical exploration

Constantin Tuleasca, Yohan Ducos, Marc Levivier, Fabrice Parker, Nozar Aghakhani

► **To cite this version:**

Constantin Tuleasca, Yohan Ducos, Marc Levivier, Fabrice Parker, Nozar Aghakhani. Arachnoid webs: preoperative differential diagnosis and surgical exploration. *Journal of Neurosurgery: Spine*, In press, <10.3171/2019.6.SPINE19671>. <hal-02289262>

HAL Id: hal-02289262

<https://hal.sorbonne-universite.fr/hal-02289262v1>

Submitted on 16 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Arachnoid webs contemporary management: particular care for preoperative differential diagnosis and surgical exploration!

Constantin Tuleasca^{1,2,3,4,5}, MD-PhD, Yohan Ducos¹, MD, Marc Levivier^{3,4}, MD, PhD,
IFAANS, Fabrice Parker¹, MD, PhD AND Nozar Aghakhani¹, MD, PhD

¹Assistance Publique-Hôpitaux de Paris, Hôpitaux Universitaires Paris Sud, Centre Hospitalier Universitaire de Bicêtre, Service de Neurochirurgie, Le Kremlin-Bicêtre, France;

²Sorbonne Université, Faculté de Médecine, Paris, France;

³Neurosurgery Service and Gamma Knife Center, Lausanne University Hospital (CHUV), Lausanne, Switzerland

⁴Université de Lausanne (Unil), Faculté de Biologie et de Médecine (FBM);

⁵Signal Processing Laboratory (LTS 5), Swiss Federal Institute of Technology (EPFL);

of abstract words : N/A

of text words : 464

of references : 9

of tables/figures :

of videos : 0

Running title: Arachnoid webs

Key words: arachnoid, webs, arachnoidolysis, shunt

Corresponding author:

Constantin Tuleasca, MD-PhD, Assistance Publique-Hôpitaux de Paris, Hôpitaux Universitaires Paris Sud, Centre Hospitalier Universitaire de Bicêtre, Service de Neurochirurgie, Le Kremlin-Bicêtre, France- 78, Rue Général Leclerc, 94270, Le Kremlin-Bicêtre, France; Sorbonne Université, Faculté de Médecine, Paris, France; Université de Lausanne (Unil), Faculté de Biologie et de Médecine (FBM); Tel: +41-21-314-26-02; Fax: +41-21-314-11-99; e-mail: constantin.tuleasca@gmail.com;

Financial Disclosures: None

Dear Editor,

We read with great interest the recent article by Nisson⁶ et al., discussing the rare presence of spine arachnoid webs, while performing a systematic review of the literature, including 43 patients. Almost half of the cases presented symptoms for more than 1 year before surgery. Arachnoid web excision was performed in 86% of cases. Following surgery, 91% reported improvement in neurological symptoms. Several aspects warrant for further consideration in this uncommon spinal pathology.

Firstly, there is a need for a better preoperative diagnosis. The anamnesis might reveal red flags in the context, including trauma, infections, previous surgery and subarachnoid hemorrhage. Preoperative MRI could be inconclusive. First neuroimaging attempts have focused on the use of direct visualization of arachnoid membranes using 3D constructive interference in steady state (CISS) sequences⁵. More recently, cardiac-gated phase-contrast (CINE) MRI in multiple axial planes has been suggested, which usually correctly localizes the SAW and demonstrates a one-way valve like the flow of CSF because of the web³.

Secondly, the clinical and radiological diagnosis should exclude other pathologies, which might mimic an SAW, in particular thoracic intradural spinal cord herniation (ISCH). Both are uncommon conditions and can result in significant neurological morbidity if left untreated⁷. The main aspect is related to the evaluation of the nature of the dorsal cord indentation, as well as the integrity of ventral subarachnoid space at the level of spinal cord deformity. The «scalpel sign» is usually referred as pathognomonic for SAW, consisting in dorsal spinal cord indentation and anterior cord displacement. More recently, the «C-shape dorsal indentation» has been further reported for spinal cord herniation⁷. In practice, differentiating between these 2 separate entities is of crucial importance from a surgical perspective and not always straightforward preoperatively⁷. Their management is different. The SAW is treated by laminectomy with intradural lysis of adhesions and resection of arachnoid bands, while the ISCH repair includes reduction of the herniation and restoring of the spinal cord to its normal position.

Thirdly, if surgical exploration is performed for SAW with no major intraoperative findings, one should also consider ISCH and further explore the presence of such condition.

Fourthly, the «wait-and-scan» strategy versus proactive treatment is, in our experience, quite clear. Conservative treatment with clinical assessment and serial imaging can be offered to asymptomatic cases. At the opposite, in symptomatic patients, treatment is always surgical.

Lastly, some of the authors consider arachnoidolysis as curative². However, we do believe that there is a small risk of readhesion, which should be considered and explained to the patient preoperatively. Approaches with arachnoid dissection and decompression have been reported, including by our group¹. In this context, laminectomy and intradural exploration became a standard of care, although minimally invasive techniques were also recently described, and propose either expandable tubular retractors⁸ or fibroscopic visualization after classical arachnoid opening⁴.

The systematic review by Nisson et al.⁶ nicely fulfills a gap of knowledge, towards a better understanding of this uncommon entity which is the arachnoid web.

Funding: Constantin Tuleasca gratefully acknowledges receipt of a ‘Young Researcher in Clinical Research Grant’ (Jeune Chercheur en Recherche Clinique) from the University of Lausanne (UNIL), Faculty of Biology and Medicine (FBM) and the Lausanne University Hospital (CHUV)

Acknowledgments: Paris-South (CHU de Bicêtre) University Hospital.

Conflict of interest: No conflict of interest.

References:

1. Aghakhani N, Baussart B, David P, Lacroix C, Benoudiba F, Tadie M, et al: Surgical treatment of posttraumatic syringomyelia. **Neurosurgery** **66**:1120-1127; discussion 1127, 2010
2. Aiyer R, El-Sherif Y, Voutsinas L: Dorsal thoracic arachnoid web presenting as neuropathic pain: 'Scalpel' sign found on MRI. **Neuroradiol J** **29**:393-395, 2016
3. Chang HS, Nagai A, Oya S, Matsui T: Dorsal spinal arachnoid web diagnosed with the quantitative measurement of cerebrospinal fluid flow on magnetic resonance imaging. **J Neurosurg Spine** **20**:227-233, 2014
4. Fujiwara Y, Manabe H, Izumi B, Shima T, Adachi N: Microscope and Fiberscope-assisted Subarachnoid-Subarachnoid (S-S) Bypass: A Novel Surgical Technique to Reestablish Cerebrospinal Fluid Flow in Treating Dorsal Spinal Arachnoid Webs, Diagnosed by Cine-MRI. **Clin Spine Surg** **31**:58-64, 2018
5. Hirai T, Korogi Y, Shigematsu Y, Sugahara T, Takahashi M, Ushio Y, et al: Evaluation of syringomyelia with three-dimensional constructive interference in a steady state (CISS) sequence. **J Magn Reson Imaging** **11**:120-126, 2000
6. Nisson PL, Hussain I, Hartl R, Kim S, Baaj AA: Arachnoid web of the spine: a systematic literature review. **J Neurosurg Spine**:1-10, 2019
7. Schultz R, Jr., Steven A, Wessell A, Fischbein N, Sansur CA, Gandhi D, et al: Differentiation of idiopathic spinal cord herniation from dorsal arachnoid webs on MRI and CT myelography. **J Neurosurg Spine** **26**:754-759, 2017
8. Vergara P, Barone DG: Minimally Invasive Excision of Thoracic Arachnoid Web. **World Neurosurg** **109**:e81-e87, 2018