



HAL
open science

Development of a standardized framework for the evaluation of biodiversity in the context of biodiversity offsets

Lucie Bezombes, Stéphanie Gaucherand, Thomas Spiegelberger, Véronique Gouraud, Christian Kerbirou

► To cite this version:

Lucie Bezombes, Stéphanie Gaucherand, Thomas Spiegelberger, Véronique Gouraud, Christian Kerbirou. Development of a standardized framework for the evaluation of biodiversity in the context of biodiversity offsets. *Ecological Indicators*, 2018, 93, pp.1244-1252. 10.1016/j.ecolind.2018.06.027 . hal-02295898

HAL Id: hal-02295898

<https://hal.sorbonne-universite.fr/hal-02295898v1>

Submitted on 27 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

1 **Development of a standardized framework for the evaluation of biodiversity in the context of**
2 **biodiversity offsets.**

3

4 Lucie Bezombes (1, 2), Stéphanie Gaucherand (1), Thomas Spiegelberger (1), Véronique Gouraud (2)
5 Christian Kerbiriou (3, 4)

6

7 (1) Université Grenoble Alpes, Irstea, UR EMGR, 2 rue de la Papeterie-BP 76, F-38402 St-Martin-
8 d'Hères, France

9 (2) EDF R&D, Laboratoire National d'Hydraulique et Environnement, 6 Quai Watier, F-78400 Chatou,
10 France

11 (3) Muséum National d'Histoire Naturelle, Centre d'Ecologie et de Sciences de la Conservation,
12 UMR7204 MNHN-UPMC-CNRS, 55 Rue Buffon, F-75005 Paris, France

13 (4) Station de biologie marine, place de la croix, F-29900 Concarneau, France

14

15 **Corresponding author**

16 Lucie Bezombes, lucie.bezombes@irstea.fr, +33476762846

17

18 **Acknowledgments**

19 We are thankful to Marie-Eve Reinert for her useful comments on an earlier draft of this article. We are
20 grateful to the researchers from IRSTEA who participated in the working group on the framework
21 development. We also thank Agnes Barillier and Frederick Jacob for relevant comments and advice on
22 the choice of indicators, and all the organizations that collected the data on the study sites. This
23 research was financed by the French government "CIFRE" grant for PhD students and Electricité de
24 France (EDF).

25

26 **Key words**

27 Biodiversity offsets, biodiversity indicators, standardized framework, ecological equivalence, offset
28 policy, operationality, scientific basis, comprehensiveness

29 **Abstract**

30 In response to biodiversity erosion caused by human activities, biodiversity offsets are widely used to
31 compensate for negative impacts with an objective of “No Net Loss” (NNL). One major challenge is to
32 evaluate what biodiversity will be offset, and select the indicators based on which the losses of
33 biodiversity due to impacts and the gains due to offsets are calculated. In most European countries
34 and in France particularly, there is no recommended and available standardized method to evaluate
35 biodiversity for offsets. This leads to heterogeneous practices that do not guarantee achieving NNL. In
36 this paper we present the development of a practical framework for biodiversity evaluation adapted to
37 the European offset policy with French specificities. We follow four steps during which filters are
38 applied so that the framework is science-based, operational and comprehensive. First all stakeholders
39 involved were rallied around the scientific vision of biodiversity in order to integrate its complexity into
40 the framework (step 1). Then the European and French legislation requirements were identified (step
41 2) and a practical framework for biodiversity evaluation integrating the outcomes of step 1 and 2 was
42 developed (step 3). Finally, a relevant set of indicators within this framework was selected. The
43 resulting framework ensures that biodiversity is evaluated at a general level before focusing on the
44 species and habitat at stake and that the landscape context is taken into account. We argue that the
45 development of the framework is innovative and can be generalized to a wide range of situations.
46 Finally, we identify perspectives for losses, gains and equivalence calculation based on the
47 framework.

48

49	<u>Summary</u>	
50	1. Introduction	5
51	2. Step 1: Rallying all the stakeholders around the scientific vision of biodiversity	6
52	3. Step 2: Identifying the characteristics of the offset policy and the legislation	
53	requirements	6
54	3.1. <i>Mandatory</i>	7
55	3.2. <i>Not mandatory but recommended</i>	7
56	4. Step 3: Developing a practical framework for biodiversity evaluation	7
57	4.1. <i>What biodiversity components at stake does the framework target?</i>	8
58	4.2. <i>At what spatial scale is biodiversity evaluated?</i>	8
59	4.3. <i>What dimensions of biodiversity are taken into account within the framework?</i>	9
60	4.4. <i>How does the framework integrate non mandatory good practices and</i>	
61	<i>recommendations?</i>	9
62	5. Step 4: Prospecting and selecting a relevant set of indicators within the	
63	framework	10
64	5.1. <i>Prospection phase</i>	10
65	5.2. <i>Selection phase</i>	11
66	5.2.1. <i>Scientific basis filter: removal of redundant indicators</i>	11
67	5.2.2. <i>Operationality filter: taking into account temporal, technical and financial</i>	
68	<i>constraints</i>	11
69	5.2.3. <i>Final result of selection</i>	12
70	6. Discussion	12
71	6.1. Innovative aspects in the framework development	12
72	6.1.1. <i>Going beyond strict legal requirements</i>	12
73	6.1.2. <i>Non-aggregation of indicators values</i>	12
74	6.1.3. <i>Combining quantitative values and expert opinion</i>	13
75	6.2. The adaptability of the framework to a wide range of situations	13
76	6.3. A basis for losses, gains and equivalence assessment	14
77	Conclusion	15
78		
79		

80 **Figures and Tables**

81 **Figure 1:** Schematic representation of the steps followed in the approach and the related filters
82 applied. Dark grey boxes represent the filters and light grey boxes the different constituent parts of the
83 methodological framework.

84 **Figure 2:** a) Schematic and b) cartographic representation of the practical framework of biodiversity
85 evaluation. Biodiversity is evaluated at two levels (General and Specific, declined in species and
86 habitats) and two spatial scales: Large Scale (grey boxes) and Site scale (white boxes). The perimeter
87 of the Larger Scale is defined at each level.

88

89 **Table 1:** Specificities of French offset policy in the European context.

90 **Table 2:** Criteria used within the framework for biodiversity evaluation.

91 **Table 3:** Details on the study sites used to test the correlations between indicators.

92 **Table 4:** Framework structure (Levels, Scales, Criteria and Indicators).

93

94 **Supplementary material**

95 **Appendix 1:** List of all preselected indicators and details of the selection for the final set of indicators.

96 **Appendix 2:** Correlation matrix used in the indicator selection phase.

97 **Appendix 3:** Qualitative features used as an operationality filter for indicator selection.

99 Biodiversity offsets are currently widely used to counteract impacts from development (Pilgrim *et al.*
100 2013). It consists in compensating biodiversity losses with equivalent gains provided by offset
101 measures, in order to achieve “no net loss” (NNL) of biodiversity (ten Kate *et al.* 2004). Implementing
102 biodiversity offsets implies that the initial state of biodiversity on both impacted and compensatory
103 sites must be evaluated. Consequently, indicators have to be chosen as surrogates for biodiversity,
104 based on which ecological equivalence between losses and gains can be calculated to demonstrate
105 NNL (Quétier & Lavorel 2011).

106 Depending on the offset policy, indicators would target different biodiversity components according to
107 local and national conservation issues (McKenney & Kiesecker 2010). In the USA, wetlands are
108 mitigated under the Clean Water Act (1972), and indicators are usually related to wetland functions
109 (State of Florida 2004; Levrel *et al.* 2015). In the state of Victoria in Australia, the Native Vegetation
110 Act (2003) imposes compensation for endemic forest, thus indicators represent vegetation cover and
111 structure (Parkes *et al.* 2003). In Europe, offset policies are complex and have been built up since the
112 1970’s based on early conservation principles (Quétier *et al.* 2014). Biodiversity offsets are currently
113 required for several biodiversity components, for which conservation outcomes are stipulated in
114 different legislations at both European and member states level. Offset measures must be
115 implemented when significant residual impacts (those remaining after avoidance and reduction) occur
116 on those biodiversity components.

117 Despite the recent enactment of new laws in European countries strengthening offset objectives and
118 implementation (e.g., in France the Recapture of Biodiversity and Landscapes law in 2016), in some
119 countries there are still no recommended and available standardized indicators or methods to size
120 offsets (e.g., France, UK, Spain, Italy). Only a few methods are starting to emerge in these countries
121 (DREFA 2013; Gayet *et al.* 2016). Therefore, there exists a gap between on one side offset policy
122 requirements and scientific recommendations, and on the other side usual practices. This makes it
123 problematic to achieve the NNL of specific target biodiversity components (Regnery *et al.* 2013).

124 In the context of biodiversity offsetting, the biodiversity components at stake (i.e. that must be offset)
125 are usually selected based upon anthropogenic considerations to address patrimonial issues (Regnery
126 *et al.* 2013) but can also address ecological issues such as ecosystem functionalities (Levrel *et al.*
127 2012). However, how these components at stake should be evaluated remains a key challenge. The
128 main reason is that biodiversity is multidimensional and can be observed at several organizational,
129 spatial and temporal scales (Noss 1990), which makes its evaluation complex. Moreover, biodiversity
130 offsets are implemented in a context of development projects where the evaluation of biodiversity is
131 subject to technical, financial and temporal constraints (Cuperus *et al.* 2001; Quigley & Harper 2006).

132 Standardized frameworks could improve the evaluation of biodiversity and the assessment of
133 ecological equivalence between losses and gains, with quantitative indicators that better combine
134 legislation requirements (McKenney & Kiesecker 2010) and biodiversity complexity (Gibbons &
135 Lindenmayer 2007; Gardner *et al.* 2013). Based on this observation, we developed a standardized
136 framework for a practical evaluation of biodiversity in the context of biodiversity offsets. The aim of this
137 paper is to present the development this framework step by step. We proceed in four steps: (1) rallying

138 all the stakeholders around the scientific vision of biodiversity; (2) identifying the characteristics of the
139 offset policy and the legislation requirements; (3) developing a practical framework for biodiversity
140 evaluation; and (4) prospecting for and selecting a relevant set of indicators within the framework,
141 based on data collected on eight sites in France, that happen to be used in an offsets context.
142 Following these four steps, different filters were applied (Fig. 1) to ensure that the framework is
143 operational, science-based and comprehensive, which are necessary attributes for the framework to
144 enable achieving NNL (Bezombes *et al.* 2017). The framework is adapted to the European context,
145 including specificities of the French legislation but could be generalized to other contexts as we
146 discuss it hereafter.+

147 **2. Step 1: Rallying all the stakeholders around the scientific vision of biodiversity**

148 The first step consisted in rallying all stakeholders involved in biodiversity offsets around the scientific
149 vision of what biodiversity is and how it should be evaluated. It aims to broaden offsets beyond the
150 strict box-ticking exercise of offset requirements. Although it can seem trivial, this step is crucial as in
151 the context of development projects, biodiversity issues are often considered by companies only as
152 legislative constraints. This first step should be a good opportunity to open up a dialogue between all
153 stakeholders and spread knowledge about biodiversity toward the ones not usually directly involved in
154 biodiversity matters.

155 Since biodiversity is multidimensional, it is important to conceptualize its intrinsic complexity as an
156 explicit common basis for its evaluation, which should enlarge the required foci for offsets. This should
157 increase the project efficiency by avoiding wasting time and money on doubtful evaluation of
158 biodiversity and give a better advertising for developers by improving their image with the society.

159 In our case, all stakeholders involved in the framework development (scientists, practitioners,
160 members of environmental authorities) agreed on the definition of biodiversity given by the Convention
161 of Biological Diversity (CBD Secretariat 1992): biodiversity is the diversity between organisms and
162 their interactions at different levels of organization, from genes to ecosystems. We completed this
163 definition with the possibility to evaluate biodiversity through three primary dimensions: the
164 composition, structure and function, as recommended by Noss (1990). These dimensions should be
165 evaluated at both spatial and temporal scales of biodiversity. Indeed, spatial patterns drive the
166 interactions between organisms at each level of organization (e.g., the flow of species from one
167 favorable habitat to another through ecological corridors). There are also temporal successions of
168 fauna and flora from the pioneer to the mature state of an ecosystem whose dynamic is influenced by
169 the perturbation regime. We agreed that evaluating biodiversity according to the vision integrating all
170 the above-mentioned elements should capture biodiversity in its complexity.

171 **3. Step 2: Identifying the characteristics of the offset policy and the legislation** 172 **requirements**

173 The second step consists in identifying the characteristics of the offset policy and the legislation
174 requirements that determine a set of biodiversity components at stake that have to be offset. This
175 involves analyzing what is mandatory and what is not mandatory but considered as good practices

176 (that contribute to improving biodiversity offsetting, and in turn the acceptance of project
177 developments; BBOP 2012b).

178 **3.1. Mandatory**

179 In Europe, species protection is based on the EU Habitats Directive and the EU Birds Directive (EEC
180 1992, 2009). The protection of these species (with particularly declining population dynamics for most
181 of them; Donald *et al.* 2007) prohibits the destruction or alteration of their habitats. When developers
182 cannot avoid negative impacts on species or their habitat (considering the economic and social issues
183 of development projects), derogation requests to legal protection can be attributed under articles 9 and
184 16 of the EU Habitats and Birds Directives. According to these articles, projects of development may
185 be authorized under the condition that offset measures maintain or restore a favorable conservation
186 status of the species populations concerned (Table 1). Thus, to effectively conserve these populations,
187 offset measures have to target the same species that are impacted (like for like).

188 In addition to the European laws, in France, Environmental Impact Assessment (EIA) procedures are
189 mandatory for projects defined in article R.122-2 of the Reform of the EIA Decree (2012). In these
190 cases, the mitigation hierarchy must be applied and any significant residual impacts on woodlands,
191 wetlands, protected species and protected natural habitats (at the European and national scales) have
192 to be compensated by appropriate offset measures (Table 1) because they are considered as
193 biodiversity components at stake (Quetier *et al.* 2014).

194 **3.2. Not mandatory but recommended**

195 In addition to these mandatory requirements, a French consultative process called “Grenelle de
196 l’Environnement” has also encouraged developers since 2007 to evaluate the impacts of their projects
197 on landscape level connectivity. Moreover, the EU 2020 Biodiversity Strategy requires that
198 ecosystems and their services be maintained and enhanced by establishing green infrastructure
199 (Commission to the European Parliament 2014). Also, other areas classified as natural areas of
200 ecological, zoological and floristic value (ZNIEFF) are considered as patrimonial areas without
201 mandatory protection, but the jurisprudence confirms that their presence must be taken into account in
202 EIA.

203 In a general way, increasing recommendations from scientists stipulate that offset measures should
204 focus on ecosystem functionalities. In practice, impacts on “ordinary biodiversity” (i.e., biodiversity
205 components not considered to be at stake) are usually declared as nonsignificant, even though
206 ordinary biodiversity supports ecosystem functionalities (Smith & Knapp 2003; Solan *et al.* 2004) and
207 is declining (Julliard *et al.* 2004; Jiguet *et al.* 2010).

208 **4. Step 3: Developing a practical framework for biodiversity evaluation**

209 The objective of the third step is to develop a practical framework for biodiversity evaluation taking into
210 account the complexity of biodiversity (step 1) but also adapted to a particular offset policy with
211 legislation requirements (step 2, Fig. 1). Those brought together make the framework comprehensive.
212 The elaboration of the framework was guided by the four following questions.

213 **4.1. What biodiversity components at stake does the framework target?**

214 To be operational, the framework should first focus on biodiversity components that must be offset by
215 legislation. In the French offset policy, EIA procedures ensure that a diagnosis of the biodiversity found
216 and possibly impacted is carried out in order to detect the components at stake that are required to be
217 offset in both European and French offset policies (see Step 2). The problem is that currently, these
218 components are evaluated quite heterogeneously in EIA since there is no standardize framework
219 recommended by authorities (Bigard *et al.* 2017). Therefore, to fit all procedures (Table 1) in a more
220 standardized and reproducible way, we suggest evaluating biodiversity at two distinct levels (general
221 and specific; see Fig. 2a):

222 -the general level (GL) answer the question “what biodiversity is found on impacted and compensatory
223 sites? “The biodiversity found is evaluated in a general way, without focusing on a particular species or
224 natural habitat. The ordinary (supporting ecosystem functionalities) and patrimonial biodiversity (i.e.
225 protected, rare, endangered or emblematic; Delzon *et al.* 2013) can be diagnosed in EIA with this
226 level. This level should be used for every site evaluation.

227 -the specific level (SL), which is further subdivided into species level (spL) and habitat level (hL). It put
228 the focus on species or habitats at stake detected at the GL, answering the question “what is the state
229 of the species or habitats on the sites (species needs, population viability, threats, habitat structure
230 and maturation etc.)? Within the European offset policy, the spL can be used to evaluate species
231 protected under the EU Habitats and Birds Directives, and the hL for natural habitats such as wetlands
232 (French specificities). This level should be used for sites where species or habitats at stake are
233 identified.

234

235 **4.2. At what spatial scale is biodiversity evaluated?**

236 In Europe, only significant residual impacts on important ecological corridors tend to be compensated,
237 but there is no obligation to integrate landscape context (general term used here to describe the
238 surrounding environment that is several kilometers around the impacted or compensatory sites) in EIA.
239 However, taking into account landscape context for linear infrastructures such as roads and railways is
240 crucial because of their highly fragmenting effects over long distances (Cuperus *et al.* 2001; Jones *et*
241 *al.* 2014; Mimet *et al.* 2016). The major difficulty in evaluating the value of the impacted or
242 compensatory site in the landscape context is the lack of available data, which makes the assessment
243 of losses and gains in the landscape context quite sensitive.

244 Therefore, we suggest evaluating not only biodiversity on the impacted or compensatory site area, but
245 also at a broader scale. To do this in a standardized way, for each of the three levels (GL, sL and hL),
246 we suggest that biodiversity is evaluated at two spatial scales, corresponding to different perimeters
247 (see Fig. 2b):

248 -the Site Perimeter (SP), which is the perimeter delimited by the developer around the impacted and
249 compensatory sites. Biodiversity is usually evaluated ate this scale in EIA. The direct influence of

250 impacts (e.g., destruction of 1 ha of wetland) or offset measures (e.g., restoration of 2 ha of wetlands)
251 can be observed at this scale, as well as species perturbation.

252 -the Expanded Perimeter (EP), which is an area surrounding the sites (Fig. 2b). This perimeter is
253 defined according to various issues: the size and shape of impacted and compensatory sites (at the
254 GL), the sites location in the catchment (at the hL), and dispersal capacity or vital area of the species
255 at stake (at the spL). The EP does not properly correspond to a landscape context because depending
256 on the project it would cover a very wide area that is too time-consuming to prospect. Evaluation at
257 this scale is a proxy for landscape level phenomena such as connectivity between favorable habitats
258 for a species, which can be fragmented by development projects or restored with offset measures.

259 **4.3. What dimensions of biodiversity are taken into account within the framework?**

260 As mentioned in step 1, we consider three main dimensions to evaluate biodiversity: composition,
261 structure and function (Noss 1990). Depending on the objectives on conservation outcomes in offset
262 policy, some dimensions seem more appropriate to evaluate the biodiversity components at stake than
263 others. For example, in the *Fish Habitat* approach (Harper & Quigley 2005), the composition, structure
264 and function of lacustrine habitats are evaluated, given that the offset policy aims to maintain lake fish
265 productivity. In Europe, as there are several objectives for biodiversity in European and French
266 legislation depending on the components considered (Table 1), we estimated that all three dimensions
267 are necessary to evaluate biodiversity in a comprehensive way.

268 To facilitate the understanding of the framework without in-depth knowledge in ecology, we broke
269 down the compositional, structural and functional dimensions of biodiversity into seven criteria more
270 often used in EIA: Diversity, Patrimonial Status, Representativeness, Vegetation structure,
271 Connectivity, Functionalities and Pressure (see Table 2 for definitions and references). Most of the
272 criteria are related to ecological processes, except for the “Patrimonial Status” criterion, which refer to
273 anthropogenic conservation choices.

274 According to Noss (1990), the term “function” involves ecological and evolutionary processes,
275 including gene flow, disturbances and nutrient cycling. Therefore, we related it to the “Pressure” and
276 “Functionalities” criteria. The “Functionalities” criterion refers to four main processes identified as the
277 most relevant for offset policy objectives: (i) capacity of reproduction of species (ii) population
278 specialization, (iii) soil quality, and (iv) maturity of the vegetal succession (Table 2).

279 **4.4. How does the framework integrate non mandatory good practices and recommendations?**

280 Some practices and recommendations may not be enacted in the legislation, but still contribute to
281 effective biodiversity offsetting and acceptance of the project. They can be mentioned in guidelines
282 recognized by the offset policy (e.g., Cuperus *et al.* 1999; BBOP 2012a, b; CGDD 2013), they can be
283 encouraged by expert panels evaluating and advising legal authorities on the application of the
284 mitigation hierarchy (e.g., National Committee for Nature Protection (CNPN) in France) and they can
285 emerge from scientific research (Evans *et al.* 2015; Gelcich *et al.* 2016).

286 In France, the obligation of achieving NNL is currently legislated, but no method detailing how to
287 measure it is imposed. Therefore biodiversity offsetting is based to a large extent on
288 recommendations, notably:

289 -Offsetting impacts on connectivity are recommended at both the European and national levels (EU
290 Strategy on Green Infrastructure and Grenelle de l'Environnement). Our framework integrates an
291 evaluation of "Connectivity" (Table 2), from a structural approach.

292 -Scientific research recognizes that the evaluation of biodiversity in terms of species lists or habitats is
293 not sufficient to evaluate losses and gains, and urges countries to assess impacts on species and
294 habitats also in terms of their functionalities. Our framework integrates some of these functionalities
295 within a dedicated criterion (Table 2).

296 **5. Step 4: Prospecting and selecting a relevant set of indicators within the framework**

297 The fourth step of the approach consists in first prospecting and then selecting a relevant set of
298 indicators for biodiversity evaluation, as a basis for the calculation of biodiversity losses and gains. In
299 the framework presented herein, indicators should aim to characterize the seven criteria chosen for
300 the evaluation of biodiversity as best possible (Table 2). As a multitude of indicators is currently
301 available, during this step we applied filters to reduce the high number of potential indicators (Heink &
302 Kowarik 2010). Two main filters can be applied (Fig. 1) so that the indicators selected are both
303 science-based and operational i.e., adapted to the temporal, technical and financial constraints that
304 developers must contend with.

305 **5.1. Prospection phase**

306 During the prospection phase, an exhaustive list of indicators was created for assessing each criterion
307 defined in step 3. We applied the scientific basis filter by investigating indicators validated in scientific
308 publications to ensure that indicators have been tested and approved as valid surrogates of the target
309 elements (e.g., Gibbons & Freudenberger 2006; Delzons *et al.* 2013). It is crucial that the indicators
310 prospected respond to disturbances, anthropogenic stresses and changes over time in a predictable
311 manner and that there is low variability in this response (Dale & Beyeler 2001).

312 In the context of offsets, the changes refer to development projects impacts and offset measures
313 benefits on biodiversity. In European countries, development projects concern mainly urbanization,
314 linear infrastructures, quarries, touristic activities (ski resorts, coastal development), industrial
315 activities...; and offset measures that are accepted consist in ecosystem restoration, creation or
316 maintenance (with preference when there is legal protection; Jacob *et al.* 2014). We searched only for
317 indicators with quantitative indicators, excluding indirect notations relative to qualitative characteristics,
318 in order to keep the evaluation direct and transparent.

319 The list of the preselected indicators after the prospection phase is detailed in Appendix 1. A total of
320 170 indicators were preselected (86 for GL, 57 for hL and 27 for sL). Indicators at specific levels were
321 classified into the indicators adapted in every situation (i.e., for the evaluation of all habitats or
322 species), and those adapted only for one type of natural habitat or species (e.g., the number of dead
323 trees for forest habitats).

324 **5.2. Selection phase**

325 During the selection phase, a final set of most relevant indicators for biodiversity evaluation is selected
326 out of the exhaustive list, by applying the scientific basis and operationality filters (Fig. 1). In a
327 preliminary step, we identified the indicators usually found in current procedures (see Appendix 1) in
328 order to give them priority in the selection of the final set of indicators, as, environmental authorities
329 assessing procedures expect certain information.

330 **5.2.1. Scientific basis filter: removal of redundant indicators**

331 To obtain a final set of indicators without redundancy in the information they provide, we analyzed the
332 correlations between indicators. In theory, they should be analyzed for all preselected indicators, but to
333 illustrate this process we restrained our analysis only to indicators from the GL (see Appendix 1).

334 The data used were collected on eight sites in France: a mitigation banking experiment in the
335 Belledonne mountains (French Alps) in the department of Isère; a compensatory site located on the
336 upper Rhine River, in the department of Haut Rhin; four impacted sites along the Romanche River and
337 two related compensatory sites downstream on the same river also in the department of Isère (see
338 Table 3 for details). All data refer to the initial states of biodiversity before impacts or offset measures
339 and various types of data were used: inventories, GIS, public data and field prospection.

340 This selection is intended to remove the indicators highly correlated to others from the exhaustive list
341 (based on Spearman rank correlation; high correlation being considered as $R^2 > 0.6$, $P < 0.05$) in order to
342 obtain a minimum set of independent indicators. The selection proceeded in three stages: a) analyzing
343 correlations between indicators not usually used in procedures; (b) analyzing correlations between
344 indicators that are usually used in procedures (from mandatory requirements) and c) analyzing
345 correlations between the indicators selected in a) and b). The three correlation matrices are presented
346 in Appendix 3. Approximately half of the preselected indicators tested were removed (39 out 75; see
347 Appendix 1) because they were highly correlated with other indicators.

348 **5.2.2. Operationality filter: taking into account temporal, technical and financial constraints**

349 We applied this filter to all three levels. Since the practical framework for biodiversity evaluation should
350 remain operational in the context of biodiversity offsets, we passed all indicators (from General and
351 Specific Levels) through an operationality filter. It aims to exclude indicators that would not be
352 appropriate in a context driven by temporal (data collection could take several years), technical
353 (software for which skills are not yet common among practitioners, e.g., “Graphab” for landscape
354 connectivity evaluation; Foltête *et al.* 2012) and financial (expensive material) constraints.

355 We categorized operationality into three features related to the above-mentioned constraints
356 (Appendix 3) based on the authors’ expertise of procedures: estimated time of data collection
357 (temporal), type of skills needed to fill in the indicator (technical) and price of the data collection
358 needed to fill in the indicator price (financial). Depending on the features’ modalities, three levels of
359 operationality were assigned to indicators: low, medium or high. The overall level of operationality was
360 evaluated as the mean levels of each feature. Indicators with an overall “low” operationality were
361 removed first, but indicators with a medium or high level of operationality were also removed if they did

362 not pass the scientific basis filter (five out of 81 for GL, 13 out of 56 for hL and four out of 27 for sPL;
363 see Appendix 1).

364 5.2.3. *Final result of selection*

365 The final set of indicators organized by levels, scales and criteria is summarized in Table 4 (41
366 indicators in the GL, 43 in the HL and 23 in the sL). This constitutes a standardized framework for
367 biodiversity evaluation adapted to the European context constructed following our methodological
368 approach.

369 6. Discussion

370 Following a step by step approach, we constructed a standardized framework for biodiversity
371 evaluation, adapted to the European and French offsets context and combining scientific basis,
372 operationality and comprehensiveness. In this section we discuss the choices we made for the
373 development of such framework and their implication in biodiversity offsetting. We then identify
374 perspectives for equivalence assessment.

375 6.1. Innovative aspects in the framework development

376

377 6.1.1. *Going beyond strict legal requirements*

378 The framework enables a general diagnosis of the biodiversity found on the sites, before making a
379 focus on components considered at stake that are required to be offset by the French legislation. Thus
380 the "ordinary" biodiversity (Doremus 2001) is visible and so will be the possible benefit from offset
381 measures, as increasingly recommended by the scientific sphere (Brownlie & Botha 2009; BenDor &
382 Stewart 2011; Moreno-Mateos *et al.* 2015). Moreover, evaluating biodiversity in an Expanded
383 Perimeter (EP) instead of restricting the evaluation to the Site Perimeter (SP) allows to contextualizes
384 the assessment in regard to the surrounding landscape, and particularly ecological corridors that
385 contribute to the success of offsets (Kiesecker *et al.* 2009; Bull *et al.* 2013). In a general way, the
386 framework places great emphasis on the evaluation of the functionalities, with a dedicated criteria, to
387 preserve both target components at stake (species and habitat) and the ecosystem's self-preservation
388 capacity in uncertain future conditions (Naeem *et al.* 2012).

389

390 6.1.2. *Non-aggregation of indicators values*

391 For more transparency and coherence in the evaluation, we chose not to aggregate indicators into
392 notes per criterion or even an overall "score of biodiversity" for an entire site (Maseyk *et al.* 2016).
393 Even though an aggregation of indicators is easily interpretable for decision making, composite indices
394 can lead to simplistic conclusions, some dimensions may be hidden by others and the necessary
395 weighting of indicators is often arbitrary (Nardo *et al.* 2005).

396 With a non-aggregation, biodiversity components showing high or low values could be detected more
397 easily and the consequences of actions on these components would be directly visible. For example,
398 impacts should be avoided on site showing an important "Patrimonial Status" revealing the presence
399 of numerous species or habitats at stake. Also, some actions can favor a species with a high

400 patrimonial value but have a negative effect on other more ordinary biodiversity components.
401 Therefore, these inevitable trade-offs have to be visible in order to inform properly decision-makers.
402 The difficulty is to make such a large amount of information intelligible; therefore the visualization of
403 indicators is all the more important for their understanding and interpretation by stakeholders.

404 6.1.3. *Combining quantitative values and expert opinion*

405 The standardization of the framework makes the evaluation of biodiversity operational and efficient
406 (Laycock *et al.* 2013). It ensures that the essential components are evaluated in a quantitative way
407 and better exploits efforts already made, for data collection notably. Within the standardized
408 framework however, expert opinion is needed to fill in certain indicators in place of precise indicators
409 not available considering the requirements described in step 4.

410 This concerns mainly the indicators within the species level (spL) such as “the area of favorable
411 habitat” (on the site before and after impacts or offset measures). Instead of analyzing each habitat
412 feature for all potential species at stake in a standardized way (over 900 species are protected under
413 the EU Habitats Directive 92/43/EEC; EEC 1992), we enabled experts to decide on the favorability
414 aspect of the habitat (i.e. what surface of the site does gather all vital elements for the target
415 species?). The result is still a quantitative value (e.g. hectares). Nonetheless, experts should argue
416 the attribution of the indicator values with documentation in order that the framework outputs remain
417 transparent.

418

419 **6.2. The adaptability of the framework to a wide range of situations**

420 The framework we developed can be used for the evaluation of the biodiversity of a wide range of
421 ecosystems and respond to the European and French offset policy requirements. It is designed to be
422 applied on a case by case basis because in Europe and France it is the most common approach.

423 The strength of this framework is that it can also be adapted to other offset policies that claim different
424 objectives and are constructed upon different conservation issues. Indeed, the changes would
425 concern the set of indicators related to the biodiversity targeted by a given legislation or at stake in a
426 given country that do not exist in the European context. We detail some adaptation hereafter
427 according to:

428 -The *offset policy*. This framework focuses on species and habitats at stake in response to the
429 importance given by EU and French offset policies in protecting species and habitat conservation. In
430 the USA for example, wetland mitigation represents a major part in offset policy (Madsen *et al.* 2010)
431 and therefore the framework would probably focus on a specific level for wetlands or a comprehensive
432 set of indicators dedicated to wetlands evaluation within the habitat level. In the UK there is willingness
433 from the government to preserve ordinary biodiversity (with less focus on biodiversity at stake; DREFA
434 2013), which could be evaluated only with an extended General level (GL).

435 -The *naturalness of ecosystems*. In European countries, biodiversity has evolved with human activities
436 for centuries (Doxa *et al.* 2012) and only very few remaining ecosystems can be considered to have
437 high naturalness (e.g., some Mediterranean primary forests, peatlands). In regions where wilderness
438 areas never impacted by human activities still exist (e.g., tropical forests in Brazil), the framework

439 should enable the evaluation of this naturalness, with a dedicated criterion for example. Moreover,
440 such cases should be more favorable (less subject to debates) for comparison of the selected
441 indicators to a reference state (i.e., high naturalness). Indicators would be relative rather than absolute
442 indicators (e.g., values expressed as percentages of the benchmark; Parkes *et al.* 2003).

443 -The *type of ecosystems*. The framework is adapted to terrestrial biodiversity (including wetlands) and
444 at the moment does not include indicators specific to aquatic or marine ecosystems. The indicators
445 would necessarily be different since these ecosystems have different properties than the terrestrial
446 ones. They notably have high temporal variability due to frequent hydrological disturbances (floods,
447 low water) (Resh *et al.* 1988; Poff *et al.* 1997), very pronounced longitudinal structuring (Kondolf *et al.*
448 2003) and higher consumer biomass than primary producers (unlike terrestrial ecosystems).

449 -The *spatial scales*. In Europe, the high population density of countries (the urbanization rate is 75%
450 according to the European Commission), and the low amount of remaining pristine habitat, influences
451 land use which becomes very strategic. Land use is managed on smaller areas compared to countries
452 like the USA, Australia, Brazil or Canada. In those areas, the Expanded Perimeter could be more
453 related to an actual landscape scale since species dynamics can be observed over wide areas
454 (Hanski 1998).

455

456 **6.3. A basis for losses, gains and equivalence assessment**

457 The indicators suggested in the framework are the basis for the assessment of the ecological
458 equivalence between biodiversity losses and gains. For this purpose, each relevant indicator should
459 be assessed four times: before impacts and offset measures (initial state of impacted and
460 compensatory sites) and after impacts and offset measures (predicted state of the sites; Bezombes *et al.*
461 2017). Consequently, losses and gains can be calculated for each indicator as the predicted values
462 minus the initial values. The ecological equivalence is logically reached when the values of losses
463 equal the values of gains. To complete these steps, the two following aspects would still need to be
464 worked on.

465

466 6.3.1. *Loss and gain predictions*

467 The ecological equivalence has to be assessed early in the project development process, in order that
468 authorities evaluate whether the design of offset measures can achieve NNL. This timing implies that
469 the values of indicators after impacts and offset measures are predictions that have to be made based
470 on the impacted and compensatory sites' initial state. The use of modelling to evaluate "natural"
471 ecosystem trajectories, i.e. without impacts or offset measures (Stringham *et al.* 2003), is an accurate
472 way of calculating losses and gains but in practice experts involved do not have access to such
473 technics (Sunderland *et al.* 2009). In any cases, the predictions are accompanied by a certain amount
474 of uncertainties (Moilanen *et al.* 2009). Predicted values of indicators should therefore be modulated
475 depending on the level of this uncertainty, which could be determined according to the target species

476 or habitat (Tischew *et al.* 2010), the type and intensity of the impacts (temporary or permanent), the
477 type of offset measures (Anderson 1995) etc.

478

479 6.3.2. *Equivalence assessment*

480 As we did not aggregate the indicators within the framework, the balance between losses and gains
481 must be observed for each indicator individually. In an ideal situation, every indicator would be at least
482 balanced or show a net gain, and consequently ecological equivalence would be reached for the
483 whole project. In practice however, some indicators will be balanced (gain equals loss), while others
484 will probably show a net gain or a net loss (Bull *et al.* 2014). To determine whether equivalence is
485 reached overall, we suggest ranking indicators according to the biodiversity issues and the offset
486 policy requirements (should we prioritize the conservation of natural areas, rare species, landscape
487 fragmentation...; Brooks *et al.* 2006). Thus, equivalence (or net gain) should be reached at least for all
488 “priority” indicators, i.e., the ones identified to represent the main biodiversity issues. The prioritization
489 of indicator will certainly vary from a project to another, but some issues may be common to an entire
490 territory.

491

492 **Conclusion**

493 Using the framework we present in this paper should ensure that the evaluation of biodiversity on
494 impacted and compensatory sites is science-based, operational and comprehensive. This framework
495 can provide an objective basis for discussion on biodiversity offset management and should be shared
496 by the different stakeholders involved.

497 **References**

- 498 Anderson P. (1995). Ecological restoration and creation: a review. *Biological Journal of the Linnean*
499 *Society*, 56, 187-211.
- 500 European Economic Community (EEC) (1992). Council Directive 92/43/EEC of 21 May 1992 on the
501 conservation of natural habitats and of wild fauna and flora. *Official Journal of the European*
502 *Union*, 206, 7-50.
- 503 European Economic Community (EEC) (2009). Directive 2009/147/EC of the European Parliament and
504 of the Council of 30 November 2009 on the conservation of wild birds on the conservation of
505 wild birds (codified version). *Official Journal L20*, 7–25.
- 506 Bekessy S.A., Wintle B.A., Lindenmayer D.B., McCarthy M.A., Colyvan M., Burgman M.A. &
507 Possingham H.P. (2010). The biodiversity bank cannot be a lending bank. *Conservation*
508 *Letters*, 3, 151-158.
- 509 BenDor T. & Stewart A. (2011). Land Use Planning and Social Equity in North Carolina's
510 Compensatory Wetland and Stream Mitigation Programs. *Environmental Management*, 47,
511 239-253.
- 512 Bezombes L., Gaucherand S., Kerbiriou C., Reinert M.-E. & Spiegelberger T. (2017). Ecological
513 equivalence assessment methods: what trade-offs between operationality, scientific basis
514 and comprehensiveness? *Environmental Management*.
- 515 Bigard C., Pioch S. & Thompson J.D. (2017). The inclusion of biodiversity in environmental impact
516 assessment: Policy-related progress limited by gaps and semantic confusion. *J. Environ.*
517 *Manage.*, 200, 35-45.
- 518 Brooks T.M., Mittermeier R.A., da Fonseca G.A., Gerlach J., Hoffmann M., Lamoreux J.F., Mittermeier
519 C.G., Pilgrim J.D. & Rodrigues A.S. (2006). Global biodiversity conservation priorities. *Science*,
520 313, 58-61.
- 521 Brownlie S. & Botha M. (2009). Biodiversity offsets: adding to the conservation estate, or 'no net
522 loss'? *Impact Assessment and project appraisal*, 27, 227-231.
- 523 Bull J.W., Milner-Gulland E.J., Suttle K.B. & Singh N.J. (2014). Comparing biodiversity offset
524 calculation methods with a case study in Uzbekistan. *Biological Conservation*, 178, 2-10.
- 525 Bull J.W., Suttle K.B., Singh N.J. & Milner-Gulland E.J. (2013). Conservation when nothing stands still:
526 moving targets and biodiversity offsets. *Front. Ecol. Environ.*, 11, 203-210.
- 527 Business and Biodiversity Offsets Programme (BBOP) (2012a). Resource Paper: No Net Loss and
528 Loss-Gain Calculations in Biodiversity Offsets. *BBOP, Washington, D.C.*
- 529 Business and Biodiversity Offsets Programme (BBOP) (2012b). Standard on Biodiversity Offset. *BBOP,*
530 *Washington, D.C.*
- 531 Business and Biodiversity Offsets Programme (BBOP) (2014). Working towards NNL of Biodiversity
532 and Beyond: Strongman Mine – A Case Study. *BBOP, Washington, D.C.*
- 533 CBD Secretariat (1992). Convention on biological diversity. In: *Convention on Biological Diversity*.
- 534 Commissariat Général du Développement Durable (CGDD). (2013). Lignes directrices nationales sur la
535 séquence éviter, réduire et compenser les impacts sur les milieux naturels. Collection «
536 Références » du Service de l'Économie, de l'Évaluation et de l'Intégration du Développement
537 Durable (SEEIDD) du Commissariat Général au Développement Durable (CGDD).
- 538 Commission to the European Parliament, the European Economic and Social Committee and the
539 Committee of the Regions (2014). *Communication from the Commission to the European*
540 *Parliament, the Council, the European Economic and Social Committee and the Committee of*
541 *the Regions. Our life insurance, our natural capital: an EU biodiversity strategy to 2020*
542 *Bryssel.*
- 543 Cuperus R., Bakermans M.M.G.J., Udo de Haes H.A. & Canters K.J. (2001). Ecological compensation in
544 Dutch highways planning. *Environmental Management*, 27, 75-89.
- 545 Cuperus R., Canters K.J., Udo de Haes H.A. & Friedman D.S. (1999). Guidelines for ecological
546 compensation associated with highways. *Biological Conservation*, 90, 41-51.

547 Curran M., Hellweg S. & Beck J. (2013). Is there any empirical support for biodiversity offset policy?
548 *Ecological Applications*, 24, 617-632.

549 Dale V.H. & Beyeler S.C. (2001). Challenges in the development and use of ecological indicators.
550 *Ecological indicators*, 1, 3-10.

551 Delzons O., Gourdain P., Siblet J.-P., Touroult J., Herard K. & Poncet L. (2013). L'IQE: Un indicateur de
552 biodiversité multi-usages pour les sites aménagés ou à aménager. *Revue d'écologie*, 68, 105-
553 119.

554 Department for Environment Food & Rural Affairs (DREFA) (2013). Biodiversity offsetting in England
555 green paper.

556 Donald P.F., Sanderson F.J., Burfield I.J., Bierman S.M., Gregory R.D. & Waliczky Z. (2007).
557 International conservation policy delivers benefits for birds in Europe. *Science*, 317, 810-813.

558 Doremus H. (2001). Biodiversity and the Challenge of Saving the Ordinary. *Idaho L. Rev.*, 38, 325.

559 Doxa A., Paracchini M.L., Pointereau P., Devictor V. & Jiguet F. (2012). Preventing biotic
560 homogenization of farmland bird communities: the role of High Nature Value farmland.
561 *Agriculture, Ecosystems & Environment*, 148, 83-88.

562 Evans D.M., Altwegg R., Garner T.W.J., Gompper M.E., Gordon I.J., Johnson J.A. & Pettorelli N. (2015). Biodiversity offsetting: what
563 are the challenges, opportunities and research priorities for animal conservation? *Animal
564 Conservation*, 18, 1-3.

565 Fennessy M.S., Jacobs A.D. & Kentula M.E. (2007). An evaluation of rapid methods for assessing the
566 ecological condition of wetlands. *Wetlands*, 27, 543-560.

567 Foltête J.-C., Clauzel C. & Vuidel G. (2012). A software tool dedicated to the modelling of landscape
568 networks. *Environmental Modelling & Software*, 38, 316-327.

569 Gardner T.A., Von Hase A., Brownlie S., Ekstrom J.M.M., Pilgrim J.D., Savy C.E., Stephens R.T.T.,
570 Treweek J., Ussher G.T., Ward G. & Ten Kate K. (2013). Biodiversity Offsets and the Challenge
571 of Achieving No Net Loss. *Conservation Biology*, 27, 1254-1264.

572 Gayet G., Baptist F., Baraille L., Caessteker P., Clément J.C., Gaillard J., Gaucherand S., Isselin-
573 Nondedeu F., Poinot C. & Quétier F. (2016). Méthode nationale d'évaluation des fonctions
574 des zones humides. *Fondements théoriques, scientifiques et techniques. Onema, MNHN*, 310.

575 Gelcich S., Vargas C., Carreras M.J., Castilla J.C. & Donlan C.J. (2016). Achieving biodiversity benefits
576 with offsets: Research gaps, challenges, and needs. *Ambio*, 1-6.

577 Gibbons P. & Freudenberger D. (2006). An overview of methods used to assess vegetation condition
578 at the scale of the site. *Ecological Management & Restoration*, 7, S10-S17.

579 Gibbons P. & Lindenmayer D.B. (2007). Offsets for land clearing: No net loss or the tail wagging the
580 dog? *Ecological Management & Restoration*, 8, 26-31.

581 Hanski I. (1998). Metapopulation dynamics. *Nature*, 396, 41-49.

582 Harper D.J. & Quigley J.T. (2005). No Net Loss of Fish Habitat: A Review and Analysis of Habitat
583 Compensation in Canada. *Environmental Management*, 36, 343-355.

584 Heink U. & Kowarik I. (2010). What criteria should be used to select biodiversity indicators?
585 *Biodiversity and conservation*, 19, 3769-3797.

586 Jacob C., Quétier F., Aronson J., Pioch S. & Levrel H. (2014). Vers une politique française de
587 compensation des impacts sur la biodiversité plus efficace: défis et perspectives. *[VertigO] La
588 revue électronique en sciences de l'environnement*, 14.

589 Jaunatre R., Buisson E. & Dutoit T. (2014). Can ecological engineering restore Mediterranean
590 rangeland after intensive cultivation? A large-scale experiment in southern France. *Ecological
591 Engineering*, 64, 202-212.

592 Jiguet F., Gregory R.D., Devictor V., Green R.E., VOŘÍŠEK P., Van Strien A. & Couvet D. (2010).
593 Population trends of European common birds are predicted by characteristics of their
594 climatic niche. *Global change biology*, 16, 497-505.

595 Jones I.L., Bull J.W., Milner-Gulland E.J., Esipov A.V. & Suttle K.B. (2014). Quantifying habitat impacts
596 of natural gas infrastructure to facilitate biodiversity offsetting. *Ecol. Evol.*, 4, 79-90.

597 Julliard R., Jiguet F. & Couvet D. (2004). Common birds facing global changes: what makes a species
598 at risk? *Global Change Biology*, 10, 148-154.

- 599 Kiesecker J.M., Copeland H., Pocewicz A., Nibbelink N., McKenney B., Dahlke J., Holloran M. & Stroud
600 D. (2009). A Framework for Implementing Biodiversity Offsets: Selecting Sites and
601 Determining Scale. *BioScience*, 59, 77-84.
- 602 Kondolf G.M., Piégay H., Schmitt L. & Montgomery D.R. (2003). Geomorphic classification of rivers
603 and streams. *Tools in fluvial geomorphology*, 133-158.
- 604 Laycock H.F., Moran D., Raffaelli D.G. & White P.C.L. (2013). Biological and operational determinants
605 of the effectiveness and efficiency of biodiversity conservation programs. *Wildlife Research*,
606 40, 142-152.
- 607 Levrel H., Frascaria-Lacoste N., Hay J., Martin G. & Pioch S. (2015). *Restaurer la nature pour atténuer*
608 *les impacts du développement: Analyse des mesures compensatoires pour la biodiversité.*
609 Editions Quae.
- 610 Levrel H., Pioch S. & Spieler R. (2012). Compensatory mitigation in marine ecosystems: which
611 indicators for assessing the “no net loss” goal of ecosystem services and ecological functions?
612 *Marine Policy*, 36, 1202-1210.
- 613 Madsen B., Moore Brands K. & Carroll N. (2010). State of biodiversity markets: offset and
614 compensation programs worldwide.
- 615 Maseyk F., Barea L., Stephens R., Possingham H., Dutson G. & Maron M. (2016). A disaggregated
616 biodiversity offset accounting model to improve estimation of ecological equivalency and no
617 net loss. *Biological Conservation*, 204, 322-332.
- 618 McKenney B. & Kiesecker J. (2010). Policy Development for Biodiversity Offsets: A Review of Offset
619 Frameworks. *Environmental Management*, 45, 165-176.
- 620 Mimet A., Clauzel C. & Foltête J.-C. (2016). Locating wildlife crossings for multispecies connectivity
621 across linear infrastructures. *Landscape Ecol*, 1-19.
- 622 Moilanen A., Van Teeffelen A.J.A., Ben-Haim Y. & Ferrier S. (2009). How Much Compensation is
623 Enough? A Framework for Incorporating Uncertainty and Time Discounting When Calculating
624 Offset Ratios for Impacted Habitat. *Restoration Ecology*, 17, 470-478.
- 625 Moreno-Mateos D., Maris V., Béchet A. & Curran M. (2015). The true loss caused by biodiversity
626 offsets. *Biological Conservation*, 192, 552-559.
- 627 Naeem S., Duffy J.E. & Zavaleta E. (2012). The functions of biological diversity in an age of extinction.
628 *Science*, 336, 1401-1406.
- 629 Nardo M., Saisana M., Saltelli A., Tarantola S., Hoffman A. & Giovannini E. (2005). Handbook on
630 constructing composite indicators.
- 631 Noss R.F. (1990). Indicators for monitoring biodiversity: a hierarchical approach. *Conservation*
632 *biology*, 355-364.
- 633 Parkes D., Newell G. & Cheal D. (2003). Assessing the quality of native vegetation: The ‘habitat
634 hectares’ approach. *Ecological Management & Restoration*, 4, S29-S38.
- 635 Pilgrim J.D., Brownlie S., Ekstrom J.M.M., Gardner T.A., von Hase A., Kate K.t., Savy C.E., Stephens
636 R.T.T., Temple H.J., Treweek J., Ussher G.T. & Ward G. (2013). A process for assessing the
637 offsetability of biodiversity impacts. *Conservation Letters*, 6, 376-384.
- 638 Poff N.L., Allan J.D., Bain M.B., Karr J.R., Prestegard K.L., Richter B.D., Sparks R.E. & Stromberg J.C.
639 (1997). The natural flow regime. *BioScience*, 47, 769-784.
- 640 Quétier F. & Lavorel S. (2011). Assessing ecological equivalence in biodiversity offset schemes: Key
641 issues and solutions. *Biological Conservation*, 144, 2991-2999.
- 642 Quetier F., Regnery B. & Levrel H. (2014). No net loss of biodiversity or paper offsets? A critical
643 review of the French no net loss policy. *Environmental Science & Policy*, 38, 120-131.
- 644 Quigley J. & Harper D. (2006). Effectiveness of Fish Habitat Compensation in Canada in Achieving No
645 Net Loss. *Environmental Management*, 37, 351-366.
- 646 Regnery B., Couvet D. & Kerbiriou C. (2013). Offsets and Conservation of the Species of the EU
647 Habitats and Birds Directives. *Conservation Biology*, 27, 1335-1343.
- 648 Resh V.H., Brown A.V., Covich A.P., Gurtz M.E., Li H.W., Minshall G.W., Reice S.R., Sheldon A.L.,
649 Wallace J.B. & Wissmar R.C. (1988). The role of disturbance in stream ecology. *Journal of the*
650 *North American benthological society*, 7, 433-455.

651 Smith M.D. & Knapp A.K. (2003). Dominant species maintain ecosystem function with non-random
652 species loss. *Ecol Lett*, 6, 509-517.
653 Solan M., Cardinale B.J., Downing A.L., Engelhardt K.A., Ruesink J.L. & Srivastava D.S. (2004).
654 Extinction and ecosystem function in the marine benthos. *Science*, 306, 1177-1180.
655 State of Florida (2004). F-DEP UMAM Chapter 62–345.
656 ten Kate K., Bishop J. & Bayon R. (2004). Biodiversity offsets: Views, experience, and the business
657 case. *IUCN, Gland, Switzerland and Cambridge, UK and Insight Investment, London, UK*.
658 Tischew S., Baasch A., Conrad M.K. & Kirmer A. (2010). Evaluating Restoration Success of Frequently
659 Implemented Compensation Measures: Results and Demands for Control Procedures.
660 *Restoration Ecology*, 18, 467-480.
661 Virah-Sawmy M., Ebeling J. & Taplin R. (2014). Mining and biodiversity offsets: A transparent and
662 science-based approach to measure "no-net-loss". *J. Environ. Manage.*, 143, 61-70.

663

664

665

666