Bone histology of *Iberosuchus macrodon* (Sebecosuchia, Crocodylomorpha)

Jorge Cubo, Meike Köhler, Vivian de Buffrénil

► To cite this version:

Jorge Cubo, Meike Köhler, Vivian de Buffrénil. Bone histology of *Iberosuchus macrodon* (Sebecosuchia, Crocodylomorpha). *Lethaia*, 2017, 19 (4), pp.495-503. 10.1111/let.12203 . hal-02295975

HAL Id: hal-02295975

<https://hal.sorbonne-universite.fr/hal-02295975>

Submitted on 24 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bone histology of *Iberosuchus macrodon* (Sebecosuchia, Crocodylomorpha).

By

Jorge Cubo^{1*}, Meike Köhler^{2,3}, Vivian de Buffrénil⁴

¹ Sorbonne-Universités, UPMC-Univ. Paris 06, CNRS, Institut des Sciences de la Terre de Paris (ISTeP), 4 place Jussieu, BC 19, 75005 Paris, France

² ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain.

³ Institut Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, Carrer de les Columnes s/n, 08193 Cerdanyola del Vallés, Spain

⁴ Museum National d'Histoire Naturelle, Centre de Recherche sur la Paléobiodiversité et les Paléoenvironnements (CR2P), 75005 Paris, France

* Corresponding author : jorge.cubo_garcia@upmc.fr

RH – Bone histology of *Iberosuchus*

ABSTRACT

Iberosuchus macrodon is a Cenozoic crocodyliform interpreted as a terrestrial, cursorial form. In order to assess if this adaptation was accompanied by a high growth rate and an elevated resting metabolic rate (two features commonly attributed to several terrestrial Triassic Crocodylomorpha based on histology) we studied bone histology in the femora of two specimens attributed to *I. macrodon*. Beyond this question is the broader problem of the possible survival to the Cretaceous-Paleogene extinction event of tachymetabolic sauropsids other than birds. At mid-diaphysis, bone cortices in *Iberosuchus* are made of a parallel-fibered tissue that turns locally to true lamellar bone. Cortical vascularization consists of simple longitudinal canals forming a network of medium density. The spacing pattern of conspicuous lines of arrested growth suggests asymptotic growth for *Iberosuchus*. This general histological structure prevails also in the metaphyseal region of the bones. It is basically similar to that encountered in certain large lizards adapted to active predation, the Varanidae and the Teiidae. In one of the two *Iberosuchus* femora, however, an intra-cortical meniscus made of a tissue displaying a global radial architecture, occurs in the region of the fourth trochanter. Histologically, the latter can be interpreted either as compacted spongiosa, or as a fibro-lamellar complex with a gross radial orientation, a tissue corresponding to fast periosteal apposition. These observations suggest that, *Iberosuchus* basically had a slow, cyclical growth indicative of an ecto-poikilothermic, lizard-like, resting metabolic rate. However it might also have retained a limited capacity for fast periosteal accretion in relation to local morphogenetic requirements as, for instance, the development of crests or trochanters.

Key-words: Crocodylomorphs, Cretaceous-Paleogene extinction event, terrestriality, bone structure, growth, metabolism.

INTRODUCTION

The presence of a four-chambered heart in crocodiles and birds (Seymour *et al.* 2004, Summers 2005) and unidirectional air flow through the lungs (Farmer & Sanders 2010) suggests that the last common ancestor of archosaurs was endothermic, and that this character state was inherited by Ornithodira (pterosaurs and dinosaurs including birds) and Pseudosuchia (taxa more closely related to crocodiles than to birds), but was lost somewhere during the evolution of the latter (Seymour *et al.* 2004). Indirect evidence for this hypothesis has been gained from inferences of bone growth rates of extinct archosaurs (Ricqlès *et al.* 2008; Cubo *et al.* 2012; Legendre *et al.* 2013), assuming a direct relationship between bone growth rates and resting metabolic rates (Montes *et al.* 2007). Recently, resting metabolic rates of extinct archosaurs and non-archosaurian Archosauromorpha were inferred using bone paleohistology (Legendre *et al.* 2016), providing more direct support for that hypothesis.

Interestingly, within Pseudosuchia, aquatic (Dyrosauridae), semiaquatic (Eusuchia) and terrestrial (Sebecidae and other Sebecosuchia such as *Iberosuchus*) forms survived the Cretaceous-Paleogene extinction event (Macleod *et al.* 1997). According to the null hypothesis, *Iberosuchus* retained resting metabolic rates similar to those of Triassic terrestrial Pseudosuchia with an upright stance, such as *Terrestrisuchus* (Ricqlès *et al.* 2003). Two alternative hypotheses will be tested: (1) The inferred cursorial locomotion (Riff & Kellner 2011) of *Iberosuchus* suggests that it might have acquired even higher resting metabolic rates than its Triassic terrestrial relatives. (2) The Cretaceous-Paleogene extinction event filtered the diversity of Pseudosuchia, so that only taxa showing low resting metabolic rates survived. Bone histology has been acknowledged for several decades as one of the major clues for assessing the gross physiological adaptations of extinct taxa (Ricqlès 1974, 1978); unfortunately, there is no description of the histological features of long bones in notosuchians up to now (only osteoderms in some taxa were studied; cf. Buffrénil *et al.* 2015). In order to settle the question, we analyzed the bone histology of *Iberosuchus*, and tentatively interpreted it in terms of growth rates and metabolism activity.

MATERIAL AND METHODS

We analyzed two partial femora assigned to *Iberosuchus macrodon* Antunes 1975 from the Paleocene of La Boixedat (Spain). These specimens belong to the paleontological collections of the Institut Català de Paleontologia (ICP), labeled IPS4930 and IPS4932. In contrast to the condition in the Eusuchia, the femora of *Sebecus* and *Iberosuchus* are (a) straighter and (b) the medial edge of the greater trochanter is a prominent, sharp, longitudinal crest (Pol *et al.* 2012). Although the specimens in this study cannot be attributed unequivocally to *Iberosuchus* (whose holotype is a skull fragment), the morphological congruence of available specimens nevertheless justifies the assumption that the scarce, non-neosuchian Meoseucrocodylia specimens from the Paleogene of the Iberian Peninsula and southern France are at least closely related, indistinguishable forms (Ortega *et al.* 1996). We follow the phylogeny of Pol *et al.* (2012) in which Neosuchia and Notosuchia are sister taxa and *Iberosuchus* is a sebecosuchian nested in Notosuchia.

After photography, the bones were embedded under vacuum in a polyester resin, and two transverse slices, 3 mm in thick, were removed from the middle of the diaphysis (Fig. 1A, B) and the base of the proximal metaphysis (Fig. 1E, F) of each femur. According to standard ground section procedures (e.g. Lamm 2013), these slices were polished on one side, glued on glass slides and grounded to a thickness $100\ \mu\text{m} \pm 20$. These thin sections were observed and photographed with a Nikon Eclipse E600POL microscope, under normal and cross-polarized light, with or without a lambda compensator.

Simple morphometric measurements were also performed on binary (black and white) images of the sections at mid-diaphysis (Fig. 1C, D) using the software Image J (Schneider *et al.* 2012). These measurements include: a) the compactness of each section, or GC (ratio, expressed in percent, of the area actually occupied by bone tissue to the total sectional area); b) compactness of the cortex proper, or CC (ratio, expressed in percent, of the actual area of the cortex occupied by bone tissue to the total cortical area, including bone plus cavities); c) the cortico-diaphyseal index, or CDi (mean thickness of bone cortex as a fraction of the mean diaphyseal radius).

RESULTS

Diaphyseal region of the femora

Both femoral diaphyses have a tubular morphology, with a free medullary cavity surrounded by a compact cortex. However, specimen IPS 4930, though smaller than IPS 4932 (38 % less cross sectional area) is clearly more compact (GC = 87.16% vs 77.52%) and has a thicker (CDi = 0.65 vs 0.54) and more compact (CC = 99.41% vs 79.06%) cortex. Perimedullary resorption is limited in this specimen; as a consequence bone layers deposited in early ontogenetic stages are preserved.

Histologically, the bone forming the diaphyseal cortex has similar characteristics in the two specimens. Most of its volume is predominantly composed of parallel-fibered tissue (Fig. 1A, B; Fig. 2A, B), displaying mass birefringence and collagen fiber bundles oriented circularly (i.e. parallel to the outer contours of the bones). In IPS 4932, this tissue can locally turn to the lamellar type (Fig. 2B). Frequent irregularities in the birefringence properties of the parallel-fibered bone suggest some degree of local variation in fiber orientation. Cell lacunae in this tissue are oriented parallel to the collagen fibers, and they display a small size, as compared to the lacunae occurring in primary or secondary endosteal deposits forming e.g. osteons (Fig. 2C).

The vascularization of the primary cortex mainly consists of simple vascular canals oriented longitudinally and, to a much lesser extent, obliquely. These canals are relatively few in the peripheral (outer) part of the cortex; especially in IPS 4932 (Fig. 2A, B). Vascular density increases in deeper cortical parts, a process much more pronounced in IPS 4930 (Fig. 1B; Fig. 2D) than in IPS 4932 (Fig. 1A). In both specimens, the cortical region bordering the medullary cavity has undergone Haversian remodeling. This process was relatively diffuse in IPS 4932, where it created sparse secondary osteons (Fig. 2C, E); it was more restricted but more intense in IPS 4930 where dense Haversian bone tissue was spatially limited (Fig. 2F). Superficial resorption/reconstruction processes of variable intensity also occurred around the medullary cavity, creating endosteal layers of secondary lamellar bone.

In both specimens the femoral cortex shows cyclical growth marks in the form of lines of arrested growth (LAGs). The latter appear as thin dark lines parallel to the contour of the bones (Fig. 2G). These lines are broadly, but unevenly, spaced in the depth of the cortex (280 μ m in the average on Fig. 2G). In IPS 4932, the spacing of the LAGs suddenly becomes more regular and much narrower in the most peripheral cortical layers, where the intervals between consecutive LAGs drop to 78 μ m in the

average (Fig. 2G). Both femora also have bundles of short Sharpey's fibers (length $50\text{ }\mu\text{m} \pm 10$) oriented radially and obliquely (Fig. 2H).

Histology of the metaphyseal region

The histological structure of primary periosteal cortices in the metaphyseal region of both specimens (Fig. 1E, F) is basically similar to that prevailing in the diaphyseal region: most of the cortical volume consists of parallel-fibered bone, the collagen fibers of which are oriented circularly (Fig. 3A, B). However, three main differences exist. 1) Bone vascularization, mainly represented by simple vascular canals and primary osteons oriented longitudinally, tends to be lower in the metaphyses than in the diaphyses, especially in IPS 4930 (Fig. 3A, B). 2) Loose networks of remodeled endosteal bone trabeculae (Fig. 1E, F; Fig. 3C) partly fill the medullary cavity (the medullary cavity is free in the diaphysis). 3) In IPS 4932, the cortex is not only composed of vascularized parallel-fibered tissue; but also contains a broad, central crescent-like meniscus formation displaying different histological features (Fig. 1F; Fig. 3D). This formation is briefly described below.

This peculiar bone layer is located under the forth trochanter of the femur and consequently, it occupies only a part of the sectional area (Fig. 1F). It is inserted in the middle of the cortex, between two layers (under and above it) of ordinary parallel-fibered bone (Fig. 3D). Histologically, the thinnest parts of this meniscus consist merely of some big longitudinal primary osteons (Fig. 3E). In its thicker part, the meniscus displays two components: 1) Multiple oblong areas made of brightly birefringent parallel-fibered tissue (blue in Fig. 3F, G) with a dominant, though variable, radial orientation (including for cell lacunae) and vascular canals also displaying a gross radial orientation. 2) Small and irregular monorefringent, or poorly birefringent areas, unevenly wedged between the birefringent ones, and formed by woven bone (red in Fig. 3F, G). They contain big multipolar cell lacunae.

This basic structure can be interpreted in two distinct ways. The first interpretation is that it represents former spongiosa, compacted by endosteal deposits. Crests, trochanters and other bone excrescences related to muscle insertion are most often associated with spongy tissues within bone cortices (e.g. Ricqlès 1976a). During growth, such spongiosae are frequently made compact by inter-trabecular filling, a situation actually observed by one of us (VB) in sections of the fourth trochanter

of two alligatorids, *Alligator mississippiensis* and *Diplocynodon ratelii*. The gross radial orientation of this compacted spongiosa in IPS 4932 reflects the strong traction stress exerted on the periosteum and the growing bone cortex by the muscle *caudofemoralis* (according to Wolf's law; cf. Currey 2002). An alternative interpretation of the meniscus structure is that it is made of a radiating fibro-lamellar bone complex, a tissue known to be characteristic of fast or very fast accretion. The monorefringent areas would represent the woven-fibered trabeculae initially deposited by the periosteum. The brightly birefringent areas would be primary osteons. The main difference between the bone forming the meniscus and typical fibro-lamellar complexes is that the highly ordered and conspicuous periosteal scaffoldings of woven-fibered trabeculae that characterize these complexes cannot be clearly identified in *Iberosuchus*. As a consequence, the limits and shape of the putative primary osteons cannot be traced precisely. For this reason, the tissue forming the thick part of the meniscus should be interpreted, with necessary caution, as an atypical form of radial fibro-lamellar bone tissue. Both interpretations, compacted spongiosa vs radiating fibro-lamellar complex, differ little from each other. In both cases, the differentiation of the fourth trochanter basically involves the sub-periosteal accretion of a spongiosa, and its subsequent compaction by inter-trabecular, centripetal deposits of endosteal lamellar or parallel-fibered tissue. Their main difference would reside in the dynamics of bone deposits: fast or very fast periosteal and endosteal deposits in the case of a radiating fibro-lamellar complex; much slower deposits in the case of a compacted spongiosa.

The basal part of the meniscus is in continuity (though histologically very distinct) with the subjacent parallel-fibered tissue. Conversely, the peripheral border of the meniscus is marked by a reversion line displaying a typical scalloped contour at high magnification (Fig. 3H). Above this line, towards bone periphery, the parallel-fibered tissue prevails again. The occurrence of this reversion line means that, during growth, the height of the basal part of the trochanter had to be reduced, through a resorption process, to become compatible with the diameter of the diaphysis into which the trochanter was sequentially relocated.

DISCUSSION

The most significant result of this study is that the femoral cortex of *Iberosuchus* is made of a parallel-fibered tissue with variable vascular density and conspicuous lines of arrested growth. This situation was observed in all bone sections, especially those sampled at mid-diaphysis, i.e. a sectional plane classically considered as a general reference for histological studies of long bones; e.g. Lamm 2013). This result strongly suggests that *Iberosuchus* was not a fast growing, tachymetabolic animal, but a slow growing crocodile with steep, cyclic decreases in growth rate. Several experimental studies indeed show that, on the one hand, bone tissue with scattered longitudinal vascular canals results from accretion speeds less than 5 $\mu\text{m}/\text{day}$ (Castanet *et al.* 1996) and that, on the other hand, parallel-fibered bone corresponds to accretion rates less than 1 $\mu\text{m}/\text{day}$ (e.g. Buffrénil & Pascal 1984). The actual apposition rate on the femoral cortex of *Iberosuchus* could have been between these values which are anyway far below those prevailing for the woven-fibered tissue, especially when it contributes to the constitution of fibro-lamellar complexes (laminar, plexiform, radiating bone tissues) typically encountered in fast growing endotherms (Castanet *et al.* 1996, 2000; Margerie *et al.* 2002; see also Cubo *et al.* 2012). In most modern crocodiles, the femoral cortex is made of a lamellar-zonal tissue (Enlow & Brown 1957, Ricqlès 1976b, Lee 2004) displaying conspicuous yearly growth cycles. The latter consist of zones made of woven-fibered tissue (that may change to the parallel-fibered type), associated with annuli made of parallel-fibered or true lamellar bone (Buffrénil 1980a, b; Hutton 1986). These two components of an annual growth cycle are deposited sequentially and reflect a progressive decrease in growth rate that may end, each year, in a total cessation of growth and the formation of a LAG (Buffrénil 1980a). Since it integrates a woven-fibered component, the lamellar-zonal tissue reflects faster deposition than the mere parallel-fibered type prevailing in *Iberosuchus*. The occurrence of lines of arrested growth in this taxon, as well as the absence of annuli, clearly shows that the maximum growth rate in *Iberosuchus* was comparable to the slowest rates of modern crocodiles, and that growth stopped completely each year. As observable in the femur, the histological features of *Iberosuchus* can be best compared to those of large predatory squamates such as the varanids or the teids (e.g. Duarte-Varela & Cabrera 2000, Buffrénil & Hémery 2002, see also Cubo *et al.* 2014). In both cases, bone cortices are made of parallel-fibered tissue (that may turn to the woven-fibered type in the inner cortex) comprising longitudinal simple vascular canals or primary osteons,

and lines of arrested growth. Therefore, if the histological structure of long bones indeed reflects growth rate, as initially proposed by Amprino (1947) and universally acknowledged today, then the growth activity of *Iberosuchus* should be considered similar to that of extant large lizards. This comparison is further substantiated by the spacing pattern of cyclical growth marks in IPS 4932. Growth mark spacing indeed suggests that a sudden and steep decrease in growth activity occurred about six years before this animal died. Its growth pattern was thus clearly asymptotic, a situation commonly encountered in squamates in which epiphyseal and metaphyseal fusion limits growth possibilities (Maisano 2008, Buffrénil *et al.* 2004). Conversely, this growth pattern is unusual in crocodiles, though it was observed in some populations of *Alligator mississippiensis* (Woodward *et al.* 2011; see also Lee *et al.* 2013). IPS 4932 is larger than IPS 4930 and displays signs of a sudden decrease in growth. These two characteristics indicate an older age for IPS 4932. Moreover, the porosity of deep cortical layers in this specimen is a feature commonly encountered in mature crocodilian females (Wink & Elsey 1986; Wink *et al.* 1987).

If the general relationship between the histological structure of primary bone cortices and their appositional rate is now strongly evidenced by experimental data (e.g. Castanet *et al.* 1996), the association between, on the one hand, bone tissue types and growth rate and, on the other hand, the resting metabolic rate of an organism, seems to be less simple and less clearly deciphered. Montes *et al.* (2007) found a direct relationship between periosteal bone growth rate and resting metabolic rate in a sample of growing amniotes. In general, long bone cortices made of parallel-fibered tissue, be it vascularized or not, are characteristically encountered in ecto-poikilothermic tetrapods (e.g. Enlow and Brown 1956-1958, Ricqlès 1976). However, the few extant large squamates displaying vascular canals (simple canals or primary osteons) within parallel-fibered bone, i.e. the largest Teiidae (Duarte-Varela & Cabrera 2000, Cubo *et al.* 2014) and the Varanidae more than 30 cm in snout-vent length (Buffrénil *et al.* 2008), can experience transitory episodes of relatively high and constant metabolic rate during either their reproductive cycle (Tattersall *et al.* 2016), or during times of intense foraging activity (synthesis in Thompson 1999). Otherwise, these animals, including the largest ones, have a typical ecto-poikilothermic physiology (Green *et al.* 1991, Christian & Conley 1994, Wikramanayake *et al.*

1999). The question is made more complex by the fact that the very same type of bone tissue (vascularized parallel-fibered bone) can be observed in long bone cortices in some large anurans, such as *Rana Catesbeiana*, *Rhinella marina*, or *Pipa pipa* (pers. obs. VB), known to have much lower metabolic rates than the squamates in general (White et al. 2006). Such observations suggest that the actual relationship between the details of bone structure and the metabolic rate of an organism result from a complex, multifactorial causality that needs to be fully deciphered. It nevertheless remains that bone cortices mainly composed of parallel-fibered tissue are very unlikely to belong to an endotherm-homeotherm animal. For this reason, the fundamental thermal regime of *Iberosuchus* should be considered to have been ectotherm and poikilotherm, as is also the case for the large extant squamates or lissamphibians mentioned above.

The particular case of *Iberosuchus* suggests that, in pseudosuchians, a terrestrial habitat is not necessarily associated with a high, sustained growth speed and the tachymetabolic regime consistent with it. Of course, this conclusion holds only for *Iberosuchus*. The Notosuchia (of which the Sebecosuchia are but one clade) represent a particularly rich and diversified lineage of terrestrial forms in which a number of bizarre and ecologically enigmatic taxa occur (e.g. Ortega *et al.* 2000). Future comparative studies should document whether this high morphological diversity also involved significant physiological discrepancies.

One element in our observations could challenge the conclusion presented above: the possible occurrence of radiating fibro-lamellar tissue in a limited area of the metaphyseal cortex of IPS 4932. This type of bone tissue is considered to result from the fastest accretion rate (Margerie *et al.* 2004) and is supposed to be an exclusive feature of fast-growing tachymetabolic tetrapods (Ricqlès 1974, 1976a). For several reasons, the occurrence of radiating fibro-lamellar tissue in the core of the metaphyseal cortex cannot be alleged to conclude that (1) episodes of fast growth occurred in the life of IPS 4932, and (2) that this taxon had the physiological competence for sustaining them at the level of the organism as a whole. As mentioned in the observations, the identification of this tissue is debatable, due to atypical histological characteristics. Nevertheless, even if the tissue forming the meniscus is really akin to fibro lamellar radiating bone, the complete lack of this tissue in diaphyseal

cortices (despite the excellent preservation of primary cortical tissues) would clearly indicate that its occurrence in the proximal metaphysis of IPS 4932 does not indicate a general trend towards fast growth in *Iberosuchus*, but a local, topographically restricted, process. This taxon might indeed have retained the potential capacity to develop fast periosteal accretion but, in the specimen studied here, this capacity would have been expressed in a strictly local context, in relation to the differentiation, growth and sequential relocation of the fourth trochanter during ontogeny.

With reference to the hypotheses listed in the introduction, our results suggest that the situation may be more complex than expected because some crocodyliforms might have preserved physiological potentialities that could have been expressed, or non-expressed, depending on morphological and/or ecological contexts. Future studies based on larger samples should settle more firmly this interesting question, and help assess if the capacity for fast growth, at least in limited skeletal regions and for a limited duration, is a frequent feature in the Notosuchia.

LITERATURE CITED

Amprino, R. 1947: La structure du tissu osseux envisagée comme expression de différences dans la vitesse de l'accroissement. *Archives de Biologie* 58, 315–330.

Antunes, M.T. 1975: *Iberosuchus*, crocodile Sébécosuchien nouveau, de l'Eocène ibérique au Nord de la Chaîne centrale, et l'origine du canyon de Nazaré. *Comunicações dos Serviços Geológicos de Portugal* 59, 285-330.

Antunes, M.T. 1986: *Iberosuchus* et *Pristichampsus*, crocodiliens de l'Eocène : données complémentaires, discussion, distribution stratigraphique. *Ciências da Terra* 8, 111-122.

Buffrénil, V. de. 1980a: Données préliminaires sur la structure des marques de croissance squelettiques chez les crocodiliens actuels et fossiles. *Bulletin de la Société Zoologique de France* 105, 355-361.

Buffrénil, V. de. 1980b: Mise en évidence de l'incidence des conditions de milieu sur la croissance de *Crocodylus siamensis* (Schneider, 1801) et valeur des marques de croissance squelettiques pour l'évaluation de l'âge individuel. *Archives de Zoologie Expérimentale et Générale* 121, 63-76.

282 Buffrénil, V. de & Pascal, M. 1984: Croissance et morphogenèse postnatales de la mandibule du
 283 vison (*Mustela vison* Schreiber) : données sur la dynamique et l'interprétation fonctionnelle des dépôts
 284 osseux mandibulaires. *Canadian Journal of Zoology* 62, 2026-2037.

285 Buffrénil, V. de & Castanet, J. 2000 : Age estimation by skeletochronology in the Nile monitor
 286 (*Varanus niloticus*), a highly exploited species. *Journal of Herpetology* 34, 414-424.

287 Buffrénil, V. de & Hémery, G. 2002: Variation in longevity, growth, and morphology in exploited
 288 Nile monitors (*Varanus niloticus*) from sahelian Africa. *Journal of Herpetology* 36, 419-426.

289 Buffrénil, V. de, Ineich, I. & Böhme, W. 2004: Comparative data on epiphyseal development in the
 290 family Varanidae. *J. Herpetology* 37, 328-335.

291 Buffrénil, V. de, Clarac, F., Fau M., Martin, S., Martin B., Pellé, E. & Laurin, M. 2015:
 292 Differentiation and growth of bone ornamentation in vertebrates: a comparative histological study
 293 among the Crocodylomorpha. *Journal of morphology* 276, 425-445.

294 Castanet, J., Grandin, A., Abourachid, A., & Ricqlès, A. de. 1996: Expression de la dynamique de
 295 croissance dans la structure de l'os périostique chez *Anas platyrhynchos*. *Comptes-Rendus de.*
 296 *L'Académie des Sciences, Paris, Sciences de la Vie* 319, 301-308.

297 Castanet, J., Curry-Rogers, K., Cubo, J. & Boisard, J-J. 2000: Periosteal bone growth rates in
 298 extant ratites (ostrich and emu). Implications for assessing growth in dinosaurs. *Comptes-Rendus de.*
 299 *L'Académie des Sciences, Paris, Sciences de la Vie* 323, 543-550.

300 Christian, K.A. & Conley, K.E. 1994: Activity and resting metabolism of varanid lizards compared
 301 with "typical" lizards. *Australian Journal of Zoology* 42, 185-193.

302 Cubo, J., Le Roy, N., Martinez-Maza, C., Montes, L. 2012: Paleohistological estimation of bone
 303 growth rate in extinct archosaurs. *Paleobiology* 38, 335-349.

304 Cubo, J., Baudin, J., Legendre, L., Quilhac, A., Buffrénil, V. de 2014: Geometric and metabolic
 305 constraints on bone vascular supply in diapsids. *Biological Journal of the Linnean Society of London*
 306 112, 668-677.

307 Duarte-Varela, C.F. & Cabrera, M.R. 2000: Testing skeletochronology in black tegu lizards
 308 (*Tupinambis merianae*) of known ages. *Herpetological Review* 31, 224-226.

309 Enlow, D.H. & Brown, S.O. 1957: A comparative histological study of fossil and recent bone tissues.
 310 Part II. *Texas Journal of Science* 9, 186-214.

311 Farmer, C.G. and Sanders, K. 2010: Unidirectional Airflow in the Lungs of Alligators. *Science*
 312 327, 338–340.

313 Green, B., King, D., Braysher, M., & Saim, A. 1991: Thermoregulation, water turnover and
 314 energetics of free-ranging Komodo dragons, *Varanus komodoensis*. *Comparative Biochemistry and*
 315 *Physiology* 99A, 97-101.

316 Hutton, J.M. 1986: Age determination of living Nile crocodiles from the cortical stratification of
 317 bone. *Copeia* 1986(2), 332-341.

318 Lamm, E.-T. 2013: Preparation and sectioning of specimens. In Padian, K., Lamm, E.-T. (eds): *Bone*
 319 *Histology of Fossil Tetrapods*, 55–160. University of California Press, Berkeley.

320 Lee, A.H. 2004: Histological organization and its relationship to function in the femur of *Alligator*
 321 *mississippiensis*. *Journal of Anatomy* 204, 197-207.

322 Legendre, L.J., Segalen, L. and Cubo, J. 2013: Evidence for high bone growth rate in *Euparkeria*
 323 obtained using a new paleohistological inference model for the Humerus. *Journal of Vertebrate*
 324 *Paleontology* 33, 1343–1350.

325 Legendre, L.J., Guénard, G., Botha-Brink, J. and Cubo, J. 2016: Paleohistological evidence for
 326 ancestral high metabolic rate in archosaurs. *Systematic Biology* 65, 989-996.

327 Macleod, N., Rawson, P.F., Forey, P.L., Banner, F.T., BoudagherFadel, M.K., Bown, P.R.,
 328 Burnett, J.A., Chambers, P., Culver, S., Evans, S.E., Jeffery, C., Kaminski, M.A., Lord, A.R., Milner,
 329 A.C., Milner, A.R., Morris, N., Owen, E., Rosen, B.R., Smith, A.B., Taylor, P.D., Urquhart, E. and
 330 Young, J.R. 1997: The Cretaceous-Tertiary biotic transition. *Journal of the Geological Society* 154,
 331 265–292.

332 Maisano, J.A. 2002: Terminal fusion of skeletal elements as indicators of maturity in squamates.
 333 *Journal of Vertebrate Paleontology*, 22, 268-275.

334 Margerie, E. de, Cubo, J. & Castanet, J. 2002: Bone typology and growth rate: testing and
 335 quantifying « Amprino's rule » in the mallard (*Anas platyrhynchos*). *Comptes-Rendus de. L'Académie*
 336 *des Sciences, Paris, Sciences de la Vie* 325, 221-230.

337 Margerie, E. de, Robin, J-P., Verrier, D., Cubo, J. & Groscolas, R. 2004: Assessing the
 338 relationship between bone microstructure and growth rate: a fluorescent labelling study in the king
 339 penguin chick (*Aptenodytes patagonicus*). *Journal of Experimental Biology* 207, 869-879.

340 Montes, L., Le Roy, N., Perret, M., Buffrénil, V. de, Castanet, J. & Cubo, J. 2007: Relationship
 341 between bone growth rate, body mass and resting metabolic rate in growing amniotes: a phylogenetic
 342 approach. *Biological Journal of the Linnean Society* 92, 63-76.

343 Ortega, F., Buscalioni, A.D. and Gasparini, Z. 1996: Reinterpretation and new denomination of
 344 *Atacisaurus crassipratoratus* (Middle Eocene; Issel, France) as cf *Iberosuchus* (Crocodylomorpha,
 345 Metasuchia). *Geobios* 29, 353–364.

346 Ortega, F., Gasparini, Z., Buscalioni, A.D. and Calvo, J.O. 2000: A new species of *Araripesuchus*
 347 (Crocodylomorpha, Mesoeucrocodylia) from the Lower Cretaceous of Patagonia (Argentina). *Journal*
 348 *of Vertebrate Paleontology* 20, 57–76.

349 Pol, D., Leardi, J.M., Lecuona, A. and Krause, M. 2012: Postcranial Anatomy of *Sebecus*
 350 *Icaeorhinus* (crocodyliformes, Sebecidae) from the Eocene of Patagonia. *Journal of Vertebrate*
 351 *Paleontology* 32, 328–354.

352 Ricqlès, A. de. 1974: Evolution of endothermy: histological evidence. *Evolutionary Theory* 1, 51-
 353 80.

354 Ricqlès, A. de. 1976a: Recherches paléohistologiques sur les os longs des tetrapodes. VII. – Sur la
 355 classification, la signification fonctionnelle et l'histoire des tissus osseux des tetrapodes. *Annales de*
 356 *Paléontologie* 62, 71-126.

357 Ricqlès, A. de. 1976b: On bone histology of fossil and living reptiles, with comments on its
 358 functional and evolutionary significance. *In* Bellairs, A. d'A. & Cox, C.B. (eds) *Morphology and*
 359 *Biology of Reptiles*, 123-150. Linnean society Symposium Series n°3, London.

360 Ricqlès, A. de. 1978: Recherches paléohistologiques sur les os longs des tetrapods. VII. – Sur la
 361 classification, la signification fonctionnelle et l'histoire des tissus osseux des tétrapodes (troisième
 362 partie). *Annales de Paléontologie* 64, 85-111.

363 Ricqlès, Ad., Padian, K. & Horner, J.R. 2003: On the bone histology of some Triassic
 364 pseudosuchian archosaurs and related taxa. *Annales de Paléontologie* 89, 67–101.

365 Ricqlès, A. de, Padian, K., Knoll, F., Horner, J.R. 2008: On the origin of high growth rates in
 366 archosaurs and their ancient relatives: Complementary histological studies on Triassic archosauriforms
 367 and the problem of a “phylogenetic signal” in bone histology. *Annales de Paléontologie* 94, 57-76.

368 Riff, D. and Armin Kellner, A.W. 2011: Baurusuchid crocodyliforms as theropod mimics: clues
 369 from the skull and appendicular morphology of *Stratiosuchus maxhechti* (Upper Cretaceous of
 370 Brazil). *Zoological Journal of the Linnean Society* 163, S37–S56.

371 Schneider, C.A. Rasband, W.S. and Eliceiri, K.W. 2012 : [NIH Image to ImageJ: 25 years of image](#)
 372 [analysis](#). *Nature methods* 9(7), 671-675.

373 Seymour, R.S., Bennett-Stamper, C.L., Johnston, S.D., Carrier, D.R. and Grigg, G.C. 2004:
 374 Evidence for endothermic ancestors of crocodiles at the stem of archosaur evolution. *Physiological*
 375 *and Biochemical Zoology* 77, 1051–1067.

376 Summers, A.P. 2005: Evolution - Warm-hearted crocs. *Nature* 434, 833–834.

377 Tattersall, G.J., Leite, C.A.C., Sanders, C.E., Cadena, V., Andrade, D.V., Albe, A.S., Milson, W.K.
 378 2016 : Seasonal reproductive endothermy in Tegu lizards. *Scientific Advances* 2016 ; e1500951.

379 Thompson, G. 1999 : Goanna metabolism : different to other lizards, and if so, what are the
 380 ecological consequences ? *In* Horn, H.-G. & Böhme, W. (eds.) *Advances in monitor research II. –*
 381 *Mertensiella* 11, 79-90.

- Wink, C.S. & Elsey, R.M. 1986: Changes in femoral morphology during egg-laying in *Alligator mississippiensis*. *Journal of Morphology* 189, 183-188.
- Wink, C.S., Elsey, R.M. & Hill, E.M. 1987: Changes in femoral robusticity and porosity during the reproductive cycle of the female Alligator (*Alligator mississippiensis*). *Journal of Morphology* 193: 317-321.
- White, C.R., Phillips, N.F., Seymour, R.S. 2006 : The scaling and temperature dependence of vertebrate metabolism. *Biology Letters* 2, 125-127.
- Wikramanayake, E.D, Ridwan, W. & Marcellini, D. 1999 : The thermal ecology of free-ranging Komodo dragons, *Varanus komodoensis*, on Komodo Island, Indonesia. In Horn, H.-G. & Böhme, W. (eds.) *Advances in monitor research II. – Mertensiella* 11, 157-166.
- Woodward, H.N., Horner, J.R. & Farlow, J.O. 2011: Osteohistological evidence for determinate growth in the American alligator. *Journal of Herpetology* 45, 339-342.

ACKNOWLEDGEMENTS

We thank Francisco Ortega for his valuable help in performing the taxonomic assignment of the samples analyzed to *Iberosuchus*. We would like to express our gratitude to an anonymous reviewer for his/her helpful comments and to an associate editor for his/her stylistic suggestions. This work was supported by the Spanish Ministry of Economy and Competitiveness: CGL2015-63777-P, PI: MK, and 2014 SGR 1207, PI: MK, and [CERCA Programme / Generalitat de Catalunya](#). We have no conflict of interest to declare.

FIGURE LEGENDS

Figure 1: Gross aspect of the sections.

A: Global aspect of the the mid-diaphyseal section in IPS 4930. B: Global aspect of the the mid-diaphyseal section in IPS 4932. C: Binary image of the mid-diaphyseal section in specimen IPS 4930.

D: Binary image of the mid-diaphyseal section in specimen IPS 4932. E: Global aspect of the metaphyseal section in IPS 4930. F: Global aspect of the metaphyseal section in IPS 4932. Scale bar: 2 mm.

Figure 2: Bone histology in the femoral diaphysis in IPS 4930 and 49832.

A: Peripheral cortex of the femur in IPS 4930. The upper part is viewed in transmitted polarized light with lambda compensator; the lower part in ordinary transmitted light. B: Peripheral cortex of the femur in IPS 4932. Upper part: transmitted polarized light with lambda compensator; lower part: ordinary light. C: Size of osteocyte lacunae in periosteal and endosteal deposits (IPS 4932). The insert is a closer view at the wall of a secondary osteon as compared to the primary periosteal cortex. D: vascular density in the deep, peri-medullary cortex of IPS 4930. E: Secondary osteons scattered in the cortex of specimen IPS 4932. Polarized light. F: Localized area of dense Haversian tissue around the medullary cavity in IPS 4930. G: Lines of arrested growth, or LAG (arrows) in the cortex of IPS 4932. Note the tight spacing of the last LAGs. H: Sharpey's fibers in the cortex of IPS 4930 (arrows). All scale bars are equal to 0.5 mm but that of the close-up in C, which equals 0.1 mm.

Figure 3: Histological characteristics of the metaphyseal region.

A: Basic appearance of the femoral cortex in the metaphyseal region of IPS 4930. Upper part: transmitted polarized light with lambda compensator; lower half: ordinary transmitted light. B: Femoral cortex in the metaphyseal region of IPS 4932. Upper part: transmitted polarized light with lambda compensator; lower half: ordinary transmitted light. C: Remodeled endosteal trabeculae occupying the medullary cavity in the metaphysis of IPS 4930. Polarized light. D: Complex cortical structure observed locally in the metaphyseal of IPS 4932. The inner (left) and the outer (right) parts of the cortex are formed by parallel-fibered bone, the collagen fibers of which are oriented circularly (yellow). In between these layers the cortex is occupied by a tissue with a dominant radial structuration. E: Longitudinal primary osteons in the thin extremities of the crescent-like meniscus. Polarized light. F: Complex histological organization in the thick parts of the meniscus. Brightly birefringent (here in blue) and poorly birefringent (here in red) areas oriented radially are unevenly inter-mixed. Polarized light with lambda compensator. G: Continuity of the basal, parallel-fibered part

434 of the cortex (bottom part of the picture) with the tissue forming the meniscus. Polarized light with
435 lambda compensator. H: Reversion line (RL) separating the meniscus and the peripheral layer of
436 parallel-fibered bone tissue. The insert shows the typical scalloped contour of the reversion line. All
437 scale bars are equal to 0.5 mm but that of the close-up in H, which equals 0.1 mm.

A

B

C

D

E

F