



HAL
open science

**The complete chloroplast genome sequence of
Dactylorhiza majalis (Rchb.) P.F. Hunt et Summerh.
(Orchidaceae)**

Michal May, Alžběta Novotná, Julita Minasiewicz, Marc-André Selosse,
Marcin Jąkaliski

► **To cite this version:**

Michal May, Alžběta Novotná, Julita Minasiewicz, Marc-André Selosse, Marcin Jąkaliski. The complete chloroplast genome sequence of *Dactylorhiza majalis* (Rchb.) P.F. Hunt et Summerh. (Orchidaceae). *Mitochondrial DNA Part B Resources*, 2019, 4 (2), pp.2821-2823. 10.1080/23802359.2019.1660282 . hal-02302282


HAL Id: hal-02302282

<https://hal.sorbonne-universite.fr/hal-02302282>

Submitted on 1 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Mitochondrial DNA Part B Resources

ISSN: (Print) 2380-2359 (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

The complete chloroplast genome sequence of *Dactylorhiza majalis* (Rchb.) P.F. Hunt et Summerh. (*Orchidaceae*)

Michał May, Alžběta Novotná, Julita Minasiewicz, Marc-Andre Selosse & Marcin Jąkowski

To cite this article: Michał May, Alžběta Novotná, Julita Minasiewicz, Marc-Andre Selosse & Marcin Jąkowski (2019) The complete chloroplast genome sequence of *Dactylorhiza majalis* (Rchb.) P.F. Hunt et Summerh. (*Orchidaceae*), *Mitochondrial DNA Part B*, 4:2, 2821-2823, DOI: [10.1080/23802359.2019.1660282](https://doi.org/10.1080/23802359.2019.1660282)

To link to this article: <https://doi.org/10.1080/23802359.2019.1660282>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 02 Sep 2019.



Submit your article to this journal [↗](#)



Article views: 57



View related articles [↗](#)



View Crossmark data [↗](#)

The complete chloroplast genome sequence of *Dactylorhiza majalis* (Rchb.) P.F. Hunt et Summerh. (*Orchidaceae*)

Michał May^a , Alžběta Novotná^b , Julita Minasiewicz^a , Marc-Andre Selosse^{a,c}  and Marcin Jąkowski^a 

^aDepartment of Plant Taxonomy and Nature Conservation, Faculty of Biology, University of Gdańsk, Gdańsk, Poland; ^bFaculty of Science, University of South Bohemia, České Budějovice, Czech Republic; ^cInstitut de Systématique, Evolution, Biodiversité (ISYEB), Muséum national d'Histoire naturelle, CNRS, Sorbonne Université, EPHE, Paris, France

ABSTRACT

The complete chloroplast genome of *Dactylorhiza majalis* (Rchb.) P.F. Hunt et Summerh. (*Orchidaceae:Orchidoideae*) was assembled and characterized using next-generation sequencing data. The plastome (154,108 bp) possesses the typical circular structure consisting of a large single-copy region (LSC; 83,196 bp), a small single-copy region (SSC; 26,580 bp), and two copies of inverted repeats (17,752 bp each). Its overall GC content is 36.99% and the plastome encodes 134 genes. Reconstruction of phylogenetic relationships using complete plastome sequences of *Orchidaceae* representatives showed that *D. majalis* was nested within the *Orchidoideae* tribe *Orchideae*. The complete plastome comprises a valuable tool in elucidating taxonomic uncertainties within the genus *Dactylorhiza*.

ARTICLE HISTORY

Received 19 July 2019
Accepted 3 August 2019

KEYWORDS

Chloroplast genome;
Dactylorhiza majalis; orchid;
Orchidaceae; next-generation sequencing

Dactylorhiza Necker ex Nevski is a temperate orchid genus known from its complex evolutionary relationships between species driven by high frequency of hybridization, introgression, and polyploidization. Reticulate evolution pattern along with relatively great morphological variability within a species is considered a major challenge in taxonomy of the genus (Hedrén 1996; Pillon et al. 2007, and references therein). *Dactylorhiza majalis* (Rchb.) P.F. Hunt et Summerh. can be found in western and central Europe, Baltic region, and northern Russia (Hultén and Fries 1986; Balao et al. 2016). It is an allotetraploid species belonging to a polyploid complex formed iteratively by crosses between *Dactylorhiza incarnata* s.l. and *Dactylorhiza maculata* s.l. with the last species being always maternal parent (Hedrén et al. 2008 and references therein). Complete, annotated plastidial genome, upon which new molecular markers can be described, would be a valuable tool in untangling evolutionary history within the genus. Chloroplast genomes provide researchers with data invaluable for resolving major phylogenetic relationships between orchid subfamilies (Givnish et al. 2015). Currently, only 15 complete chloroplast genomes are available within the subfamily *Orchidoideae* (Delannoy et al. 2011; Lin et al. 2015; Yu et al. 2015; Zhu et al. 2016; Roma et al. 2018; Lallemand et al. 2019; Oh et al. 2019). This makes this subfamily largely underrepresented among other orchids. Species from the *Dactylorhiza* genus were so far only subject to phylogenetic studies employing ITS, microsatellite loci, selected marker genes, or morphology, and results of these still often remain incongruent (Bateman et al. 2003; Shipunov et al. 2004; Balao et al. 2016; Jin et al. 2017).

Fresh leaves were collected from an individual growing in Psary, Poland (N50°22'07.4" E19°04'53.3"). Leaves dried in silica gel (voucher SG-13237, Herbarium of University of Gdansk, UGDA) were used for extraction of the total genomic DNA with Dneasy Plant Mini Kit (Qiagen, Hilden, Germany). Sequencing library was generated with Accel-NGS[®] 1S Plus DNA Library Kit (Swift Biosciences Inc., Ann Arbor, MI). Next-generation paired-end sequencing was performed with Illumina HiSeq 4000 (San Diego, CA). The obtained reads were used for genome assembly with the Geneious software version 10.2.4 (<https://www.geneious.com>) with medium-low sensitivity parameters and a subset of 25% of the reads, followed by mapping to the closest reference plastome (*Platanthera japonica*, NC_037440.1), and reassembly with medium sensitivity parameters to increase the assembly quality. Annotation was performed within Geneious as well as using GeSeq (Tillich et al. 2017), and manually corrected afterwards. Phylogenetic relationships of *D. majalis* with other orchids were inferred from maximum-likelihood analyses with RAXML-NG (Kozlov et al. 2019) using selected available complete orchid plastomes aligned with MAFFT (Katoh and Standley 2013).

The chloroplast DNA of *D. majalis* is 154,108 bp in length, presenting the overall GC content (the proportion of guanine and cytosine bases) of 36.99%. Consistent with other known orchid plastomes it is comprised two inverted repeats (IRa and IRb) with 26,580 bp in length, an 83,196 bp large single-copy region (LSC), and a 17,752 bp long small single-copy region (SSC). A total of 134 genes were annotated, of which 113 are unique. These are 4 rRNA genes, 30 tRNA genes, and




Figure 1. Phylogenetic relationships inferred from maximum-likelihood analyses with full length plastome sequences of *Orchidaceae* representatives, including the newly assembled *D. majalis* plastome. Node support values are derived from RAXML assessment with 1000 bootstraps replicates. Non-orchid monocots were used for tree rooting.

79 protein-coding genes. Twenty genes are duplicated in the IR region. Additionally, 12 protein-coding genes and 6 tRNA genes contain introns. The annotated sequence was deposited at GenBank with the accession number MK984209. Results of the phylogenetic relationships investigation between *D. majalis* and other members of the *Orchidaceae* showed its clustering together with representatives of the subtribe *Orchidinae* (Figure 1). The complete plastome sequence we provided here constitutes a valuable aid for addressing the taxonomic uncertainties within the genus *Dactylorhiza*, as well as analysing the genetic diversity of the *Orchidaceae* family.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The presented research was supported by the grant funded by Narodowe Centrum Nauki (National Science Center, Poland), grant No. [2015/18/A/NZ8/00149].

ORCID

Michał May <http://orcid.org/0000-0001-8027-1369>
 Alžběta Novotná <http://orcid.org/0000-0002-9835-0855>
 Julita Minasiewicz <http://orcid.org/0000-0002-0330-7011>
 Marc-Andre Selosse <http://orcid.org/0000-0003-3471-9067>
 Marcin Jąkowski <http://orcid.org/0000-0002-5481-9148>

References

Balao F, Tannhäuser M, Lorenzo MT, Hedrén M, Paun O. 2016. Genetic differentiation and admixture between sibling allopolyploids in the *Dactylorhiza majalis* complex. *Heredity*. 116:351–361.

Bateman RM, Hollingsworth PM, Preston J, Yi-Bo L, Pridgeon AM, Chase MW. 2003. Molecular phylogenetics and evolution of Orchidinae and selected Habenariinae (*Orchidaceae*). *Bot J Linn Soc*. 142:1–40.

Delannoy E, Fujii S, Colas Des Francs-Small C, Brundrett M, Small I. 2011. Rampant gene loss in the underground orchid *Rhizanthella gardneri* highlights evolutionary constraints on plastid genomes. *Mol Biol Evol*. 28:2077–2086.

Givnish TJ, Spalink D, Ames M, Lyon SP, Hunter SJ, Zuluaga A, Iles WJD, Clements MA, Arroyo MTK, Leebens-Mack J, et al. 2015. Orchid phylogenomics and multiple drivers of their extraordinary diversification. *Proc R Soc B Biol B*. 282:20151553.

Hedrén M. 1996. Genetic differentiation, polyploidization and hybridization in northern European *Dactylorhiza* (*Orchidaceae*): evidence from allozyme markers. *Plant Syst Evol*. 201:31–55.

Hedrén M, Nordström S, Ståhlberg D. 2008. Polyploid evolution and plastid DNA variation in the *Dactylorhiza incarnata/maculata* complex (*Orchidaceae*) in Scandinavia. *Mol Ecol*. 17:5075–5091.

Hultén E, Fries M. 1986. Atlas of north European vascular plants: north of the tropic of cancer. Germany: Koeltz Scientific Books.

Jin WT, Schuitman A, Chase MW, Li JW, Chung SW, Hsu TC, Jin XH. 2017. Phylogenetics of subtribe *Orchidinae* s.l. (*Orchidaceae; Orchidoideae*) based on seven markers (plastid matK, psaB, rbcL, trnL-F, trnH-psbA, and nuclear nrITS, Xdh): implications for generic delimitation. *BMC Plant Biol*. 17:222.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. 30:772–780.

Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019. RAXML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*. btz305, doi:10.1093/bioinformatics/btz305

Lallemant F, May M, Ichnatowicz A, Jąkowski M. 2019. The complete chloroplast genome sequence of *Platanthera chlorantha* (*Orchidaceae*). *Mitochondrial DNA B*. 4:2683–2684.

Lin CS, Chen JJW, Huang YT, Chan MT, Daniell H, Chang WJ, Hsu CT, Liao DC, Wu FH, Lin SY, et al. 2015. The location and translocation of *ndh* genes of chloroplast origin in the *Orchidaceae* family. *Sci Rep*. 5:9040.

Oh SH, Suh HJ, Park J, Kim Y, Kim S. 2019. The complete chloroplast genome sequence of a morphotype of *Goodyera schlechtendaliana* (*Orchidaceae*) with the column appendages. *Mitochondrial DNA B*. 4: 626–627.

Pillon Y, Fay MF, Hedrén M, Bateman RM, Devey DS, Shipunov AB, van der Bank M, Chase MW. 2007. Evolution and temporal diversification

- of western European polyploid species complexes in *Dactylorhiza* (*Orchidaceae*). *Taxon*. 56:1185–1208.
- Roma L, Cozzolino S, Schlüter PM, Scopece G, Cafasso D. 2018. The complete plastid genomes of *Ophrys iricolor* and *O. sphegodes* (*Orchidaceae*) and comparative analyses with other orchids. *PLoS One*. 13:e0204174.
- Shipunov AB, Fay MF, Pillon Y, Bateman RM, Chase MW. 2004. *Dactylorhiza* (*Orchidaceae*) in European Russia: combined molecular and morphological analysis. *Am J Bot*. 91:1419–1426.
- Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, Greiner S. 2017. GeSeq – versatile and accurate annotation of organellar genomes. *Nucleic Acids Res*. 45:W6–W11.
- Yu C, Lian Q, Wu K, Yu S, Xie L, Wu Z. 2015. The complete chloroplast genome sequence of *Anoectochilus roxburghii*. *Mitochondrial DNA*. 27:1–2.
- Zhu S, Niu Z, Yan W, Xue Q, Ding X. 2016. The complete chloroplast genome sequence of *Anoectochilus emeiensis*. *Mitochondrial DNA A*. 27:3565–3566.