



Factors associated with poor adherence to medication among hypertensive patients in twelve low and middle income Sub-Saharan countries

Diane Macquart de Terline, Adama Kane, Kouadio Euloge Kramoh, Ibrahim Ali Toure, Jean Bruno Mipinda, Ibrahima Bara Diop, Carol Nhavoto, Dadhi M Balde, Beatriz Ferreira, Martin Dèdonougbo Houenassi, et al.

► To cite this version:

Diane Macquart de Terline, Adama Kane, Kouadio Euloge Kramoh, Ibrahim Ali Toure, Jean Bruno Mipinda, et al.. Factors associated with poor adherence to medication among hypertensive patients in twelve low and middle income Sub-Saharan countries. PLoS ONE, 2019, 14 (7), pp.e0219266. 10.1371/journal.pone.0219266 . hal-02303488

HAL Id: hal-02303488

<https://hal.sorbonne-universite.fr/hal-02303488>

Submitted on 2 Oct 2019


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RESEARCH ARTICLE

Factors associated with poor adherence to medication among hypertensive patients in twelve low and middle income Sub-Saharan countries

Diane Macquart de Terline^{1,2,3*}, Adama Kane⁴, Kouadio Euloge Kramoh⁵, Ibrahim Ali Toure⁶, Jean Bruno Mipinda⁷, Ibrahima Bara Diop⁸, Carol Nhavoto⁹, Dadhi M. Balde¹⁰, Beatriz Ferreira⁹, Martin Dèdonougbo Houenassi¹¹, Méo Stéphane Ikama¹², Samuel Kingue¹³, Charles Kouam Kouam¹⁴, Jean Laurent Takombe¹⁵, Emmanuel Limbole¹⁶, Liliane Mfeukeu Kuate¹⁷, Roland N'guetta⁵, Jean Marc Damorou¹⁸, Zouwera Sesso¹⁸, Abdallahi Sidy Ali¹⁹, Marie-Cécile Perier², Michel Azizi^{3,20,21}, Jean Philippe Empana², Xavier Jouven^{2,3,22}, Marie Antignac^{1,2}



OPEN ACCESS

Citation: Macquart de Terline D, Kane A, Kramoh KE, Ali Toure I, Mipinda JB, Diop IB, et al. (2019) Factors associated with poor adherence to medication among hypertensive patients in twelve low and middle income Sub-Saharan countries. PLoS ONE 14(7): e0219266. <https://doi.org/10.1371/journal.pone.0219266>

Editor: Yan Li, Shanghai Institute of Hypertension, CHINA

Received: February 13, 2019

Accepted: June 19, 2019

Published: July 10, 2019

Copyright: © 2019 Macquart de Terline et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by: INSERM (Institut national de la Santé et de la Recherche Médicale), AP-HP (Assistance Publique Hôpitaux de Paris), Paris Descartes University.

Competing interests: The authors have declared that no competing interests exist.

1 Department of Pharmacy, Saint Antoine hospital, HUEP, AP-HP, Paris, France, **2** Paris Cardiovascular Research Centre, INSERM U970, European Georges Pompidou Hospital, Paris, France, **3** Paris Descartes University, Paris, France, **4** Cardiology Department, University Hospital of Aristide Le Dantec, Dakar, Senegal, **5** Institute of Cardiology of Abidjan, Abidjan, Côte d'Ivoire, **6** Internal Medicine and Cardiology Department, University Hospital of Lamorde, Niamey, Niger, **7** University hospital of Libreville, Libreville, Gabon, **8** Cardiology Department, University Hospital of Fann, Dakar, Senegal, **9** Instituto do Coração (ICOR), Maputo, Mozambique, **10** Department of Cardiology, University Hospital of Conakry, Conakry, Guinea, **11** National University hospital of Hubert K. MAGA (CNHU-HKM), Cotonou, Bénin, **12** Cardiology Department, National University Hospital of Brazzaville, Marien NGOUABI University, Brazzaville, Congo, **13** University of Yaoundé, Ministry of Public Health, Yaoundé, Cameroon, **14** Internal Medicine Department, Régional Hospital, Bafoussam, Cameroon, **15** General Hospital of Kinshasa, Kinshasa, Democratic Republic of the Congo, **16** Department of Internal Medicine of la Gombe (CMCG), Department of Internal Medicine, Ngaliema Hospital, Kinshasa; Democratic Republic of the Congo, **17** Central Hospital of Yaoundé, Yaoundé; Cameroon, **18** Cardiology department, CH Lomé, Lomé; Togo, **19** Cardiology clinics, Nouakchott, Mauritania, **20** Hypertension Unit, Assistance-Publique Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Paris, France, **21** Institut National de la Santé et de la Recherche Médicale, Centre d'Investigation Clinique, Paris, France, **22** Department of Cardiology, European Georges Pompidou Hospital, AP-HP, Paris, France

* diane.macquart-de-terline@aphp.fr

Abstract

Introduction

Over the past few decades, the prevalence of hypertension has dramatically increased in Sub-Saharan Africa. Poor adherence has been identified as a major cause of failure to control hypertension. Scarce data are available in Africa.

Aims

We assessed adherence to medication and identified socioeconomics, clinical and treatment factors associated with low adherence among hypertensive patients in 12 sub-Saharan African countries.

Method

We conducted a cross-sectional survey in urban clinics of both low and middle income countries. Data were collected by physicians on demographics, treatment and clinical data among hypertensive patients attending the clinics. Adherence was assessed by questionnaires completed by the patients. Factors associated with low adherence were investigated using logistic regression with a random effect on countries.

Results

There were 2198 individuals from 12 countries enrolled in the study. Overall, 678 (30.8%), 738 (33.6%), 782 (35.6%) participants had respectively low, medium and high adherence to antihypertensive medication. Multivariate analysis showed that the use of traditional medicine (OR: 2.28, 95%CI [1.79–2.90]) and individual wealth index (low vs. high wealth: OR: 1.86, 95%CI [1.35–2.56] and middle vs. high wealth: OR: 1.42, 95%CI [1.11–1.81]) were significantly and independently associated with poor adherence to medication. In stratified analysis, these differences in adherence to medication according to individual wealth index were observed in low-income countries ($p < 0.001$) but not in middle-income countries ($p = 0.17$). In addition, 26.5% of the patients admitted having stopped their treatment due to financial reasons and this proportion was 4 fold higher in the lowest than highest wealth group (47.8% vs 11.4%) ($p < 0.001$).

Conclusion

This study revealed the high frequency of poor adherence in African patients and the associated factors. These findings should be useful for tailoring future programs to tackle hypertension in low income countries that are better adapted to patients, with a potential associated enhancement of their effectiveness.

Introduction

High Blood Pressure (BP) is the worldwide leading global burden of disease risk factor; 16.5% of total deaths and 7.0% of global disability adjusted life years are attributed to high BP [1].

Due to the aging of the population and the obesity epidemic [2], high BP is increasing and despite the availability and effectiveness of multiple antihypertensive class of drugs, hypertension remains poorly controlled worldwide. Over the past few decades, hypertension prevalence has dramatically increased in developing countries [3], especially in Sub-Saharan Africa [1, 4] and tackling its burden is regarded now as a high priority task. Among various factors, poor adherence to antihypertensive medications is a major cause of failure to control hypertension [5] and a considerable proportion of CVD events (9% in Europe) could be attributed to poor adherence to antihypertensive medications [6]. Poor adherence reduces the effectiveness of drugs and represents a significant barrier to achieving better patient outcomes [7–9]. According to the WHO, the extent of poor adherence in developing countries is likely to be even higher given the lack and unequal access to health care and medications [10].

Given the scarcity of health resources available in developing countries, especially in Sub-Saharan Africa, only quality improvement interventions that are cost-efficient are likely to be feasible [11]. Interventions to improve adherence to medication should have multifaceted

dimension [12] and may represent a great opportunity to improve cardiovascular outcomes and reduce health care spending [13, 14] and underutilization of already limited treatment resources [10]. However, to be effective such interventions must be tailored to the particular economical, geographical, sociological and educational context of the patients. Therefore, there is a serious need of accurately assessing not only adherence, but also to identify barriers and behaviors that influences adherence to medication.

Studies on adherence to medication are scarce in sub-Saharan Africa and usually come from single-center studies using various measures of adherence [15] and do not precisely assess other factors specific to the African context.

To cover this unmet need we performed a new analysis of the EIGHT (Evaluation of Hypertension in Sub-Saharan Africa) Study [16], a large multinational study conducted in 12 sub-Saharan African countries. In this paper, we assessed adherence to medication and identify socioeconomics, clinical and treatment factors associated with poor adherence in African hypertensive patients.

Objectives

We therefore performed an analysis of the EIGHT study to assess adherence to medication and to identify socioeconomics, clinical and treatment factors associated with poor adherence among hypertensive patients in 12 sub-Saharan African countries.

Methods

Study design and setting

We conducted an observational cross sectional study during outpatient consultations in cardiology departments of 29 hospitals from 17 cities across 12 sub-Saharan African countries (Benin, Cameroon, Congo (Brazzaville), Democratic Republic of the Congo, Gabon, Guinea, Côte d'Ivoire, Mauritania, Mozambique, Niger, Senegal, Togo) between January 2014 and November 2015.

The study was conceived and designed by a multidisciplinary collaborative team of epidemiologists, cardiologists and pharmacists from Africa and France. The team had extensive prior research experience and existing collaborations with a network of physician-scientists in Africa such as in the field of rheumatic heart disease [17], sickle cell disease [18], quality of cardiovascular drugs [19, 20] which aided planning and launch of the present study.

The study was approved by the Ile-de-France III ethics committee (Number 2014-A00710-47) and was declared to the National Commission of Informatics (Number 1762715).

Participants

Patients were enrolled during outpatient consultations in the cardiology departments of the participating hospitals. Patients with hypertension diagnosis, ≥ 18 years old were eligible to participate. Each patient received an information leaflet about the study. Further, the on-site physician presented and explained the study in the regional language to all patients meeting eligibility criteria and collected patient consent. Due to the observational, non-interventional nature of the study, with anonymized data, informed consent was verbally obtained and registered by the same physician. Patients who agreed to participate completed a standard questionnaire while waiting for their appointment. Participating investigators (physicians) at each center received a training note detailing the study with standardized instructions on how they should interact with the patients while filling the questionnaire in the survey.

Questionnaire and clinical data report

Questionnaire. A dedicated questionnaire was conceived by the authors of the study. A first section was completed by patients and collected data on patient's demographics (age, gender, marital status etc.) and adherence to treatment. Adherence to medication was assessed by the 8-items Medication Adherence Scale [21]. The questionnaire is scored from 0 to 8 with scores of <6, 6 to <8, and 8 reflecting low, medium, and high adherence, respectively. Additional items on patient financial situation on adherence and reasons of non-adherence (list of options asking patients to document the main reason for not taking their treatment) were included. A second section was filled out by the physician during the consultation and collected data on patient's sociodemographic, drug regimen and clinical data.

A pilot investigation involving 90 patients who tested the questionnaire was conducted in January 2014 in Côte d'Ivoire. This pilot study confirmed patient's understanding, feasibility of completion, and resulted in few modifications which were incorporated in the questionnaire given to patients in the final survey (S1 File Survey questionnaire).

Country level income and individual wealth category. The country income was obtained from the World Bank database (<http://data.worldbank.org/country>, accessed 12 January 2015) and categorized into low income and middle income countries. Individual patient wealth categories were assessed by the treating physician and classified as low, middle and high:

- "Low" defined poor patients who have difficulties to afford medical consultations
- "Middle" defined patients who can manage with paying medical consultations
- "High" defined patients who have no difficulties to pay medical consultations.

Statistical methods

Continuous and categorical variables were expressed as mean (standard deviation) and numbers (percentages) where appropriate. Comparisons on qualitative variables were performed with the chi-squared test while those on quantitative variables were performed with the Student test.

First, the odds ratios (ORs) and 95% confidence intervals of socioeconomics, clinical and medication factors for low adherence were estimated in logistic regression. A random effect on the country was added (generalized estimated equation models) to account for inter-country variability. Then, in a multivariate analysis, models were adjusted on all factors with a pValue of less than 0.15 in the univariate analysis.

The level of adherence according to the individual wealth index was explored separately in low and middle income countries using a chi-squared test.

All analyses were performed through scripts developed in the R software (3.4.1 (2017-06-30)). The level of significance was set at $p < 0.05$.

Results

Participants

The EIGHT study enrolled 2198 individuals in 12 sub-Saharan countries between January 2014 and November 2015. Patients' characteristics are reported in Table 1. A total of 1017 patients (46.3%) were from 6 low-income countries (Benin, Democratic Republic of the Congo, Guinea, Mozambic, Niger, and Togo), and 1181 (53.7%) were from 6 middle-income countries (Cameroon, Congo [Brazzaville], Gabon, Côte d'Ivoire, Mauritania, and Senegal). A

Table 1. Characteristics of the participants according to their adherence level.

	GLOBAL	ADHERENCE TO MEDICATION			
		LOW	MEDIUM	HIGH	PValue*
N, (%)	2198 (100)	678 (30.8)	738 (33.6)	782 (35.6)	
Age, mean (sd)	58.3 (11.8)	58.0 (12.1)	58.4 (12.1)	58.4 (11.3)	0.81
Male, N (%)	874 (39.8)	258 (38.1)	299 (40.5)	317 (40.5)	0.55
Patient wealth index, N (%)					<0.001
Low	376 (17.5)	143 (21.7)	132 (18.4)	101 (13.2)	
Middle	1053 (49.2)	342 (51.9)	360 (50.3)	351 (45.7)	
High	713 (33.3)	174 (26.4)	224 (31.3)	315 (41.1)	
Countries income level (low vs middle), N (%)	1017 (46.3)	351 (51.8)	320 (43.4)	346 (44.2)	0.002
Location (Urban vs Rural), N (%)	1702 (78.9)	506 (76.6)	571 (78.8)	625 (81.1)	0.11
Recent diagnosis of hypertension (<1 year), N (%)	335 (15.6)	101 (15.1)	108 (15.0)	126 (16.5)	0.66
Body Mass Index, mean (sd)	27.9 (5.8)	28.0 (5.9)	27.6 (5.7)	28.1 (5.9)	0.27
Cardiovascular risks factors, N (%)					
None	461 (28.0)	153 (28.8)	146 (26.6)	162 (28.7)	0.77
Sedentary Lifestyle	649 (39.5)	206 (38.7)	232 (42.3)	211 (37.4)	0.23
Hypercholesterolemia	328 (19.9)	104 (19.5)	99 (18.0)	125 (22.2)	0.22
Diabetes mellitus	288 (17.5)	102 (19.2)	94 (17.1)	92 (16.3)	0.44
Hypertriglyceridemia	88 (5.3)	29 (5.5)	29 (5.3)	30 (5.3)	0.99
Tobacco use	84 (5.1)	31 (5.8)	31 (5.6)	22 (3.9)	0.27
Cardiovascular complications (none vs a least one), N (%)	1174 (55.8)	382 (59.0)	372 (52.3)	420 (56.5)	0.03
Use of traditional medicine, N (%)	512 (24.1)	213 (32.0)	166 (23.1)	133 (17.9)	<0.001
No. of antihypertensive prescribed, mean (sd)	1.97 (0.18)	1.97 (0.17)	1.96 (0.19)	1.97 (0.18)	0.59
Antihypertensive drug class, N (%)					
Calcium channel blocker	1219 (57.4)	358 (54.4)	416 (58.7)	445 (58.9)	0.17
Diuretic	1167 (55.0)	355 (54.0)	387 (54.6)	425 (56.2)	0.67
Angiotensin-converting-enzyme inhibitor	981 (46.2)	299 (45.4)	343 (48.4)	339 (44.8)	0.36
Beta-blocker	466 (22.0)	129 (19.6)	154 (21.7)	183 (24.2)	0.11
Angiotensin II receptor antagonist	321 (15.1)	85 (12.9)	103 (14.5)	133 (17.6)	0.04
Systolic BP, mean (sd)	149.1 (23.6)	152.4 (23.8)	150.3 (24.8)	145.0 (21.5)	<0.001
Diastolic BP, mean (sd)	88.4 (14.3)	90.8 (15.3)	88.3 (14.9)	86.4 (12.4)	<0.001

*p-value for a chi-squared test or t-test where appropriate.

<https://doi.org/10.1371/journal.pone.0219266.t001>

greater proportion of participants were women (60.2%). Age ranged from 19 to 95 years and mean age was 57.7 ± 12.0 years for women and 59.2 ± 11.4 years for men. A majority of patients were living in urban setting ($n = 1702$, 78.9%) compared to rural way of life ($n = 455$, 21.1%). Individual wealth index was low, midlevel, and high in 376 (17.6%), 1053 (49.2%), and 713 patients (33.3%), respectively. Most of the patients ($n = 1816$, 84.4%) were diagnosed with hypertension for more than one year prior to inclusion to the survey. Overall, 653 (29.7%), 927 (42.2%), and 543 (24.7%) of the participants were prescribed 1, 2, and ≥ 3 antihypertensive drugs, respectively; a quarter (24.1%) of patients admitted using traditional medicine.

Adherence to medication and factors associated with low adherence level. In the study population, 678 (30.8%), 738 (33.6%), 782 (35.6%) participants had respectively low, medium and high adherence to antihypertensive drugs.

As shown in Fig 1, the distribution of adherence level differed across countries ($p < 0.001$), the proportion of low adherence to medication ranged from 15.0% in Senegal to 55.2% in Democratic Republic of the Congo.




Fig 1. Adherence to medication according to countries. Squares represent OR and lines, 95% confidence interval (CI). ORs derived from logistic regression analysis adjusted on age, patient wealth index, complications, use of traditional medicine and antihypertensive drugs with a random effect on country to account for intra and inter-country variability.

<https://doi.org/10.1371/journal.pone.0219266.g001>

In univariate analysis, low adherence was not significantly associated with age, sex, BMI, or the presence of cardiovascular complications or comorbidities (Table 2). Contrariwise, low adherence was significantly associated with patient wealth index ($p < 0.001$); patients with a lower wealth index were more likely to be less adherent to their antihypertensive medications (OR: 1.83, 95%CI [1.38–2.45]).

Patients using traditional medicine were also less adherent to their medication ($p < 0.001$), with an OR reaching 2.22, 95%CI [1.78–2.78]. Patients prescribed a calcium channel blocker were more adherent to their treatment (OR 0.77, 95%CI [0.63–0.93] $p = 0.007$); this association was not observed for any other class of antihypertensive drug, nor with the number of prescribed antihypertensive drugs.

In the multivariate analysis, the use of traditional medicine and poor individual wealth index remained significantly and independently associated with low adherence to medication (Fig 2). The odds of low adherence increased to 2.28 fold (OR: 2.28, 95%CI [1.79–2.90], $p < 0.001$) in patients using traditional medicine, and 1.86 fold (OR: 1.86, 95%CI [1.35–2.56]) and 1.42 (OR = 1.42, 95%CI [1.11–1.81]) in patients with low and middle individual wealth as compared to those with high individual wealth ($p < 0.001$). Conversely, prescription of calcium channel blocker was associated with a better adherence (OR 0.8 [0.65–0.99], $p = 0.007$)).

Stratified analysis by country level income

We then further investigated the association between level of adherence and individual wealth category separately in low- and middle- income countries.

Table 2. Odds ratios for low adherence to medication according to socioeconomics, clinical and medication factors in univariate analysis.

Outcomes	OR [95% CI]	pValue
Age	0.99 [0.98–1.00]	0.08
Male	0.95 [0.79–1.16]	0.64
Patient wealth index		<0.001
Low	1.83 [1.38–2.45]	
Middle	1.42 [1.14–1.80]	
High	Ref.	
Countries income level (low vs middle)	1.18 [0.65–2.14]	0.59
Location (Urban vs Rural)	0.92 [0.71–1.20]	0.51
Recent diagnosis of hypertension (<1 year)	0.95 [0.74–1.22]	0.56
Body Mass Index	1.01 [0.99–1.03]	0.17
Cardiovascular risks factors (none vs a least one)	0.95 [0.75–1.20]	0.67
Complications (none vs a least one)	0.84 [0.69–1.03]	0.10
Use of traditional medicine	2.22 [1.78–2.78]	<0.001
No. of antihypertensive prescribed	1.01[0.59–1.74]	0.96
Antihypertensive drug class		
Calcium channel blocker	0.77 [0.63–0.93]	0.007
Diuretic	0.98 [0.81–1.19]	0.82
Angiotensin-converting-enzyme inhibitor	1.05 [0.86–1.28]	0.65
Beta-blocker	0.92 [0.72–1.16]	0.46
Angiotensin II receptor antagonist	0.77 [0.58–1.02]	0.07

Odds ratios were estimated by logistic regression analysis with a random effect on countries using adherence (medium and high adherence) as reference category.

<https://doi.org/10.1371/journal.pone.0219266.t002>

In low-income countries, the proportion of low adherence increased progressively and considerably with decreasing level of individual patient wealth and was 24.3%, 39.7% and 49.0% in patients with high, middle and low individual wealth, respectively ($p < 0.001$, Fig 3).

In contrast, in middle income countries, we observed a minor and non-significant increase of the proportion of low adherence between individual wealth categories (24.5%, 27.5% and 31.3%, in patients with high, middle and low individual wealth respectively).

Adherence and patients' affordability. A total of 563 patients (26.5%) reported having stopped their treatment due to financial reasons. This proportion was much higher in the lowest wealth group (47.8%) in contrast to the highest wealth group (11.4%) ($p < 0.001$). After forgetfulness, high cost of medications was the second main reason mentioned by patients for not taking their medications (Fig 4). Among patients categorized as good adherent, 19.2% ($n = 279$) reported having stopped their treatment due to financial reasons.

Discussion

Key results

We obtained large data on factors associated with antihypertensive drugs adherence in cardiology practices in 12 African countries. More than 60% of patients presented a suboptimal (low or medium) adherence. Furthermore, we observed several factors significantly and independently associated with low adherence including the use of traditional medicine and low wealth index.




Fig 2. Odds ratios of patient's factors significantly associated with low adherence level in multivariate analysis. Squares represent OR and lines, 95% confidence interval (CI). ORs derived from logistic regression analysis adjusted on age, patient wealth index, complications, use of traditional medicine and antihypertensive drugs with a random effect on country to account for intra and inter-country variability.

<https://doi.org/10.1371/journal.pone.0219266.g002>

Stratified analysis shows that these differences of adherence levels according to individual wealth index were observed in low-income countries and not in middle-income countries. More than one quarter of patients admitted having stopped their treatment due to financial reasons; this proportion was 4 fold higher in the lowest compared to the highest wealth group.

In our study, two thirds of participants had respectively low or medium adherence to antihypertensive medication. A systematic review of studies evaluating adherence to antihypertensive drugs in developing countries was published in 2017 [15]. Of the 22 studies selected, 6 were located in sub-Saharan African countries (3 in Nigeria, 1 in Ethiopia, 1 in Ghana, 1 in Zimbabwe). The percentage of poor adherence ranged from 35.4% in Ethiopia to 93.3% in Ghana. Two of the studies used the MMAS-8 self-questionnaire, reporting a very high percentage of poor adherence (92.5% [95% CI 89.3–95.8] in Nigeria and 93.3% [95% CI 90.9–95.8] in Ghana).

There is strong consensus that medication low adherence rates are higher among African patients with respect to others regions. A meta-analysis of non-adherence to antihypertensive therapy published in 2017 [22] showed major regional differences with a higher percentage of poor adherence levels among African patients (62.4%) compared to Asians (43.5%), Europeans (36.6%), and Americans (36.6%).

The multiplicity of reasons related to patient, clinician, and health system factors make poor adherence a challenging problem to address [23]. In our study, we identified relevant factors associated with low adherence, which could help to shed light on complex issue of low adherence in the particular context of the study.




Fig 3. Percentage of patients according to their adherence level by patient wealth index stratified by country-level income.

<https://doi.org/10.1371/journal.pone.0219266.g003>




Fig 4. Patients' main reason for not taking their treatment.

<https://doi.org/10.1371/journal.pone.0219266.g004>

Socio-economic factors are important to consider as highlighted by the present report. It has been described that low adherence and economic level of the patients are related [24]. Our study addressed for the first time the relative association of individual wealth and country level income with adherence. Although financial inequalities represent a barrier to access to treatment, the situation is potentially even more noticeable in sub-Saharan Africa. Current programs to improve access to medication in low-income and middle-income countries operate within complex health systems and reducing the wholesale price of medicines might not always or immediately translate to improved patient access [25].

Our results underlined the strong association between low adherence and the use of traditional medicine. Indeed, in developing countries, an important part of the population regularly uses traditional medicine [26, 27] define as to a set of healthcare practices that are delivered outside of the mainstream healthcare system including traditional, complementary and alternative medicine [28]. Therefore, there is a real challenge of involving traditional healers in conventional medicine. Few collaborative projects exists in sub Saharan Africa in the fields of HIV [29] and mental disorder [30] for example. Also, effort were made to conceal these two sides of medicine, in Ghana were the government began a pilot project in 17 public hospitals for integrating herbal medicine into biomedicine [31]. In conclusion, to be widely accepted, interventions and programs to improve adherence to medication in developing countries should include traditional healthcare professionals.

Channel calcium blockers are widely used in the treatment of hypertension in black African. This antihypertensive treatment is recommended as the first-line treatment for recently diagnosed black patients [32]. Recent study has shown that channel calcium blocker associated with hydrochlorothiazide or perindopril are more effective than other dual therapies for lowering blood pressure in sub-Saharan African patients [33]. From a patient point of view, this treatment presents less inconvenient side-effects such as cough with Angiotensin-converting-enzyme inhibitor or frequent urination with diuretics.

These reasons might explain a high level of confidence in calcium channel blocker treatment by health professionals and patients, and explain the association with adherence. However this interpretation might be specific to the African context, in Europe, the order of persistence rate from the highest to the lowest is angiotensin II receptor antagonist, angiotensin converting enzyme inhibitor, channel calcium blocker, beta-blocker, and diuretic [34].

In addition to these context specific factors, our results identified reasons for treatment discontinuation commonly known by health professionals in developed countries. Indeed, the most common cause for low adherence was forgetfulness, which accounted for more than 1/3 treatment discontinuity. Therefore, intervention to enhance adherence should be based upon medication counseling, patient education or any other action to avoid forgetfulness is a way to improve adherence [35].

We acknowledged the following limitations. First, self-reported questionnaire are known to overestimate adherence. In a meta-analysis of 11 studies ($n = 1684$ patients) by Shi et al [36], the pooled correlation coefficient of self-report compared with electronic drug monitoring was 0.45 [95% CI, 0.34–0.56], but self-reported rates of adherence were higher than rates of adherence measured by electronic drug monitors (84% vs 75%). Furthermore, the usual adherence questionnaires were conceived in high income countries, omitting to include discontinuation of treatment due to financial reason. In the particular context of low and middle income countries, future evaluation of adherence should include questions concerning access to medication such as patient affordability [37] and accessibility to medication [38]. However, self-reported medication adherence is a practical way to measure adherence because of its low cost and potential to be easily implemented into the clinical workflow. Also, evidence showed that self-reported adherence is predictive of clinical outcomes and especially in hypertension [6, 21].

This study has also many strengths including its multisite design, with over 2,000 patients from 29 medical centers in 17 cities from 12 countries. Furthermore, this study was supported by a strong and structured collaborative multidisciplinary network. Active involvement of African cardiologists, who are familiar with the problems of this area, helped derive specific questions and analysis.

The EIGHT study embraced 12 African countries which are usually not considered in international studies, such as meta-analysis previously cited [15, 22] or the well-known PURE study providing global data on cardiovascular epidemiology. Our analysis gave the opportunity to extend knowledge on non-adherence in the African continent. However, it is important to bear in mind that these data derived from specific urban clinics likely represent the best case scenario and the magnitude of the problem in the general population with hypertension could be underestimated.”

Conclusion

This study revealed the high level of low adherence in African patients with hypertension, and identified the associated factors specific to the regional context. These findings should be useful for tailoring future programs to tackle hypertension in low income countries, which are better adapted to patients, with a potential associated enhancement of their effectiveness.

Supporting information

S1 File. Survey questionnaire.

(PDF)

S2 File. Study dataset, individual data about patients’ characteristics and answers to the questionnaire used in the study.

(CSV)

Acknowledgments

The authors would like to thank the patients and the members of the network for their participation in this research.

Author Contributions

Conceptualization: Ibrahima Bara Diop, Xavier Jouven, Marie Antignac.

Data curation: Kouadio Euloge Kramoh.

Formal analysis: Diane Macquart de Terline, Marie-Cécile Perier, Jean Philippe Empana, Marie Antignac.

Investigation: Adama Kane, Kouadio Euloge Kramoh, Ibrahim Ali Toure, Jean Bruno Mipinda, Ibrahima Bara Diop, Carol Nhavoto, Dadhi M. Balde, Beatriz Ferreira, Martin Dèdonougbo Houenassi, Méo Stéphane Ikama, Samuel Kingue, Charles Kouam Kouam, Jean Laurent Takombe, Emmanuel Limbole, Liliane Mfeukeu Kuate, Roland N’guetta, Jean Marc Damorou, Zouwera Sesso, Abdallahi Sidy Ali.

Methodology: Diane Macquart de Terline, Ibrahima Bara Diop, Jean Philippe Empana, Xavier Jouven, Marie Antignac.

Supervision: Ibrahima Bara Diop, Xavier Jouven, Marie Antignac.

Validation: Diane Macquart de Terline, Adama Kane, Kouadio Euloge Kramoh, Ibrahim Ali Toure, Jean Bruno Mipinda, Ibrahima Bara Diop, Carol Nhavoto, Dadhi M. Balde, Beatriz Ferreira, Martin Dèdonougbo Houenassi, Méo Stéphane Ikama, Samuel Kingue, Charles Kouam Kouam, Jean Laurent Takombe, Emmanuel Limbole, Liliane Mfeukeu Kuate, Roland N'guetta, Jean Marc Damorou, Zouwera Sesso, Abdallahi Sidy Ali, Marie-Cécile Perier, Michel Azizi, Jean Philippe Empana, Xavier Jouven, Marie Antignac.

Visualization: Diane Macquart de Terline, Xavier Jouven, Marie Antignac.

Writing – original draft: Diane Macquart de Terline, Xavier Jouven, Marie Antignac.

Writing – review & editing: Diane Macquart de Terline, Michel Azizi, Jean Philippe Empana, Xavier Jouven, Marie Antignac.

References

1. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-based measurement studies with 19.1 million participants. *Lancet* (London, England). 2017; 389(10064):37–55. Epub 2016/11/20. [https://doi.org/10.1016/s0140-6736\(16\)31919-5](https://doi.org/10.1016/s0140-6736(16)31919-5) PMID: 27863813
2. Chobanian AV. Shattuck Lecture. The hypertension paradox—more uncontrolled disease despite improved therapy. *The New England journal of medicine*. 2009; 361(9):878–87. Epub 2009/08/28. <https://doi.org/10.1056/NEJMsa0903829> PMID: 19710486.
3. Beaney T, Schutte AE, Tomaszewski M, Ariti C, Burrell LM, Castillo RR, et al. May Measurement Month 2017: an analysis of blood pressure screening results worldwide. *The Lancet Global health*. 2018; 6(7): e736–e43. Epub 2018/05/21. [https://doi.org/10.1016/S2214-109X\(18\)30259-6](https://doi.org/10.1016/S2214-109X(18)30259-6) PMID: 29778399.
4. Ibrahim MM, Damasceno A. Hypertension in developing countries. *Lancet* (London, England). 2012; 380(9841):611–9. Epub 2012/08/14. [https://doi.org/10.1016/s0140-6736\(12\)60861-7](https://doi.org/10.1016/s0140-6736(12)60861-7) PMID: 22883510.
5. Bramley TJ, Gerbino PP, Nightengale BS, Frech-Tamas F. Relationship of blood pressure control to adherence with antihypertensive monotherapy in 13 managed care organizations. *Journal of managed care pharmacy: JMCP*. 2006; 12(3):239–45. Epub 2006/04/21. <https://doi.org/10.18553/jmcp.2006.12.3.239> PMID: 16623608.
6. Chowdhury R, Khan H, Heydon E, Shroufi A, Fahimi S, Moore C, et al. Adherence to cardiovascular therapy: a meta-analysis of prevalence and clinical consequences. *European heart journal*. 2013; 34(38):2940–8. Epub 2013/08/03. <https://doi.org/10.1093/eurheartj/ehd295> PMID: 23907142.
7. Bosworth HB, Granger BB, Mendys P, Brindis R, Burkholder R, Czajkowski SM, et al. Medication adherence: a call for action. *American heart journal*. 2011; 162(3):412–24. Epub 2011/09/03. <https://doi.org/10.1016/j.ahj.2011.06.007> PMID: 21884856
8. Cutler DM, Everett W. Thinking outside the pillbox—medication adherence as a priority for health care reform. *The New England journal of medicine*. 2010; 362(17):1553–5. Epub 2010/04/09. <https://doi.org/10.1056/NEJMp1002305> PMID: 20375400.
9. Osterberg L, Blaschke T. Adherence to medication. *The New England journal of medicine*. 2005; 353(5):487–97. Epub 2005/08/05. PMID: 16079372.
10. WHO. Adherence to Long-term Therapies: Evidence for action [Internet]. World Health Organization. 2003 [cited december 2018]. www.who.int/chp/knowledge/publications/adherence_report/en/.
11. Gaziano TA. Cardiovascular disease in the developing world and its cost-effective management. *Circulation*. 2005; 112(23):3547–53. <https://doi.org/10.1161/CIRCULATIONAHA.105.591792> PMID: 16330695.
12. Haynes RB, McDonald H, Garg AX, Montague P. Interventions for helping patients to follow prescriptions for medications. *The Cochrane database of systematic reviews*. 2002;(2):Cd000011. Epub 2002/06/22. <https://doi.org/10.1002/14651858.CD000011> PMID: 12076376.
13. Choudhry NK. Improving the pathway from cardiovascular medication prescribing to longer-term adherence: new results about old issues. *Circulation Cardiovascular quality and outcomes*. 2010; 3(3):223–5. Epub 2010/05/21. <https://doi.org/10.1161/CIRCOUTCOMES.110.957142> PMID: 20484200.
14. Dragomir A, Cote R, Roy L, Blais L, Lalonde L, Berard A, et al. Impact of adherence to antihypertensive agents on clinical outcomes and hospitalization costs. *Medical care*. 2010; 48(5):418–25. Epub 2010/04/16. <https://doi.org/10.1097/MLR.0b013e3181d567bd> PMID: 20393367.

15. Nielsen JO, Shrestha AD, Neupane D, Kallestrup P. Non-adherence to anti-hypertensive medication in low- and middle-income countries: a systematic review and meta-analysis of 92443 subjects. *Journal of human hypertension*. 2017; 31(1):14–21. Epub 2016/06/17. <https://doi.org/10.1038/jhh.2016.31> PMID: 27306087.
16. Antignac M, Diop IB, Macquart de Terline D, Kramoh KE, Balde DM, Dzudie A, et al. Socioeconomic Status and Hypertension Control in Sub-Saharan Africa: The Multination EIGHT Study (Evaluation of Hypertension in Sub-Saharan Africa). *Hypertension* (Dallas, Tex: 1979). 2018; 71(4):577–84. Epub 2018/01/31. <https://doi.org/10.1161/hypertensionaha.117.10512> PMID: 29378852.
17. Marijon E, Ou P, Celermajer DS, Ferreira B, Mocumbi AO, Jani D, et al. Prevalence of rheumatic heart disease detected by echocardiographic screening. *The New England journal of medicine*. 2007; 357(5):470–6. Epub 2007/08/03. <https://doi.org/10.1056/NEJMoa065085> PMID: 17671255.
18. Ranque B, Menet A, Diop IB, Thiam V, Diallo D, Diop S, et al. Early renal damage in patients with sickle cell disease in sub-Saharan Africa: a multinational, prospective, cross-sectional study. *Lancet Haematol*. 2014;e64–73. [https://doi.org/10.1016/S2352-3026\(14\)00007-6](https://doi.org/10.1016/S2352-3026(14)00007-6) PMID: 27030156
19. Antignac M, Diop BI, Macquart de Terline D, Bernard M, Do B, Ikama SM, et al. Fighting fake medicines: First quality evaluation of cardiac drugs in Africa. *Int J Cardiol*. 2017; 243:523–8. <https://doi.org/10.1016/j.ijcard.2017.04.099> PMID: 28641892.
20. Macquart de Terline D, Diop BI, Bernard M, Do B, Ikama MS, N'Guetta R, et al. Substandard drugs among five common antihypertensive generic medications: an analysis from 10 African countries. *J Hypertens*. 2017. <https://doi.org/10.1097/HJH.0000000000001560> PMID: 28957849.
21. Korb-Savoldelli V, Gillaizeau F, Pouchot J, Lenain E, Postel-Vinay N, Plouin PF, et al. Validation of a French version of the 8-item Morisky medication adherence scale in hypertensive adults. *Journal of clinical hypertension* (Greenwich, Conn). 2012; 14(7):429–34. Epub 2012/07/04. <https://doi.org/10.1111/j.1751-7176.2012.00634.x> PMID: 22747615.
22. Abegaz TM, Shehab A, Gebreyohannes EA, Bhagavathula AS, Elnour AA. Nonadherence to antihypertensive drugs: A systematic review and meta-analysis. *Medicine*. 2017; 96(4):e5641. Epub 2017/01/26. <https://doi.org/10.1097/MD.0000000000005641> PMID: 28121920
23. Kini V, Ho PM. Interventions to Improve Medication Adherence: A Review. *Jama*. 2018; 320(23):2461–73. Epub 2018/12/19. <https://doi.org/10.1001/jama.2018.19271> PMID: 30561486.
24. Bowry AD, Shrank WH, Lee JL, Stedman M, Choudhry NK. A systematic review of adherence to cardiovascular medications in resource-limited settings. *Journal of general internal medicine*. 2011; 26(12):1479–91. Epub 2011/08/23. <https://doi.org/10.1007/s11606-011-1825-3> PMID: 21858602
25. Rockers PC, Laing RO, Ashigbie PG, Onyango MA, Mukiira CK, Wirtz VJ. Effect of Novartis Access on availability and price of non-communicable disease medicines in Kenya: a cluster-randomised controlled trial. *The Lancet Global health*. 2019; 7(4):e492–e502. Epub 2019/02/26. [https://doi.org/10.1016/S2214-109X\(18\)30563-1](https://doi.org/10.1016/S2214-109X(18)30563-1) PMID: 30799142.
26. Bodeker G. Lessons on integration from the developing world's experience. *BMJ (Clinical research ed)*. 2001; 322(7279):164–7. Epub 2001/02/13. <https://doi.org/10.1136/bmj.322.7279.164> PMID: 11159579
27. Hollenberg D, Zakus D, Cook T, Xu XW. Re-positioning the role of traditional, complementary and alternative medicine as essential health knowledge in global health: do they still have a role to play? *World health & population*. 2008; 10(4):62–75. Epub 2009/06/25. PMID: 19550163.
28. WHO. World Health Organization Traditional Medicine Strategy 2014–2023. Geneva: 2014.
29. Mills E, Singh S, Wilson K, Peters E, Onia R, Kanfer I. The challenges of involving traditional healers in HIV/AIDS care. *International journal of STD & AIDS*. 2006; 17(6):360–3. Epub 2006/06/01. PMID: 16734953.
30. van der Watt ASJ, Nortje G, Kola L, Appiah-Poku J, Othieno C, Harris B, et al. Collaboration Between Biomedical and Complementary and Alternative Care Providers: Barriers and Pathways. *Qualitative health research*. 2017; 27(14):2177–88. Epub 2017/09/14. <https://doi.org/10.1177/1049732317729342> PMID: 28901831.
31. Appiah B, Amponsah IK, Poudyal A, Mensah MLK. Identifying strengths and weaknesses of the integration of biomedical and herbal medicine units in Ghana using the WHO Health Systems Framework: a qualitative study. *BMC complementary and alternative medicine*. 2018; 18(1):286. Epub 2018/10/24. <https://doi.org/10.1186/s12906-018-2334-2> PMID: 30348173
32. Kramer N, Barasa F, Damasceno A, Dzudie A, Jones E, Mondo C, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *The New England journal of medicine*. 2014; 311(5):507–20. Epub 2019/03/19. 2013/12/20. PMID: 24352797.
33. Ojji DB, Mayosi B, Francis V, Badri M, Cornelius V, Smythe W. Comparison of Dual Therapies for Lowering Blood Pressure in Black Africans. 2019. <https://doi.org/10.1056/NEJMoa1901113> PMID: 30883050.

34. Pruijm MT, Maillard MP, Burnier M. Patient adherence and the choice of antihypertensive drugs: focus on lercanidipine. *Vascular health and risk management*. 2008; 4(6):1159–66. Epub 2008/01/01. <https://doi.org/10.2147/vhrm.s3510> PMID: 19337529
35. Ruppert TM, Dunbar-Jacob JM, Mehr DR, Lewis L, Conn VS. Medication adherence interventions among hypertensive black adults: a systematic review and meta-analysis. *J Hypertens*. 2017; 35(6):1145–54. Epub 2017/01/21. <https://doi.org/10.1097/HJH.0000000000001260> PMID: 28106662.
36. Shi L, Liu J, Fonseca V, Walker P, Kalsekar A, Pawaskar M. Correlation between adherence rates measured by MEMS and self-reported questionnaires: a meta-analysis. *Health and quality of life outcomes*. 2010; 8:99. Epub 2010/09/15. <https://doi.org/10.1186/1477-7525-8-99> PMID: 20836888
37. Attaei MW, Khatib R, McKee M, Lear S, Dagenais G, Igumbor EU, et al. Availability and affordability of blood pressure-lowering medicines and the effect on blood pressure control in high-income, middle-income, and low-income countries: an analysis of the PURE study data. *The Lancet Public health*. 2017; 2(9):e411–e9. Epub 2017/12/19. [https://doi.org/10.1016/S2468-2667\(17\)30141-X](https://doi.org/10.1016/S2468-2667(17)30141-X) PMID: 29253412.
38. Wirtz VJ, Kaplan WA, Kwan GF, Laing RO. Access to Medications for Cardiovascular Diseases in Low- and Middle-Income Countries. *Circulation*. 2016; 133(21):2076–85. Epub 2016/05/25. <https://doi.org/10.1161/CIRCULATIONAHA.115.008722> PMID: 27217433