



HAL
open science

One more proof of the Alexandrov–Fenchel inequality

Dario Cordero-Erausquin, Bo'Az Klartag, Quentin Mérigot, Filippo Santambrogio

► **To cite this version:**

Dario Cordero-Erausquin, Bo'Az Klartag, Quentin Mérigot, Filippo Santambrogio. One more proof of the Alexandrov–Fenchel inequality. *Comptes Rendus. Mathématique*, 2019, 357 (8), pp.676-680. 10.1016/j.crma.2019.07.004 . hal-02321749

HAL Id: hal-02321749

<https://hal.sorbonne-universite.fr/hal-02321749>

Submitted on 21 Oct 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Functional analysis/Geometry

One more proof of the Alexandrov–Fenchel inequality

*Une autre preuve de l'inégalité d'Alexandrov–Fenchel*Dario Cordero-Erausquin^a, Bo'az Klartag^b, Quentin Merigot^c,
Filippo Santambrogio^d^a Institut de mathématiques de Jussieu, Sorbonne Université, 4, place Jussieu, 75252 Paris cedex 05, France^b Department of Mathematics, Weizmann Institute of Science, Rehovot 76100, Israel^c Laboratoire de mathématiques d'Orsay, Université Paris-Sud, CNRS, Université Paris-Saclay, 91405 Orsay, France^d Institut Camille-Jordan, Université Claude-Bernard – Lyon-1, 43, boulevard du 11-Novembre-1918, 69622 Villeurbanne cedex, France

ARTICLE INFO

Article history:

Received 18 April 2019

Accepted after revision 16 July 2019

Available online 11 September 2019

Presented by Gilles Pisier

ABSTRACT

We present a short proof of the Alexandrov–Fenchel inequalities, which mixes elementary algebraic properties and convexity properties of mixed volumes of polytopes.

© 2019 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

R É S U M É

Nous donnons une preuve courte des inégalités d'Alexandrov–Fenchel qui repose sur des propriétés algébriques élémentaires ou de convexité des volumes mixtes de polytopes.

© 2019 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

This short text has the modest goal of presenting another proof of the Alexandrov–Fenchel inequality for mixed volumes. It is similar to the proof for polytopes discussed in the recent work of Shenfeld and van Handel [5], which is mainly devoted to a short and neat treatment of the case of smooth bodies. Our presentation puts the emphasis on the basic algebraic properties of the polynomials involved in the construction of mixed volumes. Actually, once elementary and classical geometric and algebraic properties have been recalled, our argument reduces to the simple Proposition 3 below.

The history of the Alexandrov–Fenchel inequality and of its various proofs until the 1980s is described in the book by Burago and Zalgaller [1, Section 20.3]. The more recent literature contains a proof by Wang [7] that was inspired by Gromov's work [2], in addition to the proof by Shenfeld and van Handel [5]. Applications to combinatorics are described by Stanley [6].

The Minkowski sum of two sets $A, B \subseteq \mathbb{R}^n$ is $A + B = \{a + b; a \in A, b \in B\}$, and we also write $tA = \{tx; x \in A\}$ for $t \in \mathbb{R}$. Minkowski has shown that, when $K_1, \dots, K_N \subseteq \mathbb{R}^n$ are convex bodies, the function

$$\mathbb{R}_+^N \ni (t_1, \dots, t_N) \mapsto Vol_n \left(\sum_{i=1}^N t_i K_i \right) \quad (1)$$

E-mail address: dario.cordero@imj-prg.fr (D. Cordero-Erausquin).<https://doi.org/10.1016/j.crma.2019.07.004>1631-073X/© 2019 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

is a homogeneous polynomial of degree n . Particular cases were discussed earlier by the 19th-century geometer Jacob Steiner. Here, $\mathbb{R}_+^N = \{x \in \mathbb{R}^N; \forall i, x_i > 0\}$, a convex body is a compact, convex set with a non-empty interior and Vol_n is n -dimensional volume.

The *mixed volumes* are defined as the coefficients of the homogeneous polynomial in (1). Specifically, the mixed volume of n compact, convex sets $K_1, \dots, K_n \subseteq \mathbb{R}^n$ is

$$V(K_1, \dots, K_n) = \frac{1}{n!} \frac{\partial^n}{\partial t_1 \partial t_2 \dots \partial t_n} Vol_n \left(\sum_{i=1}^n t_i K_i \right) \Bigg|_{t_1 = \dots = t_n = 0}.$$

The reader is referred to Schneider’s book [4, Chapter 5] for explanations and for the basic properties of the mixed volumes. The mixed volume $V(K_1, \dots, K_n)$ is symmetric (invariant under permuting the convex bodies), continuous in the Hausdorff metric, it satisfies $V(K, \dots, K) = Vol_n(K)$, it is multi-linear with respect to Minkowski addition, and it is invariant under translating the convex bodies. It turns out that $V(K_1, \dots, K_n)$ is always non-negative and monotone increasing (with respect to inclusion) in each of its arguments. When $K_1, \dots, K_n \subseteq \mathbb{R}^n$ have a non-empty interior, their mixed volume is in fact positive.

The Alexandrov–Fenchel inequality from the 1930s states that

$$V(K_1, K_2, K_3, \dots, K_n)^2 \geq V(K_1, K_1, K_3, \dots, K_n) V(K_2, K_2, K_3, \dots, K_n) \tag{2}$$

for any convex bodies $K_1, \dots, K_n \subseteq \mathbb{R}^n$. In other words, this inequality states that the function

$$t \rightarrow V(K_1 + tK_2, K_1 + tK_2, K_3, \dots, K_n) \quad (t > 0) \tag{3}$$

is a quadratic polynomial with a non-negative discriminant. Alternatively, the Alexandrov–Fenchel inequalities express inequalities of Brunn–Minkowski type for mixed volumes. Given $m = 2, 3, \dots, n$, and $(n - m + 2)$ convex bodies $K_1, \dots, K_{n-m+2} \subset \mathbb{R}^n$, it follows from the Alexandrov–Fenchel inequality that the function

$$t \rightarrow f_m(t) := V(\underbrace{K_1 + tK_2, \dots, K_1 + tK_2}_{m \text{ times}}, K_3, \dots, K_{n-m+2})^{1/m} \quad \text{is concave on } \mathbb{R}^+. \tag{4}$$

The inequality $f_m''(0) \leq 0$ in the case $m = 2$ is precisely equivalent to the Alexandrov–Fenchel inequality (2), and it easily implies the other cases. The case $m = n$ is the Brunn–Minkowski inequality for convex sets.

We will prove, by induction on the dimension n , that the functions f_m are concave. To be precise, assuming the Alexandrov–Fenchel inequalities in dimension $\leq n - 1$, we will prove that in dimension n and for $m \geq 3$, the function f_m is concave. Crucial but elementary properties of hyperbolic polynomials allow us to proceed to the desired case $m = 2$, namely:

Lemma 1 (“ $m = 3 \Rightarrow m = 2$ ”). *In order to prove the Alexandrov–Fenchel inequality (2) in \mathbb{R}^n , it suffices to establish property (4) with $m = 3$ for all convex bodies in \mathbb{R}^n .*

Proof. We rely on an elementary linear algebra statement from the Appendix of Hörmander’s book [3]. Introduce the symmetric 3-linear form \tilde{f} on \mathbb{R}^3 , defined on the cone $(\mathbb{R}_+)^3$ by

$$\tilde{f}(x, y, z) = V(x_1 K_1 + x_2 K_2 + x_3 K_3, y_1 K_1 + y_2 K_2 + y_3 K_3, z_1 K_1 + z_2 K_2 + z_3 K_3, K_4, \dots, K_n)$$

for $x, y, z \in (\mathbb{R}_+)^3$, and the associated 3-form $f(x) = \tilde{f}(x, x, x)$, $x \in \mathbb{R}^3$. Our assumption is that, for $x, y \in (\mathbb{R}_+)^3$, the function $t \rightarrow f(x + ty)^{1/3}$ is concave on \mathbb{R}^+ , which is property (i)’ of [3, Proposition A1]. This implies property (iii) of [3, Proposition A1] which gives that

$$\tilde{f}(x, y, z)^2 \geq \tilde{f}(x, x, z) \tilde{f}(y, y, z), \quad \forall x, y, z \in (\mathbb{R}_+)^3$$

and in turn implies (and is equivalent) to the Alexandrov–Fenchel inequalities (2), for instance by picking (at the limit) $x = (1, 0, 0)$, $y = (0, 1, 0)$ and $z = (0, 0, 1)$. \square

As we said, we prove the Alexandrov–Fenchel inequality (2) by induction on the dimension n . The base case of our induction is the case $n = 2$, in which it is well known that the Alexandrov–Fenchel inequality follows from the Brunn–Minkowski inequality. We do not provide here an alternative proof for the Alexandrov–Fenchel inequality in two dimensions.

Thus, assume that we are given $n \geq 3$, and that the Alexandrov–Fenchel inequality is already proven in dimension $n - 1$. Our goal is to prove that, given $m \geq 3$ and $m + n - 2$ convex bodies, the function f_m above is concave (the case $m = 3$ suffices). By translating the convex bodies, it suffices to prove (4) under the additional assumption that the origin belongs to the relative interior of the K_i . By continuity with respect to the Hausdorff distance, it suffices to prove (4) in the case

where $K_1, \dots, K_{n-m+2} \subseteq \mathbb{R}^n$ are polytopes containing the origin in their interior. In fact, according to [4, Theorem 2.4.15], we may even assume that these polytopes are *simple strongly isomorphic*.

Let us briefly explain the definition and the basic properties of simple, strongly isomorphic polytopes, using Schneider [4, Chapter 2] as our main reference. A polytope in \mathbb{R}^n is simple if each of its vertices is contained in exactly n facets. The polytopes K_1, \dots, K_{n-m+2} are strongly isomorphic if for any $\theta \in S^{n-1}$, the (affine) dimension of the convex sets

$$\{x \in K_i; \langle x, \theta \rangle = \sup_{y \in K} \langle y, \theta \rangle\}$$

is the same for all $i = 1, \dots, n - m + 2$. Here $S^{n-1} = \{x \in \mathbb{R}^n; |x| = 1\}$ is the unit sphere in \mathbb{R}^n , and we write $\langle x, y \rangle$ for the standard scalar product between $x, y \in \mathbb{R}^n$.

Let $u_1, \dots, u_N \in S^{n-1}$ be the list of outer unit normals to the facets of K_1 , say. Then this list is also the list of outer unit normals to the facets of K_i , for all i . For $h \in \mathbb{R}_+^N$ we consider the set

$$K[h] = \{x \in \mathbb{R}^n; \langle x, u_j \rangle \leq h_j \text{ for } j = 1, \dots, N\},$$

which is a convex body containing the origin. Denote

$$C = \{h \in \mathbb{R}_+^N; K[h] \text{ is strongly isomorphic to } K_1\}.$$

It is well known, and stated in the next lemma, that C is a convex cone in \mathbb{R}^N which provides a parameterization of the space of all polytopes that are strongly isomorphic to K_1 and with zero in their interior. Moreover, this parameterization is linear with respect to the Minkowski sum (Lemma 2).

Lemma 2. *We have the following properties:*

- (i) *the set $C \subseteq \mathbb{R}_+^N$ is an open convex cone with an apex at zero;*
- (ii) *each polytope strongly isomorphic to K_1 and containing the origin in its interior takes the form $K[h]$ for a certain uniquely determined $h \in C$;*
- (iii) *for any $h, h' \in C$,*

$$K[h + h'] = K[h] + K[h']. \tag{5}$$

Proof. Begin with the proof of (ii). Each polytope strongly isomorphic to K_1 has u_1, \dots, u_N as the unit outer normal to its facets. If such polytope contains the origin in its interior, then it takes the form $K[h]$ for a certain uniquely determined $h \in \mathbb{R}_+^N$. By definition, $h \in C$, and (ii) is proven. Next, it is clear that $th \in C$ whenever $t > 0$ and $h \in C$. We need to prove that C is open and convex and that (5) holds true. First, note that when $K[h]$ is strongly isomorphic to K_1 , the polytope $K[h]$ has a facet whose outer normal is u_j , and hence

$$h_j = \sup_{x \in K[h]} \langle x, u_j \rangle \quad \text{for all } j.$$

Given $h, h' \in C$, the convex set

$$K[h] + K[h']$$

is strongly isomorphic to K_1 , according to [4, Corollary 2.4.12]. It follows from (ii) that $K[h] + K[h'] = K[h'']$ for a certain $h'' \in C$. In order to show that $h'' = h + h'$, we note that

$$h''_j = \sup_{x \in K[h] + K[h']} \langle x, u_j \rangle = \sup_{x \in K[h]} \langle x, u_j \rangle + \sup_{x \in K[h']} \langle x, u_j \rangle = h_j + h'_j.$$

Thus $h'' = h + h' \in C$, and C is convex. Moreover, $K[h] + K[h'] = K[h + h']$ and (iii) is proven. The fact that C is open follows from the simplicity of the polytope $K[h]$ for any $h \in C$, see [4, Lemma 2.4.13]. This completes the proof of (i), and the lemma is proven. \square

We conclude from Lemma 2 that each K_i takes the form $K_i = K[h]$ for a certain $h = h^{(i)} \in \mathbb{R}_+^N$. We will study the function

$$\mathcal{F}(h) := V(\underbrace{K[h], \dots, K[h]}_{m \text{ times}}, K_3, \dots, K_{n-m+2}), \quad h \in C \subset \mathbb{R}^N.$$

With Lemma 2 in hand, it is clear that when working with our simple strongly isomorphic polytopes K_i , the concavity of f_m on \mathbb{R}_+^N is equivalent to the concavity of the function $\mathcal{F}^{1/m}$ on the cone C , since

$$f_m(t) = \mathcal{F}^{1/m}(h^{(1)} + th^{(2)}).$$

For proving that f_m is concave on \mathbb{R}_+^N , it is enough to prove that the function $\mathcal{F}^{1/m}$ is concave on C . We will do it locally by analyzing its Hessian.

A crucial property of \mathcal{F} is that it is homogeneous of degree m on C ; actually, \mathcal{F} may be extended to a homogeneous polynomial of degree m in \mathbb{R}^N as is explained in [4, Chapter 5]. Other properties that follow from the geometry of polytopes will allow us to perform induction on the dimension. For a convex body $K \subseteq \mathbb{R}^n$, we write

$$F_j(K) = \left\{ x \in K ; \langle x, u_j \rangle = \sup_{y \in K} \langle y, u_j \rangle \right\}$$

for the facet whose normal is u_j . In our case, when $K = K_i$, the facets $F_1(K), \dots, F_N(K)$ have positive $(n - 1)$ -dimensional volume. It is wellknown (e.g., follows from [4, Lemma 5.1.5]) that

$$\partial^i \mathcal{F} := \frac{\partial \mathcal{F}}{\partial h_i} = \frac{m}{n} \cdot \underbrace{V(F_i(K[h]), \dots, F_i(K[h]), F_i(K_3), \dots, F_i(K_{n-m+2}))}_{m-1 \text{ times}}$$

where this is a mixed volume of $(n - 1)$ bodies in dimension $n - 1$. The facets satisfy the property of Lemma 2, that is, for $h, h' \in C$,

$$F_i(K[h + h']) = F_i(K[h]) + F_i(K[h']).$$

This follows from Lemma 2 and [4, Theorem 1.7.5]. Therefore, by the induction hypothesis (namely the concavity of a certain function $t \rightarrow f_{m-1}(t)$ for convex bodies in $\mathbb{R}^{n-1} \simeq u_i^\perp$), the function $(\partial^i \mathcal{F})^{1/(m-1)}$ is concave on C , and so in particular, $\partial^i \mathcal{F}$ is log-concave on C . Similarly,

$$\partial^{ij} \mathcal{F} := \frac{\partial^2 \mathcal{F}}{\partial h_i \partial h_j} = c_{ij} \cdot \underbrace{V(F_{ij}(K[h]), \dots, F_{ij}(K[h]), F_{ij}(K_3), \dots, F_{ij}(K_{n-m+2}))}_{m-2 \text{ times}}$$

where $F_{ij}(K) = F_i(K) \cap F_j(K)$ and where $c_{ij} = (m/n) \cdot ((m - 1)/(n - 1)) / \sqrt{1 - \langle u_i, u_j \rangle^2}$ is a non-negative coefficient. Note that $\partial^{ij} \mathcal{F}$ is non-negative for all $i \neq j$. Moreover, $\partial^{ij} \mathcal{F}(h) > 0$ whenever the facets $F_i(K[h])$ and $F_j(K[h])$ intersect in an $(n - 2)$ -dimensional face. We conclude that the Hessian matrix $\nabla^2 \mathcal{F}$ is *irreducible*: for any two indices i, j , there is a chain of indices i, i_1, \dots, i_L, j such that $\partial^{i_1} \mathcal{F}(h) > 0, \partial^{i_1 i_2} \mathcal{F}(h) > 0, \dots, \partial^{i_L j} \mathcal{F}(h) > 0$. The conclusion now follows from the following simple abstract proposition (with $p = m$ and $f = \mathcal{F}$), which is the core of our argument.

Proposition 3. *Let $C \subseteq \mathbb{R}_+^N$ be an open, convex cone and let $p \in (2, +\infty)$. Let $f : C \rightarrow (0, \infty)$ be a smooth function such that*

- (i) *the function f is p -homogeneous;*
- (ii) *$\partial^i f$ is positive and log-concave for $1 \leq i \leq N$ (in other words, $\log(\partial^i f)$ is concave);*
- (iii) *$\partial^{ij} f$ is non-negative for all distinct $i, j \in \{1, \dots, N\}$; moreover, sufficiently many of these numbers are strictly positive, so that the Hessian matrix $\nabla^2 f$ is irreducible.*

Then the function $f^{1/p}$ is concave on the convex cone C .

Proof. For any i , the function $f_i = \partial^i f$ is log-concave and $(p - 1)$ -homogeneous, hence $f_i^{1/(p-1)}$ is concave and so

$$\nabla^2 f_i \leq \frac{p - 2}{p - 1} \frac{\nabla f_i \otimes \nabla f_i}{f_i}. \tag{6}$$

Note that $\sum_i x_i \nabla^2 f_i = \sum_i x_i \partial_i (\nabla^2 f) = (p - 2) \nabla^2 f$, again by homogeneity. By multiplying (6) by x_i and summing over i ,

$$(p - 2) \nabla^2 f = \sum_i x_i \nabla^2 f_i \leq \frac{p - 2}{p - 1} \sum_{i=1}^N x_i \frac{\nabla f_i \otimes \nabla f_i}{f_i} = \frac{p - 2}{p - 1} (\nabla^2 f) D (\nabla^2 f) \tag{7}$$

where D is defined via

$$D = \text{diagonal}(x_1/f_1, \dots, x_N/f_N).$$

Since D has positive entries, we may multiply from the left and right by $D^{1/2}$, hence

$$(p - 2) D^{1/2} (\nabla^2 f) D^{1/2} \leq \frac{p - 2}{p - 1} \left[D^{1/2} (\nabla^2 f) D^{1/2} \right]^2. \tag{8}$$

We conclude that the matrix $M = D^{1/2}(\nabla^2 f)D^{1/2}$ thus has no spectrum in the interval $(0, p - 1)$. The matrix M has non-negative off-diagonal entries, it is irreducible, and it has an eigenvector $D^{-1/2}x$ with non-negative entries, corresponding to the eigenvalue $p - 1$. By the Perron–Frobenius theorem, this is the simple, maximal eigenvalue; actually, since we are dealing with a symmetric matrix, this can also be proved by elementary computations without using the Perron–Frobenius theorem. Hence,

$$M \leq (p - 1) \frac{(D^{-1/2}x) \otimes (D^{-1/2}x)}{|D^{-1/2}x|^2} = (p - 1) \frac{D^{-1/2}(x \otimes x)D^{-1/2}}{p \cdot f}.$$

Multiplying by $D^{-1/2}$ on the left and on the right, we get as desired

$$\nabla^2 f \leq (p - 1) \frac{D^{-1}(x \otimes x)D^{-1}}{p \cdot f} = \frac{p - 1}{p} \frac{\nabla f \otimes \nabla f}{f}. \quad \square$$

Remark 4. Properties (7)–(8) for our homogeneous function play the role of a second-order integration by parts formula (*à la* Bochner).

Remark 5. Had we had the Proposition in the case $p = m = 2$, we could have deduced the Alexandrov–Fenchel inequalities directly without having to use Lemma 1. For this, one needs to find a proper replacement for the condition (ii), which as such seems too weak in the case $p = 2$ (the function $\partial^i f$ is linear). We leave this as question to the interested reader.

Acknowledgement

We thank Ramon van Handel for his comments on an earlier version of this text.

References

- [1] Yu.D. Burago, V.A. Zalgaller, *Geometric Inequalities*, Translated from Russian by A.B. Sosinskiĭ, Springer, 1988.
- [2] M. Gromov, Convex sets and Kähler manifolds, in: *Advances in Differential Geometry and Topology*, World Science Publishers, 1990, pp. 1–38.
- [3] L. Hörmander, *Notions of Convexity*, Birkhäuser, 1994.
- [4] R. Schneider, *Convex Bodies: The Brunn–Minkowski Theory*, second edition, Cambridge University Press, 2014.
- [5] Y. Shenfeld, R. van Handel, *Mixed volume and the Bochner method*, preprint, arXiv:1811.08710.
- [6] R.P. Stanley, Two combinatorial applications of the Aleksandrov–Fenchel inequalities, *J. Comb. Theory* 31 (1) (1981) 56–65.
- [7] X. Wang, A remark on the Alexandrov–Fenchel inequality, *J. Funct. Anal.* 274 (7) (2018) 2061–2088.