



# **B Next-generation sequencing yields mitochondrial genome of *Coccidophilus cariba* Gordon (Coleoptera: Coccinellidae) from museum specimen**

Romain Nattier, Karen Salazar

## **► To cite this version:**

Romain Nattier, Karen Salazar. B Next-generation sequencing yields mitochondrial genome of *Coccidophilus cariba* Gordon (Coleoptera: Coccinellidae) from museum specimen. Mitochondrial DNA Part B Resources, 2019, 4 (2), pp.3780-3781. 10.1080/23802359.2019.1679684 . hal-02363704


**HAL Id: hal-02363704**

**<https://hal.sorbonne-universite.fr/hal-02363704>**

Submitted on 14 Nov 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Next-generation sequencing yields mitochondrial genome of *Coccidophilus cariba* Gordon (Coleoptera: Coccinellidae) from museum specimen

Romain Nattier & Karen Salazar

To cite this article: Romain Nattier & Karen Salazar (2019) Next-generation sequencing yields mitochondrial genome of *Coccidophilus cariba* Gordon (Coleoptera: Coccinellidae) from museum specimen, Mitochondrial DNA Part B, 4:2, 3780-3781, DOI: [10.1080/23802359.2019.1679684](https://doi.org/10.1080/23802359.2019.1679684)

To link to this article: <https://doi.org/10.1080/23802359.2019.1679684>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 25 Oct 2019.



Submit your article to this journal [↗](#)



Article views: 43



View related articles [↗](#)



View Crossmark data [↗](#)

MITOGENOME ANNOUNCEMENT



## Next-generation sequencing yields mitochondrial genome of *Coccidophilus cariba* Gordon (Coleoptera: Coccinellidae) from museum specimen

Romain Nattier<sup>a</sup> and Karen Salazar<sup>a,b</sup> 

<sup>a</sup>Institut de Systématique, Evolution, Biodiversité, ISYEB, Muséum national d'Histoire naturelle (MNHN), CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France; <sup>b</sup>Grupo de Investigación Insectos de Colombia, Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá, Colombia

### ABSTRACT

We assembled and annotated the first complete mitochondrial genome of a species from the subfamily Microweiseinae, *Coccidophilus cariba* Gordon, a predator of scale insect pest. The circular mitogenome consists of 15,343 bp in length, including 13 PCGs, 22 tRNA, and 2 rRNA genes, and exhibits the typical ladybird mitogenome structure. A phylogenetic analysis with the published mitogenomes of other 12 ladybirds is presented, which confirms the position of Microweiseinae as a sister group of Coccinellinae. The *C. cariba* mitogenome could be a useful source to future studies in the relationships inside the subfamily and the genus.

### ARTICLE HISTORY

Received 19 September 2019  
Accepted 25 September 2019

### KEYWORDS



Illumina; ladybird;  
Microweiseinae; mitoge-  
nome; phylogeny

Coccinellidae comprises around 360 genera and 6000 species (Vandenberg 2002) and has been recently divided into the subfamilies Microweiseinae and Coccinellinae (Seago et al. 2011). To date, 28 mitochondrial genomes, complete or partial have been published in Genbank, all belonging to the Coccinellinae. The Microweiseinae *Coccidophilus cariba* Gordon is a scale predator and is the only member of *Coccidophilus* living in the West Indies (Gordon 1978).

The specimen sequenced was collected in Gueloupe in 1997 (Ravine Chaude, GP 1417) and a voucher was deposited in the MNHN (MNHN-EC-7788). DNA was extracted from the whole specimen, following the non-destructive protocol described in Gilbert et al. (2007), and then quantified using a Qubit fluorometer (Life Technologies, Paisley, UK). Genomic DNA was indexed and libraries prepared using the NEBNext Ultra II library prep kit (New England BioLabs; NEB), with a modified version of Meyer and Kircher (2010) protocol. DNA was sequenced at the Genome and Transcriptome Platform of Toulouse (Genotoul, France) using Illumina HiSeq 3000 technology (150 bp paired-end). Quality and length distribution of the sequences were inspected prior and post-clean using FASTQC (Andrews 2010). Low-quality reads and adaptor contamination were then trimmed using the BBDuk plugin as implemented in Geneious prime 2019.1.3 (Biomatters Ltd., Auckland, New Zealand). To separate the mitochondrial genome sequences from the rest of the sequence data, an iterative read mapping strategy was employed using Geneious workflow Align/Assemble 'Map to Reference' and the mitochondrial sequence of *Henosepilachna pusillanima* as reference (Geneious mapper, custom sensitivity with a maximum


mismatch of 30%, fine-tuning 25 times). We repeat this operation with a mismatch of 10% and we conclude with a final de novo step (maximum mismatch of 30%). 7,514 overlapped reads (mean 109.1 bp) were used to generate a consensus sequence and to create a circular molecule. We obtain a mitogenome of 15,343 bp long, with a read coverage of 53×. The identity and position of 13 PCGs, 22 tRNA, and 2 rRNA genes were determined using the MITOS web server (Bernt et al. 2013; <http://mitos.bioinf.uni-leipzig.de>), in combination with the annotation with the mitogenome of *Henosepilachna pusillanima* in Geneious. The gene order is similar to other Coccinellidae mitochondrial genomes. The mitochondrial genome was submitted to GenBank (accession number # MN447521) with the submission tool implemented in Geneious.

The phylogenetic position of *C. cariba* was inferred from 12 mitochondrial genome sequences of Coccinellidae available in Genbank, with *Priasilpha obscura* (Priasilphidae) and *Cucujus clavipes* (Cucujidae) as outgroups. The 13 PCGs and the 2 rRNA was extracted from each genome, aligned separately with the MAFFT algorithm (Katoh and Standley 2013) implemented in MEGA version X (Kumar et al. 2018), then concatenated. Phylogenetic analysis was performed with using Iq-Tree 1.5 (Nguyen et al. 2015) as implemented in the W-Iq-Tree web server (<http://iqtree.cibiv.univie.ac.at>; Trifinopoulos et al. 2016). All PCGs and rRNA alignment were partitioned, the models of substitution were automatically selected, and an ultrafast bootstrap with 1,000 iterations was performed. The resulting topology (Figure 1) confirms the position of Microweiseinae as sister group of Coccinellinae (Seago et al. 2011; Robertson et al. 2015).

**CONTACT** Romain Nattier  [nattier@mnhn.fr](mailto:nattier@mnhn.fr)  Institut de Systématique, Evolution, Biodiversité, ISYEB, Muséum national d'Histoire naturelle (MNHN), CNRS, Sorbonne Université, EPHE, Université des Antilles, Paris, France

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1.** Phylogenetic relationships of Coccinellidae, based on 13 mitochondrial genomes, reconstructed with Iq-Tree 1.5 as implemented on the W-Iq-Tree web server. Ultrafast bootstrap support from 1000 replicates is indicated at the right of each node. For each species is showed the GenBank accession number of the mitochondrial genome.

## Acknowledgments

The authors thank to the 'Service de Systématique Moléculaire, SSM' (UMS 2700 CNRS-MNHN) where the molecular work was realised. We are grateful to Pascaline CHIFFLET-BELLE for technical support in the laboratory phase, the staff of SSM for the support in the molecular data, and Tony ROBILLARD for his advice on improving the manuscript. The financial support in the doctoral studies to KS was provided by the government agency Colciencias-Colombia (program number 756-2016).

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

This work was supported by the ATM blanche du MNHN and Departamento Administrativo de Ciencia, Tecnología e Innovación (COLCIENCIAS).

## ORCID

Karen Salazar  <http://orcid.org/0000-0002-2284-6379>

## References

- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data [accessed 2019 July 2019]. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M, Stadler PF. 2013. MITOS: improved de novo metazoan mitochondrial genome annotation. *Mol Phylogenet Evol.* 69:313–319.

- Gilbert MTP, Moore W, Melchior L, Worobey M. 2007. DNA extraction from dry museum beetles without conferring external morphological damage. *PLoS One.* 2:e272.
- Gordon RD. 1978. West Indian Coccinellidae II (Coleoptera): some scale predators with keys to genera and species. *Coleopt Bull.* 32:205–218.
- Katoh K, Standley DM. 2013. MAFFT Multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30:772–780.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol.* 35:1547–1549.
- Meyer M, Kircher M. 2010. Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb Protoc.* 2010:prot5448.
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol.* 32:268–274.
- Robertson JA, Ślipiński A, Moulton M, Shockley FW, Giorgi A, Lord NP, McKenna DD, Tomaszewska W, Forrester J, Miller KB, et al. 2015. Phylogeny and classification of Cucujoidea and the recognition of a new superfamily Coccinelloidea (Coleoptera: Cucujiformia). *Syst Entomol.* 40:745–778.
- Seago AE, Giorgi JA, Li J, Ślipiński A. 2011. Phylogeny, classification and evolution of ladybird beetles (Coleoptera: Coccinellidae) based on simultaneous analysis of molecular and morphological data. *Mol Phylogenet Evol.* 60:137–151.
- Trifinopoulos J, Nguyen LT, von Haeseler A, Minh BQ. 2016. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 44:W232–235.
- Vandenberg NJ. 2002. Coccinellidae Latreille 1807. Vol. 2. In: Arnett RH Jr., Thomas MC, Skelley PE, Frank JH, editors. *American Beetles. Polyphaga: Scarabaeoidea through Curculionioidea*. Boca Raton (FL): CRC Press; p. 371–389.