Multimodal management for benign skull base meningiomas.

Constantin Tuleasca, Lorenzo Giammattei, Roy Thomas Daniel, Marc Levivier

► To cite this version:

Constantin Tuleasca, Lorenzo Giammattei, Roy Thomas Daniel, Marc Levivier. Multimodal management for benign skull base meningiomas.. Journal of Neurosurgery, In press, 10.3171/2019.6.JNS191682 . hal-02402147

HAL Id: hal-02402147

<https://hal.sorbonne-universite.fr/hal-02402147>

Submitted on 10 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multimodal management for benign skull-base meningiomas

Constantin Tuleasca^{1,2,3,4,5,6}, MD-PhD, Lorenzo Giammattei¹, MD, Roy Thomas Daniel^{1,2}, MBBS, MCh AND Marc Levivier^{1,2}, MD, PhD, IFAANS

¹Department of Clinical Neurosciences, Neurosurgery Service and Gamma Knife Center, Lausanne University Hospital (CHUV), ²University of Lausanne (Unil), Faculty of Biology and Medicine (FBM), ³Signal Processing Laboratory (LTS 5), Ecole Polytechnique Fédérale de Lausanne (EPFL), ⁴Sorbonne Université, Faculté de Médecine, ⁵Assistance Publique-Hôpitaux de Paris, Hôpitaux Universitaires Paris-Sud, Centre Hospitalier Universitaire Bicêtre, Service de Neurochirurgie, Paris, France; ⁶Sorbonne Université, Faculté de Médecine;

of abstract words : N/A

of text words : 500

of references : 10

of tables/figures : 0 figures

of videos : 0

Running title: Combined management for skull-base benign lesions

Key words: skull-base, management, meningioma, schwannoma

Corresponding author:

Constantin Tuleasca, MD-PhD, Assistance Publique-Hôpitaux de Paris, Hôpitaux Universitaires Paris Sud, Centre Hospitalier Universitaire de Bicêtre, Service de Neurochirurgie, Le Kremlin-Bicêtre, France- 78, Rue Général Leclerc, 94270, Le Kremlin-Bicêtre, France; Sorbonne Université, Faculté de Médecine, Paris, France; Tel: +41-21-314-26-02; Fax: +41-21-314-11-99; e-mail: constantin.tuleasca@gmail.com;

Financial Disclosures: None

Dear Editor,

We read with great interest the article by Kim et al.², discussing multimodal management of petroclival meningiomas. Current authors' preference is for multimodal management, with planned subtotal resection followed by adjuvant SRS.

During the past ten years, our unit has developed an optimally invasive skull-base surgery approach for large benign tumors. The philosophy of this approach is planned subtotal resection followed by SRS using Gamma Knife (GK), with the aim to reduce morbidity and mortality. Gamma Knife is usually performed between 3 to 6 months after microsurgical resection, without a "wait-and-scan" follow-up period of the remnant tumor. In relationship with this recently published paper by Kim and co-authors, we also apply this approach to other skull-base meningiomas, such as the clinoid and cavernous sinus ones.

In the particular case of petroclival meningiomas radical removal remains challenging, primarily due to brainstem adherence or invasion, and encasement of basilar artery/perforators, cranial nerves V, VII, VIII, IX, X, XI⁵. In these cases, we prefer to use the supracerebellar infratentorial (extended retrosigmoid) approach. The extent of tumor removal frequently depends on extent of clival component, invasion of cavernous sinus, adherence to brainstem pia and basilar artery or its branches. What is deemed risky to remove is left behind for subsequent GK-SRS. In our series of combined approaches for benign skull-base lesions, we have not had any mortality or major neurological morbidity. Single fraction GKS can be safely applied on residual tumors¹, even in cases with brainstem contact, due to low prescription marginal doses of 12 Gy, which allow for high rates of tumor control⁶. In cases where limiting factor is optic apparatus, due to GK steep gradient, it is possible to keep the dose below 8 Gy^{3,4}.

In the particular case of clinoid and cavernous sinus meningiomas, surgical challenge is to preserve carotid artery and branches, recover visual function and limit oculomotor cranial nerve palsies. We don't routinely remove intracavernous sinus part for these tumors. We also perform chiasmopexy, by interposing small fat graft between optic nerve, chiasm, and residual intracavernous sinus tumor component, allowing safe GK on remnant. The same strategy is applied to pituitary stalk whenever needed, to minimize risk of pituitary insufficiency.

In conclusion, we are completely in agreement and we congratulate Kim and co-authors for a very nice paper, which will contribute, in the future, to major paradigm shift in the treatment of these difficult skull-base tumors.

Funding: Constantin Tuleasca gratefully acknowledges receipt of a ‘Young Researcher in Clinical Research Grant’ (Jeune Chercheur en Recherche Clinique) from the University of Lausanne (UNIL), Faculty of Biology and Medicine (FBM) and the Lausanne University Hospital (CHUV)

Acknowledgments: Lausanne University Hospital and University of Lausanne.

Conflict of interest: No conflict of interest.

References:

1. Faramand A, Kano H, Niranjana A, Johnson SA, Hassib M, Park KJ, et al: Cranial nerve outcomes after primary stereotactic radiosurgery for symptomatic skull base meningiomas. **J Neurooncol** **139**:341-348, 2018
2. Kim JW, Jung HW, Kim YH, Park CK, Chung HT, Paek SH, et al: Petroclival meningiomas: long-term outcomes of multimodal treatments and management strategies based on 30 years of experience at a single institution. **J Neurosurg**:1-8, 2019
3. Leavitt JA, Stafford SL, Link MJ, Pollock BE: Long-term evaluation of radiation-induced optic neuropathy after single-fraction stereotactic radiosurgery. **Int J Radiat Oncol Biol Phys** **87**:524-527, 2013
4. Pollock BE, Link MJ, Leavitt JA, Stafford SL: Dose-volume analysis of radiation-induced optic neuropathy after single-fraction stereotactic radiosurgery. **Neurosurgery** **75**:456-460; discussion 460, 2014
5. Seifert V: Clinical management of petroclival meningiomas and the eternal quest for preservation of quality of life: personal experiences over a period of 20 years. **Acta Neurochir (Wien)** **152**:1099-1116, 2010
6. Tuleasca C, Leroy HA, Regis J, Levivier M: Gamma Knife radiosurgery for cervical spine lesions: expanding the indications in the new era of Icon. **Acta Neurochir (Wien)** **158**:2235-2236, 2016