



HAL
open science

Mutation Update of the CLCN5 Gene Responsible for Dent Disease 1

Lamisse Mansour-Hendili, Anne Blanchard, Nelly Le Pottier, Isabelle Roncelin, Stéphane Lourdel, Cyrielle Treard, Wendy González, Ariela Vergara-Jaque, Gilles Morin, Estelle Colin, et al.

► **To cite this version:**

Lamisse Mansour-Hendili, Anne Blanchard, Nelly Le Pottier, Isabelle Roncelin, Stéphane Lourdel, et al.. Mutation Update of the CLCN5 Gene Responsible for Dent Disease 1. *Human Mutation*, 2015, 36 (8), pp.743-752. 10.1002/humu.22804 . hal-02453168

HAL Id: hal-02453168

<https://hal.sorbonne-universite.fr/hal-02453168>

Submitted on 24 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Mutation Update of the CLCN5 gene responsible for Dent disease 1

Journal:	<i>Human Mutation</i>
Manuscript ID:	humu-2015-0056.R1
Wiley - Manuscript type:	Mutation Update
Date Submitted by the Author:	25-Mar-2015
Complete List of Authors:	<p>MANSOUR-HENDILI, Lamisse; APHP-HEGP, Genetic; Université Paris Descartes, Blanchard, Anne; Université Paris Descartes, ; INSERM, UMR970, ; Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, centre d'investigation clinique; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA), Le Pottier, Nelly; HEGP, Dpt de Genetique Roncelin, Isabelle; HEGP, Dpt de Genetique Lourdel, Stéphane; UPMC, UMRS_872 Treard, Cyrielle; HEGP, Dpt de Genetique; INSERM, UMR970, González, Wendy; Universidad de Talca, Centro de Bioinformática y Simulación Molecular Vergara-Jaque, Ariela; Universidad de Talca, Centro de Bioinformática y Simulación Molecular Morin, Gilles; CHU Amiens Picardie, Genetics Colin, Estelle; Service de Génétique, Hôpital d'Angers, Holder-Espinasse, Muriel; Hôpital Jeanne de Flandre, CHRU, Service de Génétique Clinique; Department of Clinical Genetics, Guy's Hospital, Bacchetta, Justine; Centre de Référence des Maladies Rénales Rares, Service de Néphrologie et Rhumatologie Pédiatriques, Hôpital Femme Mère Enfant, Baudouin, Veronique; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA), ; Service de Néphrologie, Assistance Publique-Hôpitaux de Paris, Hôpital Robert Debré, Benoit, Stephane; Centre Hospitalier Universitaire de Tours, Service de Néphrologie,, Berard, Etienne; Centre Hospitalier Universitaire de Nice, Service de Néphrologie pédiatrique, Bourdat-Michel, Guylhene; Centre Hospitalier Universitaire de Grenoble. Département de Pédiatrie, Bouchireb, Karim; Assistance Publique Hôpitaux de Paris, Hôpital Necker- Enfants-malades, Service de Néphrologie Pédiatrique, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA), Burtey, Stephane; Aix-Marseille université, VRCM, centre de néphrologie et transplantation rénale, Cailliez, Mathilde; Assistance Publique Hôpitaux de Marseille, Hôpital La Timone, Unité de Néphrologie Pédiatrique,</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>Cardon, Gerard; Centre Hospitalier de Douai, Service de Néphrologie, Cartery, Claire; Assistance Publique-Hôpitaux de Paris, Hôpital Tenon, Service de Néphrologie et dialyse,</p> <p>Champion, Gerard; Centre Hospitalier Universitaire d'Angers, Département de Pédiatrie, LUNAM,</p> <p>Chauveau, Dominique; Centre Hospitalier Universitaire de Toulouse, Hôpital Rangueil, Département de Néphrologie et Transplantation d'organes,</p> <p>Cochat, Pierre; Centre de Référence des Maladies Rénales Rares Service de Néphrologie Rhumatologie Dermatologie Pédiatriques Hospices Civils de Lyon,</p> <p>Dahan, Karin; Cliniques Universitaires St Luc, Centre de Génétique Humaine</p> <p>De La Faille, Renaud; Centre Hospitalier Universitaire de Bordeaux, Service de Néphrologie Transplantation Dialyse,</p> <p>Debray, François-Guillaume; Centre Hospitalier Universitaire Sart-Tilman. Département de Génétique,</p> <p>Dehoux, Laurene; Service de Néphrologie, Assistance Publique-Hôpitaux de Paris, Hôpital Robert Debré, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA),</p> <p>Deschênes, Georges; Service de Néphrologie, Assistance Publique-Hôpitaux de Paris, Hôpital Robert Debré, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA),</p> <p>Desport, Estelle; Centre Hospitalier Universitaire de Poitiers, Service de Néphrologie,</p> <p>Devuyst, Olivier; Cliniques Universitaires Saint-Luc, Université Catholique de Louvain, ; Institute of Physiology, University of Zurich,</p> <p>Dieguez, Stella; Hospital General de Agudos Dr. Teodoro Álvarez, Nefrologia Infantil,</p> <p>Emma, Francesco; Ospedale Bambino Gesù, IRCCS, Department of Nephrology and Urology</p> <p>Fischbach, Michel; Centre Hospitalier Universitaire Hautepierre, Service de Pédiatrie,</p> <p>Fouque, Denis; Centre Hospitalier Universitaire Lyon Sud, Departement de Néphrologie,</p> <p>Fourcade, Jacques; Centre Hospitalier de Chambéry, Service de Néphrology,</p> <p>François, Helene; Assistance Publique-Hôpitaux de Paris, Hôpital Kremlin Bicêtre, Service de Néphrologie,</p> <p>Gilbert-Dussardier, Brigitte; Centre Hospitalier Universitaire de Poitiers ; EA 3808 Université de Poitiers, Service de Génétique</p> <p>Hannedouche, Thierry; Centre Hospitalier Universitaire de Strasbourg, Hôpital Civil, Service de Néphrologie, ; Faculté de Médecine, Université de Strasbourg,</p> <p>Houillier, Pascal; Assistance Publique Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Département de Physiologie, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA), ; INSERM, Université Paris Descartes, Sorbonne Paris Cité, UMR S1138, Centre de Recherche des Cordeliers, CNRS ERL 8228, F-75006,</p> <p>Izzedine, Hassan; Assistance Publique-Hôpitaux de Paris. Hôpital Pitié Salpêtrière. Service de Néphrologie,</p> <p>Janner, Marco; University of Berne Children's Hospital, Department of Paediatric Endocrinology, Diabetology and Metabolism,</p> <p>Karras, Alexandre; Assistance Publique Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Service de Néphrologie,</p> <p>Knebelmann, Bertrand; Assistance Publique Hôpitaux de Paris, Hôpital Necker-Enfants-malades, Service de Néphrologie adulte, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA),</p> <p>Lavocat, Marie Pierre; CHU Saint Etienne, Hopital Nord, Department of Paediatrics</p>
--	---

	<p>Lemoine, Sandrine; hôpital Edouard Herriot, Service d'exploration fonctionnelle rénale</p> <p>Leroy, Valerie; Centre Hospitalier Universitaire de Lille, Hôpital Jeanne de Flandre, Service de Néphrologie Pédiatrique,</p> <p>Loirat, Chantal; Service de Néphrologie, Assistance Publique-Hôpitaux de Paris, Hôpital Robert Debré, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA),</p> <p>Macher, Marie-Alice; Service de Néphrologie, Assistance Publique-Hôpitaux de Paris, Hôpital Robert Debré, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA),</p> <p>Martin-Coignard, dominique; CH le Mans, Service de Génétique Médicale</p> <p>Morin, Denis; Centre Hospitalier Universitaire de Montpellier, Unité de Néphrologie Pédiatrique</p> <p>Niaudet, Patrick; Assistance Publique Hôpitaux de Paris, Hôpital Necker-Enfants-malades, Service de Néphrologie Pédiatrique; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA),</p> <p>Nivet, Hubert; Centre Hospitalier Universitaire de Tours, Service de Néphrologie,</p> <p>Nobili, Francois; CHU Besancon, Nephrologie pediatrique</p> <p>Novo, Robert; Centre Hospitalier Universitaire de Lille, Hôpital Jeanne de Flandre, Service de Néphrologie Pédiatrique</p> <p>Faivre, Laurence; Dijon Hospital, medical Genetics;</p> <p>Rigothier, Claire; Centre Hospitalier Universitaire de Bordeaux, Service de Néphrologie Transplantation Dialyse,</p> <p>Rousseau-Kesler, Gwenaelle; Centre Hospitalier Universitaire de Nantes, Unité de Néphrologie Pédiatrie,</p> <p>Salomon, Remi; AP-HP, Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA), Service de Néphrologie Pédiatrique, Hôpital Necker-Enfants Malades</p> <p>Schleich, Andrea; Institute of Nephrology Stadtpital Waid Zuerich,</p> <p>Sellier-Leclerc, Anne-Laure; Centre de Référence des Maladies Rénales Rares, Service de Néphrologie et Rhumatologie Pédiatriques, Hôpital Femme Mère Enfant,</p> <p>Soulami, Kenza; CHU Ibn Rochd, Service de Néphrologie Dialyse Transplantation</p> <p>Tiple, Aurelien; Centre Hospitalier Universitaire Gabriel-Montpied Service de Néphrologie,</p> <p>Tim, Ulinski; Université Pierre et Marie Curie, AP-HP, Hôpital Armand Trousseau, Département de Néphrologie Pédiatrique</p> <p>Van Regemorter, Nicole; Université Libre de Bruxelles, Hôpital Erasme Département de Génétique Médicale,</p> <p>Vanhille, Philippe; Centre hospitalier de Valenciennes, Service de néphrologie et de médecine interne</p> <p>Jeunemaitre, Xavier; Université Paris Descartes, Faculté de Médecine, ; Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Service de Génétique, ; INSERM, UMR970, Paris-Cardiovascular Research Center, ; Centre de Référence des Maladies Rénales Héritaires de l'Enfant et de l'Adulte (MARHEA),</p> <p>Vargas Poussou, Rosa; HEGP, Dpt de Genetique</p>
Key Words:	CLCN5, Dent disease 1, CIC-5, low molecular weight proteinuria , renal failure

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Mutation Update of the *CLCN5* gene responsible for **Dent disease 1**

Lamisse Mansour-Hendili,^{1,2} Anne Blanchard,^{1,3,4,5} Nelly Le Pottier,² Isabelle Roncelin,² Stéphane Lourdel,^{6,7} Cyrielle Tread,^{2,3} Wendy González,⁸ Ariela Vergara-Jaque,⁸ Gilles Morin,⁹ Estelle Colin,¹⁰ Muriel Holder-Espinasse,^{11,12} Justine Bacchetta,¹³ Véronique Baudouin,^{5,14} Stéphane Benoit,⁵ Etienne Bérard,¹⁶ Guylhène Bourdat-Michel,¹⁷ Karim Bouchireb,^{5,18} Stéphane Burtey,¹⁹ Mathilde Cailliez,²⁰ Gérard Cardon,²¹ Claire Cartery,²² Gerard Champion,²³ Dominique Chauveau,²⁴ Pierre Cochat,¹³ Karin Dahan,²⁵ Renaud De la Faille,²⁶ François-Guillaume Debray,²⁷ Laurene Dehoux,^{5,14} Georges Deschenes,^{5,14} Estelle Desport,²⁸ Olivier Devuyt,^{29,30} Stella Dieguez,³¹ Francesco Emma,³² Michel Fischbach,³³ Denis Fouque,³⁴ Jacques Fourcade,³⁵ Hélène François,³⁶ Brigitte Gilbert-Dussardier,³⁷ Thierry Hannedouche,³⁸ Pascal Houillier,^{1,5,7,39} Hassan Izzedine,⁴⁰ Marco Janner,⁴¹ Alexandre Karras,⁴² Bertrand Knebelmann,^{5,43} Marie-Pierre Lavocat,⁴⁴ Sandrine Lemoine,⁴⁵ Valérie Leroy,⁴⁶ Chantal Loirat,^{5,14} Marie-Alice Macher,^{5,14} Dominique Martin-Coignard,⁴⁷ Denis Morin,⁴⁸ Patrick Niaudet,^{5,18} Hubert Nivet,¹⁵ François Nobili,⁴⁹ Robert Novo,⁴⁶ Laurence Faivre,⁵⁰ Claire Rigother,²⁶ Gwenaëlle Roussey-Kesler,⁵¹ Remi Salomon,^{1,5,18} Andreas Schleich,⁵² Anne-Laure Sellier-Leclerc,¹³ Kenza Soulami,⁵³ Aurélien Tiple,⁵⁴ Tim Ulinski,^{5,55} Philippe Vanhille,⁵⁶ Nicole Van Regemorter,⁵⁷ Xavier Jeunemaître^{1,2,3,5} and Rosa Vargas-Poussou^{2,3,5}

¹Université Paris Descartes, Faculté de Médecine, Paris, France. ²Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Service de Génétique, Paris, France. ³INSERM, UMR970, Paris-Cardiovascular Research Center, Paris, France. ⁴Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, centre d'investigation clinique, Paris, France. ⁵Centre de Référence des Maladies Rénales

1
2
3 Héritaires de l'Enfant et de l'Adulte (MARHEA), Paris, France. ⁶ Sorbonne Universités,
4
5 UPMC Université Paris 06, ⁷INSERM, Université Paris Descartes, Sorbonne Paris Cité, UMR
6
7 S1138, Centre de Recherche des Cordeliers, CNRS ERL 8228, F-75006, Paris, France;
8
9
10 ⁸Centro de Bioinformática y Simulación Molecular, Universidad de Talca, Talca, Chile ;
11
12 ⁹Centre Hospitalier Universitaire Amiens Picardie, Service de Génétique et Oncogénétique,
13
14 Amiens France. ¹⁰Centre Hospitalier Universitaire d'Angers, Département de Biochimie et
15
16 Génétique, LUNAM Angers, France. ¹¹Centre Hospitalier Universitaire de Lille, Département
17
18 de Génétique, Lille, France. ¹²Department of Clinical Genetics, Guy's Hospital, London, UK.
19
20 ¹³Centre de Référence des Maladies Rénales Rares. Service de Néphrologie Rhumatologie
21
22 Dermatologie Pédiatriques, Hospices Civils de Lyon, Lyon, France. ¹⁴Service de Néphrologie,
23
24 Assistance Publique-Hôpitaux de Paris, Hôpital Robert Debré, Paris, France. ¹⁵Centre
25
26 Hospitalier Universitaire de Tours, Service de Néphrologie, Tours, France. ¹⁶Centre
27
28 Hospitalier Universitaire de Nice, Service de Néphrologie pédiatrique, Nice, France. ¹⁷Centre
29
30 Hospitalier Universitaire de Grenoble. Département de Pédiatrie, Grenoble France.
31
32
33
34
35
36 ¹⁸Assistance Publique Hôpitaux de Paris, Hôpital Necker-Enfants-malades, Service de
37
38 Néphrologie Pédiatrique, Paris, France. ¹⁹Aix-Marseille Université, VRCM, centre de
39
40 néphrologie et transplantation rénale, Marseille, France ²⁰Assistance Publique Hôpitaux de
41
42 Marseille, Hôpital La Timone, Unité de Néphrologie Pédiatrique, Marseille, France. ²¹Centre
43
44 Hospitalier de Douai, Service de Néphrologie, Douai, France. ²²Assistance Publique-Hôpitaux
45
46 de Paris, Hôpital Tenon, Service de Néphrologie et dialyse, Paris, France. ²³Centre Hospitalier
47
48 Universitaire d'Angers, Département de Pédiatrie, LUNAM Angers, France. ²⁴Centre
49
50 Hospitalier Universitaire de Toulouse, Hôpital Rangueil, Département de Néphrologie et
51
52 Transplantation d'organes, Toulouse, France. ²⁵Institut de Pathologie et de Génétique,
53
54 Département de Génétique Humaine, Gosselies, Belgique. ²⁶Centre Hospitalier Universitaire
55
56 de Bordeaux, Service de Néphrologie Transplantation Dialyse, Bordeaux, France. ²⁷Centre
57
58
59
60

1
2
3 Hospitalier Universitaire Sart-Tilman. Département de Génétique, Liège Belgique. ²⁸Centre
4
5 Hospitalier Universitaire de Poitiers, Service de Néphrologie, Poitiers, France. ²⁹Cliniques
6
7 Universitaires Saint-Luc, Université Catholique de Louvain, Brussels, Belgium. ³⁰Institute of
8
9 Physiology, University of Zurich, Zurich, Switzerland. ³¹Hospital General de Agudos Dr.
10
11 Teodoro Álvarez, Nefrologia Infantil, Buenos Aires, Argentina. ³²Bambino Gesù Children's
12
13 Hospital – IRCCS, Division of Nephrology and Dialysis, Rome, Italy. ³³Centre Hospitalier
14
15 Universitaire Hautepierre, Service de Pédiatrie, Strasbourg, France. ³⁴Centre Hospitalier
16
17 Universitaire Lyon Sud, Département de Néphrologie, Lyon, France. ³⁵Centre Hospitalier de
18
19 Chambéry, Service de Néphrologie, Chambéry, France. ³⁶Assistance Publique-Hôpitaux de
20
21 Paris, Hôpital Kremlin Bicêtre, Service de Néphrologie, Le Kremlin-Bicêtre, France. ³⁷Centre
22
23 Hospitalier Universitaire de Poitiers, Service de Génétique, EA 3808, Université de Poitiers,
24
25 Poitiers, France. ³⁸Hôpitaux Universitaires de Strasbourg, Service de Néphrologie et Faculté
26
27 de Médecine, Université de Strasbourg, Strasbourg, France. ³⁹Assistance Publique Hôpitaux
28
29 de Paris, Hôpital Européen Georges Pompidou, Département de Physiologie, Paris, France.
30
31 ⁴⁰Assistance Publique-Hôpitaux de Paris. Hôpital Pitié Salpêtrière. Service de Néphrologie,
32
33 Paris, France. ⁴¹University of Berne Children's Hospital, Department of Paediatric
34
35 Endocrinology, Diabetology and Metabolism, Berne, Switzerland. ⁴²Assistance Publique
36
37 Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Service de Néphrologie, Paris,
38
39 France. ⁴³Assistance Publique Hôpitaux de Paris, Hôpital Necker-Enfants-malades, Service de
40
41 Néphrologie adulte, Paris, France. ⁴⁴Centre Hospitalier Universitaire de Saint Etienne, Hôpital
42
43 Nord, Département de Pédiatrie, Saint Etienne, France. ⁴⁵Hospices Civils de Lyon, Hôpital
44
45 Edouard-Herriot, Service d'Exploration Fonctionnelle Rénale, Lyon, France. ⁴⁶Centre
46
47 Hospitalier Universitaire de Lille, Hôpital Jeanne de Flandre, Service de Néphrologie
48
49 Pédiatrique, Lille, France. ⁴⁷Centre Hospitalier du Mans, Services de Génétique, Le Mans,
50
51 France. ⁴⁸Centre Hospitalier Universitaire de Montpellier, Unité de Néphrologie Pédiatrique,
52
53
54
55
56
57
58
59
60

1
2
3 Montpellier, France. ⁴⁹Centre Hospitalier Universitaire de Besançon, Unité de Néphrologie
4
5 Pédiatrie, Besançon, France. ⁵⁰Centre Hospitalier Universitaire de Dijon, Centre de
6
7 Génétique, Dijon, France. ⁵¹Centre Hospitalier Universitaire de Nantes, Unité de Néphrologie
8
9 Pédiatrie, Nantes, France. ⁵²Institute of Nephrology Statspital Waid Zuerich, Zuerich,
10
11 Switzerland. ⁵³Néphrologue Pédiatre, Casablanca, Maroc. ⁵⁴Centre Hospitalier Universitaire
12
13 Gabriel-Montpied Service de Néphrologie, Clermont-Ferrand, France. ⁵⁵Assistance Publique–
14
15 Hôpitaux de Paris, Hôpital Trousseau, Service de Néphrologie et Transplantation Rénale,
16
17 Paris, France. ⁵⁶Centre Hospitalier de Valenciennes, Service de Néphrologie et Médecine
18
19 Interne, Valenciennes, France. ⁵⁷Université Libre de Bruxelles, Hôpital Erasme Département
20
21 de Génétique Médicale, Brussels, Belgique
22
23
24
25
26
27
28

29 Contract grant sponsors: French Ministry of Health (Plan Maladies Rares), European
30
31 Community FP7EUNEFRON 201590 and EUReOmics 2012-305608. The Chilean PIA-
32
33 Conycit program: ACT-1104 grant. The Swiss National Science Foundation project grant
34
35 310030_146490; and RADIZ, Rare Diseases Initiative Zürich, a KFSP of the University of
36
37 Zurich.
38
39

40
41 Key words: *CLCN5*, *CIC-5*, Dent disease 1, low molecular weight proteinuria, renal failure
42
43
44
45

46 Corresponding author:

47 Rosa Vargas Poussou

48 Hôpital Européen George Pompidou

49 20-40 rue Leblanc

50 75015 Paris, France

51
52
53
54
55
56
57
58
59
60
rosa.vargas@egp.aphp.fr

Abstract

Dent disease is a rare X-linked tubulopathy characterised by low molecular weight proteinuria (LMWP), hypercalciuria, nephrocalcinosis and/or nephrolithiasis, progressive renal failure and variable manifestations of other proximal tubule dysfunctions. It often progresses over a few decades to chronic renal insufficiency, and therefore molecular characterization is important to allow appropriate genetic counselling. Two genetic subtypes have been described to date: **Dent disease 1** is caused by mutations of the *CLCN5* gene, coding for the chloride/proton exchanger CIC-5; and **Dent disease 2** by mutations of the *OCRL* gene, coding for the **inositol polyphosphate 5-phosphatase OCRL-1**.

Herein, we review previously reported mutations (n= 192) and their associated phenotype in 377 male patients with **Dent disease 1** and describe phenotype and novel (n=42) and recurrent mutations (n=24) in a large cohort of 117 **Dent disease 1** patients belonging to 90 families.

The novel missense and in-frame mutations described were mapped onto a three-dimensional homology model of the CIC-5 protein. This analysis suggests that these mutations affect the dimerization process, helix stability or transport. The phenotype of our cohort patients supports and extends the phenotype that has been reported in smaller studies.

Background

Dent disease (OMIM 300009) is a rare X-linked renal proximal tubulopathy clinically defined by low molecular weight proteinuria (LMWP) associated with hypercalciuria and/or its complications (nephrocalcinosis or nephrolithiasis) and progressive renal failure. LMWP is present in all affected males and in almost all obligate female carriers. **Dent disease** may also be associated with defective reabsorption of one or several of the following solutes: amino acids, glucose, phosphate, uric acid, potassium and bicarbonate; some patients may also have rickets. **Dent disease** may also present as a generalized proximal tubular dysfunction (i.e.

1
2
3 renal Fanconi syndrome). The main complication of this disease is progression to chronic
4 kidney disease during the second to the fourth decade of life in 30 to 80% of cases [Devuyst
5 et al. 2010].
6
7
8
9

10
11 **Dent disease** is genetically heterogeneous: about 50-60% of patients harbour inactivating
12 mutations of the *CLCN5* gene (**Dent disease 1**) and about 15% harbour inactivating mutations
13 in the *OCRL* gene (**Dent disease 2**). Other yet unidentified causative genes are likely involved
14 in the disease [Devuyst et al. 2010]
15
16
17
18
19

20
21 *CLCN5* (OMIM #300008) maps on chromosome Xp11.22 and encodes a 746 amino-acid
22 electrogenic $2\text{Cl}^-/\text{H}^+$ **antiporter** (ClC-5) [Lloyd et al., 1996; Scheel et al., 2005]. The *OCRL*
23 gene maps on chromosome Xq25 and encodes the **inositol polyphosphate 5-phosphatase**
24 **OCRL-1** [Attree et al., 1992].
25
26
27
28
29
30

31
32 In the human kidney, ClC-5 is expressed in the proximal tubule, the thick ascending limb and
33 the intercalated cells of the collecting duct. ClC-5 protein is a dimer with two identical
34 subunits, each of which contains a pore responsible for the selective coupling of the Cl^- flux to
35 H^+ counter-transport [Dutzler et al., 2005]. Each subunit is constituted by 18 α -helices in anti-
36 parallel orientation, with two phosphorylation sites and one N-glycosylation site. The carboxy
37 terminus of ClC-5 contains two CBS (cystathionine B-synthase) domains, a PY sorting signal
38 and a potential PDZ protein-binding module [Schwake et al., 2001; Hryciw et al., 2006].
39
40
41
42
43
44
45
46
47
48

49
50 An important number of mutations of the *CLCN5* gene affecting sites throughout the protein
51 have been reported since the identification of the gene in 1996 but only a limited number is
52 described in the databases [HGMD]. *CLCN5* mutations have been grouped into three classes
53 on the basis of functional data: mutations impairing processing and folding inducing retention
54 in the endoplasmic reticulum and degradation of the mutant protein within the cell (class 1);
55 mutations causing a delay in protein processing and reducing stability of ClC-5 (class 2); and
56
57
58
59
60

1
2
3 mutations affecting electrical activity but not trafficking of CLC-5 to the cell surface and early
4
5 endosomes (class 3) [Lourdel et al., 2012]. No correlation between genotype and phenotype
6
7 has been established.
8
9

10
11 In this study, we review the published mutations in the *CLCN5* gene (**Dent disease 1**) as well
12
13 as their phenotype and report molecular characterisation and clinical data of a cohort of 90
14
15 families analysed in our centre. The analysis of this large cohort includes the description of 42
16
17 previously non-described mutations and the proposed pathogenic mechanism for novel
18
19 missense mutations.
20
21
22
23
24
25
26

27 **Population**

28
29 The study included 90 unrelated patients diagnosed with **Dent disease** addressed for genetic
30
31 analysis to the Genetics Department of the Georges Pompidou European Hospital (Paris,
32
33 France) from January 2001 to December 2014. The study was approved by the "Comité de
34
35 Protection des Personnes, Paris-Île de France XI (Ref. 09069)" and informed consent for
36
37 genetic study was obtained from each proband or, if minor, from their parents. Genetic
38
39 investigations were usually performed because of the presence of at least two major criteria
40
41 (LMWP, hypercalciuria, renal failure, familial history of **Dent disease**) and one minor
42
43 criterion (mainly one or more defect of proximal reabsorption). In children, the presence of
44
45 only two major criteria (in most of the cases LMWP and hypercalciuria) or even persistent
46
47 LMWP of unknown aetiology is sufficient to justify molecular genetic diagnosis. Genetic
48
49 investigations were extended to relatives in 43 families.
50
51
52
53
54
55
56
57

58 **Methods of Mutation Detection and *in Silico* Prediction**

59
60

1
2
3 DNA was extracted by a saline method or with blood DNA midi kits (Qiagen columns).
4
5 *CLCN5* exons and the flanking intronic sequences were amplified by PCR and then sequenced
6
7 using BigDyeTerminator kit v3.1 cycle sequencing kits and run on an ABI Prism 3730XL
8
9 DNA Analyzer Sequencer (Perkin Elmer Applied Biosystems, Foster City, CA) as previously
10
11 described [Grand et al., 2009].
12
13

14
15
16 DNA mutations were identified using Sequencher software by comparison to the *CLCN5* gene
17
18 reference sequence: NM_000084.2. Each mutation was confirmed by sequencing a second
19
20 independent PCR product.
21
22

23
24 Missense and splicing mutations were interpreted with Alamut V.2.2 software (Interactive
25
26 Biosoftware, Rouen, France; <http://www.interactivebiosoftware.com>). Complementary
27
28 analyses were performed with SIFT (<http://www.Blocks.fhrc.org/sift/SIFT.html>), PolyPhen-
29
30 2 (<http://genetics.bwh.harvard.edu/pph/>), Mutpred (<http://mutpred.mutdb.org/about.html>),
31
32 SNPs&Go (<http://snps-and-go.biocomp.unibo.it/snps-and-go/info.htm>) and mutation taster
33
34 (<http://www.mutationtaster.org/documentation.html>).
35
36
37

38
39 VMD software (Humphrey W. et al, J Mol Graph 1996) was used to predict the dimeric
40
41 structure of the ClC-5; this structure was then optimized by means of energy minimization
42
43 (12,000 steps) using NAMD2.9 software [Phillips et al., 2005] and the CHARMM27 force
44
45 field [Mackerell et al., 2004]. The figures were produced with VMD software. The structure
46
47 of the monomer in the model was built with ICM software [Abagyan et al., 1994], employing
48
49 the crystallographic data of the StClC channel (PDB ID code 1KPL) as the template.
50
51
52
53

54 55 56 57 **Mutations**

58 59 60 Described Mutations

1
2
3 We searched for *CLCN5* mutations published between February 1996 and October 2014 in
4 PubMed, using the combination of the key words “*CLCN5*”, “Dent disease” and “Chloride
5 channel ClC-5”. We found 192 different described mutations in the *CLCN5* gene detected in
6
7
8
9
10 377 patients belonging to 334 families. These mutations are described in **Supp. Table S1**,
11
12 following HGVS nomenclature recommendations and the reference sequence NM_000084.2.
13
14 To date, functional studies have been performed for 41 mutations (31 missense, 5 nonsense, 4
15
16 frameshift or and 1 in-frame), the functional class and references are also quoted in **Supp.**
17
18 **Table S1**. All these mutations have been entered in the Leiden Open Variation Database
19
20 (<http://www.LOVD.nl/CLCN5>).
21
22
23
24

25 Described mutations include large deletions (4%), nonsense and frameshift mutations that
26
27 produce premature stop codons (17 and 28% respectively), splice site mutations that interfere
28
29 or are predicted to interfere with correct splicing (11%), missense and small in-frame
30
31 deletions affecting conserved amino acid residues (36.5 % and 2.6 % respectively).
32
33
34

35 Novel and Recurrent Mutations in our population

36
37
38
39 DNA sequence analysis of the entire coding region of the *CLCN5* gene from one affected
40
41 member of each of the 90 families revealed 66 different mutations; 42 of the mutations have
42
43 not previously been described (**Table 1 and 2** summarize the novel and recurrent mutations
44
45 detected). Eleven families of this cohort were previously described [Blanchard et al., 2008,
46
47 Grand et al., 2009]
48
49
50

51 Prediction of pathogenicity of novel Mutations (Biological relevance)

52
53
54
55 The novel mutations include: 14 frameshift, eight splicing, eight nonsense, eight missense,
56
57 one in-frame mutation and three large deletions. Most of these mutations (25 frameshift
58
59 nonsense and deletions) are predicted to result in the production of unstable mRNAs,
60
truncated or absent proteins. Six out of the eight splice site mutations disrupt the obligatory

1
2
3 consensus donor or acceptor splice sites and are considered pathogenic. For one mutation
4
5 located in -8 in the acceptor site of exon 6, MaxEntScan predicts a decreasing of splice site
6
7 score of 42% and SpliceSiteFinder-like a loss of a splice site and the creation of a cryptic
8
9 splice site. Finally, the affected nucleotide responsible for synonym p.Glu35Glu change is the
10
11 last nucleotide of the exon 2 and *in silico* analysis show a significant decreasing of site scores
12
13 by 36% and 13.2 % for MaxEntScan and SpliceSiteFinder-like respectively. Unfortunately no
14
15 mRNA was available for analysis of the transcripts for these two patients.
16
17
18

19
20 The seven missense and one in-frame mutations we report here for the first time affect highly
21
22 conserved amino acids and all are predicted to be potentially pathogenic by *in silico* tools
23
24 (Supp. Table S2). The novel missense and in-frame mutations were mapped onto a three-
25
26 dimensional homology model of the CIC-5 protein (Figure 1): in-frame
27
28 (p.Pro258delinsArgAsn) and two missense (p.Thr518Ala and p.Gly530Ser) mutations affect
29
30 helices involved in the formation of the dimer interface; four missense mutations
31
32 (p.Lys231Ieu, p.Arg239Pro, p.Gly250Arg and p.Gly470Arg) are near the transporter
33
34 interface; and the p.Glu267Asp missense mutation is located just before the “proton
35
36 glutamate” (Glu268), critical for the transport function of CIC 2Cl⁻/H⁺ exchangers.
37
38
39
40
41

42
43 Previous functional investigations in heterologous expression systems demonstrated that
44
45 *CLCN5* mutations clustering at the dimer interface can result in the retention of CIC-5 in the
46
47 endoplasmic reticulum and its rapid degradation within the cell [Lourdel et al., 2012]. CIC-5
48
49 2Cl⁻/H⁺ exchangers are found as homodimers, leading to the suggestion that the mutant
50
51 proteins are subjected to early degradation because of the impaired dimerization. This is
52
53 plausible for the in-frame (p.Pro258delinsArgAsn) and two missense (p.Thr518Ala and
54
55 p.Gly530Ser) mutations mapping to helices H and P, that are involved in the formation of the
56
57 dimer interface of CIC-5, as well as helices B, I, O and Q [Dutzler et al., 2002; Dutzler et al.,
58
59 2003; Wu et al., 2003; Feng et al., 2010].
60

1
2
3 Four other missense mutations involve residues near the transporter interface: the
4
5 p.Lys231Ileu mutation is in the loop between helices F and G, and mutations p.Arg239Pro,
6
7 p.Gly250Arg and p.Gly470Arg are in helices G and N. These mutations may have deleterious
8
9 consequences for CIC-5 by significantly affecting the stability of the helix and thus preventing
10
11 appropriate folding of the monomers. Consistent with this, other studies have reported that
12
13 such mutations result in abnormal CIC-5 proteins that are rapidly degraded [Lourdel et al.,
14
15 2012].
16
17
18

19
20 The p.Glu267Asp mutation maps to helix H, directly adjacent to the “proton glutamate”
21
22 (Glu268) that is strictly conserved on the internal side of CIC 2Cl⁻/H⁺ exchangers. This
23
24 residue plays a key role in the transport function of CIC 2Cl⁻/H⁺ exchangers by acting as an
25
26 H⁺ transfer site [Accardi et al., 2005; Neagoet al., 2010; Zdebik et al., 2008]. Neutralization
27
28 of the “proton glutamate” in CIC-5 abolishes Cl⁻ and H⁺ flux [Zdebiket al., 2008]. Because the
29
30 p.Glu267Asp substitution is very close to the H⁺ binding site, it may well severely impair the
31
32 transport cycle of CIC-5.
33
34
35
36
37
38
39
40

41 Recurrent Mutations

42
43
44 Among the 192 mutations described in the literature, 54 different were shared between two or
45
46 more families (Supp. Table S1). More frequent mutations were: p.Ser244Leu, p. p.Arg637*
47
48 and p.Arg704*, which were described in 16, 15 and 11 families respectively.
49
50
51

52
53 In our cohort, 24 previously described mutations were detected in 43 families (Table 2): the
54
55 p.Ser244Leu and the p.Arg637* mutations were detected in five different families; the
56
57 p.Arg347* nonsense mutation was detected in four different families; the p.Arg28* and the
58
59 p.Arg648* nonsense mutations and a whole gene deletion were detected in probands of three
60
families; the p.Arg704* nonsense mutation, the p.Val523del in-frame mutation and the

1
2
3 splicing mutation c.394-2A>G were found in two different families and finally, eleven
4
5 missense and two nonsense mutations, all of which had been described previously were
6
7 detected in only one family each.
8
9

10
11 Among recurrent mutations, nine nonsense (p.Arg28*, p.Arg34*, p.Trp279*, p.Arg347*,
12
13 p.Arg467*, p.Arg637*, p.Arg648*, p.Arg704* and p.Arg718*) and three missense
14
15 (p.Ser244Leu, p.Arg516Trp and p.Ser545Asn) correspond to C>T or G>A transitions arisen
16
17 in CpG dinucleotides. They have been detected in patients from different geographic origin
18
19 and can be considered as mutation hotspots.
20
21

22
23 Figure 2 summarizes novel and described mutations (n=234) by type and their location in
24
25 cDNA and protein domains. Mutations are scattered along all exons of the gene and protein
26
27 domains.
28
29

30 31 Variants of unknown clinical significance

32
33 **Table 3** summarises variants described in the literature and in our cohort as polymorphisms or
34
35 variants of unknown clinical significance (VUS). These including: four synonymous changes,
36
37 one missense variant (p.Ser386Phe) predicted *in silico* as benign and ten non-coding variants.
38
39 None of intronic variants were predicted to either create or abolish a splicing-related sequence
40
41 after *in silico* analysis with Alamut software.
42
43
44
45
46
47

48 Family studies

49
50
51 In our cohort, **initial** pedigree analysis of the 90 families with **Dent disease** showed **clinical**
52
53 familial history of X-linked disease in only eight families. We were able **to get additional**
54
55 **information (biochemical and ultrasound evaluation) and perform genetic studies in** 134
56
57 relatives from 43 families with four (n=2 families), three (n=7), two (n=31) or one (n=3)
58
59 generation **(Supp. Table S3). In these families we found co-segregation of the mutations with**
60

1
2
3 **the disease.** Thirty-seven of the 39 mothers and four of the six maternal grandmothers
4
5 analysed were heterozygous for the mutation carried by their son or grandson. In the
6
7 remaining cases mothers and grandmothers did not harbour the familial mutation for a *de*
8
9 *novo* mutation rate of 9%. These results are in accordance with the literature data available for
10
11 125 families: at a whole, 12 mothers did not harbour the mutation detected in their sons and in
12
13 3 additional cases, without familial antecedents, grandmothers did not harbour the mutation
14
15 detected in their daughters and grandsons, for a *de novo* mutation rate of 12%. Diagnosis of
16
17 **Dent disease 1** was confirmed in 28 additional males (brothers, cousins and maternal uncles);
18
19 in this group, 14 males belonging to 10 families were **clinically** asymptomatic before the
20
21 genetic diagnosis, which was performed in seven before the age of 5 years, in three between
22
23 the ages of 5 and 10 and in the remaining cases between 16 and 32 years (**Table 4** and **Supp.**
24
25 **Table S4**). No renal impairment was detected in this group but the genetic diagnosis allowed
26
27 establishing adequate follow-up by a nephrologist. In our cohort, 68 out of 84 women tested
28
29 were identified as carriers; such findings are very important for genetic counselling: i.e. early
30
31 screening of any male children from carrier mother and giving reassurance to non-carrier
32
33 mothers about the transmission of the disease to their children. The pedigrees of two large
34
35 families (44 and 66) for which four generations were studied, and the pedigrees of two
36
37 families (families 11 and 20) for which three generations were studied are given in the **Supp.**
38
39 **Figure S1.**

40
41
42
43
44
45
46
47
48
49 The geographical distribution available for 332 described families as well as for the additional
50
51 79 families of our cohort is described in **Supp. Table S5**. **Dent disease 1** is a worldwide
52
53 disease. The high number of descriptions in Japan is the result of the detection of LMWP in
54
55 children as product of nationwide program of urine screening of elementary and junior high
56
57 school students (Lloyd et al., 1997).

58
59
60 Phenotype (Clinical relevance):

1
2
3 Published clinical data from 377 male patients belonging to 334 families were analysed. Only
4
5 clinical data from articles published in English were included. We found 57 articles
6
7 describing the phenotype associated with mutations in the *CLCN5* gene. Phenotype
8
9 description in papers is very variable from qualitative description of main criteria to detailed
10
11 description in case reports. **Table 4** summarizes and compares global clinical qualitative data
12
13 from previously described large cohorts, from case-reports and from our cohort (including the
14
15 phenotypic data of fourteen patients belonging to 11 families previously described [Blanchard
16
17 et al., 2008, Grand et al., 2009]). Median age at diagnosis was 9 (0.2-67) years in the literature
18
19 and 7 (0.1-55) in our cohort. Altogether the main manifestations of the disease in our cohort
20
21 were similar to those described in worldwide literature.
22
23
24
25

26
27 In addition to the main characteristics the following clinical and biological abnormalities have
28
29 been described in patients with **Dent disease 1 (absolute value are given when we only had**
30
31 **information of the presence of a given trait on case reports; relative values correspond to the**
32
33 **number of positive/total cases when absence or presence of a given trait was clearly**
34
35 **described)** : micro or macro-haematuria (n=71), polyuria/polydipsia or urinary concentration
36
37 defect (31/43), failure to thrive including four cases with GH deficiency (n=16), hypouricemia
38
39 (13/38), proteinuria in the nephrotic range without hypoalbuminemia (n=13) [Lim et al., 2007,
40
41 Sheffer-Babila et al 2008, Frishberg et al. 2009, Fervenza 2013, Cramer et al. 2014], reduced
42
43 renal uptake of DMSA (n=7), enuresis (n=5), hypomagnesaemia (4/30), secondary
44
45 hyperaldosteronism with Bartter-like phenotype (n=3) [Besbas et al. 2005, Bogdanovic et al.
46
47 2010, Okamoto et al. 2011] , and night blindness responsive to vitamin A [Sethi et al. 2009].
48
49
50
51
52
53

54
55 We retrieved quantitative data on proteinuria, expressed in g/day, in 57 patients; the median
56
57 value was 1.28 g/24h (min 0.27 – max 4.50) (IQR 0.9-1.92). Renal biopsy was performed in
58
59 37 patients as part of diagnosis of proteinuria and/or renal impairment; in addition, renal
60
histologic analysis after necropsy was available in one patient. In three cases light microscopy

1
2
3 analysis was described as normal. In the 35 remaining patients, light microscopy analysis
4
5 showed mainly tubular atrophy and interstitial fibrosis, tubular or interstitial calcification and
6
7 nonspecific glomerular changes as sclerosis and periglomerular fibrosis. Twelve of these
8
9 cases were classified as focal global glomerulosclerosis. These results are summarized in
10
11 Supp. Table S6.
12
13

14
15 A considerable intra-familial variability in disease severity has been observed and no
16
17 genotype-phenotype correlation has been described. For a same mutation, patients from the
18
19 literature as well as from our cohort have a variable phenotype ranging from renal Fanconi
20
21 syndrome with or without rickets to the association of LMWP and hypercalciuria (with or
22
23 without nephrocalcinosis or nephrolithiasis).
24
25
26

27
28 Female carriers displayed a milder and variable phenotype, attributed to the differing amounts
29
30 of inactivation of the X chromosome that harbours the *CLCN5* mutation. We gathered
31
32 available phenotypic characteristic in 89 heterozygous women described in the literature and
33
34 data from 6 heterozygous carriers in our cohort: LMWP was found in 60% of female carriers
35
36 (52/86), 50 times lower than in affected males; median urinary β_2 microglobuline
37
38 concentration was 1.50 mg/L (IQR 0.74-3.52) mg/l in females and 52.20 (IQR 31-99) mg/l in
39
40 males. Hypercalciuria was detected in 31% (20/64), nephrolithiasis in 21% (11/51) and
41
42 nephrocalcinosis in 12% of female carriers (5/40). So far, only 1 female carrier was reported
43
44 to have end stage renal disease [Wrong 1994], but CKD has been described in 3 additional
45
46 heterozygous women: CKD stage 3a in one at 56 years [Igarashi 2000] and CKD stage 3b in
47
48 two at 65 and 78 years [Hoopes 1998, Kelleher 1998] . Supp. Figure S2 shows comparisons
49
50 of tubular proteinuria and calciuria values in male patients and heterozygous carriers.
51
52
53
54
55

56
57
58 Other forms of Dent disease:
59
60

1
2
3 **Dent disease** is genetically heterogeneous: in published cohorts about 50-60% of patients have
4
5 **Dent disease 1** caused by mutations of the *CLCN5* gene and about 15% have **Dent disease 2**
6
7 caused by mutations in the *OCRL* gene. One case of co-inheritance of mutations in both genes
8
9 has been described in a patient with a phenotype close to Lowe syndrome [Addis et al., 2013].
10
11 In our cohort of 150 distinct families, **Dent disease 1** represents 61 % and **Dent disease 2**
12
13 represents 7% of the cases; most of these cases have been previously described [Hichri et al.,
14
15 2011]. Accordingly, 25 to 35% of cases with **Dent disease** still remain without molecular
16
17 identification. *CLCN4*, *CFL1*, *SLC9A6* and *TMEM27* genes coding for ClC-4, Cofilin-1,
18
19 NHE6 and collectrin have been excluded as cause of **Dent disease** [Ludwig and Utsch 2004,
20
21 Hoopes R et al 2005, Wu et al., 2009 and Tosetto et al., 2009].
22
23
24
25
26
27

28 Animal models:

29
30
31 Two independent strains of ClC-5 knock-out (KO) **mice** have been generated [Piwon et al.
32
33 2000, Wang et al. 2000]. These two strains recapitulated the major features of **Dent disease**
34
35 1, including LMW proteinuria and other manifestations of proximal tubule dysfunction.
36
37 Nevertheless, as in human, there is a phenotype variability: Piwon's model showed no
38
39 hypercalciuria, increased urinary levels of PTH and reduced levels of 1.25(OH)₂-vitamin D₃
40
41 and Wang's model showed hypercalciuria and elevated levels of 1.25(OH)₂-vitamin D₃. *In*
42
43 *vitro* experiments showed defective acidification of vesicles isolated from ClC-5 KO mice,
44
45 supporting a role for ClC-5 in acidification of early endosomes [Günther et al. 2003, Novarino
46
47 et al. 2010]. After the discovery in 2005 that ClC-5 was in fact an electrogenic, 2Cl⁻/H⁺
48
49 exchanger, instead of being a simple Cl⁻ channel [Picollo & Pusch, 2005, Scheel et al. 2005],
50
51 a knock-in (KI) mouse model carrying a point mutation p.Glu211Ala was generated. This
52
53 change affects a glutamate residue that is essential for the gating of the ClC exchangers
54
55 converting ClC-5 into a pure uncoupled Cl⁻ conductor [Novarino et al. 2010]. The E211A
56
57 mutant ClC-5 did not affect endosomal acidification; however, the KI mice showed the same
58
59
60

1
2
3 renal phenotype than KO mice and patients with **Dent disease**, including LMW proteinuria,
4
5 glycosuria, hyperphosphaturia and hypercalciuria. Furthermore, both the KI and KO mouse
6
7 showed a similar impairment in PT endocytosis, with reduced levels of the endocytic
8
9 receptors megalin and cubilin and internalization of the sodium-phosphate cotransporter
10
11 NaPi-2a indicating a trafficking defect [Novarino et al. 2010].
12
13
14
15
16
17

18 Conclusion

19
20
21 In this paper, we have compiled, previously reported and novel mutations of the *CLCN5* gene
22
23 responsible for **Dent disease 1** and described the phenotype associated. We also describe the
24
25 characterization of 42 novel mutations of the *CLCN5* gene and propose mechanisms of
26
27 pathogenicity for the novel missense and in-frame mutations. By familial screening, we
28
29 identified 67 carrier females and 14 asymptomatic affected males, highlighting the importance
30
31 of genotyping the probands. Further studies are necessary to confirm the value of early
32
33 diagnosis and follow-up, as well as to understand the phenotypic variability and to identify the
34
35 molecular causes in the about 30% of the Dent disease patients for whom the aetiology is
36
37 unclear.
38
39
40
41
42
43
44
45
46

47 Acknowledgments

48
49
50 We thank the patients and their families who participated in this study for their cooperation.

51
52 We thank all the physicians of the French Tubulopathies Network and all the staff of the
53
54 genetic laboratory at Georges Pompidou Hospital. We would like to thank Dr. Johan den
55
56 Dunnen for his help in updating LOVD database.
57
58

59
60 The authors have no conflict of interest to declare.

References

- 1
2
3
4
5
6
7
8
9
10 Abagyan R, Frishman D, Argos P. 1994. Recognition of distantly related proteins through
11 energy calculations. *Proteins* 19(2):132-40.
- 12 Abagyan R, Totrov M. 1994. Biased probability Monte Carlo conformational searches and
13 electrostatic calculations for peptides and proteins. *J Mol Biol* 235(3):983-1002.
- 14 Accardi A, Walden M, Nguitragool W, Jayaram H, Williams C, Miller C. 2005. Separate ion
15 pathways in a Cl⁻/H⁺ exchanger. *J Gen Physiol* 126(6):563-70.
- 16 Addis M, Meloni C, Toso E, Ceol M, Cristofaro R, Melis MA, Vercelloni P, Del Prete D,
17 Marra G, Anglani F. An atypical Dent's disease phenotype caused by co-inheritance of
18 mutations at CLCN5 and OCRL genes. *Eur J Hum Genet* 21(6):687-90.
- 19 Akuta N, Lloyd SE, Igarashi T, Shiraga H, Matsuyama T, Yokoro S, Cox JP, Thakker RV.
20 1997. Mutations of CLCN5 in Japanese children with idiopathic low molecular weight
21 proteinuria, hypercalciuria and nephrocalcinosis. *Kidney Int* 52(4):911-6.
- 22 Anglani F, Bernich P, Toso E, Cara M, Lupo A, Nalesso F, D'Angelo A, Gambaro G. 2006.
23 Family history may be misleading in the diagnosis of Dent's disease. *Urol Res*
24 34(1):61-3.
- 25 Attree O, Olivos IM, Okabe I, Bailey LC, Nelson DL, Lewis RA, McInnes RR, Nussbaum
26 RL. 1992. The Lowe's oculocerebrorenal syndrome gene encodes a protein highly
27 homologous to inositol polyphosphate-5-phosphatase. *Nature* 358(6383):239-42.
- 28 Besbas N, Ozaltin F, Jeck N, Seyberth H, Ludwig M. 2005. CLCN5 mutation (R347X)
29 associated with hypokalaemic metabolic alkalosis in a Turkish child: an unusual
30 presentation of Dent's disease. *Nephrol Dial Transplant* 20(7):1476-9.
- 31 Blanchard A, Vargas-Poussou R, Peyrard S, Mogenet A, Baudouin V, Boudailliez B, Charbit
32 M, Deschesnes G, Ezzhair N, Loirat C and others. 2008. Effect of hydrochlorothiazide
33 on urinary calcium excretion in dent disease: an uncontrolled trial. *Am J Kidney Dis*
34 52(6):1084-95.
- 35 Bogdanovic R, Draaken M, Toromanovic A, Dordevic M, Stajic N, Ludwig M. A novel
36 CLCN5 mutation in a boy with Bartter-like syndrome and partial growth hormone
37 deficiency. *Pediatr Nephrol* 25(11):2363-8.
- 38 Bokenkamp A, Bockenhauer D, Cheong HI, Hoppe B, Tasic V, Unwin R, Ludwig M. 2009.
39 Dent-2 disease: a mild variant of Lowe syndrome. *J Pediatr* 155(1):94-9.
- 40 Bosio M, Bianchi ML, Lloyd SE, Thakker RV. 1999. A familial syndrome due to Arg648Stop
41 mutation in the X-linked renal chloride channel gene. *Pediatr Nephrol* 13(4):278-83.
- 42 Brakemeier S, Si H, Gollasch M, Hoffler D, Buhl M, Kohler R, Hoyer J, Eichler I. 2004.
43 Dent's disease: identification of a novel mutation in the renal chloride channel
44 CLCN5. *Clin Nephrol* 62(5):387-90.
- 45 Carballo-Trujillo I, Garcia-Nieto V, Moya-Angeler FJ, Anton-Gamero M, Loris C, Mendez-
46 Alvarez S, Claverie-Martin F. 2003. Novel truncating mutations in the ClC-5 chloride
47 channel gene in patients with Dent's disease. *Nephrol Dial Transplant* 18(4):717-23.
- 48 Cheong HI, Lee JW, Zheng SH, Lee JH, Kang JH, Kang HG, Ha IS, Lee SJ, Choi Y. 2005.
49 Phenotype and genotype of Dent's disease in three Korean boys. *Pediatr Nephrol*
50 20(4):455-9.
- 51 Cho HY, Lee BH, Choi HJ, Ha IS, Choi Y, Cheong HI. 2008. Renal manifestations of Dent
52 disease and Lowe syndrome. *Pediatr Nephrol* 23(2):243-9.
- 53
54
55
56
57
58
59
60

- 1
2
3 Claverie-Martin F, Flores C, Anton-Gamero M, Gonzalez-Acosta H, Garcia-Nieto V. 2005.
4 The Alu insertion in the CLCN5 gene of a patient with Dent's disease leads to exon 11
5 skipping. *J Hum Genet* 50(7):370-4.
6
7 Claverie-Martin F, Gonzalez-Acosta H, Flores C, Anton-Gamero M, Garcia-Nieto V. 2003.
8 De novo insertion of an Alu sequence in the coding region of the CLCN5 gene results
9 in Dent's disease. *Hum Genet* 113(6):480-5.
10
11 Cramer MT, Charlton JR, Fogo AB, Fathallah-Shaykh SA, Askenazi DJ, Guay-Woodford
12 LM. Expanding the phenotype of proteinuria in Dent disease. A case series. *Pediatr*
13 *Nephrol* 29(10):2051-4.
14
15 Copelovitch L, Nash MA, Kaplan BS. 2007. Hypothesis: Dent disease is an underrecognized
16 cause of focal glomerulosclerosis. *Clin J Am Soc Nephrol* 2(5):914-8.
17
18 Cox JP, Yamamoto K, Christie PT, Wooding C, Feest T, Flinter FA, Goodyer PR, Leumann
19 E, Neuhaus T, Reid C and others. 1999. Renal chloride channel, CLCN5, mutations in
20 Dent's disease. *J Bone Miner Res* 14(9):1536-42.
21
22 D'Antonio C, Molinski S, Ahmadi S, Huan LJ, Wellhauser L, Bear CE. Conformational
23 defects underlie proteasomal degradation of Dent's disease-causing mutants of ClC-5.
24 *Biochem J* 452(3):391-400.
25
26 Devuyst O, Thakker RV. Dent's disease. *Orphanet J Rare Dis* 5:28.
27
28 Dinour D, Davidovitz M, Levin-Iaina N, Lotan D, Cleper R, Weissman I, Knecht A,
29 Holtzman EJ. 2009. Truncating mutations in the chloride/proton ClC-5 antiporter gene
30 in Seven Jewish Israeli families with Dent's 1 disease. *Nephron Clin Pract*
31 112(4):c262-7.
32
33 Dutzler R. 2006. The ClC family of chloride channels and transporters. *Curr Opin Struct Biol*
34 16(4):439-46.
35
36 Dutzler R, Campbell EB, Cadene M, Chait BT, MacKinnon R. 2002. X-ray structure of a ClC
37 chloride channel at 3.0 Å reveals the molecular basis of anion selectivity. *Nature*
38 415(6869):287-94.
39
40 Dutzler R, Campbell EB, MacKinnon R. 2003. Gating the selectivity filter in ClC chloride
41 channels. *Science* 300(5616):108-12.
42
43 Feng L, Campbell EB, Hsiung Y, MacKinnon R. Structure of a eukaryotic ClC transporter
44 defines an intermediate state in the transport cycle. *Science* 330(6004):635-41.
45
46 Fervenza FC. A patient with nephrotic-range proteinuria and focal global glomerulosclerosis.
47 *Clin J Am Soc Nephrol* 8(11):1979-87.
48
49 Forino M, Graziotto R, Tosetto E, Gambaro G, D'Angelo A, Anglani F. 2004. Identification
50 of a novel splice site mutation of CLCN5 gene and characterization of a new
51 alternative 5' UTR end of ClC-5 mRNA in human renal tissue and leukocytes. *J Hum*
52 *Genet* 49(1):53-60.
53
54 Frishberg Y, Dinour D, Belostotsky R, Becker-Cohen R, Rinat C, Feinstein S, Navon-Elkan
55 P, Ben-Shalom E. 2009. Dent's disease manifesting as focal glomerulosclerosis: Is it
56 the tip of the iceberg? *Pediatr Nephrol* 24(12):2369-73.
57
58 Grand T, L'Hoste S, Mordasini D, Defontaine N, Keck M, Pennaforte T, Genete M, Laghmani
59 K, Teulon J, Lourdel S. Heterogeneity in the processing of CLCN5 mutants related to
60 Dent disease. *Hum Mutat* 32(4):476-83.
Grand T, Mordasini D, L'Hoste S, Pennaforte T, Genete M, Biyeyeme MJ, Vargas-Poussou R,
Blanchard A, Teulon J, Lourdel S. 2009. Novel CLCN5 mutations in patients with
Dent's disease result in altered ion currents or impaired exchanger processing. *Kidney*
Int 76(9):999-1005.
Gunther W, Piwon N, Jentsch TJ. 2003. The ClC-5 chloride channel knock-out mouse - an
animal model for Dent's disease. *Pflugers Arch* 445(4):456-62.

- 1
2
3 Hichri H, Rendu J, Monnier N, Coutton C, Dorseuil O, Poussou RV, Baujat G, Blanchard A,
4 Nobili F, Ranchin B and others. From Lowe syndrome to Dent disease: correlations
5 between mutations of the OCRL1 gene and clinical and biochemical phenotypes. *Hum*
6 *Mutat* 32(4):379-88.
- 7
8 Hoopes RR, Jr., Hueber PA, Reid RJ, Jr., Braden GL, Goodyer PR, Melnyk AR, Midgley JP,
9 Moel DI, Neu AM, VanWhy SK and others. 1998. CLCN5 chloride-channel mutations
10 in six new North American families with X-linked nephrolithiasis. *Kidney Int*
11 54(3):698-705.
- 12
13 Hoopes RR, Jr., Raja KM, Koich A, Hueber P, Reid R, Knohl SJ, Scheinman SJ. 2004.
14 Evidence for genetic heterogeneity in Dent's disease. *Kidney Int* 65(5):1615-20.
- 15
16 Hoopes RR, Jr., Shrimpton AE, Knohl SJ, Hueber P, Hoppe B, Matyus J, Simckes A, Tasic
17 V, Toenshoff B, Suchy SF and others. 2005. Dent Disease with mutations in OCRL1.
18 *Am J Hum Genet* 76(2):260-7.
- 19
20 Hryciw DH, Ekberg J, Ferguson C, Lee A, Wang D, Parton RG, Pollock CA, Yun CC,
21 Poronnik P. 2006. Regulation of albumin endocytosis by PSD95/Dlg/ZO-1 (PDZ)
22 scaffolds. Interaction of Na⁺-H⁺ exchange regulatory factor-2 with ClC-5. *J Biol*
23 *Chem* 281(23):16068-77.
- 24
25 Hryciw DH, Ekberg J, Pollock CA, Poronnik P. 2006. ClC-5: a chloride channel with
26 multiple roles in renal tubular albumin uptake. *Int J Biochem Cell Biol* 38(7):1036-42.
- 27
28 Humphrey W, Dalke A, Schulten K. 1996. VMD: visual molecular dynamics. *J Mol Graph*
29 14(1):33-8, 27-8.
- 30
31 Igarashi T, Gunther W, Sekine T, Inatomi J, Shiraga H, Takahashi S, Suzuki J, Tsuru N,
32 Yanagihara T, Shimazu M and others. 1998. Functional characterization of renal
33 chloride channel, CLCN5, mutations associated with Dent's Japan disease. *Kidney Int*
34 54(6):1850-6.
- 35
36 Igarashi T, Inatomi J, Ohara T, Kuwahara T, Shimadzu M, Thakker RV. 2000. Clinical and
37 genetic studies of CLCN5 mutations in Japanese families with Dent's disease. *Kidney*
38 *Int* 58(2):520-7.
- 39
40 Ji LN, Chen CY, Wang JJ, Cao L. A novel CLCN5 mutation in a Chinese boy with Dent's
41 disease. *World J Pediatr* 10(3):275-7.
- 42
43 Kelleher CL, Buckalew VM, Frederickson ED, Rhodes DJ, Conner DA, Seidman JG,
44 Seidman CE. 1998. CLCN5 mutation Ser244Leu is associated with X-linked renal
45 failure without X-linked recessive hypophosphatemic rickets. *Kidney Int* 53(1):31-7.
- 46
47 Langlois V, Bernard C, Scheinman SJ, Thakker RV, Cox JP, Goodyer PR. 1998. Clinical
48 features of X-linked nephrolithiasis in childhood. *Pediatr Nephrol* 12(8):625-9.
- 49
50 Lee BH, Lee SH, Choi HJ, Kang HG, Oh SW, Lee DS, Ha IS, Choi Y, Cheong HI. 2009.
51 Decreased renal uptake of (99m)Tc-DMSA in patients with tubular proteinuria.
52 *Pediatr Nephrol* 24(11):2211-6.
- 53
54 Li P, Huang JP. 2009. Phenotype and genotype of Dent's disease in three Chinese boys.
55 *Nephrology (Carlton)* 14(2):139-42.
- 56
57 Lim IS, Yun KW, Moon KC, Cheong HI. 2007. Proteinuria in a boy with infectious
58 mononucleosis, C1q nephropathy, and Dent's disease. *J Korean Med Sci* 22(5):928-31.
- 59
60 Lloyd SE, Gunther W, Pearce SH, Thomson A, Bianchi ML, Bosio M, Craig IW, Fisher SE,
Scheinman SJ, Wrong O and others. 1997. Characterisation of renal chloride channel,
CLCN5, mutations in hypercalciuric nephrolithiasis (kidney stones) disorders. *Hum*
Mol Genet 6(8):1233-9.
- Lloyd SE, Pearce SH, Fisher SE, Steinmeyer K, Schwappach B, Scheinman SJ, Harding B,
Bolino A, Devoto M, Goodyer P and others. 1996. A common molecular basis for
three inherited kidney stone diseases. *Nature* 379(6564):445-9.

- 1
2
3 Lloyd SE, Pearce SH, Gunther W, Kawaguchi H, Igarashi T, Jentsch TJ, Thakker RV. 1997.
4 Idiopathic low molecular weight proteinuria associated with hypercalciuric
5 nephrocalcinosis in Japanese children is due to mutations of the renal chloride channel
6 (CLCN5). *J Clin Invest* 99(5):967-74.
7
8 Lourdel S, Grand T, Burgos J, Gonzalez W, Sepulveda FV, Teulon J. CIC-5 mutations
9 associated with Dent's disease: a major role of the dimer interface. *Pflugers Arch*
10 463(2):247-56.
11
12 Ludwig M, Doroszewicz J, Seyberth HW, Bokenkamp A, Balluch B, Nuutinen M, Utsch B,
13 Waldegger S. 2005. Functional evaluation of Dent's disease-causing mutations:
14 implications for CIC-5 channel trafficking and internalization. *Hum Genet* 117(2-
15 3):228-37.
16
17 Ludwig M, Utsch B. 2004. Dent disease-like phenotype and the chloride channel CIC-4
18 (CLCN4) gene. *Am J Med Genet A* 128A(4):434-5.
19
20 Ludwig M, Utsch B, Balluch B, Frund S, Kuwertz-Broking E, Bokenkamp A. 2006.
21 Hypercalciuria in patients with CLCN5 mutations. *Pediatr Nephrol* 21(9):1241-50.
22
23 Mackerell AD, Jr., Feig M, Brooks CL, 3rd. 2004. Extending the treatment of backbone
24 energetics in protein force fields: limitations of gas-phase quantum mechanics in
25 reproducing protein conformational distributions in molecular dynamics simulations. *J*
26 *Comput Chem* 25(11):1400-15.
27
28 Matsuyama T, Awazu M, Oikawa T, Inatomi J, Sekine T, Igarashi T. 2004. Molecular and
29 clinical studies of Dent's disease in Japan: biochemical examination and renal
30 ultrasonography do not predict carrier state. *Clin Nephrol* 61(4):231-7.
31
32 Mo L, Xiong W, Qian T, Sun H, Wills NK. 2004. Coexpression of complementary fragments
33 of CIC-5 and restoration of chloride channel function in a Dent's disease mutation. *Am*
34 *J Physiol Cell Physiol* 286(1):C79-89.
35
36 Morimoto T, Uchida S, Sakamoto H, Kondo Y, Hanamizu H, Fukui M, Tomino Y, Nagano
37 N, Sasaki S, Marumo F. 1998. Mutations in CLCN5 chloride channel in Japanese
38 patients with low molecular weight proteinuria. *J Am Soc Nephrol* 9(5):811-8.
39
40 Nakazato H, Hattori S, Furuse A, Kawano T, Karashima S, Tsuruta M, Yoshimuta J, Endo F,
41 Matsuda I. 1997. Mutations in the CLCN5 gene in Japanese patients with familial
42 idiopathic low-molecular-weight proteinuria. *Kidney Int* 52(4):895-900.
43
44 Nakazato H, Yoshimuta J, Karashima S, Matsumoto S, Endo F, Matsuda I, Hattori S. 1999.
45 Chloride channel CLCN5 mutations in Japanese children with familial idiopathic low
46 molecular weight proteinuria. *Kidney Int* 55(1):63-70.
47
48 Neagoe I, Stauber T, Fidzinski P, Bergsdorf EY, Jentsch TJ. The late endosomal CIC-6
49 mediates proton/chloride countertransport in heterologous plasma membrane
50 expression. *J Biol Chem* 285(28):21689-97.
51
52 Norden AG, Scheinman SJ, Deschodt-Lanckman MM, Lapsley M, Nortier JL, Thakker RV,
53 Unwin RJ, Wrong O. 2000. Tubular proteinuria defined by a study of Dent's (CLCN5
54 mutation) and other tubular diseases. *Kidney Int* 57(1):240-9.
55
56 Novarino G, Weinert S, Rickheit G, Jentsch TJ. Endosomal chloride-proton exchange rather
57 than chloride conductance is crucial for renal endocytosis. *Science* 328(5984):1398-
58 401.
59
60 Okamoto T, Tajima T, Hirayama T, Sasaki S. A patient with Dent disease and features of
Bartter syndrome caused by a novel mutation of CLCN5. *Eur J Pediatr* 171(2):401-4.
Oudet C, Martin-Coignard D, Pannetier S, Praud E, Champion G, Hanauer A. 1997. A second
family with XLRH displays the mutation S244L in the CLCN5 gene. *Hum Genet*
99(6):781-4.
Park E, Choi HJ, Lee JM, Ahn YH, Kang HG, Choi YM, Park SJ, Cho HY, Park YH, Lee SJ
and others. Muscle involvement in Dent disease 2. *Pediatr Nephrol* 29(11):2127-32.

- 1
2
3 Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, Chipot C, Skeel RD, Kale
4 L, Schulten K. 2005. Scalable molecular dynamics with NAMD. *J Comput Chem*
5 26(16):1781-802.
- 6
7 Picollo A, Pusch M. 2005. Chloride/proton antiporter activity of mammalian CLC proteins
8 CIC-4 and CIC-5. *Nature* 436(7049):420-3.
- 9
10 Piwon N, Gunther W, Schwake M, Bosl MR, Jentsch TJ. 2000. CIC-5 Cl⁻-channel disruption
11 impairs endocytosis in a mouse model for Dent's disease. *Nature* 408(6810):369-73.
- 12
13 Ramos-Trujillo E, Claverie-Martin F, Garcia-Nieto V, Ariceta G, Vara J, Gonzalez-Acosta H,
14 Garcia-Ramirez M, Fons J, Cordoba-Lanus E, Gonzalez-Paredes J and others. 2013.
15 Dent's disease: Identification of seven new pathogenic mutations in the CLCN5 gene.
16 *Journal of Pediatric Genetics* 2(3):133-140.
- 17
18 Ramos-Trujillo E, Garcia-Nieto V, Gonzalez-Acosta H, Vara J, Perez-Diaz V, Nadal I,
19 Oliveros R, Claverie-Martin F. 2007. Molecular analysis of the CLCN5 gene in Dent's
20 disease: first mutation identified in a patient from South America. *Clin Nephrol*
21 68(6):367-72.
- 22
23 Ramos-Trujillo E, Gonzalez-Acosta H, Flores C, Garcia-Nieto V, Guillen E, Gracia S,
24 Vicente C, Espinosa L, Maseda MA, Santos F and others. 2007. A missense mutation
25 in the chloride/proton CIC-5 antiporter gene results in increased expression of an
26 alternative mRNA form that lacks exons 10 and 11. Identification of seven new
27 CLCN5 mutations in patients with Dent's disease. *J Hum Genet* 52(3):255-61.
- 28
29 Samardzic M, Pavicevic S, Ludwig M, Bogdanovic R. Effect of growth hormone replacement
30 therapy in a boy with Dent's disease: a case report. *J Med Case Rep* 5:400.
- 31
32 Scheel O, Zdebik AA, Lourdel S, Jentsch TJ. 2005. Voltage-dependent electrogenic
33 chloride/proton exchange by endosomal CLC proteins. *Nature* 436(7049):424-7.
- 34
35 Scheel O, Zdebik AA, Lourdel S, Jentsch TJ. 2005. Voltage-dependent electrogenic
36 chloride/proton exchange by endosomal CLC proteins. *Nature* 436(7049):424-7.
- 37
38 Scheinman SJ, Cox JP, Lloyd SE, Pearce SH, Salenger PV, Hoopes RR, Bushinsky DA,
39 Wrong O, Asplin JR, Langman CB and others. 2000. Isolated hypercalciuria with
40 mutation in CLCN5: relevance to idiopathic hypercalciuria. *Kidney Int* 57(1):232-9.
- 41
42 Scheinman SJ. 2009. Dent's disease. In: Lifton RP, Somlo S, Giebisch GH, Seldin DW,
43 editors. *Genetic Diseases of the Kidney*. 1st Elsevier, pAcademic Press Elsevier, p213-
44 226.
- 45
46 Schwake M, Friedrich T, Jentsch TJ. 2001. An internalization signal in CIC-5, an endosomal
47 Cl⁻-channel mutated in dent's disease. *J Biol Chem* 276(15):12049-54.
- 48
49 Sekine T, Komoda F, Miura K, Takita J, Shimadzu M, Matsuyama T, Ashida A, Igarashi T.
50 Japanese Dent disease has a wider clinical spectrum than Dent disease in
51 Europe/USA: genetic and clinical studies of 86 unrelated patients with low-molecular-
52 weight proteinuria. *Nephrol Dial Transplant* 29(2):376-84.
- 53
54 Sethi SK, Ludwig M, Kabra M, Hari P, Bagga A. 2009. Vitamin A responsive night blindness
55 in Dent's disease. *Pediatr Nephrol* 24(9):1765-70.
- 56
57 Sheffer-Babila S, Chandra M, Speiser PW. 2008. Growth hormone improves growth rate and
58 preserves renal function in Dent disease. *J Pediatr Endocrinol Metab* 21(3):279-86.
- 59
60 Smith AJ, Reed AA, Loh NY, Thakker RV, Lippiat JD. 2009. Characterization of Dent's
disease mutations of CLC-5 reveals a correlation between functional and cell
biological consequences and protein structure. *Am J Physiol Renal Physiol*
296(2):F390-7.
- Takemura T, Hino S, Ikeda M, Okada M, Igarashi T, Inatomi J, Yoshioka K. 2001.
Identification of two novel mutations in the CLCN5 gene in Japanese patients with
familial idiopathic low molecular weight proteinuria (Japanese Dent's disease). *Am J
Kidney Dis* 37(1):138-143.

- 1
2
3 Tanuma A, Sato H, Takeda T, Hosojima M, Obayashi H, Hama H, Iino N, Hosaka K, Kaseda
4 R, Imai N and others. 2007. Functional characterization of a novel missense CLCN5
5 mutation causing alterations in proximal tubular endocytic machinery in Dent's
6 disease. *Nephron Physiol* 107(4):p87-97.
- 7
8 Tosetto E, Addis M, Caridi G, Meloni C, Emma F, Vergine G, Stringini G, Papalia T,
9 Barbano G, Ghiggeri GM and others. 2009. Locus heterogeneity of Dent's disease:
10 OCRL1 and TMEM27 genes in patients with no CLCN5 mutations. *Pediatr Nephrol*
11 24(10):1967-73.
- 12
13 Tosetto E, Ceol M, Mezzabotta F, Ammenti A, Peruzzi L, Caruso MR, Barbano G, Vezzoli G,
14 Colussi G, Vergine G and others. 2009. Novel mutations of the CLCN5 gene
15 including a complex allele and A 5' UTR mutation in Dent disease 1. *Clin Genet*
16 76(4):413-6.
- 17
18 Tosetto E, Ghiggeri GM, Emma F, Barbano G, Carrea A, Vezzoli G, Torregrossa R, Cara M,
19 Ripanti G, Ammenti A and others. 2006. Phenotypic and genetic heterogeneity in
20 Dent's disease--the results of an Italian collaborative study. *Nephrol Dial Transplant*
21 21(9):2452-63.
- 22
23 Valina MR, Larsen CP, Kanosky S, Suchy SF, Nield LS, Onder AM. A novel CLCN5
24 mutation in a boy with asymptomatic proteinuria and focal global glomerulosclerosis.
25 *Clin Nephrol* 80(5):377-84.
- 26
27 Wang SS, Devuyt O, Courtoy PJ, Wang XT, Wang H, Wang Y, Thakker RV, Guggino S,
28 Guggino WB. 2000. Mice lacking renal chloride channel, CLC-5, are a model for
29 Dent's disease, a nephrolithiasis disorder associated with defective receptor-mediated
30 endocytosis. *Hum Mol Genet* 9(20):2937-45.
- 31
32 Wrong OM, Norden AG, Feest TG. 1994. Dent's disease; a familial proximal renal tubular
33 syndrome with low-molecular-weight proteinuria, hypercalciuria, nephrocalcinosis,
34 metabolic bone disease, progressive renal failure and a marked male predominance.
35 *Qjm* 87(8):473-93.
- 36
37 Wu F, Reed AA, Williams SE, Loh NY, Lippiat JD, Christie PT, Large O, Bettinelli A, Dillon
38 MJ, Goldraich NP and others. 2009. Mutational analysis of CLC-5, cofilin and CLC-4
39 in patients with Dent's disease. *Nephron Physiol* 112(4):p53-62.
- 40
41 Wu F, Roche P, Christie PT, Loh NY, Reed AA, Esnouf RM, Thakker RV. 2003. Modeling
42 study of human renal chloride channel (hCLC-5) mutations suggests a structural-
43 functional relationship. *Kidney Int* 63(4):1426-32.
- 44
45 Yamamoto K, Cox JP, Friedrich T, Christie PT, Bald M, Houtman PN, Lapsley MJ, Patzer L,
46 Tsimaratos M, Van THWG and others. 2000. Characterization of renal chloride
47 channel (CLCN5) mutations in Dent's disease. *J Am Soc Nephrol* 11(8):1460-8.
- 48
49 Yanagida H, Ikeoka M, Kuwajima H, Wada N, Tabata N, Sugimoto K, Okada M, Takemura
50 T. 2004. A boy with Japanese Dent's disease exhibiting abnormal calcium metabolism
51 and osseous disorder of the spine: defective megalin expression at the brushborder of
52 renal proximal tubules. *Clin Nephrol* 62(4):306-12.
- 53
54 Zdebik AA, Zifarelli G, Bergsdorf EY, Soliani P, Scheel O, Jentsch TJ, Pusch M. 2008.
55 Determinants of anion-proton coupling in mammalian endosomal CLC proteins. *J Biol*
56 *Chem* 283(7):4219-27.
- 57
58 Zhang H, Wang C, Yue H, Hu WW, Gu JM, He JW, Fu WZ, Liu YJ, Zhang Z, Zhang ZL.
59 Identification of a novel mutation in the CLCN5 gene in a Chinese family with Dent-1
60 disease. *Nephrology (Carlton)* 19(2):80-3.
- Zhu BZ, Li P, Huang JP. [Clinical and genetic analysis of Dent's disease in 6 Chinese
children with low molecular weight proteinuria]. *Zhonghua Er Ke Za Zhi* 48(5):329-
33.

1
2
3
4
5
6
7
8
9
10 Figure Legends

11
12
13 Figure 1.

14
15
16 Three-dimensional model of ClC-5 based on homology with the structure of StClC [Dutzler et
17 al., 2002] showing the location of the ClC-5 mutations. Views from the side of the membrane
18 with the extracellular medium at the top (A) and from a direction rotated by 90° from the A
19 monomer (B). The view in (B) shows the interface between the two subunits of the dimeric
20 protein. Mutated residues are shown as red spheres, representing the alpha-carbon atom of
21 each residue. The helices involved in the formation of the dimer interface are shown in
22 yellow.
23
24
25
26
27
28
29
30
31
32

33
34 Figure 2.

35
36 *CLCN5* previously described and novel mutations. A: Exon structure of the *CLCN5* gene with
37 geometric shapes indicating relative positions of different types of mutations. B: ClC-5
38 domains. C: percentages of mutations by type.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Novel mutations detected in 47 patients with **Dent disease 1** (42 mutations)

Patient	Type of mutation	Nucleotide*	Protein	Exon/intron
Dent 8-1	Large deletion	c.1-?_105+?del (E2del)	p.Met1_Glu35del (in-frame)	2
Dent 15-1	Large deletion	c.1-?_205+?del (E2-3del)	p.Met1_Ser68del (frameshift)	2-3
Dent 26-1	Large deletion	c.1-?_393+?del (E2-4del)	p.Met1_Glu131del (in-frame)	2-4
Dent 164-1	Frameshift	c.166del	p.Ser56Profs*5	3
Dent 36-1	Frameshift	c.403_404del	p.Ala135Leufs*37	5
Dent 22-1	Frameshift	c.410dup	p.Val138Serfs*35	5
Dent 111-1	Frameshift	c.597_598dup	p.Leu200Argfs*8	6
Dent 70-1	Frameshift	c.681_682del	p.His227Glnfs*9	6
Dent 176-1	Frameshift	c.699del	p.Lys234Argfs*20	6
Dent 21-1	Frameshift	c.780del	p.Gly261Glnfs*98	7
Dent 89-1	Nonsense	c.1536del	p.Val514*	10
Dent 148-1	Frameshift	c.1571dup	p.Met525Asnfs*3	10
Dent 5-1	Frameshift	c.1574del	p.Met525Serfs*4	10
Dent 149-1	Frameshift	c.1747del	p.Met583Trpfs*3	10
Dent 9-1	Frameshift	c.1751del	p.Asp584Valfs*2	10
Dent 170-1	Frameshift	c.1893dup	p.Val632Cysfs*14	10
Dent 43-1	Frameshift	c.1962del	p.Val655Leufs*23	11
Dent 66-1	Frameshift	c.2079_2080insG	p.Thr694Aspfs*48	11
Dent 50-1				
Dent 191-1				
Dent 65-1	Nonsense	c.2184del	p.Leu729*	12
Dent 69-1	Nonsense	c.308G>A	p.Trp103*	4
Dent 173-1	Nonsense	c.309G>A	p.Trp103*	4
Dent 91-1	Nonsense	c.438G>A	p.Trp146*	5
Dent 80-1	Nonsense	c.721G>T	p.Glu241*	7
Dent 103-1	Nonsense	c.1210G>T	p.Glu404*	8
Dent 83-1	Nonsense	c.1288C>T	p.Gln430*	8
Dent 87-1	Splicing	c.105G>A (exon 2 last nucleotide)	p.Glu35Glu splicing effect ¹	2
Dent 182-1	Splicing	c.206-1G>C	Loss of splice acceptor site	
Dent 11-4	Splicing	c.517-8A>G	p.? ²	6
Dent 136-1	Splicing	c.724-1G>A	Loss of splice acceptor site	7
Dent 12-3	Splicing	c.724-2A>G	Loss of splice acceptor site	7
Dent 154-1				
Dent 113-1	Splicing	c.1348-2del	Loss of splice acceptor site	9
Dent 116-1	Splicing	c.1535-1G>T	Loss of splice acceptor site	10
Dent 44-1	Splicing	c.2150+1G>A	Loss of splice donor site	11
Dent 197-1				
Dent 139-1	In-frame	c.773delinsGGAA	p.Pro258delinsArgAsn	7
Dent 155-1	Missense	c.692A>T	p.Lys231Ile	6
Dent 105-1	Missense	c.716G>C	p.Arg239Pro	6
Dent 158-1	Missense	c.748G>C	p.Gly250Arg	7
Dent 162-1	Missense	c.801A>C	p.Glu267Asp	7
Dent 134-1	Missense	c.1408G>A	p.Gly470Arg	9
Dent 159-1				
Dent 119-1	Missense	c.1552A>G	p.Thr518Ala	10
Dent 106-1	Missense	c.1588G>A	p.Gly530Ser	10
Dent 38-1	Missense	c.810C>G	p.Ser270Arg	8

* Numbering is according to the cDNA sequence (GenBank entry NM 000084.2). The A of the ATG of the Methionine initiation codon is defined as nucleotide 1.¹Splice site scores are decreased by 36% for MaxEntScan (10.1 to 6.4) and 13.2 % for SpliceSiteFinder-like (91.6 to 79.5). ²Splice site score is decreased by 42% for MaxEntScan (11.4 to 4.8). SpliceSiteFinder-like predicts the loss of a splice site and the creation of a cryptic splice site. No mRNA was available for analysis of the transcripts for these patients.

Table 2. Recurrent mutations detected in 43 patients with **Dent disease 1** (24 mutations)

Patient	Type of mutation	cDNA	Protein	Exon/ Intron	Reference
Dent 23-1 Dent 74-1 Dent180-1	Large deletion	c.(?_-30)_(*220_?)del	Whole gene deletion	1-12	Akuta et al., 1997
Dent 18-1	Frameshift	c.2108delT	p.Phe703Serfs*27	11	Blanchard et al., 2008
Dent 133-1 Dent 46-1 Dent 59-1	Nonsense	c.82C>T	p.Arg28*	2	Hoopes et al., 1998
Dent 13-1	Nonsense	c.608 C>A	p.Ser203*	6	Blanchard et al., 2008
Dent 146-1 Dent 16-1 Dent 2-1 Dent 27-1	Nonsense	c.1039C>T	p.Arg347*	8	Akuta et al., 1997 Besbas et al., 2005
Dent 171-1	Nonsense	c.1399C>T	p.Arg467*	9	Ludwig et al., 2005 Li et al., 2009
Dent 17-1 Dent 40-1 Dent 202-1 Dent 203-1 Dent 210-1	Nonsense	c.1909C>T	p.Arg637*	10	Takemura et al., 2001
Dent 31-1 Dent 128-1 Dent 181-1	Nonsense	c.1942C>T	p.Arg648*	11	Lloyd et al., 1996 Igarashi et al., 1998
Dent 126-1 Dent196-1	Nonsense	c.2110C>T	p.Arg704*	11	Lloyd et al., 1996 Nakazato et al., 1999
Dent 53-1	Nonsense	c.2152C>T	p.Arg718*	12	Carballo-Trujillo et al., 2003 Grand et al., 2009
Dent 35-1 Dent 37-1	Splicing	c.394-2A>G	Loss of splice acceptor site	4	Zhu et al., 2010
Dent 33-1 Dent 144-1	Inframe	c.1566_1568del	p.Val523del	10	Wu F et al., 2009
Dent 56-1	Missense	c.536G>A	p.Gly179Asp	6	Grand et al., 2009
Dent 96-1	Missense	c.608C>T	p.Ser203Leu	6	Grand et al., 2009
Dent 47-1	Missense	c.635G>C	p.Gly212Ala	6	Grand et al., 2009
Dent 124-1	Missense	c.661T>C	p.Cys221Arg	6	Hoopes et al., 2004 Grand et al., 2009
Dent 71-1 Dent 97-1 Dent 20-1 Dent 25-1 Dent 62-1	Missense	c.731C>T	p.Ser244Leu	7	Lloyd et al., 1996
Dent 208-1	Missense	c.781G>A	p.Gly261Arg	7	Tosetto et al., 2009
Dent 45-1	Missense	c.1406T>C	p.Leu469Pro	9	Grand et al., 2009
Dent 195-1	Missense	c.1535G>A	p.Gly512Asp	10	Tosetto et al., 2009
Dent 179-1	Missense	c.1546C>T	p.Arg516Trp	10	Akuta et al., 1997
Dent 76-1	Missense	c.1556T>A	p.Val519Asp	10	Tosetto et al., 2009
Dent 104-1	Missense	c.1636A>G	p.Lys546Glu	10	Hoopes et al., 2004
Dent 163-1	Missense	c.1639T>G	p.Trp547Gly	10	Ramos-Trujillo et al., 2007

* Numbering is according to the cDNA sequence (GenBank entry NM 000084.2). The A of the ATG of the Methionine initiation codon is defined as nucleotide 1.

Table 3. Summary of polymorphisms and variants of unknown clinical significance (VUS)

	Nucleotide	Protein	Location	RefSNP	MAF*	Geographic origin	Reference
Polymorphisms	c.993G>C	p.Leu331Leu	Exon 8	rs34122840	EA 0.06%; AA 14.84%	Dent 132-1: France Dent 165-1: Senegal Dent 200-1: Algeria	This study
	c.1452C>T	p.Thr484Thr	Exon 9			Japan	Akuta et al., 1997
	c.1704C>T	p.Pro568Pro	Exon 10	rs34173954	EA 0.07%; AA 20.65%	North America ¹ Cape Verde Dent 010-1: Comoros Dent 117-1: France	Hoopes et al., 2004 Tosetto et al., 2006 This study
	c.1839T>C	p.Ser 613Ser	Exon 10			Korea	Cheong et al., 2005
VUS	c.1157C>T	p.Ser386Phe	Exon 8			Italy ¹	Tosetto et al., 2009
	c.-47-433G>A		5'UTR			Italy (Sardinia)	Tosetto et al., 2009
	c.106-17T>G		Intron 2	rs56117808	EA 0.7%; AA 0.05%	NA	Forino et al., 2004
	c.206-6A>T		Intron 3	rs6651602	EA 0.04%; AA 5.37%	Dent 040-1: France ¹	This study
	c.724-35T>G		Intron 6			Turkey ¹	Ludwig et al., 2006
	c.1348-14G>C		Intron 8	rs7063765	EA (C) 0.02%; AA (G) 47%	Dent 117-1: France Dent 132-1: France Dent 153-1: Guadeloupe ¹ Dent 165-1: Senegal Dent 200-1: Algeria	This study
	c.1535-53G>A		Intron 9			Dent 118-1: Vietnam	This study
	c.1934-60G>A		Intron 10	rs150713901	NA	Dent 153-1: Guadeloupe ¹	This study
	c.2151-67C>T		Intron 11	rs56041343	NA	Dent 017-1: France ¹ Dent 040-1: France ¹ Dent 110-1: France	This study
	c.2151-18A>G		Intron 11			Spain ¹	Ramos-Trujillo et al., 2013
c.*22A>G		3'UTR			Dent 100-1: France	This study	

*MAF: minor allele frequency from exome sequencing project (ESP6500SIV2); EA: European American; AA: African American.

¹These patients harbour a *CLCN5* pathogenic mutation; NA: not available

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


For Peer Review

Table 4. Phenotype of male patients with **Dent disease 1** in the literature and in our cohort

Phenotype	Large published cohorts			Global literature Analysis	This study		
	Europe and North America*	Japan (Sekine et al., 2014)	Bökenkamp et al., 2009 (include literature data)		Patients with clinical diagnosis of Dent disease (n=103)	Patients with screening after diagnosis in a relative (n=14)	Total (n=117)
Age at diagnosis (years)				9 (0.2-67)** (n=311)	7 (0.1-55)**§ (n=89)	5.5 (0.2-32)** (n=14)	7 (0.1-55)** (n=103)
LMW proteinuria	100%	100%	(212/212) 100%	(365/365) 100%	(99/99) 100%	11/12	(110/111) 99%
Hypercalciuria	95%	89%	(180/200) 90%	(287/359) 80%	(89/95) 94%	6/11	(87/99) 88%
Nephrocalcinosis	74%	76%	(137/182) 75%	(156/282) 55%#	(59/91) 65%	3/11	(62/102) 61%
Aminoaciduria	76%	-	(31/75) 41%	(45/93) 48%	(23/34) 67%	1/5	(24/39) 61%
Renal Insufficiency	64%	42%	(60/203) 30%	(92/347) 26.5%	(52/97) 53%	0/13	(52/110) 47%
Hypophosphatemia of renal origin	50%	-	(35/156) 22%	(72/200) 36%	(39/65) 60%	2/11	(41/76) 54%
Hypokalemia of renal origin	35%	-	(10/67) 15%	(22/60) 37%	(33/75) 44%	1/11	(34/86) 39%
Lithiasis	49%	-	ND	(41/235) 17%#	(27/81) 33%	3/10	(30/91) 33%
Rickets	30%	33%	ND	(53/247) 21%	(14/81) 17%	0/12	(14/93) 15%
Metabolic Acidosis	-	-	(2/68) 3%	(6/66) 9%	(9/60) 15%	0/8	(9/68) 13%
Glycosuria	54%	-	(18/108) 17%	(27/105) 26%	(27/62) 43%	1/8	(28/70) 40%


*Adapted from Scheiman SJ, in Genetic diseases of the kidney

#72 additional cases were described as nephrocalcinosis or nephrolithiasis. Including these case the value for nephrocalcinosis/nephrolithiasis is 232/354 (65.5%); **Median (min-max); §for 9 probands the precise age was not available but the proximal tubulopathy was diagnosed in childhood.



Three-dimensional model of CIC-5 based on homology with the structure of StCIC [Dutzler et al., 2002] showing the location of the CIC-5 mutations. Views from the side of the membrane with the extracellular medium at the top (A) and from a direction rotated by 90° from the A monomer (B). The view in (B) shows the interface between the two subunits of the dimeric protein. Mutated residues are shown as red spheres, representing the alpha-carbon atom of each residue. The helices involved in the formation of the dimer interface are shown in yellow.


51x17mm (300 x 300 DPI)




CLCN5 previously described and novel mutations. A: Exon structure of the CLCN5 gene with geometric shapes indicating relative positions of different types of mutations. B: CLC-5 domains. C: percentages of mutations by type.

90x51mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Supp. Figure S1. Pedigrees of four families with X-linked **Dent disease 1**: the arrow indicates the proband in each case; the asterisks indicate individuals for whom DNA sequencing analysis was performed. In families 11 and 44, father-to-daughter transmission was observed for one and three daughters, respectively.



Supp. Figure S2. Urinary Beta-2 microglobulin (left panel) and urinary calcium/creatinine ratio (right panel) values in male patients with *Dent disease 1* compared with values in female carriers. Dotted lines indicate normal upper values.

Supp. Table S1. Mutations described in the *CLCN5* gene

Type of Mutation	Nucleotide	Protein	Exon/Intron	Reference
Large deletion	c.206?_516+?del	Codons 69-172 deleted	4-5	Sekine et al., 2014
Large deletion	c.205+3_725del	p.Gly69Alafs*12	4-6	Igarashi et al., 2000
Large deletion	c.394-532_516+3delins49	Codons 132-172 deleted	5	Brakemeier et al., 2004
Large deletion	c.394?_723+?del	Codons 132-241 deleted	5-6	Lloyd et al., 1996
Large deletion	c.394?_1347+?del	Codons 132-449 deleted	5-8	Morimoto et al., 1998
Large deletion	c.2151?_2241+?del	Codons 717-746 deleted	12	Park et al., 2014
Large deletion	c.(?_30)_(*220_?)del	Whole gene deletion	1-12	Lloyd et al., 1996, Akuta et al., 1997
Promoter	c.-47-433G>A		5'UTR	Tosetto et al., 2009
Frameshift	c.16del	p.Glu6Serfs*6	2	Tosetto et al., 2006
Frameshift	c.36delG	p.Thr13Profs*	2	Zhang et al., 2014
Frameshift	c.86_88dup	p.Asp29_Arg30insHis	2	Lloyd et al., 1997
Frameshift	c.92delA	p.Asp31Valfs*17	2	Fervenza et al., 2013
Frameshift	c.118del	p.Ser40Alafs*8	3	Hoopes et al., 1998
Frameshift	c.132del	p.Trp45Glyfs*3	3	Hoopes et al., 1998
Frameshift	c.161dup	p.Ser56Phefs*42	3	Dinour et al., 2009
Frameshift	c.165_169del	p.Phe55Leufs*41	3	Sekine et al., 2014
Frameshift	c.172_173del	p.Trp58Valfs*39	3	Tosetto et al., 2006
Frameshift	c.191del	p.Ile64Metfs*7	3	Sekine et al., 2014
Frameshift	c.200dup	p.Leu67Phefs*31	3	Park et al., 2014
Frameshift	c.258del	p.Glu87Lysfs*51	4	Cox et al., 1999, Dinour et al., 2009
Frameshift	c.299dup	p.His100Glnfs*7	4	Matsuyama et al., 2004
Frameshift	c.346del	p.Cys116Valfs*22	4	Nakazato et al., 1999
Frameshift	c.445dup	p.Leu149Profs*24	5	Cho et al., 2008, Lee et al., 2009
Frameshift	c.470_471del	p.Leu157Argfs*15	5	Cox et al., 1999
Frameshift	c.485del	p.Ala162Glyfs*12	5	Park et al., 2014
Frameshift	c.564del	p.Val192Leufs*15	6	Tosetto et al., 2009
Frameshift	c.687delinsTA	p.Asn230Lysfs*7	6	Akuta et al., 1997
Frameshift	c.722_723del	p.Glu241Glyfs*26	6	Cox et al., 1999
Frameshift	c.744_745ins CAGC	p.Ala249Glnfs*20	7	Addis et al., 2013
Frameshift	c.746_752del	p.Ala249Aspfs*3	7	Sekine et al., 2014
Frameshift	c.930_931delinsA	p.Phe310Leufs*49	8	Wu et al., 2009
Frameshift	c.977del	p.Gly326Alafs*33	8	Ramos-Trujillo et al., 2007
Frameshift	Not described	p.Val363fs	8	Cramer et al., 2014
Frameshift	c.1126delT	p.Tyr376Thrfs*4	8	Park et al., 2014
Frameshift	c.1157C>T/1160_1161dup	p.Ser386Phe/p.Leu388Serfs*47	8	Tosetto et al., 2009
Frameshift	c.1175_1176del	p.Cys392Trpfs*10	8	Yamamoto et al., 2000
Frameshift	c.1182del	p.Leu395Trpfs*39	8	Yamamoto et al., 2000
Frameshift	c.1208del	p.Tyr403Leufs*31	8	Nakazato et al., 1997
Frameshift	c.1245del	p.Pro416Leufs*18	8	Dinour et al., 2009
Frameshift	c.1248delT	p.P416fsX*17	8	Ramos-Trujillo et al., 2013
Frameshift	c.1331del	p.Phe444Serfs*5	8	Tosetto et al., 2006
Frameshift	c.1526del	p.Ala509Alafs*3	9	Sekine et al., 2014
Frameshift	c.1537del	p.Gly513Glyfs*2	10	Sekine et al., 2014
Frameshift	c.1540dup	p.Val514Glyfs*14	10	Nakazato et al., 1999
Frameshift	c.1550dup	p.Met517Ilefs*11 ^a	10	Ludwig et al., 2005 ^a
Frameshift	c.1558dup	p.Ser520Phefs*8	10	Nakazato et al., 1997, Park et al., 2014
Frameshift	c.1562_1563del	p.Leu521Argfs*6 ^{a,c}	10	Ludwig et al., 2005 ^{a,c}
Frameshift	c.1668del	p.Gly556Glyfs*30	10	Sekine et al., 2014
Frameshift	c.1765dup	p.Arg589Profs*4	10	Tosetto et al., 2009
Frameshift	c.1782del	p.Leu594Phefs*2	10	Li et al., 2009, Zhu et al., 2010
Frameshift	c.1974del	p.Ile659Serfs*19	11	Yamamoto et al., 2000
Frameshift	c.2061del	p.Phe688Serfs*3	11	Hoopes et al., 2004
Frameshift	c.2079dup	p.Thr694Tyrfs*48	11	Hoopes et al., 2004
Frameshift	c.2080dup	p.Thr694Asnfs*48	11	Park et al., 2014
Frameshift	c.2081_2082insC	p.Thr694Thrfs*48	11	Sekine et al., 2014

Frameshift	c.2108del	p.Phe703Serfs*27	11	Blanchard et al., 2008
Frameshift	c.2142dup	p.His715Thrfs*27	11	Cho et al., 2008
Frameshift	c.2085del	p.Met696Trpfs*4	11	Lloyd et al., 1997
Frameshift	c.2179del	p.Asp727Metfs*3	12	Frishberg et al., 2009
Frameshift	c.2183_2205del	p.Val728Glyfs*6 ^d	12	Yamamoto et al., 2000 ^d
Complex	c.[320A>C,324_330del]	p.[p.His107Pro,p.Thr109Lysfs*27]	4	Ramos-Trujillo et al., 2013
Nonsense	c.82C>T	p.Arg28*	2	Hoopes et al., 1998, Matsuyama et al., 2004, Sekine et al., 2004, Dinour et al., 2009, Cramer et al., 2014
Nonsense	c.100C>T	p.Arg34*	2	Langlois et al., 1998, Hoopes et al., 1998, Cox et al., 1999, Ludwig et al., 2006, Tosetto et al., 2006, Sekine et al., 2014, Ramos-Trujillo et al., 2013
Nonsense	c.134G>A	p.Trp45*	3	Hoopes et al., 2004, Ludwig et al., 2005
Nonsense	c.277G>T	p.Gly93*	4	Sekine et al., 2014
Nonsense	c.285G>A	p.Trp95*	4	Ludwig et al., 2006, Cho et al., 2008, Park et al., 2014
Nonsense	c.352G>T	p.Glu118*	4	Morimoto et al., 1998
Nonsense	c.366G>A	p.Trp122*	4	Frishberg et al., 2009
Nonsense	c.370C>T	p.Gln124*	4	Sekine et al., 2014
Nonsense	c.420T>G	p.Tyr140*	5	Ramos-Trujillo et al., 2007
Nonsense	c.492T>G	p.Tyr164*	5	Ramos-Trujillo et al., 2013
Nonsense	c.566G>A	p.Trp189*	6	Sekine et al., 2014
Nonsense	c.608C>A	p.Ser203*	6	Blanchard et al., 2008
Nonsense	c.663T>A	p.Cys221*	6	Hoopes et al., 2004
Nonsense	c.837G>A	p.Trp279* ^d	8	Lloyd et al., 1996 ^d , Akuta et al., 1997, Sekine et al., 2014
Nonsense	c.942G>A	p.Trp314*	8	Ramos-Trujillo et al., 2007
Nonsense	c.996G>A	p.Trp332*	8	Ludwig et al., 2005, Ludwig et al., 2006, Tosetto et al., 2006
Nonsense	c.1028G>A	p.Trp343*	8	Lloyd et al., 1997, Hoopes et al., 1998
Nonsense	c.1039C>T	p.Arg347* ^a	8	Akuta et al., 1997, Morimoto 1998, Hoopes et al., 2004, Ludwig et al., 2005 ^a , Besbas et al., 2005, Tosetto et al., 2006, Wu et al., 2009, Sekine et al., 2014
Nonsense	c.1298T>G	p.Leu433*	8	Nakazato et al., 1997, Carballo-Trujillo et al., 2003
Nonsense	c.1399C>T	p.Arg467*	9	Ludwig et al., 2005, Tosetto et al., 2006, Ramos-Trujillo et al., 2007, Dinour et al., 2009, Li et al., 2009, Zhu et al., 2010
Nonsense	c.1467G>A	p.Trp489*	9	Sekine et al., 2014
Nonsense	c.1506T>A	p.Tyr502*	9	Ramos-Trujillo et al., 2007
Nonsense	c.1641G>A	p.Trp547*	10	Tosetto et al., 2006
Nonsense	c.1701C>(A/G?)	p.Tyr567*	10	Lee et al., 2009,

				Okamoto et al., 2012
Nonsense	c.1798C>T	p.Gln600*	10	Tosetto et al., 2006
Nonsense	c.1825G>T	p.Glu609*	10	Cheong et al., 2005, Cho et al., 2008
Nonsense	c.1851C>A	p.Tyr617* ^c	10	Yamamoto et al., 2000, Mo et al. 2004 ^c
Nonsense	c.1885C>T	p.Gln629*	10	Sekine et al., 2014
Nonsense	c.1909C>T	p.Arg637*	10	Takemura et al., 2001, Cheong et al., 2005, Ludwig et al., 2005, Tosetto et al., 2006, Cho et al., 2008, Dinour et al., 2009, Li et al., 2009, Lee et al., 2009, Zhu et al., 2010, Sekine et al., 2014
Nonsense	c.1942C>T	p.Arg648* ^c	11	Lloyd et al., 1996 ^c and 1997, Igarashi et al., 1998, Bosio et al., 1999, Cox et al., 1999, Tosetto et al., 2006, Blanchard et al., 2008, Frishberg et al., 2009, Sekine et al., 2014, Mo et al., 2004 ^c
Nonsense	c.2110C>T	p.Arg704* ^c	11	Lloyd et al., 1996 ^c , Langlois et al., 1998, Igarashi et al., 1998, Nakazato et al., 1999, Cox et al., 1999, Mo et al., 2004 ^c , Ramos-Trujillo et al., 2007, Ramos-Trujillo et al., 2013, Sekine et al., 2014, Cramer et al., 2014
Nonsense	c.2152C>T	p.Arg718* ^a	12	Carballo-Trujillo et al., 2003, Hoopes et al., 2004, Wu et al., 2009, Grand et al., 2009 ^a
Nonsense	c.2186T>G	p.Leu729*	12	Sethi et al., 2009
Splicing	c.105+2T>C	skipping of exon 3, frameshift	intron 2	Wu et al., 2009
Splicing	c.106-2A>G	skipping of exon 3, frameshift	intron 2	Ramos-Trujillo et al., 2007
Splicing	c.205+2T>C	skipping of exon 3, frameshift	intron 3	Tosetto et al., 2006, Ji et al., 2014
Splicing	c.206-1G>A	p.?	intron 3	Tosetto et al., 2006, Sekine et al., 2014
Splicing	c.393+1G>A	p.?	intron 4	Sekine et al., 2014
Splicing	c.393+4A>G	p.?	intron 4	Tosetto et al., 2009
Splicing	c.394-2A>C	p.?	intron 4	Sekine et al., 2014
Splicing	c.394-2A>G	p.?	intron 4	Zhu et al., 2010, Sekine et al., 2014
Splicing	c.516+5G>T	skipping of exon 5	intron 5	Tosetto et al., 2006
Splicing	c.516+2T>G	p.?	intron 5	Lloyd et al., 1996
Splicing	c.516+1G>A	p.?	intron 5	Lloyd et al., 1996, Sekine et al., 2014
Splicing	c.517-2A>G	skipping of exon 6	intron 5	Igarashi et al., 2000
Splicing	c.517-3C>A	skipping of exon 6	intron 5	Cox et al., 1999
Splicing	c.723+1G>T	p.?	6	Ramos-Trujillo et al., 2013
Splicing	c.723+1G>A	p.?	6	Cramer et al., 2014
Splicing	c.1347+1G>T	loss of 290 bp of exon 8, frameshift	intron 8	Tosetto et al., 2009, Sekine et al., 2014
Splicing	c.1534+1G>T	p.?	9	Park et al., 2014
Splicing	c.1535-1G>A	p.?	9	Cramer et al., 2014
Splicing	c.1535-2A>C	p.?	9	Park et al., 2014

Splicing	c.1933+2_1933+3insTGGT	p.?	Intron 10	Sekine et al., 2014
Splicing	c.2150+1G>T	skipping of exon 11, frameshift	intron 11	Tosetto et al., 2009
Inframe	c.782_784	p.Gly261del	7	Tosetto et al., 2009
Inframe	c.800_802del	p.Glu268del	7	Tosetto et al., 2009, Sekine et al., 2014
Inframe	c.830_832dup	p.Thr277_Leu278insSer	8	Tosetto et al., 2009
Inframe	c.1566_1568del	p.V523del ^d	10	Wu et al., 2009 ^d , Ramos-Trujillo et al., 2013, Sekine et al., 2014, Park et al., 2014
Missense	c.64T>G	p.Trp22Gly ^c	2	Morimoto et al., 1998 ^c , Ramos-Trujillo et al., 2013
Missense	c.170G>T	p.Gly57Val ^d	3	Lloyd et al., 1997 ^d , Smith et al., 2009 ^d
Missense	c.173G>T	p.Trp58Leu	3	Tosetto et al., 2009
Missense	c.174G>C	p.Trp58Cys	3	Sethi et al., 2009
Missense	c.193G>A	p.Gly65Arg	3	Ramos-Trujillo et al., 2013
Missense	c.263G>A	p.Gly88Asp	4	Ludwig et al., 2006
Missense	c.263G>T	p.Gly88Val	4	Sekine et al., 2014
Missense	c.270C>G	p.Cys90Trp	4	Sekine et al., 2014
Missense	c.299A>G	p.His100Arg	4	Park et al., 2014
Missense	c.302G>A	p.Cys101Tyr	4	Cho et al., 2008, Lee et al., 2009
Missense	c.307T>C	p.Trp103Arg	4	Sekine et al., 2014
Missense	c.527T>A	p.Ile176Asn	6	Sekine et al., 2014
Missense	c.536G>A	p.Gly179Asp ^a	6	Grand T et al., 2009 ^a
Missense	c.599T>G	p.Leu200Arg ^a	6	Lloyd et al., 1996 ^a , Grand et al., 2009 ^a
Missense	c.608C>T	p.Ser203Leu ^a	6	Grand et al., 2009 ^a
				Sekine et al., 2014, Park et al., 2014
Missense	c.631G>C	p.Glu211Gln	6	Sekine et al., 2014
Missense	c.634G>A	p.Gly212Ser	6	Sekine et al., 2014
Missense	c.635G>C	p.Gly212Ala ^c	6	Grand et al., 2009 ^c
Missense	c.638C>T	p.Pro213Leu	6	Sekine et al., 2014
Missense	c.655T>C	p.Cys219Arg ^a	6	Ramos-Trujillo et al., 2007, Grand et al., 2009 ^a
Missense	c.661T>C	p.Cys221Arg ^a	6	Hoopes et al., 2004, Ludwig et al., 2005 ^a , D'Antonio et al., 2013 ^a
Missense	c.674T>C	p.Leu225Pro ^a	6	Ramos-Trujillo et al., 2007, Grand et al., 2011 ^a
Missense	c.731C>T	p.Ser244Leu ^c	7	Lloyd et al., 1996 ^c , Oudet et al., 1997, Kelleher et al., 1998, Hoopes et al., 1998 and 2004, Tosetto et al., 2006, Ludwig et al., 2006, Anglani et al., 2006, Blanchard et al., 2008, Wu et al., 2009, Zhu et al., 2010, Samardzic et al., 2011, Sekine et al., 2014, Grand et al., 2011 ^c
Missense	c.779G>T	p.Gly260Val ^b	7	Tosetto et al., 2006, Anglani et al., 2006, Grand et al., 2011 ^b
Missense	c.781G>A	p.Gly261Arg	7	Tosetto et al., 2009
Missense	c.782G>A	p.Gly261Glu	7	Bogdanović et al., 2010
Missense	c.796C>G	p.Leu266Val	7	Sekine et al., 2014
Missense	c.800A>C	p.Glu267Ala	7	Hoopes et al., 2004
Missense	c.808A>C	p.Ser270Arg ^a	8	Igarashi T et al., 1998 ^a ,

				Smith et al., 2009 ^a
Missense	c.808A>G	p.Ser270Gly	8	Hoopes et al., 2004
Missense	c.814T>A	p.Tyr272Asn	8	Sekine et al., 2014
Missense	c.815A>G	p.Tyr272Cys ^c	8	Tosetto et al., 2006, Sekine et al., 2014, Grand et al., 2011 ^c
Missense	c.817T>C	p.Phe273Leu	8	Ramos-Trujillo E et al., 2007
Missense	c.834G>C	p.Leu278Phe ^b	8	Igarashi T et al., 1998, Sekine et al., 2014, Grand et al., 2011 ^b
Missense	c.839G>C	p.Arg280Pro ^d	8	Lloyd et al., 1997 ^d
Missense	c.971T>G	p.Leu324Arg ^a	8	Ludwig et al., 2005 ^a
Missense	c.997G>A	p.Gly333Arg ^a	8	Tanuma et al., 2007 ^a
Missense	c.1020C>G	p.Asn340Lys ^a	8	Tosetto et al., 2006, Grand et al., 2011 ^a
Missense	c.1384G>A	p.Gly462Ser	9	Ramos-Trujillo et al., 2013
Missense	c.1385G>A	p.Gly462Asp	9	Hoopes et al., 2004
Missense	c.1385G>T	p.Gly462Val ^a	9	Ludwig et al., 2005 ^a
Missense	c.1396G>C	p.Gly466Arg	9	Valina et al., 2012
Missense	c.1397G>A	p.Gly466Asp	9	Ramos-Trujillo et al., 2013
Missense	c.1403T>C	p.Leu468Pro	9	Sekine et al., 2014
Missense	c.1406T>C	p.Leu469Pro ^a	9	Grand et al., 2009 ^a , Lee et al., 2009
Missense	c.1505A>G	p.Tyr502Cys	9	Sekine et al., 2014
Missense	c.1511T>A	p.Met504Lys	9	Sethi et al., 2009
Missense	c.1514T>G	p.Val505Gly	9	Zhu et al., 2010
Missense	c.1516G>A	p.Gly506Arg	9	Matsuyama et al., 2004, Sekine et al., 2014
Missense	c.1517G>A	p.Gly506Glu ^a	9	Lloyd et al., 1996 ^a , Scheinman et al., 2000
Missense	c.1534G>C	p.Gly512Arg ^d	9	Lloyd et al., 1997 ^d
Missense	c.1535G>A	p.Gly512Asp	10	Tosetto et al., 2009
Missense	c.1537G>A	p.Gly513Arg ^a	10	Hoopes et al., 2004, Sekine et al., 2014, Grand et al., 2011 ^a
Missense	c.1538G>A	p.Gly513Glu ^a	10	Akuta et al., 1997, Smith et al., 2009 ^a
Missense	c.1546C>T	p.Arg516Trp ^a	10	Akuta et al., 1997, Hoopes et al., 2004, Ludwig et al., 2005 ^a , Ludwig et al., 2006, Wu et al., 2009, Sekine et al., 2014, Park et al., 2014, Smith et al., 2009 ^a
Missense	c.1547G>A	p.Arg516Gln	10	Sekine et al., 2014
Missense	c.1556T>A	p.Val519Asp	10	Tosetto et al., 2009
Missense	c.1558T>C	p.Ser520Pro ^c	10	Lloyd et al., 1996 ^c
Missense	c.1561C>T	p.Leu521Phe	10	Cramer et al., 2014
Missense	c.1566_1568del	p.Val522del	10	Sekine et al., 2014
Missense	c.1571T>A	p.Ile524Lys ^a	10	Takemura et al., 2001, Yanagida et al., 2004, Sekine et al., 2014, Smith et al., 2009 ^a
Missense	c.1581A>T	p.Glu527Asp ^b	10	Lloyd et al., 1997 ^b , Smith et al., 2009 ^b
Missense	c.1634G>A	p.Ser545Asn	10	Hoopes et al., 2004, Cho et al., 2008, Lee et al., 2009
Missense	c.1636A>G	p.Lys546Glu ^b	10	Hoopes et al., 2004, Grand et al., 2011 ^b
Missense/splicing	c.1639T>G	p.Trp547Gly ^b (skipping of exons 10 and 11)	10	Ramos-Trujillo et al., 2007, Grand et al., 2011 ^b

Missense	c.1639T>C	p.Trp547Arg	10	Tosetto et al., 2009
Missense/splicing	c.1802A>T	p.Asp601Val ; (loss of 133 bp of exon 10), p.Asp601Lysfs*33	10	Yamamoto et al., 2000
Missense	c.1862C>T	p.Pro621Leu	10	Tosetto et al., 2009
Missense	c.2108T>C	p.Phe703Ser	11	Sekine et al., 2014
Missense	c.2117T>C	p.Leu706Pro	11	Sekine et al., 2014
Missense	c.2133C>G	p.Cys711Trp	11	Sekine et al., 2014
Missense	c.2173A>G	p.Lys725Glu	12	Ludwig et al., 2006, Sekine et al., 2014
	c.1948insAlu(345pb)		11	Claverie Martin et al., 2003 and 2005

* Numbering is according to the cDNA sequence (GenBank entry NM 000084.2). The A of the ATG of the Methionine initiation codon is defined as nucleotide 1. Superscript letters indicate mutations in vitro expressed, their reference and classification as follow: ^aclass 1 mutations inducing defective protein processing; ^bclass 2 mutations inducing delayed protein processing and lower stability of the mature protein; ^cclass 3 mutations inducing altered conduction without any change in subcellular distribution; ^dmutations needing further investigation to allow its classification; ^functionally tested as L521R

For Peer Review

Supp.Table S2. *In silico* pathogenicity predictions for new in-frame and missense mutations

Nucleotide change	Amino acid change	Exon	PolyPhen-2	Sift	MutationTaster	SNPs&Go	MutPred
c.692A>T	p.Lys231Ile	6	Probably damaging	Deleterious Score 0.01	Disease causing p=1	Disease ri=8	g=0.689; loss of methylation (p=0.0196); gain of helix (p=0.0496)
c.716G>C	p.Arg239Pro	6	Benign	Deleterious Score 0.02	Disease causing p=0.99	Disease ri=9	g=0.762
c.748G>C	p.Gly250Arg	7	Probably damaging p=0.993	Deleterious Score 0	Disease causing p=1	Disease ri=9	g=0.894; gain on methylation (p=0.0078)
c.801A>C	p.Glu267Asp	7	Probably damaging p=1	Deleterious Score 0	Disease causing p=1	Disease ri=9	g=0.974; loss of catalytic residue (p=0.0138); gain of methylation
c.1408G>C	p.Gly470Arg	9	Probably damaging p=1	Deleterious Score 0.01	Disease causing p=0.99	Disease ri=9	g=0.914
c.1552A>G	p.Thr518Ala	10	Probably damaging p=1	Deleterious Score 0	Disease causing p=1	Disease ri=6	g=0.854
c.1588G>A	p.Gly530Ser	10	Probably damaging p=1	Deleterious Score 0	Disease causing p=1	Disease ri=7	g=0.768
c.773delinsGGAA	p.Pro258delinsArgAsn	7	NA	NA	Disease causing	NA	NA

For SIFT: a change is predicted to be deleterious if the score <0.05

For MutationTaster: a change is predicted to be disease causing if the score p>0.5,

For SNPs&GO: a change is predicted to be disease causing if the reliability index (ri)>5,

For MutPred: a change is predicted to be deleterious if the general score g>0.75; in addition this tool gives scores (p) for 5 structural and functional properties. Combinations of high values of general scores and low values of property scores are referred to as hypotheses: scores with g>0.5 and p<0.05 are referred to as actionable hypotheses; scores with g>0.75 and p<0.05 are referred to as confident hypotheses and scores with g>0.75 and p<0.01 are referred to as very confident hypotheses.

Supp. Table S3. Genotype of 134 relatives analysed in 43 families.

Relative	Genotype	Number of generations (number of families)				Total
		1 (n=3)	2 (n=31)	3 (n=7)**	4 (n=2)***	
Mother	Carrier	-	29	6	2	37
	Wt	-	2	0	0	2
Grandmother	Carrier	-	-	3	1	4
	Wt	-	-	2	0	2
Daughter of affected male	Carrier	-	0	1	4	5
Other female	Carrier	0	8	9	5	22
	Wt	2	5	0	5	12
Other males*	Hemizygous	2	12	4	9	28
	Wt	0	9	1	2	13
Father	Wt		7		1	8
Maternal grandfather	Hemizygous	-	-	0	0	0
	Wt	-	-	1	0	1
Total						134

* Sons of carrier mothers or brothers in sporadic cases (probands are not included).

** In two families, the third generation is the generation after that of the proband.

*** The fourth generation is one or two generations after that of the proband.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Supp. Table S4. Phenotype of patients clinically asymptomatic before the genetic diagnosis

Patient	2-4	20-2	20-4	31-2	33-3	33-4	37-2	44-13	45-2	66-6	66-8	66-9	134-2	136-5
Age at diagnosis (years)	4.0	8.0	2.0	16.0	3.0	3.5	23.0	0.5	7.0	19.0	32.0	1.3	8.0	0.2
LMW proteinuria	+	+	+	+	+	+	NA	+	+	+	NA	-	+	+
Hypercalciuria	+	-	+	+	+	+	+	-	-	-	NA	NA	NA	-
Nephrocalcinosis	-	NA	+	-	+	-	+	-	-	NA	-	-	NA	-
Aminoaciduria	NA	+	NA	NA	-	-	NA	-	-	NA	NA	NA	NA	NA
Renal Insufficiency	-	-	-	-	-	-	-	-	-	-	-	-	NA	-
Hypophosphatemia of renal origin	-	-	-	-	+	-	NA	-	+	-	-	-	NA	NA
Hypokalaemia of renal origin	-	-	-	-	+	-	NA	-	-	-	-	-	NA	NA
Lithiasis	+	-	-	-	NA	-	NA	+	-	NA	+	-	NA	-
Rickets	-	-	-	-	-	-	NA	-	-	-	-	-	NA	-
Metabolic Acidosis	-	-	-	-	-	NA	NA	-	NA	NA	NA	-	NA	-
Glycosuria	-	-	NA	NA	NA	-	NA	-	-	-	NA	-	NA	+

+, present; -, absent; NA, not available

Peer Review

Supp. Table S5. Geographical Distribution

Continent	Country	Described families Number	This paper Number	Total	Percentage
Asia	Japan	96		96	37.46%
	Korea	32		32	
	China	11		11	
	Israel	11		11	
	India	4		4	
Europe	Italy	44	2	46	50.60%
	Spain	25		25	
	Germany	19		19	
	North Europe	13		13	
	United Kingdom	10		10	
	France	9	65	74	
	Turkey	5		5	
	Finland	3		3	
	Austria	2		2	
	Belgium	1	2	3	
	Montenegro	1		1	
	Portugal	1		1	
	Serbia	1	1	2	
	Switzerland	-	3	3	
Ukraine	-	1	1		
North America	United States of America	33	1	34	10.21%
	North America	7		7	
	Canada	1		1	
South America	Bolivia	1		1	0.73%
	Uruguay	1		1	
	Argentina	-	1	1	
North Africa		1	3	4	0.97%
Total		332	79	411	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Supp. Table S6. Data of renal biopsy

Reference	Wrong et al., 1994 Lloyd et al., 1996			Langlois et al., 1998		Hoopes et al., 1998		Igarashi et al., 1998					Takemura et al., 2001		Yanagida et al., 2004	Brakemeier et al., 2004	Cheong et al., 2005	Anglani et al., 2006	Lim et al., 2007	Copelovitch et al., 2007		Sheffer Babila et al., 2008		
Age at biopsy (years)	63*	62	13	5	6.5	4	2	14	6	8	9	13	13	16	2	9	9	15	3	12	9	8	11.5	
Minimal alteration in glomeruli and interstitium													+		+				+					
Periglomerular fibrosis	+	+	+	+	+																+			
Interstitial fibrosis	+	+	+		+	+	+											+		+	+			
tubular atrophy		+	+		+	+	+				+	+				+					+	+		
tubular or interstitial calcification		+	+	+	+			+	+	+	+			+			+				+			
Sclerosed glomeruli				+	⁺ 10/50	+	+			+	+	+				+					+ 9/37	+ 2/6	⁺ 3/18	
Glomerular hyalinosis	+	+	+													+		+			+			
Mesangial proliferation														+		+					C1q (++)			
FGGS ²																					+	+	+	+
EM				N	N														N	EFP	EFP	EFP		

Reference	Frishberg et al., 2009			Li et al., 2009	Valina et al., 2012	Okamoto et al., 2011	Ramos-Trujillo et al., 2013		Fervenza et al., 2013	Cramer et al., 2014					
	8	7	9	11	7	14	12		18	3	6	7	9	6	4
Age at biopsy (years)															
Minimal alteration in glomeruli and interstitium									+						
Periglomerular fibrosis					+										+
Interstitial fibrosis			+	+	+	+			+			+			+
tubular atrophy			+		+										
tubular or interstitial calcification									+						
Sclerosed glomeruli	+	+	+		+	+	+	+	+	4/21	+				+
Glomerular hyalinosis															
Mesangial proliferation	+										+				+
FGGS ²	+	+	+		+				+	+					+
EM		EFP									10% EFP	30-40% EFP	10% EFP	Minimal EFP	

*Necropsy; FGGS =: focal global glomerulosclerosis; EM: electron microscopy; N: normal; EFP: effacement of the foot processes