



HAL
open science

Modeling indirect N₂O emissions along the N cascade from cropland soils to rivers

Gilles Billen, Josette Garnier, Agnès Grossel, Vincent Thieu, Sylvain Théry,
Catherine Hénault

► **To cite this version:**

Gilles Billen, Josette Garnier, Agnès Grossel, Vincent Thieu, Sylvain Théry, et al.. Modeling indirect N₂O emissions along the N cascade from cropland soils to rivers. *Biogeochemistry*, 2020, 148 (2), pp.207-221. 10.1007/s10533-020-00654-x . hal-02553360

HAL Id: hal-02553360

<https://hal.sorbonne-universite.fr/hal-02553360v1>

Submitted on 24 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Modeling indirect N₂O emissions along the N cascade from cropland soils to rivers

Gilles Billen · Josette Garnier · Agnès Grossel · Vincent Thieu ·
Sylvain Théry · Catherine Hénault

Received: 16 October 2019 / Accepted: 6 March 2020 / Published online: 20 March 2020
© The Author(s) 2020

Abstract The frequently observed discrepancy between estimations of N₂O emissions at regional or global scale based either on field data or inventories (bottom-up) or on direct atmospheric observations (top-down) suggests that riparian areas and river surfaces play a significant role as hot spots of emission. We developed a modeling procedure to assess N₂O emissions occurring during the transfer of water masses from the subroot water pool of the watershed to the outlet of the river drainage network, including their passage through riparian wetlands. The model was applied to three river basins of increasing size located in the sedimentary geological area of the Paris basin (France) and validated by its capability to predict river N₂O concentrations and fluxes across the

river–atmosphere interface. At the scale of the Seine watershed, indirect emissions, i.e. emissions linked to agricultural practices but occurring elsewhere than directly at the field plot, are estimated to represent approximately 20% of the direct emissions from the watershed soils, in good agreement with previous estimates based on empirical accounting approaches. Denitrification in riparian zones is responsible for the largest share of these indirect emissions. The model results are very sensitive to the value of the ratio of N₂O versus (N₂ + N₂O), in the final products of denitrification in rivers and wetlands. By calibration on river N₂O concentrations, a value of 0.015 ± 0.05 is proposed for this ratio, in agreement with recent studies. This represents the main uncertainty factor of the model. In basins with conditions prone to increasing the value of this ratio, higher proportions of indirect N₂O emissions might possibly be observed.

Responsible Editor: Adam Langley

G. Billen (✉) · J. Garnier · V. Thieu
SU CNRS EPHE, Metis UMR 7619, 4 Place Jussieu,
BP 105, 75005 Paris, France
e-mail: gilles.billen@upmc.fr

A. Grossel
INRA, UR SOLS, 45075 Orléans, France

S. Théry
SU CNRS, FIRE FR3020, 4 Place Jussieu, BP 105,
75005 Paris, France

C. Hénault
INRA, UMR Agroécologie, 17 rue Sully, BP 86510,
21065 Dijon, France

Keywords Denitrification · Greenhouse gas indirect emissions · Seine river · Loir river · Riparian wetlands

Introduction

Direct N₂O emission from agricultural land is a major anthropogenic source of greenhouse gas affecting climate change (Myhre et al. 2013), as well as posing a threat to the stratospheric ozone layer (Ravishankara et al. 2009). The Intergovernmental Panel on Climate

Change (IPCC 2006) guidelines based on data compiled by Bouwman (1996) and Bouwman et al. (2002) recommend estimating these direct emissions at approximately 1.25% of applied synthetic fertilizers and manure. However, lower emission factors (EF) have been reported under various regional contexts from temperate (Gabrielle et al. 2006) to Mediterranean climates (Cayuela et al. 2016). On the other hand, there is evidence that emission response to increasing N inputs is exponential rather than linear (Shcherbak et al. 2014). Thus, quantifying direct N₂O emission from agriculture remains uncertain at both regional and global scales.

In addition to these direct agricultural emissions, “indirect” processes, occurring far away from agricultural fields, also contribute, but their estimation [roughly 0.75% of leached nitrogen according to IPCC (2006) guidelines, revised to 1.1% in the IPCC (2019) report] is still a subject of controversy. Using a top-down approach based on the observed N₂O increase in atmospheric concentration (rather than the bottom-up approach extrapolating from land-based data as recommended by the IPCC), Crutzen et al. (2008) estimated that 3–5% of anthropogenically produced reactive nitrogen at the global scale is actually emitted as N₂O, implying a much higher overall emission factor compared with the IPCC factor.

Other authors using top-down approaches at regional scales based on tall tower trace gas measurements (Corazza et al. 2011; Thomson et al. 2011; Griffis et al. 2013) came to the same conclusion. Later on, Davidson (2009) and Reay et al. (2012) tried to reconcile the top-down and bottom-up methodologies by considering that N₂O emitted by recycled nitrogen, such as manure nitrogen, as well as emissions from nitrate-contaminated semi-natural environments, contributes more to global N₂O emissions than agricultural soils themselves. Syakila and Kroeze (2011), revisiting the global nitrous oxide budget, confirmed that the observed increase in atmospheric N₂O cannot be explained by inventories based on the current IPCC guidelines, highlighting the significance of poorly constrained indirect anthropogenic emissions linked to the nitrogen cascade.

There is in fact no comprehensive and formalized vision of the cascade of processes responsible for indirect N₂O emissions. Systematic measurements at the landscape scale show the existence of N₂O emission hot spots in wetlands and riparian areas, in

places where nitrate-enriched groundwater comes into contact with the biogeochemically active upper soil layers (Clement et al. 2002; Oehler et al. 2007; Vilain et al. 2010, 2012; Anderson et al. 2014). However, given the limited areal extent of riparian wetlands, their contribution to agricultural N₂O emission, though significant, was lower than direct emissions.

Rivers and streams appear as other hotspots of N₂O emissions to the atmosphere. Garnier et al. (2009) and Outram et al. (2012) showed that surface waters in agricultural drainage networks are systematically oversaturated with respect to atmospheric concentration, particularly in small stream orders, showing that rivers in general and headwaters in particular are significant N₂O emitters. Interestingly, domestic wastewater-impacted rivers are also major emitters (Garnier et al. 2007, 2009; Tallec et al. 2006). Beaulieu et al. (2011) directly measured N₂ and N₂O emission using whole-stream ¹⁵NO₃⁻ tracer experiments in 72 headwater streams draining different land-use types across the United States. Extrapolating their results, they concluded that river network emissions can represent about 10% of global anthropogenic N₂O emissions. Turner et al. (2015) reported measurements of N₂O fluxes across the water-atmosphere interface of streams of different Strahler order in the Mississippi basin and suggested that taking into account headwater streams (including episodic zero-order microflow stream channels or tile drain collectors) can more than double the agricultural N₂O budget compared with bottom-up estimates based on IPCC guidelines.

River emission results from two processes: (i) degassing of N₂O produced in soils and aquifers once groundwater reaches surface water, and (ii) in-stream production of N₂O in oxygen-depleted sectors either in the water column or in the benthos, through both nitrification and denitrification (Mulholland et al. 2008). In contrast to the former, the latter processes are particularly significant in large river stretches and estuaries, as reported by Barnes and Owens (1999), Kroeze et al. (2005), Garnier et al. (2006, 2007, 2009), and Rodrigues et al. (2007).

Here we develop a comprehensive modeling framework embracing the whole N cascade from cropland to river waters in order to relate direct N₂O emission from agricultural soils and the indirect processes listed above. Several mechanistic models have been developed to calculate N₂O emissions from agricultural soil profiles, such as DNDC (Li et al.

1992; Beheydt et al. 2007), DayCENT (Parton et al. 1996, 2001; Del Grosso et al. 2005), NOE (Hénault et al. 2005), and STICS (Bessou et al. 2010). These models perform well for estimating emissions linked to given pedo-climatic context and agricultural practices, but their one-dimensional (vertical) character prevents them from assessing the landscape interactions responsible for indirect emissions. A number of models were developed to simulate nitrogen cycling at the landscape scale (see the review by Cellier et al. 2011): these tools are able to calculate nitrate transfers and associated N₂O emissions through the different compartments of a small terrestrial landscape (typically 1–10 km²) at very fine spatial and temporal resolution, but they require a comprehensive and detailed data set on topography, hydrology, soil conditions, and farming practices.

Our approach requires far fewer data; it is closely connected to our modeling approach of nitrate elimination at the river–watershed interface (Billen et al. 2018). It is applied here to three temperate watersheds of increasing size, characterized by intensive agriculture: the Orgeval, the Haut Loir, and the Seine. The purpose of this work is to assess, using a coherent modeling procedure, the share and location of the different processes responsible for indirect N₂O emissions in the land- and waterscape of agricultural basins of different sizes.

Modeled watersheds

The Orgeval is a small catchment (106 km²) in the Brie Laitière agricultural region located approximately 70 km east of Paris, nested in the Seine basin. Agricultural land, mainly devoted to cereal cropping, occupies 82% of the total area, while forests represent 17%, hence with very few grasslands and livestock left. Active riparian wetlands, defined as potential wetlands covered by grasslands or forests, represent 3.2% of the watershed. In all, 90% of the agricultural area is equipped with tile drains. The geological, hydrological, and land use features are described in detail by Garnier et al. (2014, 2016). Direct N₂O emissions from the main cropping systems have been measured with manual and automatic chambers by Vilain et al. (2010, 2011, 2012) and Benoit et al. (2014, 2015a, b) for 10 years. Annual mean values integrated over crop rotation cycles ranged between

0.65 and 0.9 kgN/ha/yr. N₂O concentrations in groundwater (piezometers) and surface water have been regularly measured in parallel (Vilain et al. 2011; Garnier et al. 2007, 2009). The basin is part of the OZCAR network of experimental watersheds (<https://ozcar-ri.prod.lamp.cnrs.fr/oracle/>).

The Haut Loir basin (3600 km², upstream from Chateaudun) drains two distinct agricultural regions with contrasting geological and pedological substrates. The western part of the basin is representative of the *Perche region*, with clayey soils, heavily tile-drained, and a dense tributary network, while the eastern part is representative of the Beauce region, with a very low river drainage density mainly fed by a deep groundwater system. Direct N₂O emissions were regularly measured with manual chambers at two locations. Annual mean emission was approximately 1–2 kgN/ha/yr. The main crop rotations in the whole watershed and their N budget were recorded (see Billen et al. 2018).

N₂O concentration in surface water, as well as emission fluxes at the water–atmosphere interface, were measured throughout the 2017–2018 period by Grosselet et al. (in preparation).

The Seine watershed, upstream from Caudebec (73,500 km², including the freshwater estuary), has been described in detail by Meybeck et al. (1998) and Flipo et al. (2019). Its land use comprises 57% cropland, 26% forest, 10% grassland, and 7% urban or artificialized areas. The current structure of its agricultural system is described in detail by Le Noë et al. (2017) and Billen et al. (2019). Direct and indirect N₂O emissions were estimated by Garnier et al. (2009, 2019) using bottom–up methodologies. Numerous measurements of N₂O concentrations in the drainage network are available. They show a clear stream order structure, with higher concentrations in headwaters, decreasing to near-saturation values in third- to fifth-order streams, then increasing again in higher-order rivers (Garnier et al. 2009).

The model described below was run on these three watersheds for three contrasting hydrological years: 2016, 2017, and 2018. The first is characterized by a relatively atypical flood in June, while the other two years present a more usual temperate regime, with high flow in winter and low flow from June to October. The average specific runoff over the 3 years was 175, 112, and 145 mm/yr for the Orgeval, Haut Loir, and Seine watersheds, respectively.

Model description

The model of N_2O fluxes developed in this paper closely relies on the description of nitrate flow from cropland subroot zone to river outlet, including transfer through riparian wetlands, recently published by Billen et al. (2018). In brief, this model first assesses a fixed nitrate concentration to subroot water from each land use class of the basin, based on the N balance of typical crop rotations. This concentration is assigned to pre-calculated subsurface and base flow runoff. Unless tile drainage allows some fraction of the subsurface runoff to bypass riparian zones, the nitrate flow issued from the watershed has to cross an active denitrification area before reaching surface water. The resulting nitrate reduction that occurs there is calculated taking into account the extent of riparian wetlands in each elementary watershed, their potential denitrification rate, and the seasonal variations in temperature. Other processes affecting nitrate in the river drainage include algal uptake, water column, and benthic nitrification and denitrification as described in

the RIVE model (www.fire.upmc.fr/rive), as well as inputs from urban point sources.

The same approach is considered here to model the corresponding flows of N_2O . The model: (1) assigns a N_2O concentration to subroot water for each land use class in the watershed; (2) assesses N_2O production and emission in the riparian wetlands; and (3) calculates in-stream N_2O production and the resulting emissions across the water–atmosphere interface (Fig. 1).

N_2O concentration in subroot water

Initialization of the model of N_2O transfer requires an assessment of a representative N_2O concentration to subroot water of each land use class considered in the watershed. Direct measurement of this concentration is technically extremely difficult. However, N_2O concentration could be measured in groundwater collected from piezometer or groundwater resurgences, and showed large over-saturation with respect to atmospheric partial pressure. In fact, this level of




Fig. 1 Principles of the model of N_2O production and transfer through the land–river continuum. (Cbas, Crip and Criv stand for N_2O concentration in the water phase of basin soil, riparian soil and river respectively; Sbas, Srip, Sriv are the surface area of watershed, riparian wetlands and river network.; Ebas, Erip,

Eriv stand for the N_2O emission from watershed soils, riparian wetlands and river surface, respectively. kvs is the gas transfer coefficient from soil to atmosphere; kr is the gas transfer coefficient from river water to the atmosphere)

over-saturation is the very driver of direct emission of N_2O from soils to the atmosphere. As mentioned above, this latter flux is commonly measured using manual or automatic chambers, or can be estimated at an annual time scale through empirical relationships involving agricultural practices and climatic features, such as the one proposed by Garnier et al. (2019). An average N_2O concentration of soil water (C_{bas} , in $\mu\text{gN-N}_2\text{O/l}$) can therefore be associated with the average annual N_2O direct emission from watershed soil to the atmosphere (E_{bas} , in $\text{mgN-N}_2\text{O/m}^2/\text{h}$), considering a simple soil water–atmosphere transfer coefficient (kvs):

$$E_{bas} (\text{mgN-N}_2\text{O/m}^2/\text{h}) = kvs \times (C_{bas} - C_{eq}) \quad (1)$$

where C_{eq} ($\text{mgN-N}_2\text{O/m}^3$) is the N_2O concentration at equilibrium with the atmospheric partial pressure at the mean annual temperature, i.e. $0.33 \text{ mgN-N}_2\text{O/m}^3$ à $12 \text{ }^\circ\text{C}$ (Weiss and Price 1980). The parameter kvs (m/h) is defined as an average apparent ventilation coefficient between the soil water phase and the atmosphere. It can be calibrated on situations where both the flux of soil nitrous oxide emission and the soil water concentration are known. This is the case in a few spots in the investigated basins (Table 1), leading to a kvs value of approximately 0.0025 m/h .

N_2O production and emission in riparian wetlands

During the water transfer from the watershed to the river across the riparian zone, the process of denitrification, while reducing nitrate concentration of the inflowing water discharge (Q , m^3/s), also tends to increase its N_2O concentration. The nitrate module of

the Riverstrahler model (Billen et al. 2018) explicitly calculates the reduction of nitrate concentration ($\Delta C_{\text{NO}_3\text{denrip}}$) due to denitrification. During this process, nitrate is partly converted into N_2 , and partly into N_2O as an intermediate of denitrification. Although the share of denitrification flux converted to N_2O can vary widely according to several factors, including availability of nitrate, organic matter, as well as pH (Weier et al. 1993; Saggari et al. 2013), the assumption is made of a constant fraction (p_{Nden}) expressing the $N_2O/(N_2O + N_2)$ ratio. Because the value of this ratio is not very well constrained by empirical data, it is considered in the model as an adjustment parameter and a sensitivity analysis toward the chosen value is carried out (see "Discussion" section).

The rate of production of N_2O ($P_{\text{N}_2\text{Odenrip}}$, $\text{mg/m}^3/\text{h}$) through riparian denitrification can thus be estimated from the corresponding nitrate concentration reduction ($\Delta C_{\text{NO}_3\text{denrip}}$, mg/m^3) and the discharge Q (m^3/h) flowing through the riparian zone:

$$P_{\text{N}_2\text{Odenrip}} (\text{mg/m}^3/\text{h}) = -p_{\text{Nden}} \times \Delta C_{\text{NO}_3\text{denrip}} \times Q \quad (2)$$

As a result of this additional N_2O production, a new equilibrium is established in the riparian zone, with increased emission across the riparian soil–atmosphere interface (E_{rip} , $\text{mgN-N}_2\text{O/m}^2/\text{h}$), at a rate that can be expressed as:

$$E_{\text{rip}} = kvs \times (C_{\text{rip}} - C_{eq}) \quad (3)$$

where kvs is the soil–atmosphere transfer coefficient defined above, and C_{rip} is the average N_2O concentration in the riparian zone (here considered as a perfectly mixed reactor).

Table 1 Calibration of the apparent ventilation coefficient between the soil water phase and the atmosphere. kvs is calculated as $E_{bas}/(C_{bas}-C_{eq})$

	Orgeval basin	Haut Loir basin
Soil emission E_{bas} , gN/ha/day	3.5	4.5
$\text{mgN/m}^2/\text{h}$	0.015	0.019
Method	Manual and automatic emission chambers (Benoit et al. 2015a, b)	Manual emission chambers (unpublished)
Soil water conc. C_{bas} , $\text{mgN-N}_2\text{O/m}^3$	5.37	8.8
Method	Measurements in piezometers (unpublished)	Measurements in groundwater resurgence (unpublished)
kvs , m/h	0.0029	0.0022

This new equilibrium concentration (Crip) in the riparian zone can be calculated by considering the material balance equation between input and output of the water flux Q through the riparian reactor, considering production of N_2O by specifically riparian processes ($P_{N_2Odenrip}$) as well as by nonspecifically riparian processes, the latter corresponding to the average N_2O emission in the watershed soils and thus equal to $kvs \times Srip \times (Cbas - C_{eq})$ (see Eq. 1):

$$Q \times Crip = Q \times Cbas + P_{N_2Odenrip} + kvs \times Srip \times (Cbas - C_{eq}) - kvs \times Srip \times (Crip - C_{eq}) \quad (4)$$

where $Srip$ (m^2) is the area of the riparian zone in the element of watershed considered, and $Cbas$ here is the average N_2O concentration in soil water from the watershed resulting from non-riparian processes.

From Eq. 4 it follows that:

$$Crip = Cbas + P_{N_2Odenrip} / (Q + kvs \times Srip) \quad (5)$$

Or

$$Crip = Cbas - pNden \times \Delta C_{NO_3denrip} / (1 + kvs \times Srip / Q) \quad (6)$$

This procedure also allows calculation of the emission flux from riparian soil to the atmosphere (Eq. 3), thus providing a complete budget of N_2O in the riparian zone.

In-stream processes

Point sources of N_2O linked to urban wastewater inputs must be considered in modeling N_2O in the river network. Based on a number of surveys of N_2O concentration in effluents from different types of wastewater treatment plants, as well as on detailed specific studies (Tallec et al. 2006, 2008), the model considers that wastewater inputs are at near saturation N_2O concentration, except when a tertiary treatment of nitrification and/or denitrification treatment is involved, in which case the N_2O concentration is 10 times the saturation level.

In-stream production of N_2O can also occur by nitrification and denitrification in the river water

column and in the benthic phase. These are part of the RIVE module of the Riverstrahler model, described in detail at www.fire.upmc.fr/rive, as well as by Garnier et al. (2006, 2007) for the special case of the lower Seine River. The same parameterization has been used in the present study. Regarding benthic processes, the calculation procedure used is the one resulting from the meta-model established by Billen et al. (2015), which allows calculation of the integrated benthic denitrification flux, from the calculation of the organic carbon deposition, together with the oxygen and nitrate concentrations in the water column. The resulting N_2O emission flux from the benthos is assumed to be a fraction $pNden$ of the integrated benthic denitrification.

Transfer of N_2O through the water–atmosphere interface is classically represented by a relation linking the flux $Eriv$ ($mgN-N_2O/m^2/h$) to the oversaturation of the water phase, with a transfer coefficient kr_{N_2O} (m/h) depending on the flow velocity (v , m/h), the water column depth ($dpth$, m), and the slope (dimensionless) (Raymond et al. 2012), as done by Marescaux et al. (2018) (Weiss and Price 1980):

$$Eriv = kr_{N_2O} \times (C_{N_2O} - C_{eq}) \quad (7)$$

$$\text{with } C_{eq} = 0.0002 \times T^2 - 0.0167 \times T + 0.5038 \quad (8)$$

$$kr_{N_2O}(m/h) = k_{600} \times \text{SQR}(600/\text{Schmidt}_{N_2O}) \quad (9)$$

$$\text{Schmidt}_{N_2O} = 2056 - 137 \times T + 4.317 \times T^2 - 0.05435 \times T^3 \quad (10)$$

$$k_{600}(m/h) = 5037/24 \times (v \times \text{slope})^{0.89} \times \text{dpth}^{0.54} \quad (11)$$

Results

The model was first implemented at the scale of the small watersheds Orgeval and Haut Loir, for which the direct emissions estimated from measurements with static chambers on the main land use and crop rotations were available, thus allowing an estimation of an average subroot N_2O concentration used as the

main constraint variable of the model. The simulated N₂O concentrations in the river network are calculated for different values (from 0 to 0.07) of the adjustable parameter pNden and compared with available observations at a number of stations (Fig. 2). A satisfactory agreement is obtained for a pNden fraction of approximately 0.015. Higher values lead to overestimation of N₂O concentrations in all river stations. Lower values strongly underestimate the observations (Fig. 2).

The model also provides a complete budget of N₂O emissions along the continuum from watershed soils to the outlet of the river network (Table 2) and can be mapped at the resolution used for the calculations, i.e. the detailed land-use and potential wetland data within each of the sub-watershed areas and 1-km stretches for the river network (Fig. 3).

In the case of the Haut Loir basin, a number of direct measurements of the emission flux of N₂O across the river–atmosphere interface are also available (Grossel et al., in preparation). They range between 0.002 and 0.28 mgN/m²/h (mean 0.11 mgN/m²/h), which fits reasonably with the model estimates

of 0.08 mgN/m²/h on average at the same stations for pNden = 0.015.

These results confirm that riparian wetlands as well as river surfaces can indeed be seen as hotspots of N₂O emission, with per-area rates of emission sometimes more than twice the average direct emissions from the dominant agricultural land areas in the watershed, depending on the particular structure of the land- and waterscape. However, owing to their somewhat limited share in the total watershed area, these areas contribute to only 2–6% and 0.8–2%, respectively, of the total emissions. In other words, indirect emissions from riparian and river surfaces together in fact only represent 3–7% of the direct emissions from the soils. The Haut Loir watershed, because of its lower relative areas of wetlands, has the lowest indirect emissions.

The model was also implemented at the scale of the much larger Seine watershed, using the estimates of N₂O emission from cropland, grassland, and forest in the different agricultural regions made by Garnier et al. (2019) based on the record of fertilization practices (Le Noë et al. 2017) and regional climate data. Land use was geographically distributed based on Corine Land Cover, and the riparian wetlands were




Fig. 2 Model-calculated time and space variations of N₂O concentration in river water of **a** the Haut Loir basin and, **b** the Orgeval basin, for three values of the pNden parameter (representing the fraction of total denitrification emitted as N₂O in riparian and stream processes). **c** Scatter plot of

calculated versus observed river N₂O concentrations for 3 values of pNden. The r² of the correlation and the simulation RMSE (normalized to the mean) are indicated for each group of simulations. In all figures, red curves and points correspond to pNden = 0, blue is for pNden = 0.015, green for pNden = 0.07

Table 2 Model-calculated budget of N₂O emissions along the continuum from watershed soils to the river outlet in the three watersheds investigated

	Units	Haut Loir	Orgeval	Seine
Direct watershed emissions				
Surface area concerned	km ²	3600	106	73,500
N₂O emissions				
Per area N ₂ O emission	tonN-N ₂ O/yr	540	14.2	6113
	kgN-N ₂ O/ha/yr	1.50	1.34	0.83
Riparian emissions				
Surface area concerned (% watershed area)	km ² (%)	62.3 (1.7%)	3.4 (3.2%)	5389 (7.3%)
N ₂ O emissions	tonN-N ₂ O/yr	11.8	0.7	1034
Per area N ₂ O emission	kgN-N ₂ O/ha _{rip} /yr	1.9	2.1	1.92
Fraction of direct emissions	%	2.2	4.9	17
River emissions				
Surface area concerned (% watershed area)	km ² (%)	6.6 (0.18%)	0.14 (0.13%)	287 (0.39%)
N ₂ O emissions	tonN-N ₂ O/yr	4.6	0.29	252
Per area N ₂ O emission	kgN-N ₂ O/ha _{riv} /yr	6.9	20.7	8.8
Fraction of direct emissions	%	0.85	2.0	4.1
Total emissions				
	tonN-N ₂ O/yr	556	15.2	7399
	kgN-N ₂ O/ha/yr	1.54	1.43	1.0

defined from the map of potential wetlands established on the basis of a topographical index (Curie et al. 2007; Berthier et al. 2014; <https://geowww.agrocampus-ouest.fr/web/?p=1538>), as described by Billen et al. (2018).

The calculated average N₂O concentrations for different values of the pNden parameter were compared with those measured since 2012 in water from different Strahler orders (Fig. 4). Some unpublished measurements available in springs or in piezometers were also used for this comparison, the latter being referred to as “zero order,” the former as “subroot.” The measurements collectively show a U-shaped distribution of concentrations versus stream orders (see also Garnier et al. 2009 and Marescaux et al. 2018). The high concentrations at zero order (or subroot water) rapidly decrease with increasing stream orders, down to near saturation levels at order 3–5. N₂O concentration increases again at higher orders. This U-shaped distribution is well reproduced by the model, which, however, predicts, on average, larger concentrations in riparian waters and headwaters; this quickly tends toward saturation once exposed to the rapid evasion processes across the water–atmosphere interface in surface river bodies. In-stream processes,

fostered by increased point sources of organic pollution, are responsible for the further increase in N₂O concentration at higher stream order. The results obtained by simulating N₂O distribution with different values for the N₂O/(N₂O + N₂) denitrification ratio (pNden) show again that a value of about 0.015 is the maximum compatible with available observations at the scale of the whole watershed (Fig. 4). A full map of N₂O emissions from soils, riparian wetlands, and rivers in the Seine basin, similar to that shown in Fig. 3, can be viewed at address: <https://doi.org/10.26047/PIREN.data.N2O.2019>.

The complete budget of direct and indirect N₂O emission at the scale of the Seine watershed shows that riparian emissions represent 17% of direct watershed emissions, while stream surface emission accounts for only 4%. Thus, indirect emissions represent almost 20% of total watershed emissions (Table 2).

Discussion

The approach developed in this paper pursues the attempt initiated by the introduction of a module of riparian denitrification in the Riverstrahler model




Fig. 3 Map of calculated N₂O fluxes from watershed soils, riparian wetlands, and river surfaces. **a.** Haut Loir basin. **b.** Orgeval basin. For the Haut Loir basin, a magnification of two

selected areas shows the riparian wetlands along the river, with their higher emissions than the surrounding fields

(Billen et al. 2018) for describing the N cascade along the continuum from watershed soils to the river outlet within a coherent modeling framework. The model proposed is constrained by the direct N₂O emissions from the different land use classes of watershed soils, assumed to be known on a yearly average basis, from either empirical measurements, using e.g. static chambers, or another independent modeling approach such as, for instance, the empirical relationship linking annual N₂O emission to fertilizer inputs and climate data established by Garnier et al. (2019) based on data from the literature. From this mean annual direct N₂O emission value, the average N₂O concentration in groundwater and subsurface runoff is calculated using a simple empirically calibrated transfer coefficient. Riparian denitrification is first calculated by the model in terms of nitrate reduction, based on the hydrological

behavior of each sub-basin and on a temperature-dependent denitrification potential of riparian wetlands, the extension of which is defined by topographic information. A fraction pNden of this denitrification is assumed to be converted into N₂O, which allows the model to calculate both the enhanced emission from riparian areas and the N₂O concentration flowing into surface water. Within the drainage network, the model calculates the in-stream N₂O production by benthic and water column processes, including nitrification and denitrification, as well as transfer across the water–atmosphere interface. The model is validated by its capability to predict the distribution of N₂O concentration in river water at different places in the river network, as well as the order of magnitude of N₂O emission fluxes at the river–atmosphere interface.




Fig. 4 Model-calculated average N_2O concentration in watershed and riparian water, as well as in surface water of different Strahler streams. Comparison with available measurements carried out in the Seine basin since 2012 (Garnier et al. 2009; Marescaux et al. 2018 and unpublished data). The level of N_2O saturation with respect to atmospheric concentration for two temperatures is also indicated (blue dotted lines: $0.36 \mu gN-N_2O$

The model involves the calibration of two parameters, the soil–atmosphere transfer coefficient (kvs) and the fraction of denitrification emitted as N_2O versus N_2 ($pNden$). The meaning and value chosen for these two parameters requires some discussion.

N_2O emission by soils is the result of complex and episodic processes, depending on highly transient conditions in a relatively heterogeneous medium (Saggar et al. 2013). This explains the difficulties in measuring and modeling them. The model approach developed here does not intend to describe these processes; it starts from the knowledge of annual emissions, which can be assessed either by in situ measurements or by using an empirical modeling approach (e.g., Garnier et al. 2019) and considers that the soil water pool that generates runoff is buffered to the extent that its average N_2O concentration reflects the integrated soil N_2O production processes. A soil–atmosphere transfer coefficient (kvs , in m/h) is calibrated on local situations where both the annual N_2O emission flux from the soil and the average N_2O concentration in the soil water pool are known. The emission values found in two such situations, in the Haut Loir and Orgeval basins, are consistently about 0.0025 m/h (Table 1). Combined with the mean runoff

at 10 °C, $0.21 \mu gN-N_2O$ at 25 °C)). **b** Scatter plot of calculated versus observed N_2O concentrations for 3 values of the $N_2O/(N_2O + N_2)$ ratio ($pNden$). The r^2 of the correlation and the simulation RMSE (normalized to the mean) are indicated for each group of simulations. In both figures, red curves and points correspond to $pNden = 0$, blue is for $pNden = 0.015$, green for $pNden = 0.07$

rate (100–270 mm/yr), these values imply that no more than 1–2% of the N_2O flux produced in the soil is transferred to the hydrosphere rather than to the atmosphere. This contribution to the indirect emissions is thus somewhat limited.

The larger contributor to indirect emissions, therefore, consists of denitrification processes during the transfer of water from soils to the outlet of the river system, including riparian zones and the rivers themselves. The model calculates these processes by mechanistic modules, well established and validated by their capacity to correctly simulate nitrate and ammonium concentrations (Billen et al. 2018) with, however, one additional calibrated parameter ($pNden$) representing the fraction of N_2O emitted during nitrate reduction through denitrification. This fraction is assumed to be constant in space and time for rivers and riparian wetlands.

N_2O emission from denitrification results from an imbalance between the rate of NO_3^- reduction to N_2O and its further reduction into N_2 . Decreased pH (Rochester 2003; Simek et al. 2002; Simek and Cooper 2002) or increased temperature (Benoit et al. 2015a, b) has been shown to result in higher $N_2O/(N_2 + N_2O)$ ratios in agricultural soils. The review by

Rochester (2003) of several studies, as well as the data of Hénault et al. (2005), suggests a consistent relationship between this ratio and the pH, starting at values close to 0.01 at pH 7–8 and increasing up to 0.75 at pH 4–5. The problem with this relationship is that it was established based on laboratory studies in which the pH was manipulated. Only two results included in the relationship of Rochester (2003) come from direct field measurements: they both show a $N_2O/(N_2 + N_2O)$ ratio of 0.01 at a pH of 5.8 and 8.2, respectively. Simek et al. (2002) emphasized the point that the imbalance of microbial or enzymatic activities leading to N_2O emission during denitrification can be transient until the readjustment of the rates of the two enzymatic processes (nitrate and N_2O reductions) or the adaptation of the microbial consortium. N_2O emission can therefore be a transient process linked to changes in pH or temperature rather than controlled by constant conditions of pH and temperature. We therefore limited our literature search to studies reporting pNden values based on measurements carried out under field conditions, i.e., without manipulation of the soil samples, either using ^{15}N -labelled fertilizer tracing, acetylene block technique on entire core samples or the results of emission chambers combined with estimation of denitrification. Even so, the range of values reported for the $N_2O/(N_2 + N_2O)$ ratio is considerable, varying from less than 0.01 to 0.6, without a clear relationship to pH or any other environmental variables (Denmead et al. 1979; Rudaz et al. 1999; Rochester 2003; Clement et al. 2002; Oehler et al. 2007; Scheer et al. 2009; Autret et al. 2019).

The review by Schlesinger (2009), distinguishing records of $N_2O/(N_2 + N_2O)$ for different types of ecosystems from agricultural soils, soils under natural vegetation or freshwater wetlands shows that the latter (0.08 ± 0.02) are typically much lower than the formers (0.37 ± 0.03 and 0.49 ± 0.07), as also shown by Beaulieu et al (2011). In a more recent paper, Butterbach-Bahl et al (2013) brought arguments showing that part of the data gathered by Schlesinger (2009), based on the use of the acetylene block technique, are subject to a systematic underestimation of N_2 production by denitrification, leading to overestimation of the nitrous oxide ratio. Discarding these data leads to an average $N_2O/(N_2 + N_2O)$ ratio of 0.016 ± 0.04 for wetlands and rivers (based however on 5 measurements only). The measurements by

Beaulieu et al (2011) obtained by in situ ^{15}N tracing across 53 US rivers and streams, ranged from 0.0004 to 0.056, with an interquartile range between 0.003 and 0.01.

The pNden value selected in our model calculations (0.015) is in exactly the same range. Figures 2 and 4 show the sensitivity of the model results to the value of this parameter. A more detailed analysis of the model response in terms of riparian and riverine emissions is shown for the case of the Seine watershed in Fig. 5. For pNden set to zero, the emission of riparian wetlands corresponds to the background soil processes, similar to the average of those direct emissions occurring in the watershed, while river indirect emissions consist of the evasion of N_2O transferred from watershed soils to surface waters. At increasing pNden values, additional emission occurs from both riparian wetlands and river surfaces precisely due to denitrification processes occurring there. For a pNden value of approximately 0.015, the calculated emissions fit the observed distribution of river concentrations and emission fluxes. Higher values of pNden, e.g., 5 to 10 times higher (0.07–0.15), would clearly result in much higher estimates of indirect N_2O emissions by riparian zones and river networks, which would reach values close to the watershed direct emissions themselves. However, such values are not supported by empirical observations of N_2O




Fig. 5 Sensitivity analysis of the effect of pNden value on the model estimate of N_2O emissions by riparian and riverine processes in the Seine watershed

Table 3 Budget of direct and indirect N₂O emissions from the Seine watershed, published by Garnier et al. (2009) and Marescaux et al. (2018), compared with the results of the

present study (with an estimated 25% range based on the uncertainty of pNden calibration)

Ton-N ₂ O/yr	IPCC 2006	Garnier et al. 2009	Marescaux et al. 2018	This study
Direct emissions from watershed soils	8250	10,400	9000	6113
Total indirect emissions	1350	1050–1460		1286 ± 330
% Indirect emission with respect to direct ones	16	10–14	–	21 ± 5
Riparian emissions	–	800–1000	–	1034 ± 260
River network emissions	–	100–200	240	252 ± 82
Wastewater treatment plants	–	150–260 ^a	–	3.4 ^b

^aEmission to the atmosphere from the treatment plant^bInput to the river network through wastewater discharge

concentration and emission rates available in the investigated watersheds.

The estimate of indirect N₂O emissions obtained with the model for the Seine watershed is consistent with previous figures published by Garnier et al. (2009) Marescaux et al. (2018) based on empirical budgeting approaches (Table 3). They do not contradict the default parameters suggested in the IPCC guidelines. However, our estimate of indirect emissions in the Seine watershed does not take into account the possible contribution of ponds, lakes, and other stagnant water bodies.

Our estimated contribution of indirect emissions to total emissions from the Seine watershed is much lower compared with those reported by Turner et al. (2015) for the Mississippi. Our work is based on two different watersheds (Orgeval nested in the Seine basin) sharing the same geological substrate, both characterized by carbonate-rich sedimentary rocks. We cannot exclude that under other contexts and, in particular, in more acidic soils and waters, different values of pNden would apply and lead to much higher N₂O emission in both river and riparian wetlands.

Conclusion

The modeling approach developed in this paper offers a coherent framework to model direct and indirect N₂O emissions at the scale of the whole continuum from agricultural soils to river system. Its application to three watersheds of increasing size, from a few hundred to nearly 100,000 km² in a temperate climate and sedimentary rock context, confirms the view that, in spite of high N₂O emission rates per surface area in

riparian wetlands and headwater streams, indirect emissions represent barely 20% of direct soil emissions when expressed at the scale of the whole watershed area. The largest part of these indirect emissions occurs in riparian areas where nitrate leached from the watershed soils undergoes partial denitrification. Headwaters receiving oversaturated groundwater from the watershed are also significant emitters. Downstream in the large river network, in-stream processes, possibly amplified by point source inputs, may contribute to enhanced emissions through the river–atmosphere interface.

Acknowledgements The study was conducted within the framework of the HydroGES supported by the Agency for the Environment and Mastery of Energy (ADEME). The PIREN-Seine program and the Fédération Ile-de-France de Recherche pour l'Environnement (FIRE) are also acknowledged for their support. We are grateful to all the technical staff involved in field measurements discussed in this paper: Adeline Ayzac, Christian Lelay, Didier Laloua, Lionel Cottenot at URSols, Orléans, as well as Benjamin Mercier, Anun Matinez, and Sébastien Bosc at UMR Metis, Paris.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Anderson TR, Goodale CL, Groffman PM, Walter MT (2014) Assessing denitrification from seasonally saturated soils in an agricultural landscape: a farm scale mass-balance approach. *Agr Ecosyst Environ* 189:60–69
- Autret B, Mary B, Gréhan E, Ferchaud F, Grandeau G, Raktovolonona L, Bertrand M, Beaudoin N (2019) Can alternative cropping systems mitigate nitrogen losses and improve GHG balance? Results from a 19-yr experiment in Northern France. *Geoderma* 342:20–33. <https://doi.org/10.1016/j.geoderma.2019.01.039>
- Barnes J, Owens NJP (1999) Denitrification and N₂O concentrations in the Humber Estuary, UK and adjacent coastal zones. *Mar Poll Bull* 37:247–260
- Beaulieu JJ, Tank JL, Hamilton SK, Wollheim WM, Hall RO, Mulholland PJ, Peterson BJ, Ashkenas LR, Cooper LW, Dahm CN, Dodds WK, Grimm NB, Johnson SL, McDowell WH, Poole GC, Valett HM, Arango CP, Bernot MJ, Burgin J, Crenshaw CL, Helton AM, Johnson LT, O'Brien JM, Potter JD, Sheibley RW, Sobota DJ, Thomas SM (2011) Nitrous oxide emission from denitrification in stream and river networks. *PNAS* 108:214–219. <https://doi.org/10.1073/pnas.1011464108>
- Beheydt DPB, Sleutel S, Li C, Van Cleemput O (2007) Validation of DNDC for 22 long-term N₂O field emission measurements. *Atmos Environ* 41:6196–6211. <https://doi.org/10.1016/j.atmosenv.2007.04.003>
- Benoit M, Garnier J, Anglade J, Billen G (2014) Nitrate leaching from organic and conventional arable crop farms in the Seine Basin (France). *Nutrient Cycling in Agrosystems*. <https://doi.org/10.1007/s10705-014-9650-9>
- Benoit M, Garnier J, Billen G, Tournebize J, Gréhan E, Mary B (2015a) Nitrous oxide emissions and nitrate leaching in an organic and a conventional cropping system (Seine basin, France) *Agriculture, Ecosystems and Environment* 213(2015):131–141. <https://doi.org/10.1016/j.agee.2015.07.030>
- Benoit M, Garnier J, Billen G (2015b) Temperature dependence of nitrous oxide production of a luvisolic soil in batch experiments. *Process Biochem* 50:79–85
- Berthier L, Bardy M, Chenu JP, Guzmova L, Laroche B, Lehmann S, Lemerrier B, Martin M, Mérot P, Squiquidant H, Thiry E, Walter C (2014) Enveloppes des milieux potentiellement humides de la France métropolitaine. Notice d'accompagnement. Programme de modélisation des milieux potentiellement humides de France, Ministère d'Ecologie, du Développement Durable et de l'Energie. <https://geowww.agrocampus-ouest.fr/web/?p=1538>
- Bessou C, Mary B, Leonard J, Roussel M, Grehan E, Gabrielle B (2010) Modelling soil compaction impacts on nitrous oxide emissions in arable fields. *Eur J Soil Sci* 61:348–363. <https://doi.org/10.1111/j.1365-2389.2010.01243.x>
- Billen G, Garnier J, Silvestre M (2015) A generic algorithm for modelling benthic nutrient fluxes in river systems. *Int J Limnol* 51:37–47. <https://doi.org/10.1051/limn/2014030>
- Billen G, Ramarson A, Thieu V, Théry S, Silvestre M, Pasquier C, Hénault C, Garnier J (2018) Nitrate retention at the river–watershed interface: a new conceptual modeling approach. *Biogeochemistry* 139:31–51. <https://doi.org/10.1007/s10533-018-0455-9>
- Billen G, Garnier J, Le Noë J, Viennot P, Gallois N, Puech T, Schott C, Anglade J, Mary B, Beaudoin N, Léonard J, Mignolet C, Théry S, Thieu T, Silvestre M, Passy P (2019) The Seine watershed water-agro-food system: long-term Trajectories of C, N and P metabolism. In: Labadie P, Lestel L (eds) Flipo N. The Seine River Basin, Handbook of Environmental Chemistry
- Bouwman AF (1996) Direct emissions of nitrous oxide from agricultural soils. *Nutr Cycl Agroecosyst* 46:53–70
- Bouwman AF, Boumans LJM, Batjes NH (2002) (2002) Modeling global annual N₂O and NO emissions from fertilized fields. *Global Biogeochem* 16:1080. <https://doi.org/10.1029/2001GB001812>
- Butterbach-Bahl K, Baggs EM, Dannenmann M, Kiese R, Zechmeister-Boltenstern S (2013) Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Phil Trans R Soc Lond B Biol Sci* 368:20130122. <https://doi.org/10.1098/rstb.2013.0122>
- Cayuella ML, Aguilera E, Sanz-Cobena A, Adams DC, Abalos D, Barton L, Ryals R, Silver WL, Alfaro MA, Pappa VA, Smith P, Garnier J, Billen G, Bouwman L, Bondeau A, Lassaletta L (2016) Direct nitrous oxide emissions in Mediterranean climate cropping systems: emission factors based on a meta-analysis of available. *AEE*. <https://doi.org/10.1016/j.agee.2016.10.006>
- Cellier P, Durand P, Hutchings N, Dragosits U, Theobald M, Drouet JP, Oenema O, Bleeker A, Breuer L, Dalgaard T, Duret S, Kros J, Loubet B, Olensen J, Merot P, Viaud V, de Vries W, Sutton MA (2011) Nitrogen flows and fate in rural landscapes. In: Sutton MA, Howard CM, Erisman JW, Billen G, Bleeker A, Grennfelt P, van Grinsven H, Grizzetti B (eds) The European Nitrogen Assessment. Cambridge University Press, Cambridge, pp 229–248
- Corazza M, Bergamaschi P, Vermeulen AT, Aalto T, Haszpra L, Meinhardt F, O'Doherty S, Thompson R, Moncrieff J, Popa E, Steinbacher M, Jordan A, Dlugokencky E, Brühl C, Krol M, Dentener F (2011) Inverse modelling of European N₂O emissions: assimilating observations from different networks. *Atmos Chem Phys* 11:2381–2398. <https://doi.org/10.5194/acp-11-2381-2011>
- Clément JC, Pinay G, Marmonier P (2002) Seasonal Dynamics of Denitrification along Topohydrosequences in Three Different Riparian Wetlands. *J Environ Qual* 31:1025–1037
- Crutzen P, Mosier A, Smith K, Winiwarter W (2008) N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuel. *Atmos Chem Phys* 8:389–395
- Curie F, Gaillard S, Ducharme A, Bendjoudi H (2007) Geomorphological methods to characterizes wetlands at the scale of the Seine watershed. *Sci Total Environ* 375:59–68
- Davidson EA (2009) The contribution of manure and fertilizer N to atmospheric nitrous oxide since 1860. *Nat Geosci* 2:659–662
- Del Grosso SJ, Mosier AR, Parton WJ, Ojima DS (2005) DAYCENT model analysis of past and contemporary soil N₂O and net greenhouse gas flux for major crops in the USA. *Soil Tillage Res* 83:9–24

- Denmead OT, Freney JR, Simpson JR (1979) Nitrous oxide emission during denitrification in a flooded field. *Soil Sci Soc Am J* 43:716–718
- Flipo N, Labadie P, Lestel L (2019) The Seine River Basin. *Handb Environ Chem*. https://doi.org/10.1007/698_2019_393
- Gabrielle B, Laville P, Duval O, Nicoullaud B, Germon JC, C. Hénault C (2006) Process-based modeling of nitrous oxide emissions from wheat-cropped soils at the subregional scale. *Glob Biogeochem Cycles*. <https://doi.org/10.1029/2006GB002686>
- Garnier J, Cébron A, Tallec G, Billen G, Sebilo M, Martinez A (2006) Nitrogen behaviour and nitrous oxide emission in the tidal Seine River estuary (France) as influenced by human activities in the upstream watershed. *Biogeochemistry* 77:305–326
- Garnier J, Billen G, Cébron A (2007) Modelling nitrogen transformations in the lower Seine river and estuary (France): impact of waste water release on oxygenation and N₂O emission. *Hydrobiologia* 588:291–302
- Garnier J, Billen G, Martinez A, Mounier E, Silvestre M, Vilain G, Toche F (2009) Nitrous oxide (N₂O) in the Seine river and basin: observations and budgets. *Agr Ecosyst Environ* 133:223–233
- Garnier J, Billen G, Vilain G, Benoit M, Passy P, Tallec G, Tournebize J, Anglade J, Billy C, Mercier B, Ansart P, Azougui A, Sebilo M, Kao C (2014) Curative vs. preventive management of nitrogen transfers in rural areas: Lessons from the case of the Orgeval watershed (Seine River basin, France). *J Environ Manage* 144:125–134
- Garnier J, Le Noë J, Marescaux A, Sanz-Cobena A, Lassaletta L, Silvestre M, Thieu V, Billen G (2019) Long-term changes in greenhouse gas emissions from French agriculture and livestock (1852–2014): from traditional agriculture to conventional intensive systems. *Sci Total Environ* 660:1486–1501. <https://doi.org/10.1016/j.scitotenv.2019.01.048>
- Griffis TJ, Lee X, Baker JM, Russelle MP, Zhang X, Venterea R, Millet DB (2013) Reconciling the differences between top-down and bottom-up estimates of nitrous oxide emissions for the US Corn Belt. *Global Biogeochem Cycles* 27(3):746–754. <https://doi.org/10.1002/gbc.20066>
- Grossel, A. et al (in prep). Indirect emissions of nitrous oxide in a cropland watershed presenting contrasted hydrology in Central France.
- Hénault C, Bizouard F, Laville P, Gabrielle B, Nicoullaud B, Germon JC, Cellier P (2005) Predicting in situ soil N₂O emission using NOE algorithm and soil database. *Glob Change Biol* 2005(11):115–127. <https://doi.org/10.1111/j.1365-2486.2004.00879.x>
- IPCC (2006). N₂O Emissions from Managed Soils, and CO₂ Emissions from Lime and Urea Application. AFOLU, vol.4. 54 pp (Chapter 11). , Intergovernmental Panel on Climate Change.
- IPCC (2019) Hergouale’h, K., Akiyama, H., Bernoux, M., Chirinda, N., Del Prado, A., Kasimir, A., MacDonald, D., Ogle, S., Regina, K., van der Weerden, T., 2019. 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 5, Chapter 11: N₂O emissions from managed soils, and CO₂ emissions from lime and urea application. Technical Report. Technical Report 4–88788- 032–4, Intergovernmental Panel on Climate Change.
- Kroeze C, Dumont E, Sitzinger SP (2005) New estimates of global emissions of N₂O from rivers and estuaries. *Environ Sci* 2:159–165
- Le Noë J, Billen G, Garnier J (2017) How the structure of agro-food systems shapes nitrogen, phosphorus, and carbon fluxes: the generalized representation of agro-food system applied at the regional scale in France. *Sci Total Environ* 586:42–55. <https://doi.org/10.1016/j.scitotenv.2017.02.040>
- Li C, Frolking S, Frolking TA (1992) A model of nitrous oxide evolution from soil driven by rainfall events: I. Model structure and sensitivity. *J Geophys Res* 97:9759–9776
- Marescaux A, Thieu V, Garnier J (2018) Carbon dioxide, methane and nitrous oxide emissions from the human-impacted Seine watershed in France. *Sci Total Environ* 643:247–259
- Meybeck M, de Marsily G, Fustec E (eds) (1998) *La Seine en son bassin. Fonctionnement écologique d’un système fluvial anthropisé*. Elsevier, Paris, p 749
- Mulholland PJ et al (2008) Stream denitrification across biomes and its response to anthropogenic nitrate loading. *Nature* 452(7184):202–205
- Myhre G, Shindell D, Bréon FM, Collins W, Fuglesvedt J, Huang, J, Koch D, Lamarque JF, Lee D, Mendoza B, Nakajima T, Robock A, Stephens G, Takemura T, and Zhang H.(2013). Anthropogenic and Natural Radiative Forcing, in: *Climate Change 2013: The Physical Science Basis Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press. Cambridge, UK and New York, NY, USA.
- Oehler F, Bordenave P, Durand P (2007) Variations of denitrification in a farming catchment area. *Agr Ecosyst Environ* 120(2007):313–324
- Outram F, Hiscock K (2012) Indirect nitrous oxide emissions from surface water bodies in a lowland arable catchment: a significant contribution to agricultural greenhouse gas budgets? *Environ Sci and Technol* 46:8156–8163
- Parton WJ, Mosier AR, Ojima DS et al (1996) Generalized model for N₂ and N₂O production from nitrification and denitrification. *Global Biogeochem Cycles* 10:401–412
- Parton WJ, Holland EA, Del Grosso SJ, Hartman MD, Martin E, Mosier AR, Ojima DS, Schimel DS (2001) Generalized model for NO_x and N₂O emissions from soils. *J Geophys Res* 106:17403–17419
- Ravishankara AR, Daniel JS, Portmann RW (2009) Nitrous oxide (N₂O): the dominant ozone-depleting substance emitted in the 21st century. *Science* 326(5949):123–125
- Raymond PA, Zappa CJ, Butman D, Bott TH, Potter J, Mulholland P, Laursen AE, McDowell WH, Newbold D (2012) Scaling the gas transfer velocity and hydraulic geometry in streams and small rivers. *Limnol Oceanogr* 2:41–53. <https://doi.org/10.1215/21573689-1597669>
- Reay DS, Davidson EA, Smith KA, Smith P, Melillo JM, Dentener Fand Crutzen PJ (2012) Global agriculture and nitrous oxide emissions. *Nat Clim Change*. <https://doi.org/10.1038/NCLIMATE1458>

- Rochester IJ (2003) Estimating nitrous oxide emissions from flood-irrigated alkaline grey clays. *Aust J Soil Res* 41:197–206
- Rodrigues PPGW, Barnes J, Upstill-Goddard R (2007) Simulating estuarine nitrous oxide production by means of a dynamic model. *Mar Poll Bull* 54:164–172
- Rudaz AO, Walti E, Kyburz G et al (1999) Temporal variation in N₂O and N₂ fluxes from a permanent pasture in Switzerland in relation to management, soil water content and soil temperature. *Agric Ecosyst Environ* 73:83–91
- Saggar S, Jha N, Deslippe J, Bolan NS, Luo J, Giltrap DL, Kim DG, Zaman M, Tillman RW (2013) Denitrification and N₂O:N₂ production in temperate grasslands: processes, measurements, modelling and mitigating negative impacts. *Sci Tot Envir* 465:173–194
- Scheer C, WassmannButterbach-Bahl RK et al (2009) The relationship between N₂O, NO, and N₂ fluxes from fertilized and irrigated dryland soils of the Aral Sea Basin. *Uzbekistan Plant Soil* 314:273–283
- Shcherbak I, Millar N, Robertson GP (2014) Global meta-analysis of the nonlinear response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *Proc Natl Acad Sci USA* 111(25):9199–9204
- Schlesinger WH (2009) On the fate of anthropogenic nitrogen. *Proc National Acad Sci* 106:203–208
- Simek M, Cooper JE (2002) The influence of soil pH on denitrification: progress towards the understanding of this interaction over the last 50 years. *Eur J Soil Sci* 53:345–354
- Simek M, Jisova L, Hopkins DW (2002) What is the so-called pH-optimum of denitrification? *Soil Biol Biogeochem* 34:1227–1234
- Syakila A, Kroeze C (2011) The global nitrous oxide budget revisited. *Greenh Gas Measure Manage* 1:17–26
- Talleg G, Garnier J, Billen G, Gousailles M (2006) Nitrous oxide emissions from secondary activated sludge in nitrifying conditions of urban wastewater treatment plants: effect of oxygenation level. *Water Res* 40:2972–2980
- Talleg G, Garnier J, Billen G, Gousailles M (2008) Nitrous oxide emissions from denitrifying activated sludge of urban wastewater treatment plants, under anoxia and low oxygenation. *Biores Technol* 99:2200–2209
- Thompson RL, Gerbig C, Rödenbeck C (2011) A Bayesian inversion estimate of N₂O emissions for western and central Europe and the assessment of aggregation errors. *Atmos Chem Phys* 11:3443–3458. <https://doi.org/10.5194/acp-11-3443-2011>
- Turner PA, Griffis TJ, Lee X, BakerJM VRT, Wood JD (2015) Indirect nitrous oxide emissions from streams within the US Corn Belt scale with stream order. *Proc Natl Acad Sci*. <https://doi.org/10.1073/pnas.1503598112>
- Vilain G, Garnier J, Talleg G, Cellier P (2010) Effect of slope position and land use on nitrous oxide (N₂O) emissions (Seine Basin, France). *Agr Forest Meteorol* 150:1192–1202
- Vilain G, Garnier J, Talleg G, Tournebise J (2011) Indirect N₂O emissions from shallow groundwater in an agricultural catchment (Seine Basin, France). *Biogeochemistry*. <https://doi.org/10.1007/s10533-011-9642-7>
- Vilain G, Garnier J, Passy P, Silvestre M, Billen G (2012) Budget of N₂O emissions at the watershed scale: role of land cover and topography (the Orgeval basin, France). *Biogeosciences* 9(1085–1097):2012
- Weier KL, Doran JW, Power JF, Walters DT (1993) Denitrification and the dinitrogen nitrous-oxide ratio as affected by soil–water, available carbon, and nitrate. *Soil Sci Soc Am J* 57:66–72
- Weiss RF, Price BA (1980) Nitrous oxide solubility in water and seawater. *Mar Chem* 8:347–359

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.