



**HAL**  
open science

# THE EFFECTS OF CHANGES OF REDOX POTENTIAL ON THE GROWTH OF MARINE EGGS, PARACENTROTUS LIVIDUS

Matilda M Brooks

► **To cite this version:**

Matilda M Brooks. THE EFFECTS OF CHANGES OF REDOX POTENTIAL ON THE GROWTH OF MARINE EGGS, PARACENTROTUS LIVIDUS. *Vie et Milieu* , 1954, 5 (2), pp.296-297. hal-02572732

**HAL Id: hal-02572732**

**<https://hal.sorbonne-universite.fr/hal-02572732>**

Submitted on 13 May 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE EFFECTS OF CHANGES  
OF REDOX POTENTIAL ON THE GROWTH  
OF MARINE EGGS, *PARACENTROTUS LIVIDUS*

Matilda M. BROOKS

This is a preliminary report of the experiments on *Paracentrotus lividus* which were done at Banyuls-sur-mer. They are a continuation of those done elsewhere, on the effects of various changes in the oxidation reduction potential on the growth and development of embryos of *Paracentrotus lividus*.

Larvæ at different stages of development to the gastrula stage were placed in solutions of sea water containing one of the indicators of the redox scale of the Clark series as follows :

1. 2-6 dibrom phenol indophenol;
2. Guaiacol indophenol;
3. Methylene blue;
4. Indigo monosulphonate;
5. Janus green.

The concentrations used varied from .005 % to .000002 % depending on toxicity. The time in the solutions varied from 10 minutes to 4 days before being transferred to sea water.

In the experiments at Naples the eggs were pretreated before being subjected to experimental procedure with such reagents as  $\text{SCN}^-$  or  $\text{I}^-$  or Ca free sea water. In the experiments at Banyuls no pretreatment was used.

The conclusions in general seemed to show that continuous immersion in the experimental solution containing the more negative dyes (numbers 3, 4, 5) development was stopped at the blastula stage. If returned to sea water after a certain time pluteic form in the case of 1 and 2.



Numbers 3 and 4 produce a solid mass of mesoderm (?) at the vegetal end, whereas number 1 forms an over-developed apical end.

No spines are formed in the more negative solutions.

These results are a confirmation of the theory stated elsewhere that the oxidation-reduction potential is the controlling factor in over-developing either the apical or the vegetal end of various marine eggs.

From the Marine Laboratory, Banyuls-sur-Mer, and the Department of Physiology of the University of California, Berkeley, California.

---